

LECTURE NOTES

MATH 103A — SPRING 2023

COMPLEX ANALYSIS

University of California, Santa Cruz

DEEWANG BHAMIDIPATI

(all errors introduced are my own)

Last Updated: Wednesday 21st June, 2023

Contents

1	Part I. Preliminaries	2
1.1	Construction of the (field of) Complex Numbers	2
1.2	Geometric Properties of Complex Numbers	6
1.3	Roots of Complex Numbers	15
1.4	Basic Topology of \mathbf{C}	18
2	Part II. Holomorphic Functions	22
2.1	Complex Functions	22
2.2	Limits of Functions	23
2.3	Theorems on Limits	26
2.4	Continuous Functions	30
2.5	Complex-Differentiable Functions	32
2.6	Cauchy-Riemann Equations	35
2.7	Holomorphic Functions	40
2.8	The Exponential Function	42
2.9	The Logarithmic Function	43
2.10	Power and Exponential Functions	48
2.11	Trigonometric Functions	49
3	Part III. Integration	52
3.1	Derivatives of Functions of a Real-variable	52
3.2	Integral of $\gamma : [a, b] \rightarrow \mathbf{C}$	54
3.3	Contours	55
3.4	Contour Integration	59
3.5	Estimating Contour Integrals	66
3.6	Antiderivatives & Fundamental Theorem of Contour Integrals	69
3.7	Cauchy-Goursat Theorem	75
3.8	Simply Connected Domains	79
3.9	Multiply Connected Domains	80
3.10	Cauchy's Integral Formula	85
3.11	Liouville's Theorem and the Fundamental Theorem of Algebra	91
A	Problems	94

Lecture 1

What is Complex Analysis? The main object of study is a **holomorphic** function $f : G \rightarrow \mathbf{C}$, where $G \subseteq \mathbf{C}$. Namely, a function for which the limit

$$\lim_{h \rightarrow 0} \frac{f(z+h) - f(z)}{h}$$

exists and is finite on an open set; that is, a **complex-differentiable function** on an open set. As a set, $\mathbf{C} = \mathbf{R}^2$, so one can naively expect the theory to be similar to that of real analysis, in this case the behaviour of differentiable functions. Interestingly, the requirement of holomorphicity can yield results that have no counterpart in the real case.

A prime example of this is *Louville's Theorem*. *Every bounded holomorphic function is constant.*

Discussion 0.0.1. We begin with first addressing the existence and nature of \mathbf{C} itself. Let \mathbf{R} denote the (field of) real numbers. One immediately deduces that the equation

$$x^2 + 1 = 0 \tag{*}$$

has no solution in the real numbers. The (field of) complex numbers \mathbf{C} stems from our desire to find a set containing \mathbf{R} that extends the algebraic operations of addition and multiplication of real numbers and which contains not only solutions to the polynomial equation above but solutions to all polynomial equations.

Surprisingly enough, the construction amounts to defining a symbol i that is a solution to $(*)$ and then considering all expressions of the form

$$x + iy, \quad x, y \in \mathbf{R}$$

1. Part I. Preliminaries

1.1. Construction of the (field of) Complex Numbers

Definition 1.1.1 (The set of Complex Numbers). A **complex number** z is simply an order pair $z := (x, y)$ of real numbers. Thus, the set of all complex numbers is given by

$$\mathbf{C} := \mathbf{R}^2 = \{(x, y) : x, y \in \mathbf{R}\}$$

If $z = (x, y)$ is a complex number, then we call

$$\operatorname{Re} z := x \quad \text{and} \quad \operatorname{Im} z := y$$

the **real** and **imaginary parts** of z respectively.

Two complex numbers z_1 and z_2 are equal if and only if $\operatorname{Re} z_1 = \operatorname{Re} z_2$ and $\operatorname{Im} z_1 = \operatorname{Im} z_2$.

If $\operatorname{Re} z = 0$ and $\operatorname{Im} z \neq 0$, we say that z is **purely imaginary**. The set of purely imaginary complex numbers corresponds to the y -axis and is called the **imaginary axis** in \mathbf{C} .

Definition 1.1.2 (Binary Operations on \mathbf{C}). Let $z_1 = (x_1, y_1)$ and $z_2 = (x_2, y_2)$ be complex numbers. Then their *sum* is

$$z_1 + z_2 = (x_1, y_1) + (x_2, y_2) := (x_1 + x_2, y_1 + y_2)$$

and their *product* is

$$z_1 \cdot z_2 = (x_1, y_1) \cdot (x_2, y_2) := (x_1x_2 - y_1y_2, x_1y_2 + x_2y_1)$$

Proposition 1.1.3. *There exists a subset of \mathbf{C} that is algebraically indistinguishable from \mathbf{R} .*

Proof. Consider the set (the x -axis)

$$\mathbf{R} \times \{0\} = \{(x, 0) : x \in \mathbf{R}\} \subseteq \mathbf{C}.$$

There is a bijection

$$\phi : \mathbf{R} \rightarrow \mathbf{R} \times \{0\}, x \mapsto (x, 0).$$

Moreover,

$$\phi(x) + \phi(y) = (x, 0) + (y, 0) = (x + y, 0) = \phi(x + y)$$

$$\phi(x) \cdot \phi(y) = (x, 0) \cdot (y, 0) = (xy - 0 \cdot 0, x \cdot 0 + y \cdot 0) = (xy, 0) = \phi(xy)$$

□

According to the proposition, the operations of addition and multiplication on complex numbers we have defined extend the operations of addition and multiplication of real numbers. We therefore call the x -axis, the **real axis**.

Discussion 1.1.4. We identify each complex number $(x, 0)$ with the corresponding real number x ; more than that, abusing notation, we write

$$1 = (1, 0) \quad \text{and} \quad (x, 0) = x(1, 0) = x$$

Now, define the **imaginary unit** $i := (0, 1)$. Then

$$i^2 = i \cdot i = (0, 1) \cdot (0, 1) = (0^2 - 1^2, 0 \cdot 1 + 1 \cdot 0) = (-1, 0) = -1.$$

Moreover, for any $z = (x, y) \in \mathbf{C}$ we see that

$$\begin{aligned} z &= (x, y) \\ &= (x, 0) + y(0, 1) = x + iy = \operatorname{Re} z + i \operatorname{Im} z \end{aligned}$$

Hence, with our new notation

$$\mathbf{C} = \{x + iy : x, y \in \mathbf{R}, i^2 = -1\}$$

and

$$z_1 + z_2 = (x_1 + iy_1) + (x_2 + iy_2) = (x_1 + x_2) + i(y_1 + y_2)$$

$$z_1 \cdot z_2 = (x_1 + iy_1) \cdot (x_2 + iy_2) = (x_1x_2 - y_1y_2) + i(x_1y_2 + x_2y_1)$$

Although we have expanded the real numbers and we will see that the complex numbers have several new and familiar properties. We do end up losing one property of the real numbers when working with complex numbers: total ordering (that extends the one on \mathbf{R} or is compatible with multiplication). In the world of complex numbers, it no longer makes sense to ask if $z_1 > z_2$ (see Problem 9).

In practice, the product of complex numbers can be computed by multiplying the expressions as if they were polynomials in the variable i , and using $i^2 = -1$. The fact that this works is left as Problem 3.

Example 1.1.5. Compute $(1 + i)(1 - 3i)$.

Answer. We note

$$\begin{aligned}(1 + i)(1 - 3i) &= (1 - 3i) + i(1 - 3i) \\ &= (1 - 3i) + (i - 3i^2) \\ &= (1 - 3i) + (i + 3) = 4 - 2i\end{aligned}$$

□

Lecture 2 **Proposition 1.1.6** (Algebraic Properties of $(\mathbf{C}, +, \cdot)$).

(1) Additive Identity. For every $z \in \mathbf{C}$

$$z + 0 = z = 0 + z$$

(2) Associativity of Addition. For every triple $z_1, z_2, z_3 \in \mathbf{C}$

$$z_1 + (z_2 + z_3) = (z_1 + z_2) + z_3$$

(3) Commutativity of Addition. For every pair $z_1, z_2 \in \mathbf{C}$

$$z_1 + z_2 = z_2 + z_1$$

(4) Additive Inverses. For every $z \in \mathbf{C}$, there exists a complex number, denoted $-z$, such that

$$z + (-z) = 0 = (-z) + z$$

In fact, $-z := (-1)z$, which is described in Problem 2.

(5) Multiplicative Identity. For every $z \in \mathbf{C}$

$$z \cdot 1 = z = 1 \cdot z$$

(6) Associativity of Multiplication. For every triple $z_1, z_2, z_3 \in \mathbf{C}$

$$z_1 \cdot (z_2 \cdot z_3) = (z_1 \cdot z_2) \cdot z_3$$

(7) Commutativity of Multiplication. For every pair $z_1, z_2 \in \mathbf{C}$

$$z_1 \cdot z_2 = z_2 \cdot z_1$$

(8) Multiplicative Inverses. For every $z \in \mathbf{C}^* := \mathbf{C} \setminus \{0\}$, there exists a complex number, denoted z^{-1} or $1/z$, such that

$$z \cdot z^{-1} = 1 = z^{-1} \cdot z$$

In fact, if $z = x + iy$, then $z^{-1} = \frac{1}{z} := \frac{x}{x^2 + y^2} - i \frac{y}{x^2 + y^2}$.

(9) Distributive Law. For every triple $z_1, z_2, z_3 \in \mathbb{C}$

$$(z_1 + z_2) \cdot z_3 = z_1 \cdot z_3 + z_2 \cdot z_3$$

Proof. (1) - (7) and (9) are left as Problem 4. One proves these directly by showing that the left hand side matches the right hand side.

(8) We note that

$$\begin{aligned} z \cdot \frac{1}{z} &= (x + iy) \left(\frac{x}{x^2 + y^2} - i \frac{y}{x^2 + y^2} \right) \\ &= (x + iy) \left(\frac{x}{x^2 + y^2} + i \frac{(-y)}{x^2 + y^2} \right) \\ &= \left(x \cdot \frac{x}{x^2 + y^2} - y \cdot \frac{(-y)}{x^2 + y^2} \right) + i \left(x \cdot \frac{(-y)}{x^2 + y^2} + y \cdot \frac{x}{x^2 + y^2} \right) \\ &= \left(\frac{x^2}{x^2 + y^2} + \frac{y^2}{x^2 + y^2} \right) + i \left(\frac{-yx + xy}{x^2 + y^2} \right) \\ &= \frac{x^2 + y^2}{x^2 + y^2} + i \cdot 0 \\ &= 1 \end{aligned}$$

Of course, we should comment that $z = (x, y) \neq (0, 0)$ if and only if $x^2 + y^2 \neq 0$ (one proves this by stating and proving the contrapositive). \square

Remark 1.1.7. In the language of algebra,

- (1) – (4) tells us that $(\mathbb{C}, +)$ is an abelian group.
- (5) – (8) tells us that (\mathbb{C}^*, \cdot) is an abelian group.
- (1) – (9) tells us that $(\mathbb{C}, +, \cdot)$ is a field.

Definition 1.1.8. Consider $z_1, z_2 \in \mathbb{C}$. We define *subtraction* and *division* as follows, respectively:

$$z_1 - z_2 := z_1 + (-z_2)$$

$$\frac{z_1}{z_2} := z_1 \cdot z_2^{-1} = z_1 \cdot \left(\frac{1}{z_2} \right), \quad z_2 \neq 0$$


Writing down z_1/z_2 as $x + iy$ is not easy to remember, one obtains it by a method akin to "rationalising the denominator", in this case we could call it "realifying the denominator"

$$\frac{z_1}{z_2} = \frac{x_1 + iy_1}{x_2 + iy_2} \cdot \frac{x_2 - iy_2}{x_2 - iy_2}$$


This method will be clarified soon when we talk about conjugates and absolute value.

1.2. Geometric Properties of Complex Numbers

As a set, we have $\mathbb{C} = \mathbb{R}^2$, so it's natural to visualise complex numbers as points in the **complex plane** (also called the **Argand plane**).



Geometrically, addition of complex numbers is just the addition of the corresponding vectors in the euclidean plane. We will soon see a geometric interpretation of multiplication.



Definition 1.2.1 (Modulus). The **modulus** (or **absolute value**) of a complex number $z = x + iy$, denoted $|z|$, is the length of the vector (x, y) , or equivalently its distance from the origin; namely

$$|z| := \sqrt{(\operatorname{Re} z)^2 + (\operatorname{Im} z)^2} = \sqrt{x^2 + y^2} = \|(x, y)\|$$

Notice that this extends the usual absolute value of real numbers, as the modulus of a real number is its absolute value.

We can then immediately derive a useful inequality,

$$|z|^2 = (\operatorname{Re} z)^2 + (\operatorname{Im} z)^2 \geq (\operatorname{Re} z)^2, (\operatorname{Im} z)^2,$$

giving us

$$\operatorname{Re} z \leq |\operatorname{Re} z| \leq |z| \quad \text{and} \quad \operatorname{Im} z \leq |\operatorname{Im} z| \leq |z|.$$

Definition 1.2.2 (Distance). The **distance** between two complex numbers z_1 and z_2 is

$$|z_1 - z_2| = \|(x_1, y_1) - (x_2, y_2)\| = \|(x_1 - x_2, y_1 - y_2)\|$$

That is, it's the euclidean distance between the vectors representing these complex numbers.


Discussion 1.2.3. The absolute value can be used to define various important subsets of \mathbf{C} .

- (1) • The *circle of radius $R > 0$ centered at z_0* is the set

$$C_R(z_0) = \{z \in \mathbf{C} : |z - z_0| = R\}$$

- The *open disk (or ball) of radius $R > 0$ centered at z_0* is the set

$$D_R(z_0) = \{z \in \mathbf{C} : |z - z_0| < R\}$$




- The *closed disk (or ball) of radius $R > 0$ centered at z_0* is the set

$$\overline{D}_R(z_0) = \{z \in \mathbf{C} : |z - z_0| \leq R\} = D_R(z_0) \cup C_R(z_0).$$

- (2) The *(open) annulus of inner radius $r > 0$ and outer radius $R > 0$ centered at z_0* is the set


$$A_{r,R}(z_0) = \{z \in \mathbf{C} : r < |z - z_0| < R\}$$



Proposition 1.2.4 (Triangle Inequalities). *For all $z_1, z_2 \in \mathbf{C}$, the following inequalities hold.*

- (1) $|z_1 + z_2| \leq |z_1| + |z_2|$.
- (2) $|z_1 \pm z_2| \geq ||z_1| - |z_2||$. We sometimes refer to this inequality as the **reverse triangle inequality**.

Proof.



- (1) A standard fact about triangles.
- (2) We first assume that $|z_1| \geq |z_2|$. Then, $||z_1| - |z_2|| = |z_1| - |z_2|$. Now, note that

$$\begin{aligned}
 |z_1| - |z_2| &= |z_1 \pm z_2 \mp z_2| - |z_2| \\
 &\leq |z_1 \pm z_2| + |\mp z_2| - |z_2|, \text{ triangle inequality} \\
 &= |z_1 \pm z_2| + |z_2| - |z_2| \\
 &= |z_1 \pm z_2|
 \end{aligned}$$

If we instead assume $|z_2| \geq |z_1|$, then we do the same computation with the roles of z_1 and z_2 switched. □

Lecture 3

Proposition 1.2.5 (Modulus is Multiplicative). *For all $z, w \in \mathbf{C}$ and positive integers n ,*

- (1) $|zw| = |z| |w|$.
- (2) $|z^n| = |z|^n$.

Proof.

- (1) Left as Problem 13. One proves these directly by showing that the left hand side matches the right hand side.
- (2) The proof of this is by induction. $n = 1$ is a tautology, and $n = 2$ is (1) in the case $w = z$. Assume the statement is true for $n = k$, that is $|z^k| = |z|^k$. Then, for $n = k + 1$


$$\begin{aligned}
 |z^{k+1}| &= |z^k \cdot z| = |z^k| |z|, \text{ using (1)} \\
 &= |z|^k |z|, \text{ using the induction hypothesis} \\
 &= |z|^{k+1}
 \end{aligned}$$

Therefore we have the result by the principle of mathematical induction. □

Definition 1.2.6 (Complex Conjugation). Given a complex number $z = x + iy$, its (complex) conjugate, denoted \bar{z} , is

$$\bar{z} := x - iy$$

Geometrically, \bar{z} is the reflection of z about the real axis.



Proposition 1.2.7 (Properties of Conjugation). For all pairs $z, w \in \mathbb{C}$, we have

- (1) $\overline{\bar{z}} = z$
- (2) $|\bar{z}| = |z|$
- (3) $\overline{z + w} = \bar{z} + \bar{w}$
- (4) $\overline{z\bar{w}} = \bar{z} w$
- (5) $z\bar{z} = |z|^2$
- (6) $\operatorname{Re} z = \frac{z + \bar{z}}{2}$ and $\operatorname{Im} z = \frac{z - \bar{z}}{2i}$
- (7) $z \in \mathbb{R}$ if and only if $z = \bar{z}$

Proof. (1) – (3) is clear geometrically. (4), (6) and (7) are left as Problem 14, (7) can be proved using (6) and can also be deduced geometrically. One proves these directly by showing that the left hand side matches the right hand side.

(5) Let $z = x + iy$, then

$$\begin{aligned} z\bar{z} &= (x + iy)(x - iy) \\ &= x^2 - ixy + iyx - i^2y^2 \\ &= x^2 + y^2 + i(yx - xy) = x^2 + y^2 = |z|^2 \end{aligned}$$

□

Discussion 1.2.8. Proposition 1.2.7 (5) gives us a nice formula for z^{-1} for $z \in \mathbb{C}^*$. For such a z , we have $z\bar{z} = |z|^2$, which gives us

$$z^{-1} = z^{-1} \cdot \frac{z\bar{z}}{|z|^2} = \frac{\bar{z}}{|z|^2}$$

This tells us that z^{-1} is just a scaled \bar{z} , which means, geometrically speaking, z^{-1} lies on the line passing through the origin and \bar{z} .

Lecture 4 Recall that every non-zero point $(x, y) \in \mathbf{R}^2$ can be re-written in polar coordinates (r, θ) as

$$x = r \cos \theta \quad \text{and} \quad y = r \sin \theta$$

This suggests the following definition.

Definition 1.2.9 (Polar Form). If (r, θ) are polar coordinates for a non-zero (x, y) , then the **polar form** of a non-zero complex number $z = x + iy$ is


$$z = r(\cos \theta + i \sin \theta)$$

We sometimes abbreviate $\cos \theta + i \sin \theta$ as $\text{cis } \theta$, so $z = r \text{cis } \theta$.

Evidently, (r, θ) are related to (x, y) by the equations

$$|z| = r \quad \text{and} \quad \cos \theta = \frac{x}{r} = \frac{x}{\sqrt{x^2 + y^2}}, \quad \sin \theta = \frac{y}{r} = \frac{y}{\sqrt{x^2 + y^2}}, \quad \text{so} \quad \tan \theta = \frac{y}{x}$$

We have to be careful and take into account which quadrant (x, y) belongs to, if we think of θ with respect to its formulation using \tan .



Since \sin and \cos are periodic functions, θ is not unique (you can replace θ with $\theta + 2\pi$). Each possible value of θ is called an **argument of z** , and the set of all such θ is denoted as $\arg z$. That is, if θ_0 is one solution of $\tan \theta = y/x$, then

$$\arg z = \{\theta_0 + 2k\pi : k \in \mathbf{Z}\}$$

The polar form, specifically θ is unique, as soon as we specify bounds on θ . The unique argument in the interval $(-\pi, \pi]$ is called the **principal argument** denoted $\text{Arg } z$. Precisely speaking,

Definition 1.2.10. For $z = x + iy$, we have

$$\text{Arg } z = \begin{cases} \arctan(y/x) & \text{if } x > 0 \quad (\text{quadrants I \& IV}) \\ \arctan(y/x) + \pi & \text{if } x < 0 \text{ and } y > 0 \quad (\text{quadrant II}) \\ \arctan(y/x) - \pi & \text{if } x < 0 \text{ and } y < 0 \quad (\text{quadrant III}) \end{cases}$$

Notice that we can then write

$$\arg z = \{\text{Arg } z + 2k\pi : k \in \mathbf{Z}\}$$

Definition 1.2.11 (Euler's Formula). $e^{i\theta} := \operatorname{cis} \theta = \cos \theta + i \sin \theta$. Therefore $|e^{i\theta}| = 1$.

Remark on Definition 1.2.11. This is for now a stopgap, defining $e^{i\theta}$ in this way. In a few weeks, we'll see that this is truly an equality of holomorphic functions. Euler deduced this by looking at the Taylor series expansion of these functions. We haven't built or discussed enough machinery to give this reasoning a solid foundation yet.

Using Euler's formula, one can write the polar form of a non-zero complex number, even more succinctly in its **exponential form**


$$z = re^{i\theta}$$

Example 1.2.12.

(1) Exponential form of $1 + i$,

$$|1 + i| = \sqrt{1^2 + 1^2} = \sqrt{2} \quad \text{and} \quad \operatorname{Arg} z = \arctan(1) = \frac{\pi}{4}$$

$$\text{So, } 1 + i = \sqrt{2}e^{i\pi/4}.$$




(2) Note that

$$1 = e^{i0} = e^{i2n\pi} \text{ for any } n \in \mathbf{Z}, \quad i = e^{i\pi/2}, \quad -1 = e^{i\pi} = e^{i(2n+1)\pi} \text{ for any } n \in \mathbf{Z}$$

One could write $-i = e^{i3\pi/2}$ but $3\pi/2 \neq \operatorname{Arg}(-i)$; instead we should write $-i = e^{-i\pi/2}$.

(3) The circle $C_R(z_0)$ has a nice parametrisation

$$C_R(z_0) = \{z = z_0 + Re^{i\theta} : 0 \leq \theta < 2\pi\}$$



Example 1.2.13 (in-class). Write the exponential form of $z = 1 - i$.

Answer. As a point on the plane, since $\operatorname{Re} z > 0$ and $\operatorname{Im} z < 0$, the complex number $z = 1 - i$ lies in the fourth quadrant. Thus,

$$r = |z| = \sqrt{(1)^2 + (-1)^2} = \sqrt{2}$$

$$\operatorname{Arg} z = \arctan(-1) = -\frac{\pi}{4}$$

Thus, $1 - i = \sqrt{2}e^{-i\pi/4}$. □

Proposition 1.2.14 (Properties of Exponential Form). Let $z = re^{i\theta}$ and $w = se^{i\phi}$ be non-zero complex numbers. Then

- (1) $zw = rs e^{i(\theta+\phi)}$
- (2) $z^{-1} = (1/r)e^{-i\theta}$
- (3) $z^n = r^n e^{in\theta}$, for any $n \in \mathbf{Z}$
- (4) $\bar{z} = re^{-i\theta}$
- (5) $z/w = (r/s)e^{i(\theta-\phi)}$

Proof.

- (1) Note that

$$\begin{aligned} zw &= (re^{i\theta})(se^{i\phi}) = rs(\cos \theta + i \sin \theta)(\cos \phi + i \sin \phi) \\ &= rs((\cos \theta \cos \phi - \sin \theta \sin \phi) + i(\cos \theta \sin \phi + \sin \theta \cos \phi)) \\ &= rs(\cos(\theta + \phi) + i \sin(\theta + \phi)) \\ &= rs e^{i(\theta+\phi)} \end{aligned}$$

- (2) It suffices to show that $(re^{i\theta})((1/r)e^{-i\theta}) = 1$, for which we use (1).

- (3) We first prove this result for $n \geq 0$, the result is clear for $n = 0$ and $n = 1$. Assume the result is true for $n = k$, that is $z^k = r^k e^{ik\theta}$. Then, for $n = k + 1$

$$\begin{aligned} z^{k+1} &= z^k z \\ &= (r^k e^{ik\theta})(re^{i\theta}) \text{ using the induction hypothesis} \\ &= r^{k+1} e^{ik\theta+\theta} \text{ by (1)} \\ &= r^{k+1} e^{i(k+1)\theta} \end{aligned}$$


Therefore we have the result by the principle of mathematical induction.

Suppose $n < 0$ instead, then write $n = -m$ for a positive $m > 0$. Now, we can apply the first case to $z^n := (z^{-1})^m$ to get our result.

(4) Using $z\bar{z} = |z|^2 = r^2$, we get that $\bar{z} = r^2 z^{-1}$, and the result follows from (2).

(5) Recall $z/w = zw^{-1}$, and the result follows from (2) and (1). □

Discussion 1.2.15. Proposition 1.2.14 (1) gives us a nice geometric interpretation of complex multiplication. If $z = re^{i\theta}$ and $w = se^{i\phi}$, then $zw = rs e^{i(\theta+\phi)}$. This can be interpreted as saying that zw is obtained from w by scaling w by $|z| = r$ and rotating w by an angle of $\text{Arg } z$ (or vice versa).



Example 1.2.16. Let's use Proposition 1.2.14 to compute $(1 + i)^{2023}$, then

$$\begin{aligned}
 (1 + i)^{2023} &= (\sqrt{2}e^{i\pi/4})^{2023} \\
 &= (\sqrt{2})^{2023}(e^{i\pi/4})^{2023} \\
 &= (\sqrt{2})^{2022}\sqrt{2}(e^{i\pi/4})^{2024}e^{-i\pi/4} \\
 &= 2^{1011}\sqrt{2}(e^{i506\pi})e^{-i\pi/4} \\
 &= 2^{1011}\sqrt{2}e^{-i\pi/4} \\
 &= 2^{1011}(1 - i)
 \end{aligned}$$

Example 1.2.17. Compute $(1 + i\sqrt{3})^{101}$.

Answer. We will first compute the exponential form of our complex number. Note that

$$|1 + i\sqrt{3}| = \sqrt{1^2 + (\sqrt{3})^2} = \sqrt{4} = 2,$$

and since $1 + i\sqrt{3}$ lies in the first quadrant of the complex plane

$$\text{Arg } z = \arctan(\sqrt{3}) = \frac{\pi}{3}$$

Therefore

$$1 + i\sqrt{3} = 2e^{i\pi/3}$$

and so

$$\begin{aligned}
 (1 + i\sqrt{3})^{101} &= (2e^{i\pi/3})^{101} \\
 &= (2e^{i\pi/3})^{99} (2e^{i\pi/3})^2 \\
 &= 2^{99} e^{i33\pi} (1 + i\sqrt{3})^2 \\
 &= -2^{99} (1 - 3 + 2i\sqrt{3}), \quad \text{since 33 is odd} \\
 &= -2^{99} (-2 + 2i\sqrt{3}) \\
 &= 2^{100} (1 - i\sqrt{3})
 \end{aligned}$$

□

Lecture 5 A few more interesting consequences of Proposition 1.2.14 are

(1) The *unit circle*

$$S^1 = \{z \in \mathbf{C} : |z| = 1\} = \{e^{i\theta} : \theta \in \mathbf{R}\}$$

is closed under multiplication. It's in fact an abelian group, usually denoted $U(1)$.

(2) *De Moivre's Theorem*. From Proposition 1.2.14 (4) applied to $z = e^{i\theta}$ we get

$$(\cos \theta + i \sin \theta)^n = (\cos n\theta + i \sin n\theta)$$

Proposition 1.2.18 (Arguments of Products). *Let z, w be non-zero complex numbers, then*

$$(1) \arg(zw) = \arg z + \arg w$$

$$(2) \arg w^{-1} = -\arg w$$

Note that this is *not* saying $\text{Arg}(zw) = \text{Arg } z + \text{Arg } w$, this is actually not true, we're claiming an equality of sets. (1) and (2) together give us $\arg(z/w) = \arg z - \arg w$.

Proof.

(1) Consider $\theta \in \arg z$ and $\phi \in \arg w$, so $z = re^{i\theta}$ and $w = se^{i\phi}$. By Proposition 1.2.14 (1), we have $zw = rs e^{i(\theta+\phi)}$ and therefore $\theta + \phi \in \arg(zw)$. Hence $\arg z + \arg w \subseteq \arg(zw)$.

Consider $\psi \in \arg(zw)$, and some $\theta \in \arg z$ then we claim that $\psi - \theta \in \arg w$. We have $rs e^{i\psi} = zw = re^{i\theta}w$, then by Proposition 1.2.14 (5), we get $w = sr^{i(\psi-\theta)}$. Hence $\psi - \theta \in \arg w$, and since $\psi = \theta + (\psi - \theta) \in \arg z + \arg w$, we have $\arg(zw) \subseteq \arg z + \arg w$.

Therefore $\arg(zw) = \arg z + \arg w$.

(2) Consider $\theta \in \arg z$, so $z = re^{i\theta}$. By Proposition 1.2.14 (2), we have $z^{-1} = (1/r)e^{i(-\theta)}$ and therefore $-\theta \in \arg z^{-1}$. Hence $-\arg z \subseteq \arg z^{-1}$.

Note that $w = (w^{-1})^{-1}$, applying the above result to w^{-1} gets us $-\arg w^{-1} \subseteq \arg(w^{-1})^{-1} = \arg w$ and so $\arg w^{-1} \subseteq -\arg w$.

Therefore $\arg w^{-1} = -\arg w$.

□

Remark 1.2.19. For a complex number, $\arg z$ is a set of all possible θ 's such that we can write $z = |z|e^{i\theta}$, as you know. Therefore, we will abuse notation by sometimes calling any $\theta \in \arg z$ as an argument of z , and sometimes also writing $z = |z|e^{i\arg z}$. That is, we are not, or are careless about, distinguishing the set $\arg z$ and its element when we can be agnostic about the choice of θ ; for example, the polar form of a complex number. It will be clear when we choose to care about our choice, it will be evident because we'll be then forcing θ to lie in an interval of length 2π ; for example, the principal argument $-\pi < \text{Arg } z \leq \pi$.

Example 1.2.20.

- (1) The principal argument of $z = (\sqrt{3} - i)^6$. We first note that $\text{Arg}(\sqrt{3} - i) = -\pi/6$. By Proposition 1.2.18 (1), applied inductively, we have

$$\arg(\sqrt{3} - i)^6 = \underbrace{\arg(\sqrt{3} - i) + \cdots + \arg(\sqrt{3} - i)}_{6 \text{ times}} = \{-\pi + 2k\pi : k \in \mathbf{Z}\}$$

Then $\text{Arg}(\sqrt{3} - i)^6$ is the element in the set above in the interval $(-\pi, \pi]$ which is π .

- (2) As mentioned previously, we can't just replace \arg with Arg in the statement of Proposition 1.2.18 (1). Here's a simple example: let $z = w = -1$, then $\text{Arg } z = \text{Arg } w = \pi$ and $\text{Arg } zw = \text{Arg } 1 = 0$ but $0 \neq 2\pi = \text{Arg } z + \text{Arg } w$.
- (3) Note that $\arg z + \arg z \neq 2 \arg z$.

1.3. Roots of Complex Numbers

Lemma 1.3.1. Two non-zero complex numbers z, w are equal if and only if $|z| = |w|$ and $\arg z = \arg w$.

Proof. If $|z| = |w|$ and $\arg z = \arg w$, then clearly $z = w$.

Suppose $z = w$, then we immediately get $|z| = |w|$. Consider $\theta \in \arg z$ and $\phi \in \arg w$, then we get $e^{i\theta} = e^{i\phi}$ which is equivalent to saying $\cos(\theta - \phi) + i \sin(\theta - \phi) = e^{i(\theta - \phi)} = 1$. This gives us

$$\sin(\theta - \phi) = 0.$$

The solution to this is $\theta - \phi = 2k\pi$ for some $k \in \mathbf{Z}$. This gives us $\arg z = \arg w$. □

Lecture 6

Definition 1.3.2 (Roots). Let α be a non-zero complex number. An n^{th} root of α is a solution to the polynomial equation $z^n - \alpha = 0$.

The set of all n^{th} roots of α is denoted by $\alpha^{1/n}$, we reserve the symbol $\sqrt[n]{\cdot}$ for the unique positive n^{th} root of a positive real number.

Proposition 1.3.3 (Distinct Roots). There are precisely n distinct n^{th} roots of α , namely

$$\beta_k = \sqrt[n]{|\alpha|} e^{i\left(\frac{\text{Arg } \alpha}{n} + \frac{2k\pi}{n}\right)}, \quad k = 0, \dots, n-1$$

Proof. Let $z = re^{i\theta}$ and $\alpha = |\alpha| e^{i \operatorname{Arg} \alpha}$, we solve

$$r^n e^{in\theta} = z^n = \alpha = |\alpha| e^{i \operatorname{Arg} \alpha}.$$

By Lemma 1.3.1, this equality is true if and only if $r^n = |\alpha|$ and $n\theta = \operatorname{Arg} \alpha + 2k\pi$ for some $k \in \mathbf{Z}$. Therefore

$$z = \sqrt[n]{|\alpha|} e^{i\left(\frac{\operatorname{Arg} \alpha}{n} + \frac{2k\pi}{n}\right)}, \quad k \in \mathbf{Z}$$

We obtain distinct n complex numbers for $k = 0, \dots, n-1$ since they have distinct arguments, and they necessarily give us the n distinct n^{th} roots of α . \square

Discussion 1.3.4. With the notation of Proposition 1.3.3, the n^{th} principal root of α is

$$\beta_0 = \sqrt[n]{|\alpha|} e^{i\frac{\operatorname{Arg} \alpha}{n}}$$

If we introduce the notation $\zeta_n = e^{\frac{2\pi i}{n}}$, then

$$\zeta_n^k = e^{\frac{2k\pi i}{n}}$$

According to the proposition, the complex numbers


$$1, \zeta_n, \zeta_n^2, \dots, \zeta_n^{n-1}$$

are the distinct solutions to $z^n - 1 = 0$, the n^{th} roots of unity, making ζ_n the primitive n^{th} root of unity as it generates all n^{th} other roots of unity.

Then we can write the roots of α in terms of the principal root and the primitive root of unity

$$\begin{aligned} \beta_k &= \sqrt[n]{|\alpha|} e^{i\left(\frac{\operatorname{Arg} \alpha}{n} + \frac{2k\pi}{n}\right)} \\ &= \sqrt[n]{|\alpha|} e^{i\frac{\operatorname{Arg} \alpha}{n}} e^{\frac{2k\pi i}{n}} = \beta_0 \zeta_n^k \end{aligned}$$

That is, β_k 's all lie on the circle of radius $\sqrt[n]{|\alpha|}$ centered at the origin, and all of them are obtained by rotating β_0 by an angle of $2k\pi/n$. That is, they all lie on the vertices of an inscribed regular n -gon.




Example 1.3.5.

- (1) We compute explicitly the 4th roots of $\alpha = -16$. As a negative real number, $\text{Arg}(-16) = \pi$, so

$$\begin{aligned}\beta_k &= \sqrt[4]{16} e^{i(\frac{\pi}{4} + \frac{2k\pi}{4})} = 2 e^{i\frac{\pi}{4}} e^{\frac{ki\pi}{2}} \\ &= 2 e^{i\frac{\pi}{4}} \left(e^{\frac{i\pi}{2}} \right)^k \\ &= 2 \left(\cos \frac{\pi}{4} + i \sin \frac{\pi}{4} \right) \left(\cos \frac{\pi}{2} + i \sin \frac{\pi}{2} \right)^k = 2 \left(\frac{1}{\sqrt{2}} + i \frac{1}{\sqrt{2}} \right) i^k = \sqrt{2}(1+i)i^k\end{aligned}$$

Therefore



$$\beta_0 = \sqrt{2}(1+i), \quad \beta_1 = \sqrt{2}(-1+i), \quad \beta_2 = \sqrt{2}(-1-i), \quad \beta_3 = \sqrt{2}(1-i)$$


- (2) In the course of the previous example, we have computed the 4th roots of unity, since they are

$$e^{\frac{2ki\pi}{4}} = e^{\frac{ki\pi}{2}}, \quad k = 0, 1, 2, 3$$

as $\text{Arg } 1 = 0$. Letting $\zeta_4 = e^{i\pi/2} = i$, the 4th roots of unity are $\zeta_4^0, \zeta_4^1, \zeta_4^2, \zeta_4^3$, which are nothing but $\pm 1, \pm i$. Furthermore, note that i is the primitive 4th root of unity.

Lecture 7

Example 1.3.6. We compute the 3rd roots of unity, also called the cube roots of unity where we denote $\omega = \zeta_3$, explicitly.



Let the primitive root be $\omega = \zeta_3$, then the cube roots of unity are

$$1, \omega, \omega^2$$

where we have

$$\omega = e^{\frac{2\pi i}{3}} = \left(\cos \frac{2\pi}{3} + i \sin \frac{2\pi}{3} \right) = -\frac{1}{2} + i\frac{\sqrt{3}}{2}$$

$$\omega^2 = e^{\frac{4\pi i}{3}} = \left(\cos \frac{4\pi}{3} + i \sin \frac{4\pi}{3} \right) = -\frac{1}{2} - i\frac{\sqrt{3}}{2}$$

1.4. Basic Topology of \mathbb{C}

Our purpose now is to define the kind of subsets of \mathbb{C} that are suitable for doing complex analysis, namely *non-empty open connected sets*.


Definition 1.4.1 (Open Disks or Neighbourhoods). Let $\varepsilon > 0$. Recall the **open disk** (of radius ε centered at z_0) is the set

$$D_\varepsilon(z_0) = \{z \in \mathbb{C} : |z - z_0| < \varepsilon\}.$$

We also refer to such an open disk as an **ε -neighbourhood** or simply a **neighbourhood**.

A **deleted** (or **punctured**) **open disk** (or **neighbourhood**) is a set of the form

$$D_\varepsilon(z_0) \setminus \{z_0\} = \{z \in \mathbb{C} : 0 < |z - z_0| < \varepsilon\}.$$




Points belonging to the same ε -neighbourhood are considered "close" to each other, in the sense that they are within a distance of 2ε from each other.

Definition 1.4.2 (Various kinds of Points). Consider a $S \subseteq \mathbb{C}$.

- A point $z \in S$ is an **interior point of S** if there exists an $\varepsilon > 0$ such that $D_\varepsilon(z) \subseteq S$.
- A point $z \notin S$ is an **exterior point of S** if there exists an $\varepsilon > 0$ such that $D_\varepsilon(z) \cap S = \emptyset$.
- A point $z \in \mathbb{C}$ is a **boundary point of S** if it's neither an interior nor an exterior point of S . Equivalently, if every neighbourhood of z contains both a point in S and not in S .
- A point $z \in \mathbb{C}$ is a **accumulation** (or **cluster**) **point of S** if for every $\varepsilon > 0$ we have

$$D_\varepsilon(z) \setminus \{z\} \cap S \neq \emptyset.$$

- A point $z \in S$ is an **isolated point of S** if there exists an $\varepsilon > 0$ such that $D_\varepsilon(z) \setminus \{z\} \cap S = \emptyset$. Isolated points are examples of boundary point



Here z_1 is a boundary point, z_2 an interior point, z_3 an exterior point, and z_4 is an isolated point (and a boundary point).

Remark 1.4.3. The idea is that if we don't move too far from an interior point of S then we remain in S ; a similar idea holds for an exterior point. But at a boundary point we can make an arbitrarily small move and get to a point inside S , and we can also make an arbitrarily small move and get to a point outside S . An accumulation point is one where it has other points from S within any arbitrarily small distance, i.e. points "accumulate" near it; an isolated point is the exact opposite.

Definition 1.4.4 (Open and Closed Sets). Consider a $S \subseteq \mathbb{C}$.

- The **interior of S** is the set of all interior points of S , denoted S° .
- S is said to be **open** if $S = S^\circ$.
- The **boundary of S** is the set of all boundary points of S , denoted ∂S .
- S is said to be **closed** if $\partial S \subseteq S$. Equivalently, if its complement is open.
- The **closure of S** is the set $S \cup \partial S$, denoted \bar{S} .


Example 1.4.5.

- (1) The open disks $D_R(z_0)$ are truly open sets, and the closed disks $\bar{D}_R(z_0)$ are truly closed sets. The closure of the open disk $D_R(z_0)$ is $\bar{D}_R(z_0)$. The boundary of $D_R(z_0)$ is the circle $C_R(z_0)$.
- (2) Consider the upper half-plane

$$\mathbf{H} = \{z \in \mathbb{C} : \text{Im } z > 0\},$$

then we have $\mathbf{H}^\circ = \mathbf{H}$. Since by definition $\mathbf{H}^\circ \subseteq \mathbf{H}$, it's enough to prove $\mathbf{H} \subseteq \mathbf{H}^\circ$. Consider any $z \in \mathbf{H}$, then $\text{Im } z > 0$. Let $\varepsilon = (\text{Im } z)/2$, we claim that

$$D_\varepsilon(z) \subseteq \mathbf{H}$$



Lecture 8

Let $w \in D_\varepsilon(z)$, then

$$|w - z| < \varepsilon = \frac{\text{Im } z}{2}$$

The end of Discussion 1.2.1 tells us

$$\begin{aligned} \frac{\text{Im } z}{2} &> |w - z| \geq |\text{Im}(w - z)| \\ &= |\text{Im } w - \text{Im } z| \end{aligned}$$

The later is simply the absolute value of a real number, which gives

$$-\frac{\text{Im } z}{2} < \text{Im } w - \text{Im } z < \frac{\text{Im } z}{2}$$

Adding $\text{Im } z$ throughout the inequality, we get from the inequality on the left hand side

$$\text{Im } w > \frac{\text{Im } z}{2} > 0.$$


Therefore $w \in \mathbf{H}$, and hence $D_\varepsilon(z) \subseteq \mathbf{H}$. Thus $\mathbf{H}^\circ = \mathbf{H}$.

The points exterior to \mathbf{H} are points z such that $\text{Im } z < 0$. That is, the exterior of the upper half-plane is the (open) lower half-plane. The boundary of \mathbf{H} consists of precisely points z whose $\text{Im } z = 0$. That is, $\partial\mathbf{H} = \mathbf{R}$.

The closure of \mathbf{H} is $\overline{\mathbf{H}} = \{z \in \mathbf{C} : \text{Im } z \geq 0\}$. While $\mathbf{H} \cup \{0\}$ is neither open nor closed.

Definition 1.4.6 (Bounded Sets). A set $S \subseteq \mathbf{C}$ is **bounded** if $S \subseteq D_M(0)$ for some $M > 0$. That is, there exists an $M > 0$ such that $|z| \leq M$ for every $z \in S$.

Definition 1.4.7 (Connected Sets). A set $S \subseteq \mathbf{C}$ is said to be **connected** if each pair of points z_1 and z_2 in S can be joined by a *polygonal line*, consisting of a finite number of line segments joined end to end, that lies entirely in S . Otherwise, we say it is **disconnected**.



Definition 1.4.8 (Domain). $S \subseteq \mathbf{C}$ is called a **domain** if it's a non-empty open and connected set. A **region** is a domain together with some or all of its boundary points.

Remark 1.4.9. Domains and regions are sets we will find most suitable for stating elegant results about certain functions in a complex variable.

Example 1.4.10. \mathbf{H} is a domain since it's non-empty, open and any two points in \mathbf{H} can be connected by a straight line. It's an unbounded set. An example of a region is $\mathbf{H} \cup \{0\}$.

2. Part II. Holomorphic Functions

2.1. Complex Functions

Definition 2.1.1. A function $f : G \rightarrow \mathbf{C}$ is a rule that assigns to each $z \in G$ a unique number $f(z) \in \mathbf{C}$.

The set G is called the *domain (of definition)*. If $S \subseteq G$, then

$$f(S) := \{f(z) : z \in S\}$$

is called the *image of S under f* .

The set $f(G)$ is called the *image (or range) of f* . Points in $f(G)$ are called *values of f* .

Given a function f , we define its conjugate \bar{f} by the rule $\bar{f}(z) := \overline{f(z)}$.

Lecture 9

Discussion 2.1.2. If $f : G \rightarrow \mathbf{C}$ is a function, then the value $f(x + iy) = u + iv$ depends on a pair $(x, y) \in \mathbf{R}^2$. Collecting all values, we decompose f into its **real** and **imaginary parts**

$$f(z) = f(x + iy) = u(x, y) + i v(x, y); \quad \operatorname{Re} f = u \quad \text{and} \quad \operatorname{Im} f = v,$$

where $u, v : \mathbf{R}^2 \rightarrow \mathbf{R}$ are real-valued functions in two real variables.

In practice, as the examples below tell us, this means replace your $z = x + iy$ and do the required operations to the output $f(x + iy)$. The resulting complex number will be, as a complex number, of the form $u + iv$. The real part is u , which you will obtain in terms of x and y , and the imaginary part is v , which you will also obtain in terms of x and y .

Example 2.1.3 (Some Complex Functions).

(1) $f(z) = z^2 = (x + iy)^2 = (x^2 - y^2) + i(2xy)$. So,

$$u(x, y) = x^2 - y^2 \quad \text{and} \quad v(x, y) = 2xy.$$

(2) $f(z) = \bar{z} = x - iy$. So,

$$u(x, y) = x \quad \text{and} \quad v(x, y) = -y.$$

(3) (in-class) $f(z) = z\bar{z} = |z|^2 = x^2 + y^2$. So,

$$u(x, y) = x^2 + y^2 \quad \text{and} \quad v(x, y) = 0.$$

Such a function is *real-valued*.

(4) *Polynomials of degree n* are functions of the form

$$p(z) = a_0 + a_1 z + \cdots + a_n z^n,$$

where $a_i \in \mathbf{C}$ and $a_n \neq 0$.

A polynomial of degree 0 is simply a non-zero complex number, sometimes also referred to as a *constant polynomial*.

(5) *Rational functions (or polynomials)* are functions of the form

$$\frac{p(z)}{q(z)}$$

where $p(z)$ and $q(z)$ are polynomials. The domain of definition is wherever $q(z) \neq 0$. For example,

$$f : \mathbb{C}^* \rightarrow \mathbb{C}, z \mapsto \frac{1}{z}$$

(6) If we express z in its polar form, then a function f , when we restrict its domain of definition within \mathbb{C}^* , can be written as

$$f(z) = f(re^{i\theta}) = u(r, \theta) + i v(r, \theta)$$

For example,

$$f : \mathbb{C}^* \rightarrow \mathbb{C}, z = re^{i\theta} \mapsto \frac{1}{z} = \frac{1}{r} e^{-i\theta} = \frac{\cos \theta}{r} - i \frac{\sin \theta}{r}.$$

Here $u(r, \theta) = \frac{\cos \theta}{r}$ and $v(r, \theta) = -\frac{\sin \theta}{r}$.

(7) (in-class) Let's consider the function $f(z) = \bar{z}^2$, in polar form we have

$$\begin{aligned} f(re^{i\theta}) &= (\overline{re^{i\theta}})^2 \\ &= (re^{-i\theta})^2, \quad \text{by Proposition 1.2.14 (4)} \\ &= r^2 e^{-i2\theta}, \quad \text{by Proposition 1.2.14 (3)} \\ &= r^2 (\cos(-2\theta) + i \sin(-2\theta)) \\ &= r^2 \cos(2\theta) - i \sin(2\theta) \end{aligned}$$

Therefore, here $u(r, \theta) = r^2 \cos(2\theta)$ and $v(r, \theta) = -r^2 \sin(2\theta)$.

(8) Consider $f(z) = z^{1/n}$, where n is a non-zero integer. For no $n \neq 1$ is this a function! We have seen previously that $z^{1/n}$ has n -distinct values. Such a "function" is called multi-valued.

We can make this into a (single-valued) function by assigning a single value of $z^{1/n}$ to each z ; taking the *principal n^{th} root* of z , for instance. More on such functions soon.


2.2. Limits of Functions

Definition 2.2.1 (Limit of a Function). Consider a function $f : G \rightarrow \mathbb{C}$, and an accumulation point z_0 of G .

We say that **limit** of f , as z approaches z_0 , is $w_0 \in \mathbb{C}$ if for all $\varepsilon > 0$ there exists a $\delta > 0$ such that

$$\text{if } 0 < |z - z_0| < \delta, \quad \text{then } |f(z) - w_0| < \varepsilon$$

Equivalently, if $z \in D_\delta(z_0) \setminus \{z_0\}$, then $f(z) \in D_\varepsilon(w_0)$.



In this case we write $\lim_{z \rightarrow z_0} f(z) = w_0$ or $f(z) \rightarrow w_0$, as $z \rightarrow z_0$.

Intuitively, the limit of f at z_0 is w_0 if

" f is arbitrarily close to w_0 eventually, that is sufficiently, near z_0 ".

How close? Within an error of ε . How near, eventually? Within a distance of δ .

Lecture 10 **Example 2.2.2.** Let's show that $\lim_{z \rightarrow i} z^2 = -1$ using the definition.

Proof. Let $\varepsilon > 0$ be arbitrary. Note that $|z^2 - (-1)| = |z - i| |z + i|$. We make an initial estimate, suppose $0 < |z - i| < 1$, then

$$\begin{aligned} |z + i| &= |z - i + 2i| \\ &\leq |z - i| + |2i| \\ &< 1 + 2 \\ &= 3 \end{aligned}$$

Now, if we choose $\delta = \min \left\{ \frac{\varepsilon}{3}, 1 \right\}$, then if $0 < |z - i| < \delta$ we get

$$0 < |z - i| < 1 \text{ and } \frac{\varepsilon}{3}$$

So,

$$\begin{aligned} |z^2 - (-1)| &= |z - i| |z + i| \\ &< 3 |z - i|, \quad \text{since } |z - i| < 1 \\ &< 3 \cdot \frac{\varepsilon}{3}, \quad \text{since } |z - i| < \frac{\varepsilon}{3} \\ &= \varepsilon \end{aligned}$$

Therefore $\lim_{z \rightarrow i} z^2 = -1$. □

Theorem 2.2.3. If f has a limit at z_0 , then it is unique.

Proof. Assume

$$\lim_{z \rightarrow z_0} f(z) = \alpha \quad \text{and} \quad \lim_{z \rightarrow z_0} f(z) = \beta$$

Consider an arbitrary $\varepsilon > 0$, then we can find a $\delta_1 > 0$ such that

$$\text{if } 0 < |z - z_0| < \delta, \quad \text{then } |f(z) - \alpha| < \frac{\varepsilon}{2}$$

and $\delta_2 > 0$ such that

$$\text{if } 0 < |z - z_0| < \delta, \quad \text{then } |f(z) - \beta| < \frac{\varepsilon}{2}$$

Define $\delta := \min \{\delta_1, \delta_2\} \leq \delta_1, \delta_2$, then if $0 < |z - z_0| < \delta$ we have

$$\begin{aligned} |\alpha - \beta| &= |f(z) - f(z) + \alpha - \beta| \\ &\leq |\alpha - f(z)| + |f(z) - \beta| \\ &= |f(z) - \alpha| + |f(z) - \beta| \\ &< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \\ &= \varepsilon \end{aligned}$$

We have proven that $|\alpha - \beta| < \varepsilon$ for any $\varepsilon > 0$. Now, suppose $\alpha \neq \beta$, then for $\varepsilon = |\alpha - \beta| > 0$ we get $|\alpha - \beta| < |\alpha - \beta|$, which is preposterous. Hence $\alpha = \beta$, and thus the limit is unique. \square

Remark 2.2.4. The reason we require that z_0 be an accumulation point of the domain of f is just that we need to be sure that there are points z of the domain that are arbitrarily close to z_0 . That is, there are indeed points satisfying $0 < |z - z_0| < \delta$.

Our definition (i.e., the part that says $0 < |z - z_0|$) does not require z_0 to be in the domain of f , and if z_0 is in the domain of f , the definition explicitly ignores the value of $f(z_0)$.

Uniqueness of limits can be used to show that a limit does not exist.

Example 2.2.5. The function $f(z) = \frac{\bar{z}}{z}$ has no limit at 0.

Discussion of Example 2.2.5. Let $z = x + iy$, then

$$f(z) = \frac{x - iy}{x + iy}$$

Along the real axis, $\text{Im } z = 0$, and so $z = x$, giving us $f(z) = \frac{x}{x} = 1$.

Along the imaginary axis, $\text{Re } z = 0$, and so $z = iy$, giving us $f(z) = \frac{-y}{y} = -1$.

Taking the limit along these axes gives us different values of the limit, 1 and -1 . Hence, by the uniqueness of limits, the limit doesn't exist. \square

2.3. Theorems on Limits

Theorem 2.3.1 (Limit in terms of Real and Imaginary parts of a Function). *Suppose that*

$$f(z) = f(x + iy) = u(x, y) + i v(x, y)$$

Then

$$\lim_{x+iy \rightarrow x_0+iy_0} f(x + iy) = u_0 + i v_0$$

if and only if

$$\lim_{(x,y) \rightarrow (x_0,y_0)} u(x, y) = u_0 \quad \text{and} \quad \lim_{(x,y) \rightarrow (x_0,y_0)} v(x, y) = v_0$$

Proof. (\Rightarrow) Consider an arbitrary $\varepsilon > 0$, then there exists a $\delta > 0$ such that if

$$0 < |(x + iy) - (x_0 + iy_0)| < \delta$$

$$\text{then } |f(x + iy) - (u_0 + i v_0)| = |(u(x, y) + i v(x, y)) - (u_0 + i v_0)| < \varepsilon$$

We first note that, by definition

$$\|(x, y) - (x_0, y_0)\| = |(x + iy) - (x_0 + iy_0)|$$

and the end of Discussion 1.2.1 tells us that

$$|u(x, y) - u_0| \leq |(u(x, y) + i v(x, y)) - (u_0 + i v_0)| < \varepsilon$$

$$|v(x, y) - v_0| \leq |(u(x, y) + i v(x, y)) - (u_0 + i v_0)| < \varepsilon$$

That is, we have that

$$\text{if } 0 < \|(x, y) - (x_0, y_0)\| < \delta, \quad \text{then } |u(x, y) - u_0| < \varepsilon \quad \text{and} \quad |v(x, y) - v_0| < \varepsilon$$

Therefore,

$$\lim_{(x,y) \rightarrow (x_0,y_0)} u(x, y) = u_0 \quad \text{and} \quad \lim_{(x,y) \rightarrow (x_0,y_0)} v(x, y) = v_0$$

(\Leftarrow) Consider an arbitrary $\varepsilon > 0$, then there exists a $\delta_1 > 0$ such that

$$\text{if } 0 < \|(x, y) - (x_0, y_0)\| < \delta_1, \quad \text{then } |u(x, y) - u_0| < \frac{\varepsilon}{2}$$

and there exists a $\delta_2 > 0$ such that

$$\text{if } 0 < \|(x, y) - (x_0, y_0)\| < \delta_2, \quad \text{then } |v(x, y) - v_0| < \frac{\varepsilon}{2}$$

Define $\delta := \min \{\delta_1, \delta_2\} \leq \delta_1, \delta_2$. Now, if

$$0 < |(x + iy) - (x_0 + iy_0)| = \|(x, y) - (x_0, y_0)\| < \delta$$

then

$$\begin{aligned}
|f(x + iy) - (u_0 + iv_0)| &= |(u(x, y) + i v(x, y)) - (u_0 + iv_0)| \\
&= |(u(x, y) - u_0) + i(v(x, y) - v_0)| \\
&\leq |(u(x, y) - u_0)| + |i(v(x, y) - v_0)|, \text{ by triangle identity} \\
&= |(u(x, y) - u_0)| + |i| |v(x, y) - v_0| \\
&= |(u(x, y) - u_0)| + |v(x, y) - v_0| \\
&< \frac{\varepsilon}{2} + \frac{\varepsilon}{2} \\
&= \varepsilon
\end{aligned}$$

Therefore,

$$\lim_{x+iy \rightarrow x_0+iy_0} f(x + iy) = u_0 + iv_0$$

□

Theorem 2.3.2 (Limit Laws). Suppose

$$\lim_{z \rightarrow z_0} f(z) = \alpha \quad \text{and} \quad \lim_{z \rightarrow z_0} g(z) = \beta$$

Then

- (1) $\lim_{z \rightarrow z_0} (f(z) + g(z)) = \alpha + \beta$
- (2) $\lim_{z \rightarrow z_0} (f(z) g(z)) = \alpha \beta$
- (3) $\lim_{z \rightarrow z_0} \frac{f(z)}{g(z)} = \frac{\alpha}{\beta}$, provided $\beta \neq 0$.

Proof. The proof follows from Theorem 2.3.1 and limit laws from Calculus.

□

Example 2.3.3. Let $p(z)$ be a polynomial, then

$$\lim_{z \rightarrow z_0} p(z) = p(z_0)$$


Write $p(z) = a_0 + a_1 z + \cdots + a_n z^n$, then by Theorem 2.3.2 we have

$$\begin{aligned}
\lim_{z \rightarrow z_0} p(z) &= \lim_{z \rightarrow z_0} (a_0 + a_1 z + \cdots + a_n z^n) \\
&= \lim_{z \rightarrow z_0} a_0 + \lim_{z \rightarrow z_0} a_1 z + \cdots + \lim_{z \rightarrow z_0} a_n z^n, \text{ by Theorem 2.3.2 (1)} \\
&= \lim_{z \rightarrow z_0} a_0 + \lim_{z \rightarrow z_0} a_1 \cdot \lim_{z \rightarrow z_0} z + \cdots + \lim_{z \rightarrow z_0} a_n \cdot \lim_{z \rightarrow z_0} z^n, \text{ by Theorem 2.3.2 (2)} \\
&= a_0 + a_1 z_0 + \cdots + a_n z_0^n, \text{ by Theorem 2.3.2 (2) and } \lim_{z \rightarrow z_0} z = z_0 \\
&= p(z_0)
\end{aligned}$$

□

Definition 2.3.4 (Extended Complex Plane or the Riemann Sphere). The **Extended Complex Plane** is the set \mathbf{C} together with a symbol ∞ called the *point at infinity*, denoted $\hat{\mathbf{C}}$ or \mathbf{C}_∞ .

There is a bijection between the extended complex plane and the unit sphere given by the *stereographic projection*, and therefore the extended complex plane is also called the **Riemann Sphere**.



The point N (the north pole) corresponds to ∞ , and any point p on the sphere corresponds uniquely to a point $z \in \mathbf{C}$ which is the unique point of intersection of the complex plane with the line passing through N and p .

Definition 2.3.5 (Neighbourhood of Infinity). Let $\varepsilon > 0$, the set

$$\left\{ z \in \mathbf{C} : |z| > \frac{1}{\varepsilon} \right\}$$

is called a *neighbourhood of ∞* . Geometrically, a neighbourhood at infinity is the exterior of a circle centered at the origin, which corresponds to a neighbourhood of N on the unit sphere.

Discussion 2.3.6. We can now easily give meaning to limits

$$\lim_{z \rightarrow z_0} f(z) = w_0$$

where z_0 and w_0 are allowed to be ∞ . We replace the appropriate neighbourhood in Definition 2.2.1 with neighbourhoods of ∞ .

Theorem 2.3.7 (Limits involving Infinity).

- (1) $\lim_{z \rightarrow z_0} \frac{1}{f(z)} = 0$ if and only if $\lim_{z \rightarrow z_0} f(z) = \infty$.
- (2) $\lim_{z \rightarrow \infty} f(z) = \lim_{z \rightarrow 0} f\left(\frac{1}{z}\right)$, provided the limit exist.

Combining (1) and (2), we get

$$\lim_{z \rightarrow 0} \frac{1}{f\left(\frac{1}{z}\right)} = 0 \quad \text{if and only if} \quad \lim_{z \rightarrow \infty} f(z) = \infty.$$

Bottom line, we can simplify limits involving ∞ to limits involving 0.

Proof. The proofs are based on the simple observation that

$$\frac{1}{a} < b \quad \text{if and only if} \quad \frac{1}{b} < a$$

for non-zero real numbers a and b .

(1) Now $\lim_{z \rightarrow z_0} \frac{1}{f(z)} = 0$ if and only if for every $\varepsilon > 0$ there exists $\delta > 0$ such that

$$\text{if } 0 < |z - z_0| < \delta, \quad \text{then} \quad \frac{1}{|f(z)|} = \left| \frac{1}{f(z)} - 0 \right| < \varepsilon$$

if and only if for every $\varepsilon > 0$ there exists $\delta > 0$ such that

$$\text{if } 0 < |z - z_0| < \delta, \quad \text{then} \quad |f(z)| > \frac{1}{\varepsilon}$$

if and only if $\lim_{z \rightarrow z_0} f(z) = \infty$.

(2) $\lim_{z \rightarrow \infty} f(z) = \alpha$ if and only if for every $\varepsilon > 0$ there exists $\delta > 0$ such that

$$\text{if } |z| > \frac{1}{\delta}, \quad \text{then} \quad |f(z) - \alpha| < \varepsilon$$

if and only if for every $\varepsilon > 0$ there exists $\delta > 0$ such that

$$\text{if } 0 < \left| \frac{1}{z} \right| < \delta, \quad \text{then} \quad |f(z) - \alpha| < \varepsilon$$

if and only if, by replacing z with $1/z$, $\lim_{z \rightarrow 0} f\left(\frac{1}{z}\right) = \alpha$.

□

Example 2.3.8. We want to show $\lim_{z \rightarrow \infty} \frac{2z^4 + 1}{z^3 + 1} = \infty$. This is equivalent to showing

$$\lim_{z \rightarrow 0} \frac{1}{f(1/z)} = \lim_{z \rightarrow 0} \frac{(1/z)^3 + 1}{2(1/z)^4 + 1} = 0, \quad \text{for } f(z) = \frac{2z^4 + 1}{z^3 + 1}$$

Note that,

$$\begin{aligned} \lim_{z \rightarrow 0} \frac{(1/z)^3 + 1}{2(1/z)^4 + 1} &= \lim_{z \rightarrow 0} \frac{\frac{1+z^3}{z^3}}{\frac{2+z^4}{z^4}} \\ &= \lim_{z \rightarrow 0} z \cdot \frac{1+z^3}{2+z^4} \\ &= 0 \cdot \frac{1}{2} \\ &= 0 \end{aligned}$$

Therefore $\lim_{z \rightarrow \infty} \frac{2z^4 + 1}{z^3 + 1} = \infty$.

Example 2.3.9 (in-class). Show $\lim_{z \rightarrow \infty} \frac{2 + z^5}{z^2 + 3} = \infty$.

Answer. This is equivalent to showing

$$\lim_{z \rightarrow 0} \frac{1}{f(1/z)} = \lim_{z \rightarrow 0} \frac{(1/z)^2 + 3}{2 + (1/z)^5} = 0, \quad \text{for } f(z) = \frac{2 + z^5}{z^2 + 3}$$

Note that,

$$\begin{aligned} \lim_{z \rightarrow 0} \frac{(1/z)^2 + 3}{2 + (1/z)^5} &= \lim_{z \rightarrow 0} \frac{\frac{1+3z^2}{z^2}}{\frac{2z^5+1}{z^5}} \\ &= \lim_{z \rightarrow 0} z^3 \cdot \frac{1+3z^2}{2z^5+1} \\ &= 0^3 \cdot \frac{1}{1} \\ &= 0 \end{aligned}$$

Therefore $\lim_{z \rightarrow \infty} \frac{2 + z^5}{z^2 + 3} = \infty$. □

2.4. Continuous Functions

Definition 2.4.1 (Continuous Functions). A function $f : G \rightarrow \mathbf{C}$ is *continuous* at $z_0 \in G$ if either z_0 is an isolated point or

$$\lim_{z \rightarrow z_0} f(z) = f(z_0) = f\left(\lim_{z \rightarrow z_0} z\right)$$

That is, for all $\varepsilon > 0$, there exists a $\delta > 0$ such that

$$\text{if } 0 < |z - z_0| < \delta, \quad \text{then } |f(z) - f(z_0)| < \varepsilon.$$

A function is **continuous** if it is continuous at every point in its domain.

By the limit laws (Theorem 2.3.2), sum, product and quotient of continuous functions are continuous (whenever and wherever defined).

Theorem 2.4.2 (Composition of Continuous Functions). Suppose we have two functions $f : G_1 \rightarrow \mathbf{C}$ and $g : G_2 \rightarrow \mathbf{C}$ such that $f(G_1) \subseteq G_2$. If f is continuous at z_0 and g is continuous at $f(z_0)$, then $g \circ f$ is continuous at z_0 . That is,

$$\lim_{z \rightarrow z_0} g(f(z)) = g(f(z_0)) = g\left(\lim_{z \rightarrow z_0} f(z)\right) = g\left(f\left(\lim_{z \rightarrow z_0} z\right)\right)$$

Therefore, if f and g are continuous, so is $g \circ f$.

Proof. By continuity of g at $f(z_0)$, for an arbitrary $\varepsilon > 0$, there exists a $\delta_1 > 0$ such that

$$\text{if } 0 < |w - f(z_0)| < \delta_1, \quad \text{then } |g(w) - g(f(z_0))| < \varepsilon.$$

Now, by continuity of f at z_0 , for $\delta_1 > 0$, there exists a $\delta > 0$ such that

$$\text{if } 0 < |z - z_0| < \delta, \quad \text{then } |f(z) - f(z_0)| < \delta_1.$$

With these two statements, we have that

$$\text{if } 0 < |z - z_0| < \delta, \quad \text{then } |g(f(z)) - g(f(z_0))| < \varepsilon.$$

Therefore $g \circ f$ is continuous at z_0 . □

Lecture 12

Theorem 2.4.3. Suppose $f : G \rightarrow \mathbf{C}$ is continuous at z_0 and $f(z_0) \neq 0$, then there exists a $\delta > 0$ such that $f(z) \neq 0$ for all $z \in D_\delta(z_0)$. That is, $|f(z)| > 0$ for all $z \in D_\delta(z_0)$.

Proof. Since f is continuous and non-zero at z_0 , for $\varepsilon = \frac{|f(z_0)|}{2} > 0$ there exists a $\delta > 0$ such that

$$\text{if } z \in D_\delta(z_0), \quad \text{then } |f(z) - f(z_0)| < \frac{|f(z_0)|}{2}.$$

For such a z , the reverse triangle inequality gives us

$$||f(z)| - |f(z_0)|| \leq |f(z) - f(z_0)| < \frac{|f(z_0)|}{2}; \quad \text{so,} \quad -\frac{|f(z_0)|}{2} < |f(z)| - |f(z_0)| < \frac{|f(z_0)|}{2}$$

since the former is the absolute value of real numbers. Therefore, adding $|f(z_0)|$ to this inequality gives us

$$|f(z)| > \frac{|f(z_0)|}{2} > 0$$

as needed. □

Theorem 2.4.4 (Continuity in terms of Real and Imaginary parts of a Function). Suppose that

$$f(z) = f(x + iy) = u(x, y) + i v(x, y).$$

Then f is continuous at $z_0 = x_0 + iy_0$ if and only if u and v are continuous at (x_0, y_0) .

Proof. This directly follows from Theorem 2.3.1. □

Definition 2.4.5 (Compact Sets). A subset of \mathbf{C} is said to be **compact** if it is closed and bounded.

Definition 2.4.6 (Bounded Functions). A function $f : G \rightarrow \mathbf{C}$ is said to be a **bounded function** if the image $f(G)$ is bounded. Equivalently, if there exists $M > 0$ such that $|f(z)| \leq M$ for every $z \in G$.

Theorem 2.4.7 (Extreme Value Theorem). Suppose $K \subseteq \mathbf{C}$ is compact, and $f : K \rightarrow \mathbf{C}$ is continuous. Then f is bounded, that is there exists an $M > 0$ such that $|f(z)| \leq M$ for all $z \in K$, and there exists a $z_0 \in K$ such that $|f(z_0)| = M$.

Proof. Since $f = u + iv$ is continuous, so are $u, v : \mathbf{R}^2 \rightarrow \mathbf{R}$ by Theorem 2.4.4. Hence, so is

$$|f(z)| = |f(x + iy)| = \sqrt{u(x, y)^2 + v(x, y)^2}$$

as it's obtained as a sum, product and composition of continuous functions. This result then follows from standard Calculus, since $|f|$ is a real-valued function. □

2.5. Complex-Differentiable Functions

Definition 2.5.1 (Derivative). Consider a function $f : G \rightarrow \mathbf{C}$, the **derivative** of f at $z_0 \in G$ is the limit

$$\frac{d}{dz}(f(z_0)) = f'(z_0) := \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}$$

If the limit exists, we say f is *differentiable* at z_0 .

A function is **differentiable** if it is differentiable at every point in its domain.

Letting $h = \Delta_{z_0} z = z - z_0$, the limit can also be written as

$$f'(z_0) = \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h}$$

Example 2.5.2. Consider $f(z) = z^2$, then

$$\begin{aligned} f'(z) &= \lim_{h \rightarrow 0} \frac{f(z+h) - f(z)}{h} = \lim_{h \rightarrow 0} \frac{(z+h)^2 - z^2}{h} \\ &= \lim_{h \rightarrow 0} \frac{2zh + h^2}{h} \\ &= \lim_{h \rightarrow 0} 2z + h \\ &= 2z \end{aligned}$$

Lecture 13

Example 2.5.3. Where is $f(z) = |z|^2$ differentiable?

Consider $z \in \mathbf{C}$ and an arbitrary $h \in \mathbf{C}$, then we compute

$$\begin{aligned} f(z+h) - f(z) &= |z+h|^2 - |z|^2 \\ &= (z+h)\overline{(z+h)} - z\bar{z} \\ &= z\bar{z} + z\bar{h} + \bar{z}h + h\bar{h} - z\bar{z} \\ &= z\bar{h} + \bar{z}h + h\bar{h} \end{aligned}$$

Then

$$\frac{f(z+h) - f(z)}{h} = \frac{z\bar{h} + \bar{z}h + h\bar{h}}{h} = z\frac{\bar{h}}{h} + \bar{z} + \bar{h}$$

Along the real axis, $h = \bar{h}$, we have

$$\frac{f(z+h) - f(z)}{h} = z + \bar{z} + h;$$

therefore, as $h \rightarrow 0$, the limit is $z + \bar{z}$. Along the imaginary axis, $h = -\bar{h}$, we have

$$\frac{f(z+h) - f(z)}{h} = -z + \bar{z} - h;$$

therefore, as $h \rightarrow 0$, the limit is $-z + \bar{z}$.

Since limits are unique, if $f'(z)$ exists, then $z + \bar{z} = -z + \bar{z}$, which gives us $z = 0$. That is, if $f'(z)$ exists, it only exists for $z = 0$. So, does $f'(0)$ exist?

$$f'(0) = \lim_{h \rightarrow 0} \frac{f(h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{h\bar{h}}{h} = \lim_{h \rightarrow 0} \bar{h} = 0$$

Proposition 2.5.4 (Differentiable Functions are Continuous). *If f is differentiable at z_0 , then f is continuous at z_0 .*

Proof. Suppose f is differentiable at z_0 , then

$$\lim_{z \rightarrow z_0} f(z) - f(z_0) = \left(\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} \right) \left(\lim_{z \rightarrow z_0} z - z_0 \right) = f'(z_0) \cdot 0 = 0$$

Therefore $\lim_{z \rightarrow z_0} f(z) = f(z_0)$, and hence f is continuous at z_0 . \square

Theorem 2.5.5 (Differentiation Laws). *Suppose f and g are differentiable at z . Then,*

$$(1) (c)' = 0, \text{ for every } c \in \mathbb{C}.$$

$$(2) (c \cdot f)'(z) = c \cdot f'(z), \text{ for every } c \in \mathbb{C}. \quad (\text{Constant Rule})$$

$$(3) (z^n)' = nz^{n-1}, \text{ for every } n \in \mathbb{Z} \text{ (assume } z \neq 0 \text{ for } n < 0). \quad (\text{Power Rule})$$

$$(4) (f + g)'(z) = f'(z) + g'(z). \quad (\text{Sum Rule})$$

$$(5) (fg)'(z) = f'(z)g(z) + f(z)g'(z). \quad (\text{Product Rule})$$

$$(6) \left(\frac{f}{g} \right)'(z) = \frac{f'(z)g(z) - f(z)g'(z)}{g(z)^2}, \text{ provided } g(z) \neq 0 \quad (\text{Quotient Rule})$$

Proof. (1) and (4) are proved directly using the limit definition, (2) can be proved directly or using (1) and (5), while (3) can be proven inductively using (5) for positive n and (6) for negative n .

(5) We first compute

$$\begin{aligned} f(z+h)g(z+h) - f(z)g(z) &= f(z+h)g(z+h) - f(z)g(z) + f(z+h)g(z) - f(z+h)g(z) \\ &= f(z+h)(g(z+h) - g(z)) + g(z)(f(z+h) - f(z)) \end{aligned}$$

So,

$$\begin{aligned} (fg)'(z) &= \lim_{h \rightarrow 0} \frac{f(z+h)g(z+h) - f(z)g(z)}{h} \\ &= \lim_{h \rightarrow 0} \frac{f(z+h)(g(z+h) - g(z))}{h} + \lim_{h \rightarrow 0} \frac{g(z)(f(z+h) - f(z))}{h} \\ &= \lim_{h \rightarrow 0} f(z+h) \cdot \lim_{h \rightarrow 0} \frac{g(z+h) - g(z)}{h} + g(z) \lim_{h \rightarrow 0} \frac{f(z+h) - f(z)}{h} \\ &= f(z)g'(z) + g(z)f'(z) \end{aligned}$$

(6) We first compute

$$\begin{aligned}\frac{1}{g(z+h)} - \frac{1}{g(z)} &= \frac{g(z) - g(z+h)}{g(z)g(z+h)} \\ &= -\frac{g(z+h) - g(z)}{g(z)g(z+h)}\end{aligned}$$

So,

$$\begin{aligned}\left(\frac{1}{g}\right)'(z) &= \lim_{h \rightarrow 0} \frac{\frac{1}{g(z+h)} - \frac{1}{g(z)}}{h} \\ &= \lim_{h \rightarrow 0} -\frac{g(z+h) - g(z)}{g(z)g(z+h)} \cdot \frac{1}{h} \\ &= -\lim_{h \rightarrow 0} \frac{g(z+h) - g(z)}{h} \cdot \lim_{h \rightarrow 0} \frac{1}{g(z)g(z+h)} \\ &= -\frac{g'(z)}{g(z)^2}\end{aligned}$$

(6) then follows from the computation above and using (5) on $\frac{f(z)}{g(z)} = f(z) \cdot \frac{1}{g(z)}$. □

Proposition 2.5.6 (Chain Rule). Suppose we have two functions $f : G_1 \rightarrow \mathbf{C}$ and $g : G_2 \rightarrow \mathbf{C}$ such that $f(G_1) \subseteq G_2$. If f is differentiable at z_0 and g is differentiable at $f(z_0)$, then $g \circ f$ is differentiable at z_0 and

$$(g \circ f)'(z_0) = g'(f(z_0)) \cdot f'(z_0)$$

Proof. Let's start by defining an auxiliary function on G_2

$$\phi(w) = \begin{cases} \frac{g(w) - g(f(z_0))}{w - f(z_0)} - g'(f(z_0)) & w \neq f(z_0) \\ 0 & w = f(z_0) \end{cases}$$

Since g is differentiable at $f(z_0)$, then $\lim_{w \rightarrow f(z_0)} \phi(w) = 0 = \phi(f(z_0))$ and therefore ϕ is continuous at $f(z_0)$. Furthermore, since f is differentiable at z_0 , it is continuous at z_0 . So $\lim_{z \rightarrow z_0} \phi(f(z)) = \phi(f(z_0)) = 0$ by Theorem 2.4.2.

Rewriting the above expression, we get the following expression which is valid on all of G_2 .

$$g(w) - g(f(z_0)) = (w - f(z_0))(\phi(w) + g'(f(z_0)))$$

Now, for $w = f(z) \in f(G_1)$, we have

$$\begin{aligned}\frac{g(f(z)) - g(f(z_0))}{z - z_0} &= \frac{(f(z) - f(z_0))(\phi(f(z)) + g'(f(z_0)))}{z - z_0} \\ &= (\phi(f(z)) + g'(f(z_0))) \cdot \frac{f(z) - f(z_0)}{z - z_0}\end{aligned}$$

Therefore,

$$\begin{aligned}
 (g \circ f)'(z_0) &= \lim_{z \rightarrow z_0} \frac{g(f(z)) - g(f(z_0))}{z - z_0} \\
 &= \lim_{z \rightarrow z_0} (\phi(f(z)) + g'(f(z_0))) \cdot \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0} \\
 &= g'(f(z_0)) \cdot f'(z_0), \text{ since } \lim_{z \rightarrow z_0} \phi(f(z)) = 0
 \end{aligned}$$

□

2.6. Cauchy-Riemann Equations

Lecture 14

Theorem 2.6.1 (Cauchy-Riemann Equations). *Suppose that*

$$f(z) = f(x + iy) = u(x, y) + i v(x, y)$$

is differentiable at $z_0 = x_0 + iy_0$. Then

(a) *the first order partial derivatives of u and v exist at (x_0, y_0) and satisfy the **Cauchy-Riemann Equations***

$$\begin{aligned}
 u_x(x_0, y_0) &= v_y(x_0, y_0) \\
 u_y(x_0, y_0) &= -v_x(x_0, y_0)
 \end{aligned} \tag{CR}$$

(b) $f'(z_0) = u_x(x_0, y_0) + i v_x(x_0, y_0) = v_y(x_0, y_0) - i u_y(x_0, y_0)$.

Proof. Since f is differentiable at z_0 , we have, where we let $h = s + it$

$$\begin{aligned}
 f'(z_0) &= \lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{f((x_0 + s) + i(y_0 + t)) - f(x_0 + iy_0)}{h} \\
 &= \lim_{h \rightarrow 0} \frac{u(x_0 + s, y_0 + t) - u(x_0, y_0)}{h} + i \cdot \lim_{h \rightarrow 0} \frac{v(x_0 + s, y_0 + t) - v(x_0, y_0)}{h}
 \end{aligned}$$

As we know by now, we must get the same result if we restrict h to be on the real axis and if we restrict it to be on the imaginary axis. In the former case, $t = 0$, giving us

$$\begin{aligned}
 f'(z_0) &= \lim_{s \rightarrow 0} \frac{u(x_0 + s, y_0) - u(x_0, y_0)}{s} + i \cdot \lim_{s \rightarrow 0} \frac{v(x_0 + s, y_0) - v(x_0, y_0)}{s} \\
 &= u_x(x_0, y_0) + i v_x(x_0, y_0)
 \end{aligned}$$

In the latter case, $s = 0$, giving us

$$\begin{aligned}
 f'(z_0) &= \lim_{t \rightarrow 0} \frac{u(x_0, y_0 + t) - u(x_0, y_0)}{it} + i \cdot \lim_{t \rightarrow 0} \frac{v(x_0, y_0 + t) - v(x_0, y_0)}{it} \\
 &= \frac{1}{i} \cdot \lim_{t \rightarrow 0} \frac{u(x_0, y_0 + t) - u(x_0, y_0)}{t} + \lim_{t \rightarrow 0} \frac{v(x_0, y_0 + t) - v(x_0, y_0)}{t} \\
 &= -i u_y(x_0, y_0) + v_y(x_0, y_0)
 \end{aligned}$$

Therefore

$$u_x(x_0, y_0) + i v_x(x_0, y_0) = f'(z_0) = v_y(x_0, y_0) - i u_y(x_0, y_0),$$

and hence $u_x(x_0, y_0) = v_y(x_0, y_0)$ and $u_y(x_0, y_0) = -v_x(x_0, y_0)$. \square

The Cauchy-Riemann equations (CR) are a *necessary* condition for f' to exist. We can use them to locate possible points where the derivative does not exist but not necessarily conclude where and if the derivative exists.

Example 2.6.2.

- (1) Consider $f(z) = |z|^2 = x^2 + y^2$, so $u(x, y) = x^2 + y^2$ and $v(x, y) = 0$. The partial derivatives at (x, y) are

$$u_x = 2x \qquad v_x = 0$$

$$u_y = 2y \qquad v_y = 0$$

Therefore, the Cauchy-Riemann equations (CR) are only satisfied at $(x, y) = (0, 0)$. Hence f is not differentiable at any $z \neq 0$. Again, note that this does not say anything about the existence of $f'(0)$.

- (2) Consider $f(z) = \bar{z} = x - iy$, so $u(x, y) = x$ and $v(x, y) = -y$. The partial derivatives at (x, y) are

$$u_x = 1 \qquad v_x = 0$$

$$u_y = 0 \qquad v_y = -1$$

Note that $u_x \neq v_y$ for all (x, y) and therefore the Cauchy-Riemann equations (CR) are satisfied for no (x, y) . Hence f is nowhere complex-differentiable.

- (3) (in-class) Consider $f(z) = (z + i\bar{z})^2$, let's simplify f to identify its real and imaginary parts $u(x, y)$ and $v(x, y)$.

$$\begin{aligned} f(z) &= (z + i\bar{z})^2 \\ &= z^2 - \bar{z}^2 + 2iz\bar{z} \\ &= (z + \bar{z})(z - \bar{z}) + 2i|z|^2 \\ &= (2\operatorname{Re} z)(2i\operatorname{Im} z) + 2i|z|^2 = 2i(|z|^2 + \operatorname{Re} z \cdot \operatorname{Im} z) \\ f(x + iy) &= 2i(x^2 + y^2 + 2xy) \\ &= 2i(x + y)^2 \end{aligned}$$

Therefore $u(x, y) = 0$ and $v(x, y) = 2(x + y)^2$. The partial derivatives at (x, y) are

$$u_x = 0 \qquad v_x = 4(x + y)$$

$$u_y = 0 \qquad v_y = 4(x + y)$$

Therefore, the Cauchy-Riemann equations (CR) are satisfied if and only if $4(x + y) = 0$, if and only if $y = -x$. Hence f is not differentiable any $z \in \mathbb{C}$ such that $\operatorname{Im} z \neq -\operatorname{Re} z$.

As commented, the Cauchy-Riemann equations (CR) are not a *sufficient* condition for the existence of the derivative as the example below shows. Problem 49 gives another example.

Example 2.6.3. Consider

$$f(z) = \begin{cases} \frac{\bar{z}^2}{z} = \frac{\bar{z}^3}{|z|^2} & z \neq 0 \\ 0 & z = 0 \end{cases}$$

Then,

$$u(x, y) = \begin{cases} \frac{x^3 - 3xy^2}{x^2 + y^2} & (x, y) \neq (0, 0) \\ 0 & (x, y) = (0, 0) \end{cases} \quad \text{and} \quad v(x, y) = \begin{cases} \frac{y^3 - 3x^2y}{x^2 + y^2} & (x, y) \neq (0, 0) \\ 0 & (x, y) = (0, 0) \end{cases}$$

We show that u and v satisfy the Cauchy-Riemann equations (CR) at $(0, 0)$.

$$\begin{aligned} u_x(0, 0) &= \lim_{s \rightarrow 0} \frac{u(s, 0) - u(0, 0)}{s} = \lim_{s \rightarrow 0} \frac{\frac{s^3}{s^2} - 0}{s} = 1 \\ u_y(0, 0) &= \lim_{t \rightarrow 0} \frac{u(0, t) - u(0, 0)}{t} = \lim_{t \rightarrow 0} \frac{0 - 0}{t} = 0 \\ v_x(0, 0) &= \lim_{s \rightarrow 0} \frac{v(s, 0) - v(0, 0)}{s} = \lim_{s \rightarrow 0} \frac{0 - 0}{s} = 0 \\ v_y(0, 0) &= \lim_{t \rightarrow 0} \frac{v(0, t) - v(0, 0)}{t} = \lim_{t \rightarrow 0} \frac{\frac{t^3}{t^2} - 0}{t} = 1 \end{aligned}$$

Therefore $u_x(0, 0) = 1 = v_y(0, 0)$ and $u_y(0, 0) = 0 = -v_x(0, 0)$, and hence the Cauchy-Riemann equations (CR) are satisfied. But $f'(0)$ does not exist, as seen in Problem 44.

Imposing certain existence and continuity conditions on the first order partial derivatives of u and v , the Cauchy-Riemann equations (CR) can be upgraded to a sufficient condition for differentiability.

Theorem 2.6.4 (Sufficient Conditions for Differentiability). *Consider a function*

$$f(z) = f(x + iy) = u(x, y) + i v(x, y)$$

and a z_0 in the domain of f , such that

- (a) the first order partial derivatives of u and v exist and are continuous in an open disk centered at z_0 ;*
- and*
- (a) the Cauchy-Riemann equations (CR) are satisfied at (x_0, y_0) .*

Then $f'(z_0)$ exists and is given by $u_x(x_0, y_0) + i v_x(x_0, y_0) = v_y(x_0, y_0) - i u_y(x_0, y_0)$.

Proof. We skip the proof. You can find a proof in [BC09, Section 22, Page 66]. □

Example 2.6.5. Let's revisit examples from Example 2.6.2 and 2.6.3.

- (1) Consider $f(z) = |z|^2 = x^2 + y^2$, we noted that $u(x, y) = x^2 + y^2$ and $v(x, y) = 0$. We have seen that the only point where $f(z)$ can be differentiable is $z = 0$. The partial derivatives in a neighbourhood of $(0, 0)$ are

$$\begin{aligned} u_x &= 2x & v_x &= 0 \\ u_y &= 2y & v_y &= 0 \end{aligned}$$

which clearly exist and are continuous. We have also seen that the Cauchy-Riemann equations (CR) are satisfied at $(0, 0)$, trivially. Therefore $f'(0)$ exists and

$$f'(0) = u_x(0, 0) + i v_x(0, 0) = 0.$$

- (2) Consider $f(z) = (z + i\bar{z})^2$, we noted that $u(x, y) = 0$ and $v(x, y) = 2(x + y)^2$. We have seen that the only point where $f(z)$ can be differentiable are $z = x + iy \in \mathbf{C}$ such that $y = \text{Im } z = -\text{Re } z = -x$. That is, at points of the form $(x, -x)$. The partial derivatives in a neighbourhood of $(x, -x)$ are

$$\begin{aligned} u_x &= 0 & v_x &= 4(x + y) \\ u_y &= 0 & v_y &= 4(x + y) \end{aligned}$$

which clearly exist and are continuous. Note the Cauchy-Riemann equations (CR) are satisfied at $(x, -x)$ trivially, since

$$u_x(x, -x) = u_y(x, -x) = v_x(x, -x) = v_y(x, -x) = 0.$$

Therefore $f'(z)$ exists, for $z = x - ix$, and

$$f'(z) = u_x(x, -x) + i v_x(x, -x) = 0.$$

- (3) The reason Example 2.6.3 doesn't contradict Theorem 2.6.4 is because, u_x , in particular, is not continuous at $(0, 0)$. Note that we have

$$u(x, y) = \begin{cases} \frac{x^3 - 3xy^2}{x^2 + y^2} & (x, y) \neq (0, 0) \\ 0 & (x, y) = (0, 0) \end{cases}$$

For $(x, y) \neq (0, 0)$, we compute $u_x(x, y)$ using the quotient rule, while we have already computed $u_x(0, 0) = 1$ in Example 2.6.3, giving us

$$u_x(x, y) = \begin{cases} \frac{x^4 + 6x^2y^2 - 3y^4}{(x^2 + y^2)^2} & (x, y) \neq (0, 0) \\ 1 & (x, y) = (0, 0) \end{cases}$$

Suppose $u_x(x, y)$ is continuous at $(0, 0)$, then we have

$$\lim_{(x, y) \rightarrow (0, 0)} u_x(x, y) = \lim_{(x, y) \rightarrow (0, 0)} \frac{x^4 + 6x^2y^2 - 3y^4}{(x^2 + y^2)^2} = u_x(0, 0) = 1$$

Restricting the limit along the y -axis, where $x = 0$, we get

$$1 = \lim_{(0, y) \rightarrow (0, 0)} \frac{-3y^4}{(y^2)^2} = \lim_{y \rightarrow 0} \frac{-3y^4}{y^4} = -3,$$

a contradiction. Hence, $u_x(x, y)$ is not continuous at $(0, 0)$.

Lecture 15 **Example 2.6.6** (Complex Exponential). Define, for any $z = x + iy \in \mathbb{C}$

$$\exp(z) = e^z := e^x e^{iy} = e^x (\cos y + i \sin y)$$

the *complex exponential function*. Note that e^x is the usual real exponential and e^{iy} is given by Euler's formula (Definition 1.2.11). Here,

$$u(x, y) = e^x \cos y \quad \text{and} \quad v(x, y) = e^x \sin y$$

We then see that

$$u_x = e^x \cos y = v_y,$$

$$v_x = -e^x \sin y = -u_y;$$

so \exp satisfies the Cauchy-Riemann equations (CR) everywhere. Furthermore, u_x, u_y, v_x and v_y are everywhere defined and continuous. Hence \exp is everywhere complex-differentiable, an *entire* function. Furthermore $\exp(z)' = u_x + iv_x = e^x \cos y + ie^x \sin y = \exp(z)$.

Discussion 2.6.7 (Polar Cauchy-Riemann Equations). Recall that if the domain of a function f is contained in \mathbb{C}^* or restricted to within \mathbb{C}^* , one can express in polar coordinates at $z = re^{i\theta}$ as

$$f(z) = f(re^{i\theta}) = u(r, \theta) + i v(r, \theta)$$

Then, the Cauchy-Riemann equations (CR) at a point (r_0, θ_0) can be expressed in polar coordinates, **Polar Cauchy-Riemann Equations** (see Problem 51)

$$\begin{aligned} ru_r &= v_\theta \\ u_\theta &= -rv_r \end{aligned} \tag{Polar CR}$$

and a differentiable function at $z_0 = r_0 e^{i\theta_0}$ is then expressed as

$$f'(z_0) = f'(r_0 e^{i\theta_0}) = e^{-i\theta_0} (u_r(r_0, \theta_0) + i v_r(r_0, \theta_0)).$$

Example 2.6.8. Consider the function

$$f(z) = f(re^{i\theta}) = \sqrt{r} e^{i\frac{\theta}{2}},$$

where $r > 0$ and $-\pi < \theta < \pi$. This is the function that outputs the principal square root of z . We compute $f'(z)$ at $z = re^{i\theta}$ using the polar form of Theorem 2.6.4. We first note that

$$f(z) = \underbrace{\sqrt{r} \cos\left(\frac{\theta}{2}\right)}_{u(r, \theta)} + i \underbrace{\sqrt{r} \sin\left(\frac{\theta}{2}\right)}_{v(r, \theta)}$$

Now, we compute

$$ru_r = r \frac{1}{2\sqrt{r}} \cos\left(\frac{\theta}{2}\right) = \frac{\sqrt{r}}{2} \cos\left(\frac{\theta}{2}\right) = v_\theta$$

$$u_\theta = -\frac{\sqrt{r}}{2} \sin\left(\frac{\theta}{2}\right) = -r \frac{1}{2\sqrt{r}} \sin\left(\frac{\theta}{2}\right) = -rv_r$$

Clearly the first order partial derivatives exist everywhere and the Polar Cauchy-Riemann equations (Polar CR) are also satisfied everywhere. Hence $f'(z)$ exists and

$$\begin{aligned}
 f'(z) &= e^{-i\theta}(u_r(r, \theta) + i v_r(r, \theta)) \\
 &= e^{-i\theta} \left(\frac{1}{2\sqrt{r}} \cos\left(\frac{\theta}{2}\right) + i \frac{1}{2\sqrt{r}} \sin\left(\frac{\theta}{2}\right) \right) \\
 &= \frac{e^{-i\theta}}{2\sqrt{r}} \left(\cos\left(\frac{\theta}{2}\right) + i \sin\left(\frac{\theta}{2}\right) \right) \\
 &= \frac{1}{2\sqrt{r}} \cdot e^{-i\theta} \cdot e^{i\frac{\theta}{2}} \\
 &= \frac{1}{2\sqrt{r}e^{i\frac{\theta}{2}}} \\
 &= \frac{1}{2f(z)}
 \end{aligned}$$

2.7. Holomorphic Functions

Definition 2.7.1 (Holomorphic Functions). A function f is *holomorphic on an open set* U if $f'(z)$ exists for every $z \in U$.

We say f is *holomorphic at a point* z_0 if it is holomorphic on some open disk $D_\varepsilon(z_0)$ for an $\varepsilon > 0$. We say f is **holomorphic** if it is holomorphic at every point in its domain.

A function that is holomorphic on all of \mathbf{C} is said to be **entire**.

Example 2.7.2.

- (1) $f(z) = \frac{1}{z}$ is holomorphic on any open set not containing 0, in particular on \mathbf{C}^* .
- (2) $f(z) = |z|^2$ is nowhere holomorphic since we have already seen that f is only complex-differentiable at $z = 0$ and at no other point.
- (3) Polynomials are entire.
- (4) $f(z) = \bar{z}$ is nowhere holomorphic, since it's nowhere differentiable.

Discussion 2.7.3. Let G be a domain (open and connected subset of \mathbf{C}). We know several necessary and sufficient conditions for $f = u + iv$ to be holomorphic on G .

(Necessary) (1) f is continuous on G .

(2) Cauchy-Riemann equations (CR) are satisfied on G .

(Sufficient) (1) First order partial derivatives of u and v exist and continuous on G , and the Cauchy-Riemann equations (CR) are satisfied on G .

- (2) Differentiation Laws. If f and g are holomorphic on G , then so are $f + g$, fg and f/g (if $g \neq 0$ on G).
- (3) Composition of holomorphic functions is holomorphic.

Theorem 2.7.4 (Sufficient Condition for Constantness). Suppose G is a domain and $f'(z) = 0$ for all $z \in G$. Then $f(z)$ is constant on G .

Proof. Write $f(z) = f(x + iy) = u(x, y) + i v(x, y)$, so we have

$$0 = f'(z) = u_x + i v_x = v_y - i u_y$$

Therefore $u_x = u_y = 0$ and $v_x = v_y = 0$. We consider points $p, q \in G$ such that there's a line segment L in G connecting them. Let $\vec{w} = (a, b)$ be a unit vector parallel to L , then the directional derivative of u along L is

$$(\text{grad } u) \cdot \vec{w} = au_x + bu_y = 0.$$

So, u is constant along L . Since G is a domain, any two points can be connected by a polygon line. Applying the above argument along constituent line segments, we see that u has the same value along the endpoints of any polygon line. This shows that u is constant on G , say $u(x, y) = c$. A similar argument works for v , giving us $v(x, y) = d$. Hence

$$f(z) = c + id,$$

that is, f is constant. □

Theorem 2.7.4 has many interesting consequences.

Proposition 2.7.5. Suppose f and \bar{f} are holomorphic on a domain G . Then f is constant on G .

Proof. We write

$$f(z) = f(x + iy) = u(x, y) + i v(x, y)$$

$$\bar{f}(z) = \overline{f(x + iy)} = u(x, y) - i v(x, y)$$

Since f and \bar{f} are holomorphic, they satisfy the Cauchy-Riemann equations (CR)

$$\text{for } f: \begin{cases} u_x = v_y \\ u_y = -v_x \end{cases}$$

$$\text{for } \bar{f}: \begin{cases} u_x = (-v)_y = -v_y \\ u_y = -(-v)_x = v_x \end{cases}$$

This gives us $v_y = -v_y$ and $v_x = -v_x$, and therefore $u_x = v_x = 0$. Hence $f'(z) = u_x + i v_x = 0$, giving us that f is constant by Theorem 2.7.4. □

Corollary 2.7.6. Suppose f is holomorphic on a domain G and always real-valued. Then f is constant on G .

Proof. Since f is always real-valued, we have $f = \bar{f}$. Therefore \bar{f} is holomorphic on G as well, and hence f is constant by Proposition 2.7.5. □

Corollary 2.7.7. Suppose f is holomorphic on a domain G and $|f|$ is constant on it. Then f is also constant on G .

Proof. By assumption $|f(z)| = c$, for all $z \in G$, for some $c \in \mathbf{C}$. This gives us

$$f(z)\overline{f(z)} = |f(z)|^2 = c^2 \quad (*)$$

Suppose $c = 0$, then $|f(z)| = 0$ and therefore $f(z) = 0$. Suppose $c \neq 0$, then necessarily $f(z) \neq 0$ for every $z \in G$ by (*). Hence

$$\overline{f(z)} = \frac{c^2}{f(z)},$$

and thus \bar{f} is holomorphic. Therefore both f and \bar{f} are holomorphic and hence f is constant by Proposition 2.7.5. \square

Example 2.7.8. We apply Corollary 2.7.7 to $f(z) = \frac{\bar{z}}{z}$ to conclude that it's not holomorphic.

We first note that, for any $z \in \mathbf{C}$,

$$|f(z)| = \left| \frac{\bar{z}}{z} \right| = \frac{|\bar{z}|}{|z|} = 1;$$

that is, $|f|$ is constant. Suppose f was holomorphic on \mathbf{C} (this argument can be specialised to any domain G), then f would be a holomorphic function such that $|f|$ is constant. Therefore, by Corollary 2.7.7, f is constant on \mathbf{C} . That's a contradiction, since f is non-constant, as $f(1) = 1$ and $f(i) = -1$.

Example 2.7.9 (in-class). Is the function $f(z) = \operatorname{Re} z$ holomorphic?

Answer. Note that $f(z) = \operatorname{Re} z$ is a real-valued function, for any $z \in \mathbf{C}$. Suppose f was holomorphic on \mathbf{C} (this argument can be specialised to any domain G), then f would be a holomorphic function such that f is always real-valued. Therefore, by Corollary 2.7.6, f is constant on \mathbf{C} . That's a contradiction, since f is non-constant, as $f(1) = 1$ and $f(i) = 0$. \square

We now discuss a large class of holomorphic functions, which are complex versions of functions you may have seen in your Calculus classes

2.8. The Exponential Function

Definition 2.8.1 (The Exponential Function). The (complex) exponential function e^z (or $\exp(z)$) is defined on all of \mathbf{C} as follows

$$e^z := e^{\operatorname{Re} z} e^{i \operatorname{Im} z} = e^{\operatorname{Re} z} (\cos(\operatorname{Im} z) + i \sin(\operatorname{Im} z)).$$

That is, writing $z = x + iy$, we have

$$e^z = e^x e^{iy} = e^x (\cos y + i \sin y).$$

Since $x \in \mathbf{R}$, e^x is the usual real exponential function, while e^{iy} is given by Euler's formula.

Furthermore, the definitions give us $\overline{e^z} = e^{\bar{z}}$.

Note that when $z = x \in \mathbf{R}$, we have $e^z = e^x$, since then $\operatorname{Im} z = 0$.

Proposition 2.8.2 (Properties of the Exponential). Consider $z, w \in \mathbf{C}$.

$$(1) |e^z| = e^{\operatorname{Re} z} \text{ and } \arg e^z = \{\operatorname{Im} z + 2k\pi : k \in \mathbf{Z}\}.$$

$$(2) e^{z+w} = e^z e^w.$$

$$(3) e^{z-w} = \frac{e^z}{e^w}.$$

$$(4) e^z \text{ is entire, and } (e^z)' = e^z.$$

$$(5) e^z \text{ is periodic: } e^{z+2k\pi i} = e^z \text{ for all } k \in \mathbf{Z}.$$

Proof.

(1) Write $z = x + iy$, then $|e^z| = |e^x| |\cos x + i \sin x| = |e^x|$. Which tells us

$$\arg e^z = \{y + 2k\pi : k \in \mathbf{Z}\}.$$

(2) Write $z = x + iy$ and $w = u + iv$, then

$$\begin{aligned} e^{z+w} &= e^{(x+u)+i(y+v)} \\ &= e^{x+u} e^{i(y+v)} \\ &= e^x e^u e^{iy} e^{iv} \\ &= e^x e^{iy} e^u e^{iv} = e^z e^w \end{aligned}$$

(3) From (2) we get $e^{z-w} e^w = e^z$.

(4) This was seen in Example 2.6.6.

(5) From (2) we have $e^{z+2k\pi i} = e^z e^{2k\pi i} = e^z$.

□

2.9. The Logarithmic Function

Discussion 2.9.1. The complex logarithmic function arises, just the like the usual real logarithmic function, from trying to solve the following equation for w

$$e^w = z \quad (z \neq 0)$$

Write $z = re^{i\theta}$ and $w = u + iv$, then

$$e^u e^{iv} = e^w = z = re^{i\theta}.$$

So, $e^u = r$, giving us $u = \ln r = \ln |z|$, and $v = \theta + 2k\pi$ for some $k \in \mathbf{Z}$, that is the possible values of v are exactly $\arg z = \operatorname{Arg} z + 2k\pi$, $k \in \mathbf{Z}$.

Therefore,

$$\begin{aligned} w &= \ln |z| + i \arg(z) \\ &= \ln |z| + i \operatorname{Arg}(z) + 2k\pi i, \quad k \in \mathbf{Z} \end{aligned}$$

Essentially, w is not unique, as v is not unique. This is to be expected, since e^z is not injective as it is periodic.

Multiple functions satisfy the equation we considered, which we package into a *multi-valued function* using $\arg z$.

Definition 2.9.2 (The Logarithmic Function). We define the **logarithmic function** $\log z$ for any $z \neq 0$, following the discussion above, as

$$\log z := \ln |z| + i \arg(z)$$

Note that $\log z$ is not really a function but a *multi-valued function*, as $\arg z$ is not single-valued.

The **principal logarithm**, denoted $\text{Log } z$, is defined by taking the principal argument of z

$$\text{Log } z := \ln |z| + i \text{Arg } z, \quad -\pi < \text{Arg } z \leq \pi$$

The principal branch of \log is a single-valued function.

Proposition 2.9.3 (Properties of the Logarithm). Consider $z \in \mathbf{C}$.

- (1) $e^{\log z} = z$.
- (2) $\log e^z = z + 2k\pi i$, $k \in \mathbf{Z}$.
- (3) $\log z = \text{Log } z + 2k\pi i$, $k \in \mathbf{Z}$.
- (4) If $z = x \in \mathbf{R}_{>0}$, then $\text{Log } z = \ln x$.

Proof.

- (1) Note that

$$\begin{aligned} e^{\log z} &= e^{\ln |z| + i \arg z} \\ &= e^{\ln |z|} e^{i(\text{Arg } z + 2k\pi)}, \quad k \in \mathbf{Z} \\ &= e^{\ln |z|} e^{i \text{Arg } z} e^{2k\pi i}, \quad k \in \mathbf{Z} \\ &= |z| e^{i \text{Arg } z} \\ &= z \end{aligned}$$

- (2) Note that

$$\begin{aligned} \log e^z &= \ln |e^z| + i \arg(e^z) \\ &= \ln e^{\text{Re } z} + i(\text{Im } z + 2k\pi), \quad k \in \mathbf{Z} \\ &= \text{Re } z + i \text{Im } z + 2k\pi i, \quad k \in \mathbf{Z} \\ &= z + 2k\pi i, \quad k \in \mathbf{Z} \end{aligned}$$

- (3) Note that

$$\begin{aligned} \log z &= \log e^{\text{Log } z}, \text{ by (1)} \\ &= \text{Log } z + 2k\pi i, \quad k \in \mathbf{Z}, \text{ by (2)} \end{aligned}$$

(4) Note that if $z = x \in \mathbf{R}_{>0}$, then $\text{Arg } z = 0$, therefore

$$\text{Log } z = \ln |z| + i \text{Arg } z = \ln x.$$

□

Lecture 17 **Example 2.9.4.**

$$\begin{aligned} (1) \quad \log(1 + i\sqrt{3}) &= \ln |1 + i\sqrt{3}| + i \arg(1 + i\sqrt{3}) \\ &= \ln 2 + i \left(\frac{\pi}{3} + 2k\pi \right), \quad k \in \mathbf{Z} \end{aligned}$$

$$\text{Log}(1 + i\sqrt{3}) = \ln 2 + \frac{\pi i}{3}$$

$$\begin{aligned} (2) \quad \log 1 &= \ln |1| + i \arg 1 \\ &= 0 + i(0 + 2k\pi), \quad k \in \mathbf{Z} \\ &= 2k\pi i, \quad k \in \mathbf{Z} \end{aligned}$$

$$\text{Log } 1 = 0$$

$$\begin{aligned} (3) \quad \log -1 &= \ln |-1| + i \arg -1 \\ &= \ln 1 + i(\pi + 2k\pi), \quad k \in \mathbf{Z} \\ &= (2k + 1)\pi i, \quad k \in \mathbf{Z} \end{aligned}$$

$$\text{Log } -1 = \pi i$$

(4) Familiar properties of logarithms that you know may not hold.

$$(a) \quad \text{Log}(-1 + i)^2 \neq 2 \text{Log}(-1 + i)$$

$$\begin{aligned} \text{Log}(-1 + i)^2 &= \text{Log}(-2i) = \ln |-2i| + i \text{Arg}(-2i) \\ &= \ln 2 + i \left(-\frac{\pi}{2} \right) \\ &= \ln 2 - \frac{\pi i}{2} \end{aligned}$$

$$\begin{aligned} 2 \text{Log}(-1 + i) &= 2 \ln |-1 + i| + 2i \arg(-1 + i) \\ &= 2 \ln \sqrt{2} + 2i \left(\frac{3\pi}{4} \right) \\ &= \ln 2 + \frac{3\pi i}{2} \end{aligned}$$

$$(b) \quad \log i^2 \neq 2 \log i$$

$$\log i^2 = \log -1 = (2k + 1)\pi i, \quad k \in \mathbf{Z}$$

$$\begin{aligned} 2 \log i &= 2 \ln |i| + 2i \arg i \\ &= 0 + 2i \left(\frac{\pi}{2} + 2k\pi \right), \quad k \in \mathbf{Z} \\ &= (4k + 1)\pi i, \quad k \in \mathbf{Z} \end{aligned}$$

Proposition 2.9.5. For all $z, w \in \mathbf{C}^*$

$$(1) \log zw = \log z + \log w$$

$$(2) \log w^{-1} = -\log w$$

One treats this as an equality of sets. (1) and (2) also gives you $\log z/w = \log z - \log w$.

Proof.

(1) We have

$$\begin{aligned} \log z + \log w &= \ln |z| + i \arg z + \ln |w| + i \arg w \\ &= \ln |z| |w| + i(\arg z + \arg w) \\ &= \ln |zw| + i \arg zw, \text{ by Proposition 1.2.18 (1)} \\ &= \log zw \end{aligned}$$

(2) We have

$$\begin{aligned} \log w^{-1} &= \ln |w^{-1}| + i \arg w^{-1} \\ &= \ln |w|^{-1} + i(-\arg w), \text{ by Proposition 1.2.18 (2)} \\ &= -(\ln |w| + i \arg w) \\ &= -\log w \end{aligned}$$

This statement does not hold if we replace $\log z$ with $\text{Log } z$. □

Definition 2.9.6 (Branch of a Multi-Valued Functions). A **branch** of a multi-valued function f is a single-valued function F such that

- F is holomorphic on some domain G ; and
- F assigned to each $z \in G$ precisely one value $F(z)$ of $f(z)$.

A portion of a line or curve in the complex plane is called a **branch cut** for f if a branch f is defined on its complement. A point belonging to *every* branch cut of f is a **branch point**.

Proposition 2.9.7 (Branches of log). *Let $\alpha \in \mathbf{R}$. The function*

$$L_\alpha(z) = L_\alpha(re^{i\theta}) = \ln r + i\theta, \quad \alpha < \theta < \alpha + 2\pi$$

is a branch of $f(z) = \log z$. Note that $\text{Re } L_\alpha = u(r, \theta) = \ln r$ and $\text{Im } L_\alpha = v(r, \theta) = \theta$.

Proof. We first remark that if we were to define L_α also on the ray $\theta = \alpha$, it would not be continuous there. For if z is a point on that ray, as one notes that $\lim_{\theta \rightarrow \alpha^-} \theta = \alpha$ but $\lim_{\theta \rightarrow \alpha^+} \theta \neq \alpha$ as the points close to the ray to the right have arguments near $\alpha + 2\pi$.

It is clear that $L_\alpha(z)$ is single-valued and, for each z , $L_\alpha(z)$ is a value of $\log z$. We need to show L_α is holomorphic. Note that $u(r, \theta) = \ln r$ and $v(r, \theta) = \theta$ have continuous partial derivatives on the domain of definition

$$\begin{aligned} u_r &= \frac{1}{r} & v_r &= 0 \\ u_\theta &= 0 & v_\theta &= 1 \end{aligned}$$

Clearly, the Polar Cauchy Riemann equations ([Polar CR](#)) are satisfied, and therefore L_α is holomorphic. In fact,


$$L'_\alpha(z) = e^{-i\theta}(u_r + iv_r) = e^{-i\theta} \left(\frac{1}{r} \right) = \frac{1}{z}$$

In particular, $\text{Log } z$ for those z such that $-\pi < \text{Arg } z < \pi$ is a branch of $\log z$, called the **principal branch of the logarithm** and

$$(\text{Log } z)' = \frac{1}{z}$$

□

Remark 2.9.8. The branch cut for $\log z$ in Proposition [2.9.7](#) is the ray $r > 0, \theta = \alpha$



The branch cut for $\text{Log } z$ is the ray $r > 0, \theta = \pi$, i.e., the negative real axis. The origin is a branch point of $\log z$.

Example 2.9.9 (Integer Powers and Roots). The logarithmic function can be used to compute integer powers and roots (as previously seen and defined).

$$(1) \quad z^n = e^{n \log z}$$

$$(2) \quad z^{1/n} = e^{\frac{\log z}{n}}$$

Proof. We note that

$$\begin{aligned} e^{n \log z} &= e^{n(\ln|z| + i \arg z)} & e^{\frac{\log z}{n}} &= e^{\frac{1}{n}(\ln|z| + i \arg z)} \\ &= e^{n \ln|z|} \cdot e^{in \arg z} & &= e^{\frac{1}{n} \ln|z|} \cdot e^{i \left(\frac{\arg z}{n} \right)} \\ &= |z|^n \cdot (e^{i \arg z})^n & &= e^{\frac{1}{n} \ln|z|} \cdot e^{i \left(\frac{\text{Arg } z + 2k\pi}{n} \right)} \\ &= (|z| e^{i \arg z})^n & &= \sqrt[n]{|z|} \cdot e^{i \left(\frac{\text{Arg } z + 2k\pi}{n} \right)} \\ &= z^n & &= z^{1/n} \end{aligned}$$

Recall that z^n is single-valued, but $z^{1/n}$ is multi-valued, as complex numbers have n distinct n^{th} roots (Proposition [1.3.3](#)). In fact, using the the principal logarithm, the complex number

$$e^{\frac{\text{Log } z}{n}}$$

gives the principal n^{th} root of z .

□

2.10. Power and Exponential Functions

Lecture 18

Definition 2.10.1 (Power Function). The **power function** z^c for a fixed $c \in \mathbb{C}$ is the *multi-valued* function

$$z^c := e^{c \log z}, \quad z \neq 0$$

Proposition 2.10.2 (Branches of z^c). A branch of z^c is determined by specifying a branch of $\log z$

$$\log z = \ln |z| + i \arg z, \quad z \neq 0, \quad \alpha < \arg z < \alpha + 2\pi$$

Moreover,

$$(z^c)' = cz^{c-1},$$

whenever $z \neq 0$, $\alpha < \arg z < \alpha + 2\pi$.

Proof. We only need to verify that z^c is holomorphic, once a branch of $\log z$ has been specified. Since $z^c = e^{c \log z}$ is a composition of two holomorphic functions, z^c itself is holomorphic. Moreover, by the chain rule

$$\begin{aligned} (z^c)' &= (e^{c \log z})' = e^{c \log z} (c \log z)' \\ &= e^{c \log z} \cdot \frac{c}{z} \\ &= c \cdot \frac{e^{c \log z}}{e^{\log z}} = c \cdot e^{(c-1) \log z} = cz^{c-1} \end{aligned}$$

□

Discussion 2.10.3. The **principal branch** of z^c is defined by specifying the principal branch $\text{Log } z$ of $\log z$. The principal branch of z^c reduces to the usual power function when $z = x \in \mathbb{R}$.

Definition 2.10.4 (Exponential Function with Base c). The **exponential function with base c** , where $c \in \mathbb{C}^*$, is the *multi-valued* function

$$c^z := e^{z \log c}$$

Discussion 2.10.5. Once a branch of $\log z$ has been specified, c^z is an entire function. In that case, using chain rule we have

$$\begin{aligned} (c^z)' &= (e^{z \log c})' = e^{z \log c} (z \log c)' \\ &= e^{z \log c} \cdot \log c \\ &= c^z \log c \end{aligned}$$

What happens if we take $c = e$? Specifying the principal branch $\text{Log } z$ we see

$$e^z = e^{z \text{Log } e} = e^{z(\ln e + i \text{Arg } e)} = e^{z(1+0)} = e^z$$

Example 2.10.6.

(1) We compute

$$\begin{aligned}
 i^i &= e^{i \log i} \\
 &= e^{i(\ln|i| + i \arg i)} \\
 &= e^{i(\ln 1 + i(\frac{\pi}{2} + 2k\pi))}, k \in \mathbf{Z} \\
 &= e^{i^2(\frac{\pi}{2} + 2k\pi)}, k \in \mathbf{Z} \\
 &= e^{-\frac{\pi}{2}} e^{-2k\pi}, k \in \mathbf{Z}
 \end{aligned}$$

(2) We compute

$$\begin{aligned}
 (-1)^{\frac{1}{\pi}} &= e^{\frac{1}{\pi} \log -1} \\
 &= e^{\frac{1}{\pi}(\ln|-1| + i \arg -1)} \\
 &= e^{\frac{1}{\pi}(\ln 1 + i(\pi + 2k\pi))}, k \in \mathbf{Z} \\
 &= e^{\frac{1}{\pi}(\pi i(2k+1))}, k \in \mathbf{Z} \\
 &= e^{i(2k+1)}, k \in \mathbf{Z}
 \end{aligned}$$

2.11. Trigonometric Functions

Discussion 2.11.1. Recall that for any $z \in \mathbf{C}$,

$$\operatorname{Re} z = \frac{z + \bar{z}}{2} \quad \text{and} \quad \operatorname{Im} z = \frac{z - \bar{z}}{2i}$$

Therefore, for $x \in \mathbf{R}$,

$$\begin{aligned}
 \cos x &= \operatorname{Re}(e^{ix}) & \sin x &= \operatorname{Im}(e^{ix}) \\
 &= \frac{e^{ix} + \overline{e^{ix}}}{2} & &= \frac{e^{ix} - \overline{e^{ix}}}{2i} \\
 &= \frac{e^{ix} + e^{-ix}}{2} & &= \frac{e^{ix} - e^{-ix}}{2i}
 \end{aligned}$$

This suggests a way to extend the domain of definition of sine and cosine functions to all of \mathbf{C} .

Definition 2.11.2 (Sine and Cosine). The **(complex) sine** and **cosine functions** are defined as

$$\cos z = \frac{e^{iz} + e^{-iz}}{2} \quad \text{and} \quad \sin z = \frac{e^{iz} - e^{-iz}}{2i}$$

respectively. Moreover, this gives us $e^{iz} = \cos z + i \sin z$. And our calculations above tell us that these functions reduce to the usual sine and cosine for $z = x \in \mathbf{R}$.

Proposition 2.11.3 (Holomorphicity of sin and cos).

- (1) $\sin z$ and $\cos z$ are entire.
- (2) $(\sin z)' = \cos z$ and $(\cos z)' = -\sin z$.

Proof.

- (1) Since $\sin z$ and $\cos z$ are linear combinations of entire functions, they themselves are entire functions.

(2) We note that

$$\begin{aligned}
 (\sin z)' &= \frac{(e^{iz})' - (e^{-iz})'}{2i} & (\cos z)' &= \frac{(e^{iz})' + (e^{-iz})'}{2} \\
 &= \frac{ie^{iz} - (-i)e^{-iz}}{2i} & &= \frac{ie^{iz} - ie^{-iz}}{2} \\
 &= \frac{ie^{iz} + ie^{-iz}}{2i} & &= i \cdot \frac{e^{iz} - e^{-iz}}{2} \\
 &= \frac{e^{iz} + e^{-iz}}{2} & &= -\frac{e^{iz} - e^{-iz}}{2i} \\
 &= \cos z & &= -\sin z
 \end{aligned}$$

□

Discussion 2.11.4 (Trigonometric Identities). Various familiar identities hold, here are a few.

- | | |
|---|--------------------------------|
| (1) $\sin(-z) = -\sin z$ | (5) $\sin(z + 2\pi) = \sin z$ |
| (2) $\cos(-z) = \cos z$ | (6) $\cos(z + 2\pi) = \cos z$ |
| (3) $\sin(z + w) = \sin z \cos w + \cos z \sin w$ | (7) $\sin(\pi/2 - z) = \cos z$ |
| (4) $\cos(z + w) = \cos z \cos w - \sin z \sin w$ | (8) $\sin^2 z + \cos^2 z = 1$ |

To define other trigonometric functions, we need to understand the zeros of $\sin z$ and $\cos z$.

Theorem 2.11.5 (Zeros of Sine and Cosine). *The zeros of $\sin z$ and $\cos z$ are precisely the zeros of sine and cosine functions in a real variable:*

$$\begin{aligned}
 \sin z = 0 & \quad \text{if and only if} \quad z = k\pi, \quad k \in \mathbf{Z} \\
 \cos z = 0 & \quad \text{if and only if} \quad z = k\pi + \frac{\pi}{2}, \quad k \in \mathbf{Z}
 \end{aligned}$$

Proof. We immediately note that

$$\sin z = \sin k\pi = 0 \quad \text{and} \quad \cos z = \cos\left(k\pi + \frac{\pi}{2}\right) = 0$$

since the inputs are real numbers and sine and cosine reduce to the usual real sine and cosine for real inputs.

Conversely, suppose

$$\frac{e^{iz} - e^{-iz}}{2i} = \sin z = 0,$$

this gives us $e^{iz} = e^{-iz}$, and therefore $e^{2iz} = 1$. Applying log gives us

$$2iz + 2m\pi i = 2n\pi i, \quad \text{for } m, n \in \mathbf{Z}$$

by Proposition 2.9.3 (2) and Example 2.9.4 (2). Giving us $z = (n - m)\pi = k\pi$ for any $k \in \mathbf{Z}$.

Suppose $\cos z = 0$. By Discussion 2.11.4 (1) and (7), we have

$$\sin\left(z - \frac{\pi}{2}\right) = -\cos z = 0$$

Hence, $z - \frac{\pi}{2} = k\pi$, $k \in \mathbf{Z}$. □

Definition 2.11.6 (Other Trigonometric Functions). The (complex) **tangent**, **cotangent**, **secant** and **cosecant** functions are defined in terms of sine and cosine.

$$\begin{aligned} \tan z &:= \frac{\sin z}{\cos z}, & z &\neq k\pi + \frac{\pi}{2} & \sec z &:= \frac{1}{\cos z}, & z &\neq k\pi + \frac{\pi}{2} \\ \cot z &:= \frac{\cos z}{\sin z}, & z &\neq k\pi & \csc z &:= \frac{1}{\sin z}, & z &\neq k\pi \end{aligned}$$

These functions are holomorphic in their stated domains of definition since $\sin z$ and $\cos z$ are. They also all reduce to the usual real trigonometric functions when z is real, since $\sin z$ and $\cos z$ do. The derivatives are exactly as expected.

3. Part III. Integration

Lecture 19

We now want to develop a theory of integration of complex-valued functions in a single complex variable. Integrals will be defined over suitable curves (contours) in the complex plane. This theory of integration is a surprisingly powerful tool in the study of holomorphic functions.

Using this theory, we will obtain powerful characterisations of holomorphic functions. Roughly speaking we will prove the following: let G be a domain and $f : G \rightarrow \mathbf{C}$ a function. The following are equivalent.

- (1) f is holomorphic on G .
- (2) For all $n \in \mathbf{Z}_{>0}$, $f^{(n)}$ exists and is holomorphic on G .
- (3) In each *simply connected* subdomain D of G , there exists a holomorphic function $F : D \rightarrow \mathbf{C}$ such that $F' = f|_D$.
- (4) f is continuous on G and

$$\int_C f(z) dz = 0$$

for every *contour* C lying in a *simply connected* subdomain.

- (5) If C is a *simple closed contour* in G and z_0 is interior to C , then

$$f'(z_0) = \frac{1}{2\pi i} \int_C \frac{f(z)}{z - z_0} dz.$$

Additionally, as an application of the theory, we will prove

- *Liouville's theorem*. Every bounded holomorphic function is constant.
- *Fundamental Theorem of Algebra*. Every polynomial of degree $n \geq 1$ has at least one complex root.

3.1. Derivatives of Functions of a Real-variable

To define an integral of a complex-valued functions in a single complex variable, we need to understand how to differentiate a complex-valued function in a single real variable

$$\gamma : [a, b] \rightarrow \mathbf{C},$$

where $[a, b] \subseteq \mathbf{R}$.

Definition 3.1.1. For $\gamma : [a, b] \rightarrow \mathbf{C}$, writing $\gamma(t) = u(t) + i v(t)$, where $u, v : [a, b] \rightarrow \mathbf{R}$, we define the *derivative* of γ to be

$$\gamma'(t) = u'(t) + i v'(t),$$

provided that $u'(t)$ and $v'(t)$ exist. In this case, we say γ is differentiable.

Proposition 3.1.2. Suppose $\gamma_1(t) = u_1(t) + i v_1(t)$ and $\gamma_2(t) = u_2(t) + i v_2(t)$ are differentiable, then

$$(1) (\gamma_1 + \gamma_2)'(t) = \gamma_1'(t) + \gamma_2'(t)$$

$$(2) (\gamma_1 \gamma_2)'(t) = \gamma_1'(t) \gamma_2(t) + \gamma_1(t) \gamma_2'(t)$$

Proof.

$$\begin{aligned} (1) (\gamma_1 + \gamma_2)' &= ((u_1 + u_2) + i(v_1 + v_2))' \\ &= (u_1 + u_2)' + i(v_1 + v_2)' \\ &= (u_1' + u_2') + i(v_1' + v_2') \\ &= (u_1' + i v_1') + (u_2' + i v_2') = \gamma_1' + \gamma_2' \end{aligned}$$

$$\begin{aligned} (2) (\gamma_1 \gamma_2)' &= ((u_1 + i v_1)(u_2 + i v_2))' \\ &= ((u_1 u_2 - v_1 v_2) + i(u_1 v_2 + u_2 v_1))' \\ &= (u_1 u_2 - v_1 v_2)' + i(u_1 v_2 + u_2 v_1)' \\ &= (u_1 u_2)' - (v_1 v_2)' + i(u_1 v_2)' + i(u_2 v_1)' \\ &= (u_1' u_2 + u_1 u_2') - (v_1' v_2 + v_1 v_2') + i(u_1' v_2 + u_1 v_2') + i(u_2' v_1 + u_2 v_1') \\ &= (u_1' u_2 - v_1' v_2) + i(u_1' v_2 + u_2 v_1') + (u_1 u_2' - v_1 v_2') + i(u_1 v_2' + u_2' v_1) \\ &= (u_1' + i v_1')(u_2 + i v_2) + (u_1 + i v_1)(u_2' + i v_2') \\ &= \gamma_1' \gamma_2 + \gamma_1 \gamma_2' \end{aligned}$$

□

Example 3.1.3. We will often encounter the function $\gamma : [a, b] \rightarrow \mathbf{C}$, where

$$\gamma(t) = e^{z_0 t}, \quad z_0 \in \mathbf{C}$$

Let's compute $\gamma'(t)$, for which we first need to express it as $u(t) + i v(t)$. Let $z_0 = x_0 + i y_0$,

$$\begin{aligned} \gamma(t) &= e^{z_0 t} = e^{(x_0 + i y_0)t} \\ &= e^{x_0 t + i y_0 t} \\ &= e^{x_0 t} e^{i y_0 t} = e^{x_0 t} (\cos(y_0 t) + i \sin(y_0 t)) \end{aligned}$$

Therefore, $u(t) = e^{x_0 t} \cos(y_0 t)$ and $v(t) = e^{x_0 t} \sin(y_0 t)$. We note,

$$\begin{aligned} u'(t) &= (e^{x_0 t})'(\cos(y_0 t)) + (e^{x_0 t})(\cos(y_0 t))' & v'(t) &= (e^{x_0 t})'(\sin(y_0 t)) + (e^{x_0 t})(\sin(y_0 t))' \\ &= x_0 e^{x_0 t} \cos(y_0 t) - y_0 e^{x_0 t} \sin(y_0 t) & &= x_0 e^{x_0 t} \sin(y_0 t) + y_0 e^{x_0 t} \cos(y_0 t) \end{aligned}$$

Hence,

$$\begin{aligned} \gamma'(t) &= u'(t) + i v'(t) = x_0 e^{x_0 t} \cos(y_0 t) - y_0 e^{x_0 t} \sin(y_0 t) + i x_0 e^{x_0 t} \sin(y_0 t) + i y_0 e^{x_0 t} \cos(y_0 t) \\ &= x_0 e^{x_0 t} (\cos(y_0 t) + i \sin(y_0 t)) + i y_0 e^{x_0 t} (\cos(y_0 t) + i \sin(y_0 t)) \\ &= (x_0 e^{x_0 t} + i y_0 e^{x_0 t})(\cos(y_0 t) + i \sin(y_0 t)) \\ &= (x_0 + i y_0) e^{x_0 t} e^{i y_0 t} \\ &= z_0 e^{z_0 t} \end{aligned}$$

To summarise, for $\gamma(t) = e^{z_0 t}$, we have $\gamma'(t) = z_0 e^{z_0 t}$.

3.2. Integral of $\gamma : [a, b] \rightarrow \mathbf{C}$

Definition 3.2.1 (Definite Integral of γ). Consider a function $\gamma : [a, b] \rightarrow \mathbf{C}$ with

$$\gamma(t) = u(t) + iv(t),$$

where $u, v : [a, b] \rightarrow \mathbf{R}$. The **definite integral of γ** is defined as

$$\int_a^b \gamma(t) dt := \int_a^b u(t) dt + i \int_a^b v(t) dt$$

provided the integrals of u and v exist.

Improper integrals can be defined in a similar manner.

Example 3.2.2. We illustrate this definition by integrating $\gamma(t) = e^{it}$ on $[0, \pi]$.

$$\begin{aligned} \int_0^\pi e^{it} dt &= \int_0^\pi \cos t dt + i \int_0^\pi \sin t dt \\ &= \left[\sin t \right]_0^\pi + i \left[-\cos t \right]_0^\pi \\ &= (\sin \pi - \sin 0) + i(-\cos \pi + \cos 0) \\ &= 2i \end{aligned}$$

Definition 3.2.3 (Piecewise Continuity). A function $u : [a, b] \rightarrow \mathbf{R}$ is **piecewise continuous on $[a, b]$** if it is continuous on $[a, b]$ except at a finite number of points, where despite its discontinuity on those points, both one sided limits exist.

We call $\gamma(t) = u(t) + iv(t)$ *piecewise continuous* if both u and v are.

Remark 3.2.4. The existence of the integrals

$$\int_a^b u(t) dt \quad \text{and} \quad \int_a^b v(t) dt$$

is guaranteed when γ is piecewise continuous.

Proposition 3.2.5 (Properties of the Integral of γ). Suppose γ and γ_1 are piecewise continuous on $[a, b]$, then

$$(1) \int_a^b z_0 \gamma(t) dt = z_0 \int_a^b \gamma(t) dt, \text{ for any } z_0 \in \mathbf{C}.$$

$$(2) \int_a^b \gamma(t) + \gamma_1(t) dt = \int_a^b \gamma(t) dt + \int_a^b \gamma_1(t) dt.$$

$$(3) \int_a^b \gamma(t) dt = \int_a^c \gamma(t) dt + \int_c^b \gamma(t) dt, \text{ for any } c \in [a, b].$$

$$(4) \int_b^a \gamma(t) dt = - \int_a^b \gamma(t) dt.$$

Proof. These properties follow from the properties of regular real integrals applied to the real and imaginary part of γ and γ_1 . \square

Proposition 3.2.6 (Extension of Fundamental Theorem of Calculus). *Suppose that $\gamma(t) = u(t) + iv(t)$ is continuous on $[a, b]$ and $\Gamma(t) = U(t) + iV(t)$ is differentiable such that $\Gamma'(t) = \gamma(t)$ on $[a, b]$. Then*

$$\int_a^b \gamma(t) dt = \Gamma(b) - \Gamma(a)$$

Proof. By assumption $\Gamma' = \gamma$, therefore $U'(t) = u(t)$ and $V'(t) = v(t)$, therefore

$$\begin{aligned} \int_a^b \gamma(t) dt &= \int_a^b u(t) dt + i \int_a^b v(t) dt \\ &= U(b) - U(a) + i(V(b) - V(a)), \text{ by the Fundamental Theorem of Calculus} \\ &= U(b) + iV(b) - (U(a) + iV(a)) \\ &= \Gamma(b) - \Gamma(a) \end{aligned}$$

\square

Example 3.2.7. We use this proposition to integrate e^{it} on $[0, \pi]$. For this, we first note that

$$\left(\frac{e^{it}}{i} \right)' = \frac{1}{i} (e^{it})' = \frac{i}{i} e^{it} = e^{it}.$$

Therefore,

$$\begin{aligned} \int_0^\pi e^{it} dt &= \left[\frac{e^{it}}{i} \right]_0^\pi \\ &= \left[-ie^{it} \right]_0^\pi \\ &= -ie^{i\pi} + ie^{i \cdot 0} = i + i = 2i \end{aligned}$$

3.3. Contours

So far, we have only defined the integral of a complex-valued function in a single real variable over an interval. Integrals of complex-valued functions in a single complex variable are defined over suitable curves in the complex plane called *contours*.


Definition 3.3.1 (Arcs).

- (1) An **arc**, or **curve**, is a collection of points


$$C = \{z(t) = x(t) + iy(t) : t \in [a, b]\},$$

where $x, y : [a, b] \rightarrow \mathbf{R}$ are continuous functions (which also makes $z : [a, b] \rightarrow \mathbf{C}$ a continuous function). The function $z(t)$ is called a **parametrization of C** .


- (2) An arc (or curve) C is called **simple** or a **Jordan arc** if it does not cross itself, which is equivalent to saying the function $z(t)$ is injective; that is, if $z(t_1) = z(t_2)$ then $t_1 = t_2$.
- (3) If C is simple except for the fact that $z(a) = z(b)$, then C is called a **simple closed curve** or a **Jordan curve**.
- (4) A simple closed curve is **positively oriented** if it is transversed counter-clockwise as t increases from a to b . It is called **negatively oriented** if it is transversed clockwise.




a not simple arc




a simple arc



*a simple closed curve
with positive orientation*



*a simple closed curve
with negative orientation*




*a not simple closed
non-orientable curve*

Lecture 20

Example 3.3.2. The most frequently encountered arcs and curves are line segments and circles.


- (1) The circle of radius R centered at z_0 with positive orientation has as a parametrisation

$$z(t) = z_0 + Re^{it}, \quad t \in [0, 2\pi]$$




(2) The circle of radius R centered at z_0 with negative orientation has as a parametrisation

$$z(t) = z_0 + Re^{-it}, \quad t \in [0, 2\pi]$$



(3) The line segment from z_0 to z_1 in \mathbf{C} has as a parametrisation

$$z(t) = z_0 + (z_1 - z_0)t = (1 - t)z_0 + tz_1, \quad t \in [0, 1]$$



Definition 3.3.3 (Reparametrisation of an arc). Suppose an arc C is parametrised by $z : [a, b] \rightarrow \mathbf{C}$. A map

$$w : [c, d] \rightarrow \mathbf{C}$$

is called an **orientation-preserving reparametrisation** of C if there exists a surjective function

$$\phi : [c, d] \rightarrow [a, b]$$

with continuous derivative such that $\phi(c) = a$ (preserves initial point), $\phi(d) = b$ (preserves final point), $\phi'(s) > 0$ and $w(s) = z(\phi(s))$ (w and z trace out the same arc C).

Example 3.3.4. Note that $z(t) = e^{it}$ for $t \in [0, 2\pi]$ is a parametrisation of the unit circle. Now, consider

$$w : [0, \pi] \rightarrow \mathbf{C}, s \mapsto e^{2is},$$

this is, in fact, an orientation-preserving reparametrisation of the unit circle. To conclude this, we produce the following surjective map

$$\phi : [0, \pi] \rightarrow [0, 2\pi], s \mapsto 2s,$$

we note that $\phi(0) = 0$ and $\phi(\pi) = 2\pi$, furthermore $\phi'(s) = 2 > 0$ which is clearly continuous. Lastly, $z(\phi(s)) = z(2s) = e^{2is} = w(s)$.

Remark 3.3.5. Suppose an arc C is parametrised by $z : [a, b] \rightarrow \mathbf{C}$, a map $w : [c, d] \rightarrow \mathbf{C}$ is called an **orientation-reversing reparametrisation** of C if there exists a surjective function

$$\psi : [c, d] \rightarrow [a, b]$$

with continuous derivative such that $\psi(c) = b$ and $\psi(d) = a$ (swaps initial and final points), $\psi'(s) < 0$ and $w(s) = z(\psi(s))$ (w and z trace out the same arc C).

Consider the unit circle, which has parametrisation $z(t) = e^{it}$, $t \in [0, 2\pi]$. Then $w(t) = e^{-it}$ for $0 \leq t \leq 2\pi$ is an orientation-reversing parametrisation. To see this, we consider the surjective function

$$\psi : [0, 2\pi] \rightarrow [0, 2\pi], s \mapsto 2\pi - s;$$

we note that $\psi(0) = 2\pi$ and $\psi(2\pi) = 0$, furthermore $\psi'(s) = -1 < 0$ and

$$z(\psi(s)) = z(2\pi - s) = e^{2\pi i - is} = e^{-is} = w(s),$$

since $e^{2\pi i} = 1$.

Definition 3.3.6 (Arc length and Smooth arcs).

- (1) If C is parametrised by $z(t) = x(t) + iy(t)$ and $x'(t)$, $y'(t)$ exist and are continuous on $[a, b]$, then C is called a **differentiable arc**.
- (2) The **arc length** of such a differentiable arc C is


$$L(C) = \int_a^b |z'(t)| dt = \int_a^b \sqrt{x'(t)^2 + y'(t)^2} dt$$

- (3) A differentiable curve parametrised by $z(t)$ is called **smooth** if $z'(t) \neq 0$ on $[a, b]$.

Definition 3.3.7 (Contours). A **contour** is an arc consisting of a finite number of smooth arcs joined end to end.

A **simple closed contour** is a contour that does not cross itself except that the initial and final points are the same.

Discussion 3.3.8 (Jordan Curve Theorem). A deep theorem known as the *Jordan Curve theorem* tells us that every simple closed contour C is the boundary of two distinct domains called the **interior of C** , which is bounded, and the **exterior of C** , which is unbounded.



The theorem is geometrically evident but the proof is not easy. We will assume its truth so that we can refer to the interior of a simple closed contour.

3.4. Contour Integration

Definition 3.4.1 (Contour Integral). Suppose $f : G \rightarrow \mathbf{C}$ is a complex function and C is a contour lying in G . If $z(t)$, $t \in [a, b]$, is a parametrisation of C and $f(z(t))$ is piecewise continuous, then the **contour integral of f over C** is

$$\int_C f(z) dz := \int_a^b f(z(t)) z'(t) dt$$

Remark 3.4.2. As C is a contour, $z'(t)$ is piecewise continuous and thus the above integral exists.

Proposition 3.4.3 (Integral is parametrisation-independent). Suppose $z : [a, b] \rightarrow \mathbf{C}$ parametrises C and $w : [c, d] \rightarrow \mathbf{C}$ is an orientation-preserving reparametrisation of C , then

$$\int_C f(z) dz = \int_C f(w) dw$$


Proof. By definition of an orientation-preserving reparametrisation, there exists a surjective map $\phi : [c, d] \rightarrow [a, b]$ such that $\phi(c) = a$, $\phi(d) = b$, $\phi'(s) > 0$ and $w(s) = \phi(z(s))$. Then

$$\begin{aligned} \int_C f(w) dw &= \int_c^d f(w(s)) w'(s) ds \\ &= \int_c^d f(z(\phi(s))) \phi'(z(s)) z'(s) ds && \text{apply chain rule to } w(s) = \phi(z(s)) \\ &= \int_a^b f(z(t)) z'(t) dt && \text{set } t = \phi(s) \\ &= \int_C f(z) dz \end{aligned}$$



□

Discussion 3.4.4 (Notation for Contours).

- (1) Suppose C is a contour, then $-C$ denotes the same set of points as C but with opposite orientation. If $z : [a, b] \rightarrow \mathbf{C}$ is a parametrisation of C , then $w : [-b, -a] \rightarrow \mathbf{C}$ defined as $w(t) := z(-t)$ is a parametrisation of $-C$.



- (2) If C_1 is a contour from z_1 to z_2 and C_2 is a contour from z_2 to z_3 , then their **sum** $C = C_1 + C_2$ is the contour obtained by transversing C_1 and then C_2 .



If C_1 and C_2 have the same final point, then we can consider the sum of C_1 and $-C_2$ and is written as $C_1 - C_2 := C_1 + (-C_2)$.

Proposition 3.4.5 (Properties of Contour Integral). Assume f, g are piecewise continuous on the contours we consider below.

- (1) $\int_C z_0 f(z) dz = z_0 \int_C f(z) dz$, for any $z_0 \in \mathbf{C}$.
- (2) $\int_C f(z) + g(z) dz = \int_C f(z) dz + \int_C g(z) dz$.
- (3) $\int_{-C} f(z) dz = - \int_C f(z) dz$.
- (4) $\int_C f(z) dz = \int_{C_1} f(z) dz + \int_{C_2} f(z) dz$ if $C = C_1 + C_2$.

Proof.

(1) Suppose C is parametrised by $z : [a, b] \rightarrow \mathbf{C}$

$$\begin{aligned} \int_C z_0 f(z) dz &= \int_C z_0 f(z(t)) z'(t) dz \\ &= z_0 \int_a^b f(z(t)) z'(t) dz && \text{by Proposition 3.2.5 (1)} \\ &= z_0 \int_C f(z) dz \end{aligned}$$

(2) This will follow from Proposition 3.2.5 (2).

(3) Suppose C is parametrised by $z : [a, b] \rightarrow \mathbf{C}$, then, as we note before, a parametrisation of $-C$ is $w : [-b, -a] \rightarrow \mathbf{C}$ where $w(t) = z(-t)$. Then

$$\begin{aligned} \int_{-C} f(w) dw &= \int_{-b}^{-a} f(w(t)) w'(t) dt \\ &= - \int_{-b}^{-a} f(z(-t)) z'(-t) dt && \text{apply chain rule to } w(t) = z(-t) \\ &= - \int_{-a}^{-b} f(z(-t)) z'(-t) dt && \text{by Proposition 3.2.5 (4)} \\ &= \int_a^b f(z(s)) z'(s) ds && \text{set } s = -t \\ &= \int_C f(z) dz \end{aligned}$$

(4) We leave this as an exercise for the motivated student. □


Example 3.4.6.

(1) Integrate $f(z) = \frac{1}{z}$ over the following contours:

- C_1 : upper semicircle of the unit circle, from 1 to -1 .
- C_2 : lower semicircle of the unit circle, from 1 to -1 .
- C_3 : $C_1 - C_2$.


For C_1 , parametrise C_1 as $z(t) = e^{it}$, $0 \leq t \leq \pi$. Then

$$\int_{C_1} \frac{1}{z} dz = \int_0^\pi \frac{1}{e^{it}} i e^{it} dt = i \int_0^\pi dt = \pi i$$




For C_2 , parametrise C_2 as $z(t) = e^{-it}$, $0 \leq t \leq \pi$. Then

$$\int_{C_1} \frac{1}{z} dz = \int_0^\pi \frac{1}{e^{-it}} (-ie^{-it}) dt = -i \int_0^\pi dt = -\pi i$$



Note


$$\begin{aligned} \int_{C_3} \frac{1}{z} dz &= \int_{C_1 - C_2} \frac{1}{z} dz = \int_{C_1} \frac{1}{z} dz + \int_{-C_2} \frac{1}{z} dz \\ &= \int_{C_1} \frac{1}{z} dz - \int_{C_2} \frac{1}{z} dz \\ &= \pi i - (-\pi i) \\ &= 2\pi i \end{aligned}$$



This example shows that the integral may depend on the path taken and not just on the endpoints. Also, the integral over a closed contour may be non-zero.

(2) Integrate $f(z) = z$ over *any* contour C connecting a point w_1 to a point w_2 .

First, suppose C is a smooth arc joining w_1 and w_2 with parametrisation $z : [a, b] \rightarrow \mathbf{C}$.



Since,

$$\left(\frac{z(t)^2}{2} \right)' = \frac{z'(t)z(t) + z(t)z'(t)}{2} = z(t)z'(t).$$

Therefore,

$$\begin{aligned} \int_C f(z) dz &= \int_C z dz = \int_a^b z(t) z'(t) dt \\ &= \frac{z(b)^2}{2} - \frac{z(a)^2}{2}, \text{ by Proposition 3.2.6} \\ &= \frac{w_2^2 - w_1^2}{2} \end{aligned}$$


Now, if C is a contour, we can write $C = C_1 + \cdots + C_n$, where C_i is a smooth arc joining z_i to z_{i+1} with $z_1 = w_1$ and $z_{n+1} = w_2$. Then,

$$\begin{aligned}\int_C z \, dz &= \sum_{i=1}^n \int_{C_i} z \, dz, \text{ by Proposition 3.4.5 (4)} \\ &= \sum_{i=1}^n \frac{z_{i+1}^2 - z_i^2}{2} \\ &= \frac{z_{n+1}^2 - z_1^2}{2} \\ &= \frac{w_2^2 - w_1^2}{2}\end{aligned}$$

This example shows that some integrals do depend only on the end points and not the path taken. Also, for any contour C is closed, that is, when $w_2 = w_1$, we have shown hence that

$$\int_C z \, dz = 0.$$

(3) Integrate $f(z) = z^m \bar{z}^n$, for $m, n \in \mathbf{Z}$, over the unit circle C .



Parametrise C as $z(t) = e^{it}$, $0 \leq t \leq 2\pi$. Then,

$$\begin{aligned}\int_C f(z) \, dz &= \int_C z^m \bar{z}^n \, dz \\ &= \int_0^{2\pi} (e^{it})^m (\overline{e^{it}})^n i e^{it} \, dt \\ &= i \int_0^{2\pi} (e^{it})^m (e^{-it})^n e^{it} \, dt \\ &= i \int_0^{2\pi} e^{imt} e^{-int} e^{it} \, dt \\ &= i \int_0^{2\pi} e^{(m-n+1)it} \, dt\end{aligned}$$

Case I. $m = n - 1$

$$\int_C f(z) dz = i \int_0^{2\pi} e^{(m-n+1)it} = i \int_0^{2\pi} dt = 2\pi i$$

Case II. $m \neq n - 1$


$$\begin{aligned} \int_C f(z) dz &= i \int_0^{2\pi} e^{(m-n+1)it} = i \left[\frac{e^{(m-n+1)it}}{i(m-n+1)} \right]_0^{2\pi} \\ &= \frac{1}{m-n+1} (e^{2(m-n+1)\pi i} - e^0) \\ &= \frac{1}{m-n+1} (1 - 1) \\ &= 0 \end{aligned}$$

Lecture 22 Some examples involving a branch of a multi-valued function.

(4) Integrate the branch of square root

$$f(z) = z^{1/2} = e^{(1/2)\log z}, \quad |z| > 0, \quad 0 < \arg z < 2\pi$$

over the contour



$$C : z(t) = Re^{it}, \quad R > 0, \quad 0 \leq t \leq \pi$$

Note that $f(z)$ is not defined at the initial point $z = R$ of the contour C as $\arg R = 0$, that is, $f(z(t))$ is not defined for $t = 0$. The integral

$$\int_C f(z) dz = \int_0^\pi f(z(t)) z'(t) dt$$

nevertheless exists as the integrand $f(z(t)) z'(t)$ is piecewise continuous on $[0, \pi]$. To see this, we note that for $0 < t \leq \pi$

$$\begin{aligned} f(z(t)) z'(t) &= e^{(1/2)\log Re^{it}} Rie^{it} = iRe^{(\ln R + it)/2} e^{it} \\ &= iR(R^{1/2} e^{it/2}) e^{it} \\ &= iR^{3/2} e^{3it/2} \\ &= iR^{3/2} \left(\cos \frac{3t}{2} + i \sin \frac{3t}{2} \right) = R^{3/2} \left(-\sin \frac{3t}{2} + i \cos \frac{3t}{2} \right) \end{aligned}$$


The right hand limits of the real and imaginary parts of $f(z(t)) z'(t)$ at $t = 0$ exist, and equal 0 and $R^{3/2}$. Therefore, $f(z(t)) z'(t)$ is continuous on $[0, \pi]$ with its value at $t = 0$ defined as $iR^{3/2}$. Hence,

$$\begin{aligned} \int_C f(z) dz &= \int_0^\pi f(z(t)) z'(t) dt = \int_0^\pi iR^{3/2} e^{3it/2} dt \\ &= iR^{3/2} \int_0^\pi e^{3it/2} dt \\ &= iR^{3/2} \left[\frac{e^{3it/2}}{3i/2} \right]_0^\pi \\ &= \frac{2}{3} R^{3/2} (e^{3\pi i/2} - e^0) \\ &= \frac{2}{3} R^{3/2} (-i - 1) \\ &= -\frac{2}{3} R^{3/2} (1 + i) \end{aligned}$$

(5) Integrate the principal branch of

$$f(z) = z^{i-1} = e^{(i-1)\text{Log} z}, \quad |z| > 0, \quad -\pi < \text{Arg} z < \pi$$

over the contour



$$C : z(t) = e^{it}, \quad R > 0, \quad -\pi \leq t \leq \pi$$

Since the curve crosses the branch cut, we need to check if integrand $f(z(t)) z'(t)$ is piecewise continuous on $[-\pi, \pi]$. To see this, we note that for $-\pi < t \leq \pi$

$$\begin{aligned} f(z(t)) z'(t) &= e^{(i-1)\text{Log} e^{it}} i e^{it} = i e^{(i-1)(\ln 1 + it)} e^{it} \\ &= i e^{(i-1)it} e^{it} \\ &= i e^{(i-1)it + it} \\ &= i e^{i^2 t} = i e^{-t} \end{aligned}$$

The right hand limits of the real and imaginary parts of $f(z(t)) z'(t)$ at $t = \pi$ exist, and equal 0 and $e^{-\pi}$. Therefore, $f(z(t)) z'(t)$ is continuous on $[-\pi, \pi]$ with its value at $t = -\pi$ defined

as $ie^{-\pi}$. Hence,

$$\begin{aligned}
 \int_C f(z) dz &= \int_{-\pi}^{\pi} f(z(t)) z'(t) dt = \int_{-\pi}^{\pi} ie^{-t} dt \\
 &= i \int_{-\pi}^{\pi} e^{-t} dt \\
 &= i \left[-e^{-t} \right]_{-\pi}^{\pi} \\
 &= i \left(-e^{-\pi} - (-e^{-(-\pi)}) \right) \\
 &= i (e^{\pi} - e^{-\pi})
 \end{aligned}$$

3.5. Estimating Contour Integrals

Lemma 3.5.1 (Triangle Inequality for Integrals). *Suppose $\gamma : [a, b] \rightarrow \mathbf{C}$ is piecewise continuous. Then*

$$\left| \int_a^b \gamma(t) dt \right| \leq \int_a^b |\gamma(t)| dt$$

Proof. Let's first assume

$$\int_a^b \gamma(t) dt = 0,$$

then the lemma holds as $|\gamma(t)| \geq 0$ for all $t \in [a, b]$ and so its integral is non-negative. Otherwise, let

$$r_0 e^{it_0} = \int_a^b \gamma(t) dt \neq 0.$$

Then,

$$\begin{aligned}
 \left| \int_a^b \gamma(t) dt \right| &= |r_0 e^{it_0}| = r_0 = \operatorname{Re} r_0 = \operatorname{Re}(r_0 e^{it_0} e^{-it_0}) \\
 &= \operatorname{Re} \left(e^{-it_0} \int_a^b \gamma(t) dt \right) \\
 &= \operatorname{Re} \left(\int_a^b e^{-it_0} \gamma(t) dt \right) \\
 &= \int_a^b \operatorname{Re}(e^{-it_0} \gamma(t)) dt \\
 &\leq \int_a^b |e^{-it_0} \gamma(t)| dt, \text{ using Discussion 1.2.1} \\
 &= \int_a^b |e^{-it_0}| |\gamma(t)| dt \\
 &= \int_a^b |\gamma(t)| dt
 \end{aligned}$$

□

Theorem 3.5.2 (Bound for Contour Integrals). Suppose that C is a contour of length L and f is piecewise continuous on C . Then

$$\left| \int_C f(z) dz \right| \leq \max_{z \in C} |f(z)| \cdot L(C)$$

Proof. Suppose $z : [a, b] \rightarrow \mathbf{C}$ parametrises C . By assumption $f(z(t))$ is piecewise continuous on $[a, b]$. Hence, $\max_{z \in C} |f(z)| = \max_{t \in [a, b]} |f(z(t))|$ is finite as $f(z(t))$ is continuous on a closed and bounded interval. Thus,

$$\begin{aligned} \left| \int_C f(z) dz \right| &= \left| \int_a^b f(z(t)) z'(t) dz \right| \\ &\leq \int_a^b |f(z(t)) z'(t)| dz, \text{ by Lemma 3.5.1} \\ &= \int_a^b |f(z(t))| |z'(t)| dz \\ &\leq \int_a^b \max_{t \in [a, b]} |f(z(t))| |z'(t)| dz \\ &= \max_{t \in [a, b]} |f(z(t))| \int_a^b |z'(t)| dz = \max_{t \in [a, b]} |f(z(t))| \cdot L(C) = \max_{z \in C} |f(z)| \cdot L(C) \quad \square \end{aligned}$$

Lecture 23 Example 3.5.3.

(1) Finding a bound for

$$\int_C \frac{z^2 + 1}{z^3 + 2} dz,$$

where C is the semicircle $z(t) = 2e^{it}$, $0 \leq t \leq \pi$.

All we need to find is an $M > 0$ such that, for all $z \in C$

$$\left| \frac{z^2 + 1}{z^2 + 3} \right| \leq M, \quad \text{because then} \quad \max_{z \in C} \left| \frac{z^2 + 1}{z^2 + 3} \right| \leq M$$

Suppose $z \in C$, then $|z| = 2$, and therefore

$$|z^2 + 1| \leq |z|^2 + 1 = 5;$$

also,

$$|z^3 + 2| \geq ||z|^3 - 2| = |2^3 - 2| = 6.$$

Together, we get, for any $z \in C$

$$\left| \frac{z^2 + 1}{z^2 + 3} \right| \leq \frac{5}{6}.$$

Hence,

$$\left| \int_C \frac{z^2 + 1}{z^3 + 2} dz \right| \leq \max_{z \in C} \left| \frac{z^2 + 1}{z^2 + 3} \right| \cdot L(C) \leq \frac{5}{6} \cdot L(C) = \frac{5}{6} \cdot 2\pi = \frac{5\pi}{3}$$

(2) Show that

$$\lim_{R \rightarrow \infty} \int_{C_R} \frac{z^2 + z}{z^4 + 2z^2 + 1} dz = 0,$$

where C_R is the semicircle $z(t) = Re^{it}$, $0 \leq t \leq 2\pi$. Note that $L(C) = 2\pi R$.

Let $z \in C_R$, then $|z| = R$, and therefore

$$|z^2 + z| \leq |z|^2 + |z| = R^2 + R;$$

also,

$$|z^4 + 2z^2 + 1| \geq |(z^2 + 1)| = |z^2 + 1|^2 \geq ||z|^2 - 1|^2 = |R^2 - 1|^2 = (R^2 - 1)^2.$$

Together, we get, for any $z \in C$ and $R > 1$

$$\left| \frac{z^2 + z}{z^4 + 2z^2 + 1} \right| \leq \frac{R^2 + R}{(R^2 - 1)^2}.$$

Hence,

$$\left| \int_{C_R} \frac{z^2 + z}{z^4 + 2z^2 + 1} dz \right| \leq \frac{R^2 + R}{(R^2 - 1)^2} \cdot 2\pi R \rightarrow 0, \text{ as } R \rightarrow \infty$$

Therefore,

$$\lim_{R \rightarrow \infty} \int_{C_R} \frac{z^2 + z}{z^4 + 2z^2 + 1} dz = 0,$$

by the Sandwich theorem.

Example 3.5.4. Finding a bound for

$$\int_C \frac{z^2 - 1}{z^4 + 2} dz,$$

where C is the sector $z(t) = 5e^{it}$, $\pi/4 \leq t \leq 3\pi/4$.

Answer. Let's first compute $L(C)$. We first note that $z'(t) = i5e^{it}$, therefore,

$$\begin{aligned} L(C) &= \int_{\pi/4}^{3\pi/4} |z'(t)| dt \\ &= \int_{\pi/4}^{3\pi/4} |5ie^{it}| dt \\ &= \int_{\pi/4}^{3\pi/4} 5 dt \\ &= 5 \int_{\pi/4}^{3\pi/4} dt = 5 \left(\frac{3\pi}{4} - \frac{\pi}{4} \right) = \frac{5\pi}{2} \end{aligned}$$

Now, suppose $z \in C$, then $|z| = 5$, and therefore

$$|z^2 - 1| \leq |z^2| + |-1| = |z|^2 + 1 = 26;$$

also,

$$|z^4 + 2| \geq ||z^4| - |2|| = ||z|^4 - 2| = 623.$$

Together, we get, for any $z \in C$

$$\left| \frac{z^2 - 1}{z^4 + 2} \right| \leq \frac{26}{623}, \quad \text{hence } \max_{z \in C} \left| \frac{z^2 - 1}{z^4 + 2} \right| \leq \frac{26}{623}$$

Hence,

$$\left| \int_C \frac{z^2 - 1}{z^4 + 2} dz \right| \leq \max_{z \in C} \left| \frac{z^2 - 1}{z^4 + 2} \right| \cdot L(C) \leq \frac{26}{623} \cdot L(C) = \frac{26}{623} \cdot \frac{5\pi}{2} = \frac{65\pi}{623}$$

□


3.6. Antiderivatives & Fundamental Theorem of Contour Integrals

Discussion 3.6.1. Suppose C is a contour joining z_1 to z_2 . In general, the value of the integral

$$\int_C f(z) dz$$

depends on C . For example, we have seen that

$$\int_{C_1} \frac{1}{z} dz = \pi i \quad \text{and} \quad \int_{C_2} \frac{1}{z} dz = -\pi i$$



But on the other hand we have also seen that

$$\int_C z dz = \frac{z_2^2 - z_1^2}{2}$$

for any contour C with initial point z_1 and end point z_2 .

The difference between these functions turns out to be that $f(z) = z$ has an antiderivative on \mathbf{C} while $g(z) = 1/z$ does not any domain containing C_1 and C_2 .

Definition 3.6.2 (Antiderivative). Suppose that f is a continuous function on a domain G . Any holomorphic function $F : G \rightarrow \mathbf{C}$ is called an **antiderivative** of f if $F'(z) = f(z)$ for every $z \in G$.

Definition 3.6.3 (Independence of Path). Let $f : G \rightarrow \mathbf{C}$ be a continuous function on a domain G and fix $z_1, z_2 \in G$. If

$$\int_{C_1} f(z) dz = \int_{C_2} f(z) dz$$

for any pair of contours C_1 and C_2 joining z_1 to z_2 , then the integral of f from z_1 to z_2 is *independent of path* and we denote the unique value by

$$\int_{z_1}^{z_2} f(z) dz.$$

So, for instance, we would write

$$\int_{z_1}^{z_2} z dz = \frac{z_2^2 - z_1^2}{2},$$

since we have already proved the integral of $f(z) = z$ from z_1 to z_2 , for any $z_1, z_2 \in \mathbf{C}$, is independent of path.

Theorem 3.6.4 (Fundamental Theorem of Contour Integrals). Suppose f is continuous on a domain G . The following are equivalent.

- (1) f has an antiderivative $F : G \rightarrow \mathbf{C}$.
- (2) For all $z_1, z_2 \in G$, the integral of f from z_1 to z_2 are independent of path.
- (3) If C is any closed contour lying in G , then

$$\int_C f(z) dz = 0$$

If any of these conditions hold, then the unique value of the integral in (2) is given as

$$\int_{z_1}^{z_2} f(z) dz = F(z_2) - F(z_1)$$

where F is the antiderivative given in (1).

Proof.

- (1) \Rightarrow (2) Suppose f has an antiderivative $F : G \rightarrow \mathbf{C}$. Let $z_1, z_2 \in G$ and let C be any contour with initial point z_1 to z_2 and lying in G .

First assume C is a smooth arc parametrised by $z : [a, b] \rightarrow \mathbf{C}$; therefore, in particular, $z(a) = z_1$ and $z(b) = z_2$. Then we first note

$$(F \circ z)'(t) = F'(z(t))z'(t) = f(z(t))z'(t)$$

That is, we have found an antiderivative of $f(z(t))z'(t)$, the function $F \circ z$. Hence,


$$\begin{aligned} \int_C f(z) dz &= \int_a^b f(z(t))z'(t) dt \\ &= F(z(a)) - F(z(b)), \quad \text{by Proposition 3.2.6} \\ &= F(z_2) - F(z_1) \end{aligned}$$

Now, assume C is a contour; that is, we can write $C = C_1 + \cdots + C_n$, where C_i 's are smooth arcs with initial point w_i and end point w_{i+1} . In particular, $w_1 = z_1$ and $w_{n+1} = z_2$. Then,

$$\begin{aligned}\int_C f(z) dz &= \sum_{i=1}^n \int_{C_i} f(z) dz \\ &= \sum_{i=1}^n F(w_{i+1}) - F(w_i) \\ &= F(w_{n+1}) - F(w_1) \\ &= F(z_2) - F(z_1)\end{aligned}$$

Since $F(z_2) - F(z_1)$ only depends on z_1 and z_2 and not the contour itself, we have proved the claim.

(2) \Rightarrow (3) Let C be any closed contour lying in G , and choose two distinct points z_1 and z_2 on C . Let C_1 and C_2 be contours from z_1 to z_2 such that $C = C_1 - C_2$.



By assumption, the integral of f from z_1 to z_2 is independent of path, therefore

$$\int_{C_1} f(z) dz = \int_{C_2} f(z) dz$$

Hence,

$$\int_C f(z) dz = \int_{C_1 - C_2} f(z) dz = \int_{C_1} f(z) dz - \int_{C_2} f(z) dz = 0,$$

as claimed.

(3) \Rightarrow (2) Suppose

$$\int_C f(z) dz = 0$$

for any closed contour C lying in G . Let $z_1, z_2 \in G$ and C_1 and C_2 are two contours with initial point z_1 and end point z_2 . Then $C_1 - C_2$ is a closed contour, and therefore by assumption

$$0 = \int_{C_1 - C_2} f(z) dz = \int_{C_1} f(z) dz - \int_{C_2} f(z) dz$$

Hence,

$$\int_{C_1} f(z) dz = \int_{C_2} f(z) dz,$$

as claimed.

(2) \Rightarrow (1) Assume (2) (and also (3), since we've shown them to be equivalent). We need to show that f has an antiderivative on G . Fix any point $z_0 \in G$ and define

$$F(w) = \int_{z_0}^w f(z) dz,$$

which is well defined by (2). We need to show $F'(w) = f(w)$ for any $w \in G$. That is,

$$\lim_{h \rightarrow 0} \frac{F(w+h) - F(w)}{h} = f(w)$$

Let $\varepsilon > 0$ and consider an $z \in G$. Since f is continuous at z , we can find $\delta > 0$ such that

$$\text{if } |z - w| < \delta, \quad \text{then } |f(z) - f(w)| < \varepsilon$$


For $w \in G$, since G is a domain and so in particular an open set, we can find a $d > 0$ such that $D_d(w) \subseteq G$. Pick a $h \in \mathbf{C}$ such that $0 < |h| < \min\{d, \delta\}$. Then $0 < |h| < d$ and $0 < |h| < \delta$. In particular, $w + h \in D_d(w) \subseteq G$; then,

$$F(w+h) - F(w) = \int_{z_0}^{w+h} f(z) dz - \int_{z_0}^w f(z) dz = \int_w^{w+h} f(z) dz$$

Since our integrals are path-independent, we assume that the integral above is over a line segment from w to $w + h$, which lies in G , since $D_d(w)$ is convex. Also,

$$f(w) = \frac{f(w)h}{h} = \frac{1}{h} f(w) \int_w^{w+h} dz = \frac{1}{h} \int_w^{w+h} f(w) dz$$

Also, since $|h| < \delta$, then $|z - w| < \delta$ for any point z lying on ℓ , the line segment joining w to $w + h$. Therefore, $|f(z) - f(w)| < \varepsilon$ for any $z \in \ell$, that is, $\max_{z \in \ell} |f(z) - f(w)| < \varepsilon$.



Using the preceding computations we have

$$\begin{aligned}
 \left| \frac{F(w+h) - F(w)}{h} - f(w) \right| &= \left| \frac{1}{h} \int_w^{w+h} f(z) dz - \frac{1}{h} \int_w^{w+h} f(w) dz \right| \\
 &= \frac{1}{h} \left| \int_w^{w+h} f(z) - f(w) dz \right| \\
 &\leq \frac{1}{h} \max_{z \in \ell} |f(z) - f(w)| \cdot L(\ell) \\
 &< \frac{\varepsilon}{h} \cdot L(\ell) \\
 &= \varepsilon, \quad \text{since } L(\ell) = h
 \end{aligned}$$

We have shown that given an $\varepsilon > 0$, there exists a $\delta > 0$ such that


$$\text{if } |h| < \delta, \quad \text{then } \left| \frac{F(w+h) - F(w)}{h} - f(w) \right| < \varepsilon$$

That is, $F'(w) = f(w)$, for all $w \in G$.

□

Lecture 24 **Example 3.6.5.**

- (1) The function $f(z) = 1/z$ has no antiderivative on \mathbf{C}^* . In fact, it has no antiderivative on any domain G containing a deleted neighbourhood of 0. Take a circle $C_\varepsilon = C_\varepsilon(0)$ with radius $\varepsilon > 0$ such that it lies in our domain G .



Then,

$$\begin{aligned}
 \int_{C_\varepsilon} \frac{1}{z} dz &= \int_0^{2\pi} \frac{1}{\varepsilon e^{it}} i \varepsilon e^{it} dt \\
 &= \int_0^{2\pi} i dt \\
 &= 2\pi i
 \end{aligned}$$

By Theorem 3.6.4, $f(z)$ does not have an antiderivative on such a domain, as the integral over the closed contour C_ε was non-zero. The problem is as follows: it is true that that a branch of the logarithm $F(z) = \log z$ is such that

$$F'(z) = \frac{1}{z},$$


but it is only holomorphic on the complement of the branch cut. Since our domain contains a deleted neighbourhood of 0, it has a non-empty intersection with any branch cut we take, and therefore F is not holomorphic on G . This argument, in particular, holds for the domain \mathbf{C}^* .

- (2) The function $f(z) = \cos z$ is entire on \mathbf{C} , so is $F(z) = \sin z$. Moreover $F'(z) = \cos z = f(z)$, so f has an antiderivative on \mathbf{C} . So, for instance

$$\int_0^{\pi i} \cos z \, dz = \sin \pi i - \sin 0 = \sin \pi i$$

- (3) Although $f(z) = 1/z$ has no antiderivative on any domain containing a deleted neighbourhood of 0, we can integrate f over a circle C by using two different antiderivatives.

Let C_1 be parametrised by $z(t) = e^{it}$, $t \in [-\pi/2, \pi/2]$, a contour from $-i$ to i .




On $\mathbf{C} \setminus \mathbf{R}_{\leq 0}$, $f(z)$ has an antiderivative, namely the principal branch of the logarithm

$$\text{Log } z = \ln |z| + i \text{Arg } z, \quad -\pi < \text{Arg } z < \pi$$

Then, by Theorem 3.6.4

$$\begin{aligned} \int_{C_1} \frac{1}{z} \, dz &= \text{Log } i - \text{Log }(-i) \\ &= (\ln |i| + i \text{Arg } i) - (\ln |-i| + i \text{Arg }(-i)) \\ &= \left(\ln 1 + i \frac{\pi}{2} \right) - \left(\ln 1 - i \frac{\pi}{2} \right) \\ &= \pi i \end{aligned}$$

Let C_2 be parametrised by $z(t) = e^{it}$, $t \in [\pi/2, 3\pi/2]$, a contour from i to $-i$.



On $\mathbb{C} \setminus \mathbb{R}_{\geq 0}$, $f(z)$ has an antiderivative, namely the following branch of the logarithm

$$\log z = \ln |z| + i \arg z, \quad 0 < \arg z < 2\pi$$

Then, by Theorem 3.6.4

$$\begin{aligned} \int_{C_2} \frac{1}{z} dz &= \log(-i) - \log i \\ &= (\ln |i| + i \arg(-i)) - (\ln |-i| + i \arg i) \\ &= \left(\ln 1 + i \frac{3\pi}{2} \right) - \left(\ln 1 + i \frac{\pi}{2} \right) \\ &= \pi i \end{aligned}$$

Hence,

$$\int_C \frac{1}{z} dz = \int_{C_1} \frac{1}{z} dz + \int_{C_2} \frac{1}{z} dz = \pi i + \pi i = 2\pi i$$

3.7. Cauchy-Goursat Theorem

Discussion 3.7.1. The Cauchy-Goursat theorem gives a sufficient condition for the integral of a function over a simple closed curve to be zero. The theorem has powerful implications, ultimately it leads to

- The Cauchy Integral formula.
- The theory of residues for computing contour integrals.
- A method to evaluate real-valued functions in a real variable, using contour integration.

Historically, a weaker version of the theorem was first proved by Cauchy. We prove this first.

We first note the following.

- (1) Contour integrals are related to line (or path) integrals. We note this by writing our function $f(z) = f(x + iy) = u(x, y) + i v(x, y)$ and formally writing $dz = dx + i dy$. Then formally,

$$\begin{aligned}\int_C f(z) dz &= \int_C (u + iv)(dx + i dy) \\ &= \int_C u dx - v dy + i \int_C u dy + v dx\end{aligned}$$

- (2) **Green's Theorem.** Suppose C is a simple closed contour in \mathbf{R}^2 and let R be the region enclosed by C and including C . If $P(x, y)$ and $Q(x, y)$ have continuous partial derivatives on R . Then

$$\int_C P dx + Q dy = \iint_R \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} dA = \iint_R (Q_x - P_y) dA$$

Theorem 3.7.2 (Weak Cauchy Integral Theorem). *Let C be a simple closed contour, and let R denote the region consisting of C and its interior. If f is holomorphic on R and f' continuous on R , then*

$$\int_C f(z) dz = 0.$$

Proof. If $f(z) = u(x, y) + i v(x, y)$ is holomorphic on R , then the Cauchy-Riemann equations hold, and so $u_x = v_y$ and $u_y = -v_x$ on R , and $f'(z) = u_x + i v_x = v_y - i u_y$.

Since f' is continuous, so are u_x , u_y , v_x and v_y . Hence,

$$\begin{aligned}\int_C f(z) dz &= \int_C u dx - v dy + i \int_C u dy + v dx \\ &= \iint_R (-v_x - u_y) dA + i \iint_R (u_x - v_y) dA, \quad \text{by Green's theorem} \\ &= 0, \quad \text{using Cauchy-Riemann equations}\end{aligned}$$

□

Goursat was the first to prove that the assumption on the continuity of f' can be omitted. This turns out to be essential for the theory of holomorphic functions. The problem is that it may be difficult to prove the derivative of holomorphic function is continuous.


Theorem 3.7.3 (Cauchy-Goursat Theorem). *Let C be a simple closed contour, and let R denote the region consisting of C and its interior. If f is holomorphic on R , then*

$$\int_C f(z) dz = 0.$$

Proof (skipped in class). **For simplicity, we will assume C is a square.** The idea is to “divide and conquer”. We break the curve into a finite number of smaller squares on which we can estimate the integral. We first construct a sequence of positively oriented curves $S^{(k)}$, each of which is the boundary of a square region $R^{(k)}$.

To begin with, set $S^{(0)} = C$. Then, inductively, after the first k squares have been chosen, we

define $(k+1)^{\text{th}}$ square as follows. Divide $S^{(k)}$ into four congruent squares with positive orientation: $S_1^{(k)}, S_2^{(k)}, S_3^{(k)}, S_4^{(k)}$.



Note that the integral of f along the shared boundaries of these squares cancel. Hence,

$$\sum_{i=1}^4 \int_{S_i^{(k)}} f(z) dz = \int_{S^{(k)}} f(z) dz$$

We choose $S^{(k+1)}$ to be one of the squares $S_j^{(k)}$ such that

$$\left| \int_{S^{(k+1)}} f(z) dz \right| = \left| \int_{S_j^{(k)}} f(z) dz \right| = \left| \max_{i=1}^4 \int_{S_i^{(k)}} f(z) dz \right|$$

At this point, we have a sequence $S^{(0)}, \dots, S^{(k)}, \dots$. Note that, by triangle inequality

$$\left| \int_{S^{(k)}} f(z) dz \right| \leq \sum_{i=1}^4 \left| \int_{S_i^{(k)}} f(z) dz \right| \leq 4 \left| \int_{S^{(k+1)}} f(z) dz \right|$$

So, inductively we get

$$\left| \int_C f(z) dz \right| = \left| \int_{S^{(0)}} f(z) dz \right| \leq 4^n \left| \int_{S^{(n)}} f(z) dz \right| \quad (*)$$

We record some more facts. Denote by $d^{(n)}$ the length of the diagonal of the n^{th} square $S^{(n)}$ and denote by $P^{(n)}$ its perimeter. Then,

$$d^{(n)} = \frac{1}{2^n} \cdot d^{(0)}$$

$$P^{(n)} = \frac{1}{2^n} \cdot P^{(0)}$$

Also, $d^{(n)}, P^{(n)} \rightarrow 0$, as $n \rightarrow \infty$.

Next, consider the associated sequence of regions

$$R = R^{(0)} \supseteq R^{(1)} \supseteq \dots \supseteq R^{(k)} \supseteq \dots$$

Each $R^{(k)}$ is compact (closed and bounded) and hence, using a fact from topology, there exists a unique point

$$z_0 \in \bigcap_{i \geq 0} R^{(i)}.$$

Since $z_0 \in R^{(0)} = R$, f is holomorphic at z_0 . So, we define the following function on R

$$\psi(z) = \begin{cases} \frac{f(z) - f(z_0)}{z - z_0} - f'(z_0) & \text{if } z \neq z_0 \\ 0 & \text{if } z = z_0 \end{cases}$$

and we note

$$\lim_{z \rightarrow z_0} \psi(z) = f'(z_0) - f'(z_0) = 0 = \psi(z_0),$$

and therefore ψ is continuous at z_0 . We can write

$$f(z) = f(z_0) + (z - z_0)(\psi(z) + f'(z_0)) = f(z_0) + f'(z_0)(z - z_0) + \psi(z)(z - z_0)$$

Note that $f(z_0)$ and $f'(z_0)(z - z_0)$ have antiderivatives on \mathbf{C} , hence, by Theorem 3.6.4, we have

$$\begin{aligned} \int_{S^{(n)}} f(z) dz &= \int_{S^{(n)}} f(z_0) dz + \int_{S^{(n)}} f'(z_0)(z - z_0) dz + \int_{S^{(n)}} \psi(z)(z - z_0) dz \\ &= 0 + 0 + \int_{S^{(n)}} \psi(z)(z - z_0) dz \\ &= \int_{S^{(n)}} \psi(z)(z - z_0) dz \end{aligned}$$

Consider $\varepsilon > 0$. Since ψ is continuous at z_0 with $\psi(z_0) = 0$, choose $\delta > 0$ such that

$$\text{if } |z - z_0| < \delta, \quad \text{then } |\psi(z)| < \varepsilon$$

Since $d^{(n)} \rightarrow 0$, as $n \rightarrow \infty$, we choose an $N \in \mathbf{Z}_{>0}$ such that $|d^{(n)}| < \delta$ for every $n \geq N$. Thus, if $z \in S^{(N)}$, then $|z - z_0| < |d^{(N)}| < \delta$ and therefore $|\psi(z)| < \varepsilon$ for every $z \in S^{(N)}$. Hence,

$$\max_{z \in S^{(N)}} |z - z_0| < d^{(N)} \quad \text{and} \quad \max_{z \in S^{(N)}} |\psi(z)| < \varepsilon$$

Hence, we obtain

$$\begin{aligned} \left| \int_{S^{(N)}} f(z) dz \right| &= \left| \int_{S^{(N)}} \psi(z)(z - z_0) dz \right| \\ &\leq \max_{z \in S^{(N)}} |\psi(z)| |z - z_0| \cdot L(S^{(N)}) \\ &< \varepsilon \cdot d^{(N)} \cdot L(S^{(N)}) \\ &= d^{(N)} p^{(N)} \varepsilon \\ &= \frac{1}{4^N} d^{(0)} p^{(0)} \varepsilon \end{aligned}$$

By (*), we have

$$\begin{aligned} \left| \int_C f(z) dz \right| &\leq 4^N \left| \int_{S^{(N)}} f(z) dz \right| \\ &< 4^N \cdot \frac{1}{4^N} d^{(0)} p^{(0)} \varepsilon \\ &= d^{(0)} p^{(0)} \varepsilon \end{aligned}$$

Since $\varepsilon > 0$ is arbitrary, we necessarily get that

$$\left| \int_C f(z) dz \right| \leq 0$$

Thus,

$$\int_C f(z) dz = 0$$

□

3.8. Simply Connected Domains

Definition 3.8.1 (Simply Connected Domain). A domain G is called **simply connected** if it has the following property: if C is any simple closed contour lying in G and z is interior to C , then $z \in G$.

Intuitively, a simply connected domain is a domain that has no “holes”.


Open disks, complex plane, interior of any simple closed contour etc. are all examples of simply connected domains. While deleted open disks, $\mathbb{C} \setminus \{p\}$ etc. are examples of non-simply connected domains.

A result similar to Theorem 3.7.3 holds for closed contours, not necessarily simple, provided they lie in a simply connected domain.

Theorem 3.8.2 (Cauchy-Goursat Theorem for Simply Connected Domain). Suppose f is holomorphic on a simply connected G . If C is any closed contour lying in G , then

$$\int_C f(z) dz = 0.$$

Proof. We are presented with two cases: C has finitely many self-intersections, or infinitely many self-intersections. Let's focus on the first cases, where the proof is a consequence of Theorem 3.7.3.



Suppose C has n -many self-intersections, then those points of self-intersections allow us to write

$$C = C_1 + C_2 + \cdots + C_{n+1},$$

where each C_i is a simple closed contour that all, necessarily, lie in G .

Therefore f is holomorphic at each point interior of and on C_i , hence by Theorem 3.7.3 we get

$$\int_{C_i} f(z) dz = 0$$

Finally, we have

$$\int_C f(z) dz = \sum_{i=1}^n \int_{C_i} f(z) dz = 0$$

as claimed.

The proof in the case the contour has infinitely many self-intersections is subtle, so we assume validity without a proof. \square

Corollary 3.8.3 (Antiderivatives of Holomorphic Functions). *If f is holomorphic on a simply connected domain G , then f has an antiderivative on G .*

Proof. By Theorem 3.8.2,

$$\int_C f(z) dz = 0$$

for any closed contour C lying in G . By Theorem 3.6.4, this is equivalent to f having an antiderivative on G . \square

Corollary 3.8.4 (Entire Functions have Antiderivatives). *Suppose f is entire, then f has an antiderivative on \mathbb{C} which is necessarily also entire.*

Proof. \mathbb{C} is simply connected, the result follows from Corollary 3.8.3. \square

3.9. Multiply Connected Domains

Definition 3.9.1 (Multiply Connected Domain). A domain G is called **multiply connected** if it is not simply connected.

We can generalise Theorem 3.8.2 to a multiply connected domain with finitely many holes.


Theorem 3.9.2 (Generalised Cauchy-Goursat Theorem). *Suppose that*

- (1) C is a simple closed positively oriented contour.
- (2) C_1, \dots, C_n are simple closed negatively oriented contours enclosing regions R_1, \dots, R_n . Further assume that the regions are pairwise disjoint and interior to C .

Lecture 25

If f is holomorphic on each contour and the region consisting of all points interior to C but exterior to each C_i , then

$$\int_C f(z) dz + \sum_{i=1}^n \int_{C_i} f(z) dz = 0$$



Proof. We prove this using induction.

Base Case. $n = 1$. Assume C and C_1 are contours satisfying the hypotheses. Let z_1, z_2 be points on C while w_1, w_2 be points on C_1 . Join z_1 to w_1 with a polygon line L_1 , and also join z_2 to w_2 with a polygon line L_2 .


Define contour Γ_1 and Γ_2 as follows.

Γ_1 : Start with z_1 and follow to w_1 along L_1 , then w_1 to w_2 along C_1 (we'll call this C_{11}), then w_2 to z_2 along L_2 , and finally z_2 to z_1 along C (we'll call this C'). So,

$$\Gamma_1 = L_1 + C_{11} + L_2 + C'$$

Γ_2 : Start with z_2 and follow to w_2 along $-L_2$, then w_2 to w_1 along C_1 (we'll call this C_{12}), then w_1 to z_1 along $-L_1$, and finally z_1 to z_2 along C (we'll call this C''). So,

$$\Gamma_2 = -L_2 + C_{12} - L_1 + C''$$



We note $C' + C'' = C$ and $C_{11} + C_{12} = C_1$.

Then f is holomorphic in the interior of and on the simple closed curves Γ_1 and Γ_2 , so by Theorem 3.7.3 we have

$$\int_{\Gamma_1} f(z) dz = \int_{\Gamma_2} f(z) dz = 0$$

So, this gives us

$$\begin{aligned} 0 &= \int_{\Gamma_1} f(z) dz + \int_{\Gamma_2} f(z) dz \\ &= \left(\int_{L_1} f(z) dz + \int_{C_{11}} f(z) dz + \int_{L_2} f(z) dz + \int_{C'} f(z) dz \right) \\ &\quad + \left(- \int_{L_2} f(z) dz + \int_{C_{12}} f(z) dz - \int_{L_1} f(z) dz + \int_{C''} f(z) dz \right) \\ &= \int_{C'} f(z) dz + \int_{C''} f(z) dz + \int_{C_{11}} f(z) dz + \int_{C_{12}} f(z) dz \\ &= \int_C f(z) dz + \int_{C_1} f(z) dz \end{aligned}$$

Inductive Step. Assume the statement holds for $n = k$, that is

$$\int_C f(z) dz + \sum_{i=1}^k \int_{C_i} f(z) dz = 0$$

for any k -many contours satisfying the hypotheses.

Now, let C_1, \dots, C_k, C_{k+1} be any $k+1$ -many contours. Introduce a polygon line L that separates C_1, \dots, C_k from C_{k+1} , say with end points z_1 and z_2 . We define Γ_1 and Γ_2 as follows.


Γ_1 : Start with z_1 and follow to z_2 along C (we'll call this C'), then z_2 to z_1 along $-L$. So,

$$\Gamma_1 = C' - L$$

Γ_2 : Start with z_1 and follow to z_2 along L , then z_2 to z_1 along C (we'll call this C''). So,

$$\Gamma_2 = C'' + L$$

We note $C' + C'' = C$.



We note that

$$\begin{aligned}
 \int_{\Gamma_1} f(z) dz + \int_{\Gamma_2} f(z) dz &= \left(\int_{C'} f(z) dz - \int_L f(z) dz \right) + \left(\int_{C''} f(z) dz + \int_L f(z) dz \right) \\
 &= \int_{C'} f(z) dz + \int_{C''} f(z) dz \\
 &= \int_C f(z) dz
 \end{aligned} \tag{†}$$

By the inductive hypothesis

$$\int_{\Gamma_1} f(z) dz + \sum_{i=1}^k \int_{C_i} f(z) dz = 0 \tag{1}$$

and by the computation in the base case we have


$$\int_{\Gamma_2} f(z) dz + \int_{C_{k+1}} f(z) dz = 0 \tag{2}$$

Adding (1) and (2) and using (†) we have

$$0 = \int_{\Gamma_1} f(z) dz + \sum_{i=1}^k \int_{C_i} f(z) dz + \int_{\Gamma_2} f(z) dz + \int_{C_{k+1}} f(z) dz = \int_C f(z) dz + \sum_{i=1}^{k+1} \int_{C_i} f(z) dz$$

Thus, we have proved our result using the principle of mathematical induction. \square

Corollary 3.9.3 (Principle of Deformation of Paths). *Suppose C_1 and C_2 are positively oriented simple closed contours with C_1 interior to C_2 .*



If f is holomorphic on the region consisting of C_1 and C_2 and all the points between them, then

$$\int_{C_1} f(z) dz = \int_{C_2} f(z) dz$$

Proof. Applying Theorem 3.9.2 to C_2 and $-C_1$, we get

$$\int_{C_2} f(z) dz + \int_{-C_1} f(z) dz = 0.$$

Therefore,

$$\int_{C_1} f(z) dz = \int_{C_2} f(z) dz \quad \square$$

Among other things, the principle of deformation of paths is useful for integrating over complicated contours. Often, we can just replace this contour with a circle.


Example 3.9.4. Let C be any simple closed contour whose interior contains 0. We show that

$$\int_C \frac{1}{z} dz = 2\pi i.$$

Since 0 is interior to C , we can choose an $\varepsilon > 0$ small enough such that $C_\varepsilon = C_\varepsilon(0)$ is contained in the interior of C . The region containing C and C_ε and points between them does not contain 0, so $1/z$ is holomorphic there. By Corollary 3.9.3,

$$\begin{aligned} \int_C f(z) dz &= \int_{C_\varepsilon} f(z) dz \\ &= \int_0^{2\pi} \frac{1}{\varepsilon e^{it}} i e^{it} dt = \int_0^{2\pi} i dt = 2\pi i \end{aligned}$$

Definition 3.9.5 (Singularities). Suppose f is not holomorphic at z_0 , but every neighbourhood of z_0 contains a point at which f is holomorphic, then z_0 is called a **singular point** (or **singularity**) of f .



Example 3.9.6.

- (1) $f(z) = \frac{1}{z}$ has a singularity at 0.
- (2) $f(z) = |z|^2$ has no singular points, as f is only differentiable at 0 but is nowhere holomorphic.
- (3) $f(z) = \frac{z^2 + 3}{(z + 1)(z^2 + 5)}$ has singularities at those z where

$$(z + 1)(z^2 + 5) = 0.$$

That is, at -1 , $i\sqrt{5}$ and $-i\sqrt{5}$.

Remark 3.9.7. More generally, the generalised Generalised Cauchy-Goursat Theorem (Theorem 3.9.2) and its Corollary 3.9.3 provide a technique for integrating functions over contours whose interior contains singularities of that function. The idea is to introduce small circles around the singular points, and apply the theorem (or corollary). It is usually easy to integrate over a circle.

3.10. Cauchy's Integral Formula

Discussion 3.10.1. Cauchy's Integral Formula is a remarkable theorem. It asserts that if a function is holomorphic inside and on C , a simple closed contour, then its values interior to C are completely determined by its values on C .

Theorem 3.10.2 (Cauchy's Integral Formula). *Let C be a simple closed contour, with positive orientation, and let f be a function that is holomorphic at all points on and interior to C . Then for any $z_0 \in \text{int}(C)$, we have*

$$f(z_0) = \frac{1}{2\pi i} \int_C \frac{f(z)}{z - z_0} dz$$

Proof. Our strategy is to show that for all $\varepsilon > 0$, we get

$$\left| \int_C \frac{f(z)}{z - z_0} dz - f(z_0) \cdot 2\pi i \right| < \varepsilon$$


because then

$$\int_C \frac{f(z)}{z - z_0} dz - f(z_0) \cdot 2\pi i = 0$$

Let $\varepsilon > 0$, and since, by assumption, f is holomorphic on C , it's continuous on C . So, there exists a $\delta > 0$ such that

$$\text{if } |z - z_0| < \delta, \text{ then } |f(z) - f(z_0)| < \frac{\varepsilon}{2\pi}$$

Let $\rho > 0$ be small enough such that the circle $C_\rho = C_\rho(z_0)$ centered at z_0 of radius ρ lies in the interior of C ; assume C_ρ has positive orientation.



We may assume $\rho < \delta$, then for every point $z \in C_\rho$, since $|z - z_0| = \rho < \delta$, we have

$$|f(z) - f(z_0)| < \frac{\varepsilon}{2\pi}, \text{ therefore } \max_{z \in C_\rho} |f(z) - f(z_0)| < \frac{\varepsilon}{2\pi}$$

Now, note that

$$\frac{f(z)}{z - z_0}$$

is holomorphic on the region consisting of C , C_ρ and all points that are interior to C but exterior to C_ρ . So, by Corollary 3.9.3, we have

$$\int_C \frac{f(z)}{z - z_0} dz = \int_{C_\rho} \frac{f(z)}{z - z_0} dz$$

and then

$$\begin{aligned}
\left| \int_C \frac{f(z)}{z - z_0} dz - f(z_0) \cdot 2\pi i \right| &= \left| \int_{C_\rho} \frac{f(z)}{z - z_0} - f(z_0) \cdot 2\pi i \right| \\
&= \left| \int_{C_\rho} \frac{f(z)}{z - z_0} - f(z_0) \int_{C_\rho} \frac{1}{z - z_0} dz \right| \\
&= \left| \int_{C_\rho} \frac{f(z) - f(z_0)}{z - z_0} dz \right| \\
&\leq \max_{z \in C_\rho} \left| \frac{f(z) - f(z_0)}{z - z_0} \right| \cdot L(C_\rho) \\
&= \max_{z \in C_\rho} \frac{|f(z) - f(z_0)|}{\rho} \cdot (2\pi\rho) \\
&= \frac{1}{\rho} \max_{z \in C_\rho} |f(z) - f(z_0)| \cdot (2\pi\rho) \\
&< \frac{\varepsilon}{2\pi} \cdot 2\pi \\
&= \varepsilon
\end{aligned}$$

and the claim follows. □

Among other things, Cauchy's Integral formula is useful for computing integrals.

Lecture 26

Example 3.10.3.

- (1) Let's compute $\int_C \frac{\cos z}{z(z^2 + 2)} dz$, where C is the unit circle, positively oriented.

Consider

$$f(z) = \frac{\cos z}{z^2 + 2}$$

Then f is holomorphic on all points on and interior to C , as they don't include $\pm 2i$ and 0 is in the interior of C . Therefore, by Cauchy's Integral Formula (Theorem 3.10.2) we have


$$\int_C \frac{\cos z}{z(z^2 + 2)} dz = \int_C \frac{f(z)}{z - 0} dz = 2\pi i \cdot f(0) = \pi i.$$

- (2) Let's compute $\int_C \frac{e^{z^2}}{z - 1} dz$, where C is a positively oriented circle with radius 2.


Consider $f(z) = e^{z^2}$, then f is entire, and therefore holomorphic on all points on and interior to C . Therefore, by Cauchy's Integral Formula (Theorem 3.10.2) we have

$$\int_C \frac{e^{z^2}}{z - 1} dz = 2\pi i f(1) = 2\pi i e.$$

(3) Let's compute $\int_C \frac{z^2 + 1}{z^2 - 1} dz = \int_C \frac{z^2 + 1}{(z - 1)(z + 1)} dz$, where C is as follows



The contour C is not simple but it can be decomposed as a sum of simple closed contours $C = C_1 - C_2$



So,

$$\int_C \frac{z^2 + 1}{(z - 1)(z + 1)} dz = \int_{C_1} \frac{z^2 + 1}{(z - 1)(z + 1)} dz - \int_{C_2} \frac{z^2 + 1}{(z - 1)(z + 1)} dz$$

For C_1 , consider

$$f(z) = \frac{z^2 + 1}{z + 1},$$

then f is holomorphic on all points on and interior to C_1 , as they don't include -1 , and 1 is in the interior of C_1 . Therefore by Cauchy's Integral Formula (Theorem 3.10.2) we have

$$\int_{C_1} \frac{z^2 + 1}{(z - 1)(z + 1)} dz = \int_{C_1} \frac{f(z)}{z - 1} dz = 2\pi i \cdot f(1) = 2\pi i$$

For C_2 , consider

$$g(z) = \frac{z^2 + 1}{z - 1},$$

then g is holomorphic on all points on and interior to C_2 , as they don't include 1, and -1 is in the interior of C_2 . Therefore by Cauchy's Integral Formula (Theorem 3.10.2) we have

$$\int_{C_2} \frac{z^2 + 1}{(z-1)(z+1)} dz = \int_{C_2} \frac{g(z)}{z+1} dz = 2\pi i \cdot g(-1) = -2\pi i$$

Hence,

$$\int_C \frac{z^2 + 1}{(z-1)(z+1)} dz = \int_{C_1} \frac{z^2 + 1}{(z-1)(z+1)} dz - \int_{C_2} \frac{z^2 + 1}{(z-1)(z+1)} dz = 2\pi i + 2\pi i = 4\pi i.$$

Theorem 3.10.4 (Generalised Cauchy's Integral Formula). *Let C be a simple closed contour, with positive orientation, and let f be a function that is holomorphic at all points on and interior to C . The for any $z_0 \in \text{int}(C)$, we have that $f^{(n)}(z_0)$ exists and*

$$f^{(n)}(z_0) = \frac{n!}{2\pi i} \int_C \frac{f(z)}{(z-z_0)^{n+1}} dz$$

Proof. We prove by induction, with the base case $n = 0$ being just Theorem 3.10.2. Assume the statements holds for $n = k$, we need to prove that

$$f^{(k+1)}(z_0) := \lim_{h \rightarrow 0} \frac{f^{(k)}(z_0 + h) - f^{(k)}(z_0)}{h} = \frac{(k+1)!}{2\pi i} \int_C \frac{f(z)}{(z-z_0)^{(k+1)+1}} dz$$

We assume $|h|$ is small enough such that $z + h \in \text{int}(C)$, then by the inductive hypothesis

$$f^{(k)}(z_0 + h) = \frac{k!}{2\pi i} \int_C \frac{f(z)}{(z - (z_0 + h))^{k+1}} dz$$

$$f^{(k)}(z_0) = \frac{k!}{2\pi i} \int_C \frac{f(z)}{(z - z_0)^{k+1}} dz$$

Recall the algebraic identity, that for $a, b \in \mathbb{C}$ we have

$$a^{k+1} - b^{k+1} = (a-b)(a^k + a^{k-1}b + \dots + ab^{k-1} + b^k),$$

We will apply this to $a = \frac{1}{z - z_0 - h}$ and $b = \frac{1}{z - z_0}$, and we also note $\lim_{h \rightarrow 0} a = b$. Then,

$$\begin{aligned} & \lim_{h \rightarrow 0} \frac{f^{(k)}(z_0 + h) - f^{(k)}(z_0)}{h} \\ &= \lim_{h \rightarrow 0} \frac{k!}{2\pi i} \int_C \frac{f(z)}{h} \left(\frac{1}{(z - z_0 - h)^{k+1}} - \frac{1}{(z - z_0)^{k+1}} \right) dz \\ &= \lim_{h \rightarrow 0} \frac{k!}{2\pi i} \int_C \frac{f(z)}{h} \left(\frac{1}{z - z_0 - h} - \frac{1}{z - z_0} \right) (a^k + a^{k-1}b + \dots + ab^{k-1} + b^k) dz \\ &= \lim_{h \rightarrow 0} \frac{k!}{2\pi i} \int_C \frac{f(z)}{h} \left(\frac{h}{(z - z_0 - h)(z - z_0)} \right) (a^k + a^{k-1}b + \dots + ab^{k-1} + b^k) dz \end{aligned}$$

$$\begin{aligned}
&= \lim_{h \rightarrow 0} \frac{k!}{2\pi i} \int_C \left(\frac{f(z)}{(z - z_0 - h)(z - z_0)} \right) (a^k + a^{k-1}b + \dots + ab^{k-1} + b^k) dz \\
&= \frac{k!}{2\pi i} \int_C \lim_{h \rightarrow 0} \left(\frac{f(z)}{(z - z_0 - h)(z - z_0)} \right) (a^k + a^{k-1}b + \dots + ab^{k-1} + b^k) dz \\
&= \frac{k!}{2\pi i} \int_C \left(\frac{f(z)}{(z - z_0)^2} \right) (b^k + b^{k-1}b + \dots + b \cdot b^{k-1} + b^k) dz \\
&= \frac{k!}{2\pi i} \int_C \frac{f(z)}{(z - z_0)^2} \cdot (k+1) \cdot b^k dz \\
&= \frac{(k+1)!}{2\pi i} \int_C \frac{f(z)}{(z - z_0)^2} \cdot \frac{1}{(z - z_0)^k} dz \\
&= \frac{(k+1)!}{2\pi i} \int_C \frac{f(z)}{(z - z_0)^{(k+1)+1}} dz
\end{aligned}$$

Thus we have our result by the principle of mathematical induction. \square

Example 3.10.5. Compute the integral

$$\frac{1}{2\pi i} \int_C \frac{(1+z)^n}{z^{k+1}} dz$$

where C is any simple closed positively oriented contour whose interior contains 0 and $0 \leq k \leq n$.

Let $f(z) = (1+z)^n$, since f is entire, f is holomorphic on all points on and interior to C . Since 0 is in the interior of C , then generalised Cauchy's Integral formula (Theorem 3.10.4) gives us

$$\frac{1}{2\pi i} \int_C \frac{(1+z)^n}{z^{k+1}} dz = \frac{1}{k!} \left(\frac{k!}{2\pi i} \int_C \frac{(1+z)^n}{(z-0)^{k+1}} dz \right) = \frac{1}{k!} \cdot f^{(k)}(0)$$

We have,

$$f^{(k)}(z) = n(n-1) \cdots (n-(k-1))(1+z)^{n-k},$$

and therefore

$$f^{(k)}(0) = n(n-1) \cdots (n-(k-1)) = \frac{n!}{(n-k)!}$$

Hence,

$$\frac{1}{2\pi i} \int_C \frac{(1+z)^n}{z^{k+1}} dz = \frac{1}{k!} \cdot f^{(k)}(0) = \frac{n!}{k!(n-k)!} = \binom{n}{k}$$

Theorem 3.10.6 (Derivatives of Holomorphic functions are Holomorphic). Suppose that f is holomorphic at $z_0 \in \mathbf{C}$, then for all $n \in \mathbf{Z}_{>0}$, $f^{(n)}$ is also holomorphic at z_0 .

Proof. Suppose f is holomorphic at $z_0 \in \mathbf{C}$. Choose an open disk $D_\varepsilon(z_0)$ on which f is differentiable. To conclude f' exists and is holomorphic at z_0 , it's enough to find a neighbourhood of z_0 where $f''(w)$ exists for all w in that neighbourhood. Let C be the positive oriented circle of radius

$\varepsilon/2$ centered at z_0 , then f is holomorphic on all points on and interior to C . So, by generalised Cauchy's Integral formula (Theorem 3.10.4),

$$f''(w) = \frac{2!}{2\pi i} \int_C \frac{f(z)}{(z-w)^3} dz$$

for any w in the interior of C . Thus, f' is differentiable in the open set $D_{\varepsilon/2}(z_0)$, and hence f' is holomorphic at z_0 . Induction then gives us that $f^{(n)}$ is holomorphic at z_0 for any $n \in \mathbf{Z}_{>0}$. \square

Corollary 3.10.7. *If $f(z) = u(x, y) + iv(x, y)$ is holomorphic at $z = x + iy$, then u and v have continuous partial derivatives of all orders at (x, y) .*

Lecture 27 **Theorem 3.10.8** (Morera's Theorem). *Suppose f is continuous on a domain G . If*

$$\int_C f(z) dz = 0$$

for every closed contour $C \subseteq G$, then f is holomorphic on G .

Proof. By Theorem 3.6.4, there exists a holomorphic function $F : G \rightarrow \mathbf{C}$ such that $F'(z) = f(z)$ for all $z \in G$. But by Theorem 3.10.6, F' is holomorphic on G , and therefore so is f . \square

Remark 3.10.9. When G is simply connected, Morera's theorem (Theorem 3.10.8) is just the converse of Cauchy-Goursat Theorem for simply connected domains (Theorem 3.8.2).

Theorem 3.10.10 (Cauchy's Inequalities). *Suppose that f is holomorphic on all points on and interior to $C_R = C_R(z_0)$, a positively oriented circle of radius R centered at some $z_0 \in \mathbf{C}$. Then,*

$$|f^{(n)}(z_0)| \leq \frac{n!}{R^n} \max_{z \in C_R(z_0)} |f(z)|$$

Proof. By Theorem 3.10.4,

$$f^{(n)}(z_0) = \frac{n!}{2\pi i} \int_{C_R} \frac{f(z)}{(z-z_0)^{n+1}} dz$$

Hence,

$$\begin{aligned} |f^{(n)}(z_0)| &= \left| \frac{n!}{2\pi i} \int_{C_R} \frac{f(z)}{(z-z_0)^{n+1}} dz \right| = \frac{n!}{2\pi i} \left| \int_{C_R} \frac{f(z)}{(z-z_0)^{n+1}} dz \right| \\ &\leq \frac{n!}{2\pi i} \max_{z \in C_R} \left| \frac{f(z)}{(z-z_0)^{n+1}} \right| \cdot L(C_R) \\ &= \frac{n!}{2\pi i} \max_{z \in C_R} \frac{|f(z)|}{R^{n+1}} \cdot 2\pi R \\ &= \frac{n!}{R^n} \max_{z \in C_R} |f(z)| \end{aligned}$$

\square

3.11. Liouville's Theorem and the Fundamental Theorem of Algebra

As an application, we will prove that every non-constant polynomial with complex coefficients has a root in \mathbf{C} . In the language of algebra, we will provide a proof for the fact that \mathbf{C} is *algebraically closed*. Thus, the statement is "purely algebraic" while no "purely algebraic" proof exists. The proof relies on the following wonderful theorem.

Theorem 3.11.1 (Liouville's Theorem). *Every bounded entire function is constant.*

Proof. We show that $f'(z) = 0$ for all $z \in \mathbf{C}$, then it follows that f is constant since \mathbf{C} is a domain by Theorem 2.7.4.

Consider any $z_0 \in \mathbf{C}$. Since f is bounded, we can find a $M > 0$ such that $|f(z)| \leq M$ for all $z \in \mathbf{C}$. Let $C_R(z_0)$ be a circle of radius R centered at z_0 , then f is holomorphic at all points on and interior to $C_R(z_0)$. Hence, by Theorem 3.10.10,

$$\begin{aligned} |f'(z_0)| &\leq \frac{1}{R} \max_{z \in C_R(z_0)} |f(z)| \\ &\leq \frac{M}{R} \rightarrow 0, \text{ as } R \rightarrow \infty \end{aligned}$$

Thus $|f'(z_0)| = 0$, giving us $f'(z_0) = 0$. Since z_0 was arbitrary, the result follows. \square

Theorem 3.11.2 (Fundamental Theorem of Algebra). *For any polynomial $p(z) = a_0 + a_1z + \cdots + a_nz^n$ where $a_n \neq 0$, $a_i \in \mathbf{C}$ and $n \geq 1$, there exists an $\alpha \in \mathbf{C}$ such that $p(\alpha) = 0$. That is, every non-constant polynomial $p(z)$ has at least one root in \mathbf{C} .*

Proof. Suppose otherwise that $p(z)$ has no root in \mathbf{C} , then $p(z) \neq 0$ for every $z \in \mathbf{C}$. Hence $1/p(z)$ is an entire function. We show that $1/p(z)$ is bounded.

For a non-zero $z \in \mathbf{C}$, consider the complex number

$$w_z := \frac{a_0}{z^n} + \frac{a_1}{z^{n-1}} + \cdots + \frac{a_{n-1}}{z}$$

Note that $p(z) = (w_z + a_n) z^n$, and by triangle inequality we have

$$|w_z| \leq \frac{|a_0|}{|z|^n} + \frac{|a_1|}{|z|^{n-1}} + \cdots + \frac{|a_{n-1}|}{|z|} = \sum_{k=0}^{n-1} \frac{|a_k|}{|z|^{n-k}}$$

For each $0 \leq k \leq n-1$ we note that $\frac{|a_k|}{|z|^{n-k}} \rightarrow 0$ as $z \rightarrow \infty$.

Then, for $\varepsilon = \frac{|a_n|}{2n} > 0$, we can find an $R > 0$ such that whenever $|z| > R$, we get

$$\frac{|a_k|}{|z|^{n-k}} = \left| \frac{|a_k|}{|z|^{n-k}} - 0 \right| < \varepsilon = \frac{|a_n|}{2n}$$

for any $k = 0, \dots, n-1$. This then gives us

$$|w_z| \leq \sum_{k=0}^{n-1} \frac{|a_k|}{|z|^{n-k}} < \sum_{k=0}^{n-1} \frac{|a_n|}{2n} = n \cdot \frac{|a_n|}{2n} = \frac{|a_n|}{2}$$

Now, by the reverse triangle inequality we have

$$|a_n + w_z| \geq ||a_n| - |w_z|| > \left| |a_n| - \frac{|a_n|}{2} \right| = \frac{|a_n|}{2}$$

Thus,

$$\begin{aligned} |p(z)| &= |(w_z + a_n) z^n| \\ &= |w_z + a_n| |z^n| > \frac{|a_n|}{2} R^n \end{aligned}$$

Therefore, for any $z \in \mathbf{C}$ such that $|z| > R$, we have

$$\left| \frac{1}{p(z)} \right| \leq \frac{2}{R^n |a_n|}$$


So, $1/p(z)$ is bounded outside the closed disk $\overline{D}_R(0)$.

Now, the closed disk $\overline{D}_R(0)$ is compact (closed and bounded) and $1/p(z)$ is continuous on $\overline{D}_R(0)$. Hence $1/p(z)$ is bounded on $\overline{D}_R(0)$ by Theorem 2.4.7.

Thus, $1/p(z)$ is bounded on all of \mathbf{C} . Hence, by Theorem 3.11.1, $1/p(z)$ is constant, and therefore so is $p(z)$. We have arrived a contradiction, since $p(z)$ was non-constant by assumption. \square

Lemma 3.11.3 (Maximum Modulus Principle). *Suppose that $|f(z)| \leq |f(z_0)|$ at each point z in a neighbourhood $D_\varepsilon(z_0)$ where f is holomorphic. Then $f(z) = f(z_0)$ on $D_\varepsilon(z_0)$. That is, if a holomorphic function on an open disk achieves its maximum on it, then it is constant on the open disk.*

Proof. Let $z_1 \in D_\varepsilon(z_0)$ such that $z_1 \neq z_0$. Set $\rho := |z_1 - z_0| > 0$, and consider $C_\rho = C_\rho(z_0)$, the circle of radius $\rho > 0$ centered at z_0 , which is interior to $D_\varepsilon(z_0)$. We parametrise C_ρ as $z(t) = z_0 + \rho e^{it}$ for $0 \leq t \leq 2\pi$.



By Theorem 3.10.2,

$$\begin{aligned}
|f(z_0)| &= \left| \frac{1}{2\pi i} \int_{C_\rho} \frac{f(z)}{z - z_0} dz \right| = \frac{1}{2\pi} \left| \int_{C_\rho} \frac{f(z)}{z - z_0} dz \right| \\
&= \frac{1}{2\pi} \left| \int_0^{2\pi} \frac{f(z_0 + \rho e^{it})}{\rho e^{it}} i \rho e^{it} dt \right| \\
&= \frac{1}{2\pi} \left| \int_0^{2\pi} f(z_0 + \rho e^{it}) dt \right| \\
&\leq \frac{1}{2\pi} \int_0^{2\pi} |f(z_0 + \rho e^{it})| dt \\
&\leq \frac{1}{2\pi} \int_0^{2\pi} |f(z_0)| dt, \quad \text{by assumption} \\
&\leq |f(z_0)|
\end{aligned}$$

This tells us that

$$|f(z_0)| = \frac{1}{2\pi} \int_0^{2\pi} |f(z_0 + \rho e^{it})| dt \quad (\dagger)$$

Since, $|f(z_0)| = \frac{1}{2\pi} \int_0^{2\pi} |f(z_0)| dt$. Rewriting (\dagger) , we have

$$\frac{1}{2\pi} \int_0^{2\pi} |f(z_0)| - |f(z_0 + \rho e^{it})| dt = 0$$

By assumption $|f(z_0)| - |f(z_0 + \rho e^{it})| \geq 0$; suppose $|f(z_0)| - |f(z_0 + \rho e^{it})| > 0$, then necessarily

$$\frac{1}{2\pi} \int_0^{2\pi} |f(z_0)| - |f(z_0 + \rho e^{it})| dt > 0 \quad (*)$$

since the integrand in $(*)$ is continuous in the variable t , giving us a contradiction. Thus,

$$|f(z_0)| - |f(z_0 + \rho e^{it})| = 0$$

Therefore $|f(z)| = |f(z_0)|$ for every $z \in C_\rho(z_0)$. Varying the radius $\rho > 0$, we may then obtain $|f(z)| = |f(z_0)|$ for every $z \in D_\varepsilon(z_0)$.

Thus, $|f|$ is a holomorphic function on $D_\varepsilon(z_0)$, and thus by Corollary 2.7.7, we have f is constant on $D_\varepsilon(z_0)$ and $f(z) = f(z_0)$ for every $z \in D_\varepsilon(z_0)$. \square

A. Problems

Problem 1. Consider the set of matrices

$$X := \left\{ \begin{pmatrix} x & -y \\ y & x \end{pmatrix} : x, y \in \mathbf{R} \right\}.$$

One can check (and you should if you're unconvinced) straightforwardly that X is closed under matrix addition and matrix multiplication; that is, if $A, B \in X$, then $A + B, AB \in X$.

(a) Let \mathbf{C} denote the set of complex numbers. Show that the map $\phi : X \rightarrow \mathbf{C}$ defined by

$$\phi : X \rightarrow \mathbf{C}, \quad \begin{pmatrix} x & -y \\ y & x \end{pmatrix} \mapsto x + iy$$

is a bijection.

(b) Let $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ be the identity matrix. Consider $A, B \in X$, show that ϕ has the following properties.

(i) $\phi(A + B) = \phi(A) + \phi(B)$

(ii) $\phi(AB) = \phi(A)\phi(B)$

(iii) $\phi(I) = 1$

(c) Find a matrix J satisfying $J^2 = -I$ and show that $\phi(J) = i$.

Remark A.0.1. This indicates that one could very well define \mathbf{C} to be X . The algebraic operations on \mathbf{C} then seem less artificial, since product and sum of complex numbers correspond to the corresponding operations of matrices. Even taking the inverse and modulus is captured by X as taking inverse and the determinant of matrices. The copy of \mathbf{R} corresponds to the set of diagonal matrices in X . One obtains X by considering the linear operator of multiplying by $x + iy$ on the \mathbf{R} -vector space \mathbf{C} with basis 1 and i .

Problem 2. Using the definition of complex multiplication prove that

$$(a, 0) \cdot (x, y) = (ax, ay).$$

That is, $a(x + iy) = ax + iay$.

Problem 3. Consider complex numbers $z_1 = (x_1, y_1) = x_1(1, 0) + y_1(0, 1)$ and $z_2 = (x_2, y_2) = x_2(1, 0) + y_2(0, 1)$. Using the identity $(0, 1)^2 = (-1, 0)$. Prove that

$$(x_1(1, 0) + y_1(0, 1)) \cdot (x_2(1, 0) + y_2(0, 1)) = (x_1x_2 - y_1y_2, x_1y_2 + x_2y_1),$$

where the former is computed distributively.

Problem 4. Prove properties (1) - (7) and (9) listed in Proposition 1.1.6.

Problem 5. Prove that if $z_1 z_2 = 0$, then $z_1 = 0$ or $z_2 = 0$.

Problem 6. Show that

(a) $\operatorname{Re} iz = -\operatorname{Im} z$;

(b) $\operatorname{Im} iz = \operatorname{Re} z$

Problem 7.

(a) Verify that $z = 1 \pm i$ satisfies the equation

$$z^2 - 2z + 2 = 0.$$

(b) Solve the equation

$$z^2 + z + 1 = 0$$

for $z = x + iy$ by solving a pair of simultaneous equations in x and y .

Problem 8. Let $p(z) = az^2 + bz + c$ be a polynomial with complex coefficients ($a \neq 0$).

(a) By completing the square, show that the solution to $p(z) = 0$ is

$$z = \frac{-b \pm \Delta^{1/2}}{2a},$$

where $\Delta := b^2 - 4ac$ is called the discriminant.

Remark. There's a subtlety with taking roots that we will address later in class.

(b) Consider the polynomial $p(z) = iz^2 - 1$

(i) Compute Δ .

(ii) For the Δ obtained in (b), compute $\Delta^{1/2}$ by solving a pair of simultaneous equations in x and y obtained by considering the equation

$$x^2 - y^2 + 2ixy = (x + iy)^2 = \Delta.$$

(iii) Finally, write down the roots of $p(z)$ in the form $u + iv$.

Problem 9. Suppose \mathbf{C} had total ordering that extends the ordering on \mathbf{R} , arrive at a contradiction by comparing i and 0 .

Problem 10. Locate the numbers $z_1 + z_2$, $z_1 - z_2$ and $z_1 z_2$ in the complex plane when

$$(a) z_1 = 2i, z_2 = \frac{2}{3} - i.$$

$$(c) z_1 = (-\sqrt{3}, 1), z_2 = (\sqrt{3}, 0).$$

$$(b) z_1 = (-3, 1), z_2 = (1, 4).$$

$$(d) z_1 = x_1 + iy_1, z_2 = x_1 - iy_1.$$

Problem 11. Verify that $\sqrt{2}|z| \geq |\operatorname{Re} z| + |\operatorname{Im} z|$.

Problem 12. Let $z_0 \neq z_1 \in \mathbf{C}$ and let $\lambda > 0$.

(a) Show that if $\lambda \neq 1$, then the set of points

$$|z - z_0| = \lambda |z - z_1| \quad (\star)$$

is a circle of radius $R = \frac{\lambda}{|1 - \lambda^2|} |z_0 - z_1|$ centered at $w = \frac{z_0 - \lambda^2 z_1}{1 - \lambda^2}$.

(b) Show that every circle in the complex plane can be written in the form of (\star) for some $\lambda > 0$, $\lambda \neq 1$ and $z_0 \neq z_1 \in \mathbf{C}$.

(c) If $\lambda = 1$, show that (\star) defines a line. In fact, argue that the resulting line is perpendicular to and bisects the line segment joining z_0 and z_1 , by producing the equation of this line as a subset of \mathbf{R}^2 .

(d) Characterise points on the real (resp. imaginary) axis using (c). That is, find $z_0 \neq z_1 \in \mathbf{C}$ such that the points on the real (resp. imaginary) axis satisfy (\star) for $\lambda = 1$.

(e) Consider the map

$$M(z) = \frac{z - 3}{1 - 2z}.$$

For which values of $c \in \mathbf{R}$ is the image of the circle $|z - 1| = c$ under M a line? What is the equation of the line when considered as a subset of the plane \mathbf{R}^2 ?

Problem 13. Prove Proposition 1.2.5 (1).

Problem 14. Prove the properties, other than (5), listed in Proposition 1.2.7.

Problem 15. Prove that z is either real or pure imaginary if and only if $z^2 = \bar{z}^2$.

Problem 16. Prove that $|z| = 1$ if and only if $\bar{z} = \frac{1}{z}$.

Problem 17. Follow the steps below to give an algebraic derivation of the triangle inequality (Proposition 1.2.4 (a))

(a) Show that

$$|z_1 + z_2|^2 = (z_1 + z_2)(\bar{z}_1 + \bar{z}_2) = z_1\bar{z}_1 + (z_1\bar{z}_2 + \overline{z_1\bar{z}_2}) + z_2\bar{z}_2.$$

(b) Argue why

$$z_1\bar{z}_2 + \overline{z_1\bar{z}_2} = 2\operatorname{Re}(z_1\bar{z}_2) \leq 2|z_1||z_2|.$$

(c) Use (a) and (b) to obtain $|z_1 + z_2|^2 \leq (|z_1| + |z_2|)^2$. Finally note how the triangle inequality follows from this.

Problem 18. Let $z, w \in \mathbf{C}$.

(a) Prove the formula

$$|z + w|^2 = |z|^2 + 2\operatorname{Re} z\bar{w} + |w|^2$$

(b) Use (a) to deduce the *parallelogram law*

$$|z + w|^2 + |z - w|^2 = 2|z|^2 + 2|w|^2$$

Give a geometric interpretation of this formula.

Problem 19. Suppose p is a polynomial with *real coefficients*. Prove that

(a) $\overline{p(z)} = p(\bar{z})$.

(b) $p(z) = 0$ if and only if $p(\bar{z}) = 0$.

Problem 20. Find the principal argument $\operatorname{Arg} z$ when

(a) $-i(3 + 3i)^{-1}$.

(b) $(1 - i\sqrt{3})^6$.

Problem 21. Prove that

$$\arg z + \arg w = \{(\operatorname{Arg} z + \operatorname{Arg} w) + 2k\pi : k \in \mathbf{Z}\}$$

Combining this with Proposition 1.2.18 we get that $\operatorname{Arg} zw = \operatorname{Arg} z + \operatorname{Arg} w + 2k\pi$ for some $k \in \mathbf{Z}$ such that $-\pi < \operatorname{Arg} z + \operatorname{Arg} w + 2k\pi \leq \pi$. That is, to find $\operatorname{Arg} zw$, just add $\operatorname{Arg} z$ and $\operatorname{Arg} w$ and then add or subtract a suitable multiple of 2π to get it between $-\pi$ and π .

Problem 22. Prove that for any complex number z , we have $\operatorname{Arg} \bar{z} = \operatorname{Arg} z^{-1} = -\operatorname{Arg} z$.

Problem 23.

(a) Show that if $\operatorname{Re} z_1 > 0$ and $\operatorname{Re} z_2 > 0$, then $\operatorname{Arg}(z_1 z_2) = \operatorname{Arg} z_1 + \operatorname{Arg} z_2$.

(b) Show that if $\operatorname{Re} z > 0$, then $\operatorname{Arg}(-z) = -\pi + \operatorname{Arg} z$ if $\operatorname{Im} z > 0$ or $\operatorname{Arg}(-z) = \pi + \operatorname{Arg} z$ if $\operatorname{Im} z < 0$.

- (c) Using (a) and (b), find an expression for $\text{Arg } zw$ for any non-zero complex numbers z and w , in terms of $\text{Arg } z$, $\text{Arg } w$ and specific multiples of π .

Problem 24. Compute the 6th roots of unity, explicitly. Show that the principal 6th root of unity is $\zeta_6 = -\omega$, where ω is as in Example 1.3.6.

Problem 25.

- (a) Let $z \in \mathbf{C}$. Using the principle of mathematical induction, show that the following formula holds for all integers $n \geq 1$

$$1 + z + z^2 + \cdots + z^n = \frac{1 - z^{n+1}}{1 - z}.$$

- (b) Use (a) to derive *Lagrange's Trigonometric Identity*.

$$1 + \cos \theta + \cos^2 \theta + \cdots + \cos^n \theta = \frac{2 \sin((2n+1)\theta/2)}{2 \sin(\theta/2)}, \quad 0 < \theta < 2\pi.$$

- (c) If ζ_1, \dots, ζ_n are the *distinct* n^{th} roots of unity, show that, using (a), $\sum_{i=1}^n \zeta_i = 0$.

- (d) We compute the following sum of real numbers

$$\cos \frac{\pi}{7} + \cos \frac{3\pi}{7} + \cos \frac{5\pi}{7} \tag{†}$$

- (i) Let $w = e^{\frac{\pi i}{7}}$. What is $\text{Re } w$ and w^7 ? Furthermore, rewrite (†) as

$$\text{Re}(w^{a_1} + w^{a_2} + w^{a_3}), \quad \text{for some } 0 \leq a_i < 7.$$

- (ii) Replacing z by $-z$ in (a), find a formula for

$$\frac{z^7 + 1}{z + 1}.$$

Use this to deduce an identity involving w and its powers.

- (iii) Using the identity you found in (iii), conclude that

$$w^{a_1} + w^{a_2} + w^{a_3} = \frac{1}{1 - w}$$

where the a_i 's are the numbers you found in (ii).

- (iv) Finally compute (†).

Problem 26.

- (a) Recall that a set is open if every point of the set is an interior point. Prove that a set $U \subseteq \mathbf{C}$ is open if and only if it does not contain any of its boundary points; that is, $\partial U \cap U = \emptyset$. Then deduce that the complement of a closed set is open.
- (b) Prove that an open disk $D_\varepsilon(z_0) = \{z \in \mathbf{C} : |z - z_0| < \varepsilon\}$ is a domain; that is, a non-empty open and connected subset of \mathbf{C} .

Problem 27. Sketch the sets defined by the following constraints and determine whether they are open, closed, or neither; bounded; connected. What are their boundaries?

- (a) $|z + 3| < 2$.
 (b) $|\operatorname{Im}(z)| < 1$.
 (c) $0 < |z - 1| < 2$.
 (d) $|z - 1| + |z + 1| = 2$.
 (e) $|z - 1| + |z + 1| < 3$.
 (f) $|z| \geq \operatorname{Re}(z) + 1$.

Problem 28. Let G be the set of points $z \in \mathbf{C}$ satisfying either z is real and $-2 < z < -1$, or $|z| < 1$, or $z = 1$ or $z = 2$.

- (a) Sketch the set G , being careful to indicate exactly the points that are in G .
 (b) Determine the interior points of G .
 (c) Determine the boundary points of G .
 (d) Determine the isolated points of G .
 (e) G can be written in three different ways as the union of two disjoint nonempty disconnected subsets. Describe them.

Problem 29. For each of the functions below, describe the domain of definition that is understood.

- (a) $f(z) = \frac{1}{1 + z^2}$
 (b) $f(z) = \operatorname{Arg}\left(\frac{1}{z}\right)$
 (c) $f(z) = \frac{z}{z + \bar{z}}$
 (d) $f(z) = \frac{1}{1 - |z|^2}$

Problem 30.

- (a) Write the function $f(z) = z^3 + z + \bar{z} + 1$ in the form

$$f(z) = u(x, y) + i v(x, y).$$

- (b) Suppose that $f(z) = x^2 - y^2 - 2y + i(2x - 2xy)$, where $z = x + iy$. Use Proposition 1.2.7 (6) to write $f(z)$ in terms of z , and simplify the result.

- (c) Write the function

$$f(z) = z + \frac{1}{z} \quad (z \neq 0)$$

in the form $f(z) = u(r, \theta) + i v(r, \theta)$.

Problem 31. Let $f : G \rightarrow \mathbf{C}$ be a complex function, and suppose z_0 is an accumulation point of G . Show that

$$\lim_{z \rightarrow z_0} f(z) = w_0 \quad \text{if and only if} \quad \lim_{z \rightarrow z_0} |f(z) - w_0| = 0.$$

Thereby deduce that

$$\lim_{z \rightarrow z_0} \bar{f}(z) = \bar{w}_0 \quad \text{if and only if} \quad \lim_{z \rightarrow z_0} f(z) = w_0.$$

Problem 32. Let $f : G \rightarrow \mathbf{C}$ be a complex function, and suppose z_0 is an accumulation point of G . Show that

$$\text{if } \lim_{z \rightarrow z_0} f(z) = w_0, \quad \text{then } \lim_{z \rightarrow z_0} |f(z)| = |w_0|.$$

Hint. Use the reverse triangle inequality.

Problem 33. Let $f : G \rightarrow \mathbf{C}$ be a complex function, and suppose z_0 is an accumulation point of G . Writing $h = z - z_0$, show that

$$\lim_{z \rightarrow z_0} f(z) = w_0 \quad \text{if and only if} \quad \lim_{h \rightarrow 0} f(z_0 + h) = w_0.$$

Problem 34. Compute the following limits and prove your claim by using only the ε - δ definition.

(a) $\lim_{z \rightarrow i} \bar{z}$

(d) $\lim_{z \rightarrow 1-i} \bar{z}^2 - 1$

(b) $\lim_{z \rightarrow 1+i} z^2$

(e) $\lim_{z \rightarrow 1} z - \bar{z}$

(c) $\lim_{z \rightarrow 1} z^3$

(f) $\lim_{z \rightarrow i} \bar{z} + z$

Problem 35. Evaluate the following limits or explain why they don't exist.

(a) $\lim_{z \rightarrow i} \frac{iz^3 - 1}{z + i}$

(b) $\lim_{z \rightarrow 1-i} (x + i(2x + y))$

Problem 36. Define

$$f(z) = \frac{x^2 y}{x^4 + y^2} \quad \text{where } z = x + iy \neq 0.$$

Show that the limits of f at 0 along all straight lines through the origin exist and are equal, but $\lim_{z \rightarrow 0} f(z)$ does not exist.

Hint: Consider the limit along the parabola $y = x^2$.

Problem 37. Let $M(z) = \frac{z-3}{1-2z}$. Prove that

$$\lim_{z \rightarrow \infty} M(z) = -\frac{1}{2} \quad \text{and} \quad \lim_{z \rightarrow 1/2} M(z) = \infty$$

Problem 38. Let

$$M(z) = \frac{az + b}{cz + d}, \quad ad - bc \neq 0.$$

Prove that

(a) $\lim_{z \rightarrow \infty} M(z) = \infty$ if $c = 0$.

(b) $\lim_{z \rightarrow \infty} M(z) = \frac{a}{c}$ and $\lim_{z \rightarrow -d/c} M(z) = \infty$, if $c \neq 0$.

Problem 39. Example 2.3.3 tells us that polynomials are continuous.

- (a) Prove that the complex conjugation function $\sigma(z) := \bar{z}$ is continuous.
 (b) Prove that a polynomial in \bar{z} is continuous. That is, prove that a polynomial given as

$$p(\bar{z}) = a_n \bar{z}^n + \cdots + a_1 \bar{z} + a_0, \quad a_i \in \mathbb{C}, a_n \neq 0$$

is continuous.

- (c) Prove that the following functions are continuous by writing them as a sum or product of polynomials $p(z)$ and $q(\bar{z})$
- (i) $R(z) := \operatorname{Re} z$
 - (ii) $I(z) := \operatorname{Im} z$
 - (iii) $N(z) := |z|^2$

Problem 40. Show that the function $f : \mathbb{C} \rightarrow \mathbb{C}$ given by

$$f(z) = \begin{cases} \frac{\bar{z}}{z} & \text{if } z \neq 0 \\ 1 & \text{if } z = 0 \end{cases}$$

is continuous on \mathbb{C}^* .

Problem 41. Consider the function

$$f : \mathbb{C}^* \rightarrow \mathbb{C}, z \mapsto \frac{1}{z}.$$

Apply the definition of the derivative to give a direct proof that $f'(z) = -\frac{1}{z^2}$.

Problem 42. Find the derivative of the function

$$M(z) := \frac{az + b}{cz + d}, \quad ad - bc \neq 0.$$

When is $M'(z) = 0$?

Problem 43. Using Example 2.2.5 as an inspiration, show that $f'(z)$ does not exist for any z for the functions

- (a) $f(z) = \operatorname{Re} z$
- (b) $f(z) = \operatorname{Im} z$

Problem 44. Show that the function $f : \mathbf{C} \rightarrow \mathbf{C}$ given by

$$f(z) = \begin{cases} \frac{\bar{z}^2}{z} & \text{if } z \neq 0 \\ 0 & \text{if } z = 0 \end{cases}$$

is not differentiable at 0.

Problem 45.

- (a) Show that a polynomial of degree n , $p(z) = a_0 + a_1z + a_2z^2 + \cdots + a_nz^n$, where $a_n \neq 0$, is differentiable everywhere, with

$$p'(z) = a_1 + 2a_2z + \cdots + na_nz^{n-1}$$

- (b) Furthermore, show that for $p(z)$, as given in (a), we have

$$a_i = \frac{p^{(i)}(0)}{i!}$$

for $i = 0, \dots, n$. Where $p^{(0)}(z) = p(z)$ and $p^{(i)}(z)$, for $i > 0$, is the i^{th} derivative of $p(z)$.

Problem 46. Let G be a domain and $f : G \rightarrow \mathbf{C}$ a function that is differentiable at every point in G . Consider the domain

$$G^* = \{z \in \mathbf{C} : \bar{z} \in G\}$$

and the function

$$f^* : G^* \rightarrow \mathbf{C}, z \mapsto \overline{f(\bar{z})}$$

Show that f^* is differentiable at every point in G^* .

Problem 47. For each function, determine all points at which the derivative exists. When the derivative exists, find its value. Use Example 2.5.3 as an inspiration.

- (a) $f(z) = z + i\bar{z}$
- (b) $g(z) = (z + i\bar{z})^2$
- (b) $h(z) = z \operatorname{Im} z$

Problem 48. By definition, a function $f : G \rightarrow \mathbf{C}$ is differentiable at $z_0 \in G$ if the limit

$$f'(z_0) = \lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}$$

exists. Unpacking the limit definition, we see that f is differentiable at z_0 if and only if for every $\varepsilon > 0$, there exists a $\delta > 0$ such that

$$\text{if } 0 < |z - z_0| < \delta, \quad \text{then} \quad \left| \frac{f(z) - f(z_0)}{z - z_0} - f'(z_0) \right| < \varepsilon.$$

By appealing only to the definition, we show that $\sigma : \mathbf{C} \rightarrow \mathbf{C}$ defined by $\sigma(z) = \bar{z}$ is not differentiable anywhere by completing the following steps.

- (i) Let $z_0 \in \mathbf{C}$ and assume that $f'(z_0)$ exists. Choose $\delta > 0$ according to the definition using $\varepsilon = 1/2$ and write down the resulting statement.
- (ii) Consider $z = z_0 + \delta/2$ and conclude from (a) that $|1 - f'(z_0)| < \varepsilon$.
- (iii) Consider $z = z_0 + i\delta/2$ and conclude from (a) that $|1 + f'(z_0)| < \varepsilon$.
- (iv) Using the triangle inequality together with (ii) and (iii), obtain a contradiction.

Problem 49. Define

$$f(z) = \begin{cases} 0 & \text{if } \operatorname{Re}(z) \cdot \operatorname{Im}(z) = 0, \\ 1 & \text{if } \operatorname{Re}(z) \cdot \operatorname{Im}(z) \neq 0 \end{cases}.$$

Show that f satisfies the Cauchy–Riemann equation at $z = 0$, yet f is not differentiable at $z = 0$.

Problem 50. Show that when $f(z) = x^3 + i(1 - y)^3$, it makes sense to write

$$f'(z) = u_x + iv_x = 3x^2$$

only when $z = i$.

Problem 51. Show that $f'(z)$ does not exist at any point if

- (a) $f(z) = z - \bar{z}$
- (b) $f(z) = 2x + ixy^2$

Problem 52. Show that $f'(z)$ and its derivative $f''(z)$ exist everywhere, and find $f''(z)$ when

- (a) $f(z) = iz + 2$
- (b) $f(z) = e^{-x}e^{-iy}$

Problem 53. Let $f : G \rightarrow \mathbf{C}$ be a function, such that $G \subseteq \mathbf{C}^*$, then we can write

$$f(z) = f(x + iy) = u(x, y) + i v(x, y) \quad \text{or} \quad f(z) = f(re^{i\theta}) = u(r, \theta) + i v(r, \theta)$$

Using the fact that $x = r \cos \theta$ and $y = r \sin \theta$ and the chain rule from calculus, write u_r and u_θ in terms of u_x and u_y . Assuming f is differentiable, rewrite the CR-equations and $f'(z)$ in terms of u_r and u_θ .

Problem 54. Prove that the function

$$f(z) = e^{-\theta} \cos(\ln r) + ie^{-\theta} \sin(\ln r)$$

is differentiable when $r > 0$ and $0 < \theta < 2\pi$, and find $f'(z)$ in terms of $f(z)$.

Problem 55. Let $f = u + iv$ be a complex-valued function defined on an open set $G \subseteq \mathbf{C}$. Suppose that the first-order partial derivatives of $\operatorname{Re} f = u$ and $\operatorname{Im} f = v$ exist and are continuous on G .

(a) Recall that if $z = x + iy$, then

$$x = \frac{z + \bar{z}}{2} \quad \text{and} \quad y = \frac{z - \bar{z}}{2i}$$

Treat $f = f(x, y)$ as a function in two real-variables, and *formally* apply the chain rule in Calculus to obtain the expressions

$$\frac{\partial f}{\partial z} = \frac{1}{2} \left(\frac{\partial f}{\partial x} - i \frac{\partial f}{\partial y} \right) \quad \text{and} \quad \frac{\partial f}{\partial \bar{z}} = \frac{1}{2} \left(\frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right)$$

(b) Define $\frac{\partial f}{\partial x} := \frac{\partial u}{\partial x} + i \frac{\partial v}{\partial x}$, and similarly for $\frac{\partial f}{\partial y}$.

Prove that f is holomorphic on G if and only if $\frac{\partial f}{\partial \bar{z}} = 0$.

(c) (i) If f is holomorphic on G , prove that $f'(z) = \frac{\partial f}{\partial z}$.

(ii) The *Jacobian* of $(x, y) \mapsto (u(x, y), v(x, y))$ is the determinant of the matrix

$$\begin{pmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{pmatrix}$$

If f is holomorphic on G , prove that the Jacobian equals $|f'(z)|^2 \geq 0$.

Problem 56. Suppose f is entire and can be written as

$$f(z) = u(x) + i v(y),$$

that is, the real part of f depends only on $x = \operatorname{Re}(z)$ and the imaginary part of f depends only on $y = \operatorname{Im}(z)$.

Prove that $f(z) = az + b$ for some $a \in \mathbf{R}$ and $b \in \mathbf{C}$.

Problem 57. Suppose f is entire, with real and imaginary parts u and v satisfying

$$u(x, y) v(x, y) = 3$$

for all $z = x + iy$. Show that f is constant.

Problem 58. Prove that, if $G \subseteq \mathbf{C}$ is a domain and $f : G \rightarrow \mathbf{C}$ is a complex-valued function with $f''(z)$ defined and equal to 0 for all $z \in G$, then $f(z) = az + b$ for some $a, b \in \mathbf{C}$.

Problem 59. Show that

- (a) $\exp(2 \pm 3\pi i) = -e^2$
- (b) $\exp\left(\frac{2 + \pi}{4}\right) = \sqrt{\frac{e}{2}}(1 + i)$
- (c) $\exp(z + \pi i) = -\exp z$.

Problem 60. Prove that

- (a) $f(z) = \exp \bar{z}$ is nowhere holomorphic.
- (b) $f(z) = \exp z^2$ is entire. What is its derivative?

Problem 61. Show that

- (a) $|\exp(2z + i) + \exp(iz^2)| \leq e^{2x} + e^{-2xy}$.
- (b) $|\exp(z^2)| \leq \exp(|z|^2)$.
- (c) $|\exp(-2z)| < 1$ if and only if $\operatorname{Re} z > 0$.

Problem 62. Find all values of z such that

- (a) $\exp z = -2$
- (b) $\exp z = 1 + i\sqrt{3}$
- (c) $\exp(2z - 1) = 1$.

Problem 63. Find all solutions to the equation $e^{2z} - 2ie^z = 1$.

Problem 64. Let $G \subseteq \mathbf{C}^*$ be an open set and let f be a function that is continuous on G with the property

$$e^{f(z)} = z, \quad z \in G.$$

Show that f is holomorphic on G .

Remark A.0.2. This shows that a *continuously* defined logarithm on an open set is immediately holomorphic.

Problem 65. Find the all possible values of

- (a) $\log(-5)$
- (b) $\log(-2 + 2i)$
- (c) $\log(\sqrt{2} + i\sqrt{6})$
- (d) $\log(-ei)$

(e) $\log(1 + i)$

(f) $\log(-\sqrt{3} + i)$

Problem 66. Compute

(a) $\text{Log}(6 - 6i)$

(d) $\text{Log}((1 + i\sqrt{3})^5)$

(b) $\text{Log}(-e^2)$

(e) $\text{Log}(3 - 4i)$

(c) $\text{Log}(-12 + 5i)$

(f) $\text{Log}((1 + i)^4)$

Problem 67.

(a) Show that if $\text{Re } z_1 > 0$ and $\text{Re } z_2 > 0$, then

$$\text{Log}(z_1 z_2) = \text{Log } z_1 + \text{Log } z_2.$$

(b) Show that for any two non-zero complex numbers z_1 and z_2 ,

$$\text{Log}(z_1 z_2) = \text{Log } z_1 + \text{Log } z_2 + 2N\pi i,$$

where $N \in \{0, \pm 1\}$.

Problem 68. Example 2.9.4 (4) tells us that it's not necessarily true that $\log z^n = n \log z$, for $n \in \mathbf{Z}_{>0}$.

Writing $z = re^{i \text{Arg } z}$, show that, where $n \in \mathbf{Z}_{>0}$

$$\log(z^{1/n}) = \frac{1}{n} \ln r + i \left(\frac{\text{Arg } z + 2(pn + k)\pi}{n} \right), \quad k = 0, \dots, n-1.$$

Now, after writing

$$\frac{1}{n} \log z = \frac{1}{n} \ln r + i \left(\frac{\text{Arg } z + 2q\pi}{n} \right), \quad q \in \mathbf{Z},$$

show that we have equality of sets

$$\log(z^{1/n}) = \frac{1}{n} \log z$$

Problem 69. Find a domain in which the given function f is holomorphic; then find the derivative f' .

(a) $f(z) = 3z^2 - e^{2iz} + i \text{Log } z$

(b) $f(z) = (z + 1) \text{Log } z$

(c) $f(z) = \frac{\text{Log}(2z - i)}{z^2 + 1}$

(d) $f(z) = \text{Log}(z^2 + 1)$

Problem 70. Find the all possible values of

- | | |
|-----------------------|-----------------------|
| (a) $(-1)^{3i}$ | (e) $(-i)^i$ |
| (b) $3^{2i/\pi}$ | (f) $(ei)^{\sqrt{2}}$ |
| (c) $(1+i)^{1-i}$ | (g) $(-1)^{1/\pi}$ |
| (d) $(1+i\sqrt{3})^i$ | (h) $i^{i/\pi}$ |

Problem 71. Compute the principal value of the given complex powers.

- | | |
|--------------------------|---|
| (a) $(-1)^{3i}$ | (e) $i^{i/\pi}$ |
| (b) $3^{2i/\pi}$ | (f) $(1+i)^{2-i}$ |
| (c) 2^{4i} | (g) $\left(\frac{e}{2}(-1-i\sqrt{3})\right)^{3\pi i}$ |
| (d) $(1+i\sqrt{3})^{3i}$ | (h) $(1-i)^{4i}$ |

Problem 72.

- (a) Verify that $(z^\alpha)^n = z^{n\alpha}$ for $z \neq 0$ and $n \in \mathbf{Z}$.
- (b) Find a counterexample to the statement: $(z^\alpha)^\beta = z^{\alpha\beta}$, where $z \neq 0$ and $\alpha, \beta \in \mathbf{C}$.

Problem 73. Let z^α represent the principal value of the complex power. Find the derivative of the given function at the given point.

- | | |
|--------------------------------------|----------------------------------|
| (a) $z^{3/2}; \quad z = 1+i$ | (c) $z^{2i}; \quad z = i$ |
| (b) $z^{1+i}; \quad z = 1+i\sqrt{3}$ | (d) $z^{\sqrt{2}}; \quad z = -i$ |

Problem 74. Let $z \in \mathbf{C}$.

- (a) Prove that $|1^z|$ is single-valued if and only if $\text{Im } z = 0$.
- (b) Find a necessary and sufficient condition for $|i^z|$ to be single-valued.
- (c) Find a counterexample to the statement: 1^z is single-valued if and only if $\text{Im } z = 0$.

Problem 75. Express the value of the given trigonometric function in the form $x + iy$.

- | | |
|--------------------|--|
| (a) $\sin(4i)$ | (d) $\sin\left(\frac{\pi}{4} + i\right)$ |
| (b) $\cos(-3i)$ | (e) $\tan(2i)$ |
| (c) $\cos(2 - 4i)$ | (f) $\cot(\pi + 2i)$ |

(g) $\sec\left(\frac{\pi}{2} - i\right)$

(h) $\csc(1 + i)$

Problem 76. Find all complex values z satisfying the given equation.

(a) $\sin z = i$

(c) $\sin z = \cos z$

(b) $\cos z = 4$

(d) $\cos z = i \sin z$

Problem 77. Prove the properties stated in Discussion [2.11.4](#).

Problem 78.

(a) Prove that $\overline{\cos z} = \cos \bar{z}$.

(b) What is $\operatorname{Re} \cos z$ and $\operatorname{Im} \cos z$?

(c) Using the identity $e^{iz} = \cos z + i \sin z$, prove $\overline{\sin z} = \sin \bar{z}$ and find $\operatorname{Re} \sin z$ and $\operatorname{Im} \sin z$.

References

- [BC09] J. Brown and R. Churchill, *Complex Variables and Applications*, Brown and Churchill series, McGraw-Hill Education, 2009.
- [BMPS] M. Beck, G. Marchesi, D. Pixton, and L. Sabalka, *A First Course in Complex Analysis*, [Available online](#), [Version 1.54].
-