

Kinetic Measurement of Leukemia- Cell Proliferation Rate by Deuterium Labeling Predicts Time to Initial Treatment

Abstract 829

**Murphy EJ, Neuberg D, Rai KR, Rassenti L, Emson C, Li K, Kay
NE, Wierda WG, Hayes GM, Brown JR, Byrd JC, McConnel C,
Barrientos JC, Redd RA, Turner S, Busch R, Greaves A,
Hellerstein M, Chiorazzi N, Kipps TJ**

Cell Kinetics in CLL

- Traditionally thought of as a disease of failed cell death
- On average, birth rate of CLL cells < birth normal B cells
- It has also been shown that
 - Significant heterogeneity of birth rates between patients
 - Some individuals have significantly higher birth rates than normal B cells
 - Different clones have significantly different birth rates
- It is likely that defects in both cell production and destruction are present in CLL

CLL Cell Birth Rate as a Prognostic Marker


- Hypothesis: birth rate could be used as a prognostic marker for disease progression in CLL
- Study goals:
 - Assess CLL cell birth rates in untreated patients (Rai stage 0, 1, 2; diagnosis within 3 years)
 - Compare birth rate to other established prognostic markers
 - Assess all markers as predictors need for treatment (progression) during 2 years of patient follow-up

Demographics

		Total n = 97
Sex		
Female		39 (40%)
Male		58 (60%)
Age		
Median (range)		57 (40-85)
Rai stage at enrollment		
0		36 (37%)
I		44 (45%)
II		17 (18%)
Enrollment site		
Dana-Farber Cancer Institute		5 (5%)
North Shore - Long Island Jewish Health System		19 (20%)
Mayo Clinic		18 (19%)
MD Anderson Cancer Center		14 (14%)
Ohio State University		10 (10%)
University of California San Diego		31 (32%)
Months from CLL diagnosis to study entry		
Median (range)		12 (1-36)


Measurement of CLL Cell Birth Rate

Heavy Water Study Protocol




Heavy Water Studies

Body Water Enrichment and CLL cell incorporation




CLL Cell Birth Rates (n = 97)

Median Birth Rate 0.32% new cells per day
Range 0.07%-1.31% per day




Birth Rate in Relation to Other Prognostic Factors


IGHV Mutation Status




ZAP-70 Expression



CD38 Expression



Cytogenetics




Treated Patients


	Total n = 97	Treated CLL	
		No n = 64	Yes n = 33
Sex			
Female	39 (40%)	26 (41%)	13 (39%)
Male	58 (60%)	38 (59%)	20 (61%)
Age			
Median (range)	57 (40-85)	58 (40-85)	56 (41-78)
Rai stage at enrollment			
0	36 (37%)	32 (50%)	4 (12%)
I	44 (45%)	24 (38%)	20 (61%)
II	17 (18%)	8 (12%)	9 (27%)
Enrollment site			
Dana-Farber Cancer Institute	5 (5%)	3 (5%)	2 (6%)
North Shore - Long Island Jewish Health System	19 (20%)	11 (17%)	8 (24%)
Mayo Clinic	18 (19%)	12 (19%)	6 (18%)
MD Anderson Cancer Center	14 (14%)	9 (14%)	5 (15%)
Ohio State University	10 (10%)	0 (14%)	1 (3%)
University of California San Diego	31 (32%)	20 (31%)	11 (33%)
Months from CLL diagnosis to study entry			
Median (range)	12 (1-36)	11 (1-36)	12 (1-36)

Birth Rate vs Time to Initial Therapy

Birth Rate $>0.35\%$ per day ¹




Birth Rate $>0.49\%$ per day



1. Messmer BT, et al. *J Clin Invest.* 2005;115(3):755-764.

Murphy EK, et al. *Blood.* 2014;124: Abstract 829.

Time to Treatment Stratified by *IGHV* and Birth Rate



Conclusions

- Increased CLL cell birth rate in early-stage disease is a strong predictor of earlier treatment
- In a multivariable model, only birth rate and *IGHV* mutational status contributed significantly
- Within the groups of patients with mutated or unmutated *IGHV*, birth rate allowed further discrimination in a time-to-treatment analysis
- This direct measure of CLL-cell proliferation represents a new marker for prognostication for subjects with early-stage CLL