

The Role of Pathology in the Era of Targeted Therapy

Keith Kerr, MD, FRCPath


Aberdeen Royal Infirmary
Aberdeen, United Kingdom

www.prIMEoncology.org




It Is Worthwhile Finding an Actionable Genetic Alteration in Lung Cancer


Using Multiplexed Assays of Oncogenic Drivers in Lung Cancers to Select Targeted Drugs¹



Comparison of Survival for Patients With Lung Adenocarcinoma in Japan Before and After Gefetinib Approval²




Comparison of Survival for Patients With Lung Adenocarcinoma in Second Line Before and After Crizotinib Approval³



1. Kris MG, et al. *JAMA*. 2014;311(19):1998-2006. 2. Takano T, et al. *J Clin Oncol*. 2008;26(34):5589-5595. 3. Shaw AT, et al. *Lancet Oncol*. 2011;12(11):1004-1012.

Oncogene “Drivers” in Adenocarcinoma




NTRK1 fusion

MPRIIP-NTRK1 and *CD74-NTRK1*
3.3% of ‘onco-negative’
adenocarcinomas
Trk inhibitors exist
Vaishnavi A, et al. *Nat Med.*
2013;19(11):1469-1472.

CD74-NRG1 fusion

Search in ‘onco-negative’
adenocarcinomas
ERBB3 and PI3K-AKT pathway
activation
Mucinous adenocarcinomas
Potential therapeutic target
Fernandez-Cuesta L, et al. *Cancer
Discov.* 2014;4(4):415-422.

Squamous Cell Carcinoma of the Lung: Molecular Subtypes and Therapeutic Opportunities



EGFR

TKI vs MoAb

Mutations – rarity (VIII – 8%)

Targeting the receptor

IGFR1

Figitumumab


Some effect in squamous

Toxicity

MoAB, monoclonal antibody; TKI, tyrosine kinase inhibitor

Perez-Moreno P, et al. *Clin Cancer Res.* 2012;18(9):2443-2451.


Most Lung Cancer Samples Are Small Biopsies or Cytology-Type Samples



Surgically resected tumour



Small biopsy samples



Cytology samples

Is There Enough Material for These Studies?




Two biopsy fragments <1 mm

- Morphologic diagnosis
- Immunohistochemistry (IHC)
- Molecular testing
- Conserve tissue
- Don't waste

**On average, only 20%
of this tissue is tumour**


Immunohistochemical (IHC) Subtyping of NSCLC



- Predictive IHC has 'levelled the playing field'
- Better diagnosis possible on poorer specimens

NSCLC-NOS, non-small cell lung cancer not otherwise specified

Tumour Histology, Genotype and Treatment of Lung Cancer




SCLC, small cell lung cancer

Whom Should We Be Testing for *EGFR* Mutation and *ALK* Rearrangement?

- *All nonsquamous tumours in patients with advanced/recurrent disease should be tested for EGFR mutation and ALK rearrangement*
- *Selected squamous tumours (from patients with minimal or remote smoking history) should strongly be considered for testing*

Which Tumours Do We Test for *EGFR* Mutation and *ALK* Arrangement?



What Do We Use for the Test?


- Whatever is available – we need tumour cells!!!
- Tissue or cytology cell block sections
 - Maximise tumour cells &.....
 - Minimise nontumour cells in material submitted for DNA extraction
 - >10%, >50%.....
 - >100 cells?
 - For FISH – at least 50 assessable cells for *ALK*
 - For IHC?

Pathologic Assessment for Molecular Testing

- Tumour present?
- Prepared appropriately?
- Is there is enough tumour?
- The molecular lab knows what it is getting?




- Integration of the molecular results
 - Are they meaningful?
 - Are they reliable?



**Tissue from sections:
a standard source
of tumour DNA**



**EGFR c.2573T>G; p.Leu858Arg
(exon21 L858R)**




EGFR Mutation Testing Methodology

Increasing Sensitivity ↓	Technique	Limit of Detection, % Mutant DNA	Mutations Identified
	Direct sequencing	10 - 20	Known and new
	TaqMan PCR	10	Known only
	Loop-hybrid mobility shift assay	10	Known only
	Pyrosequencing	5	Known and new
	PCR-SSCP	5	Known and new
	dHPLC (WAVE surveyor)	3-5	Known and new
	Cycleave PCR	5	Known only
	PCR-RFLP and length analysis	5	Known only
	MALDI-TOF MS-based genotyping	5	Known only
	High resolution melting (HRM)	3-5	Known and new
	Scorpion ARMS	1	Known only
	PNA-LNA PCR clamp	1	Known only
	<i>Single molecule sequencing</i>	<i>0.1</i>	<i>Known and new</i>
	<i>Mutant-enriched sequencing</i>	<i>0.1</i>	<i>Known only</i>
	<i>SMAP</i>	<i>0.1</i>	<i>Known only</i>


ARMS, amplification refractory mutation system; MALDI-TOF, matrix-assisted laser desorption/ionization time-of-flight; PCR, polymerase chain reaction; PNA-LNA, peptide nucleic acid-locked nucleic acid; RFLP, restriction fragment length polymorphisms; SMAP, smart amplification process; SSCP, single-strand conformation polymorphism

Pao W, Ladanyi M. *Clin Cancer Res*. 2007;13(17):4954-4955.


All *EGFR* Mutations Are Not Equal



The 'break apart'
FISH test




Multiplex PCR



Ou SH, et al. *Oncologist*. 2012;
17(11):1351-1375.

IHC



Change in DNA
Sequence
ALK
Translocation

**ALK Fusion Gene
mRNA Transcript**

ALK PROTEIN


**Biologic Activity
Oncogenesis
Drug target**

Transcription



Translation

ALK: Which Test?

ALK Negative



ALK Positive



The Protein Does the Job

The Protein Is the Target of the Drug

Cases with *ALK* gene fusion

Cases with ALK protein excess

**FISH positive,
IHC negative**

Lower Response Rate?

**FISH negative,
IHC positive**

Reports of Response to ALK TKI

Who Orders the Test?

Reflex vs Bespoke Testing

Reflex—pathologist driven

- Fast
- Becomes 'routine'
- Ready for tumour board decision
- *Potential for waste*
 - Time
 - Tissue
 - Money

Bespoke—to order from oncologist

- Only when needed
- Preserves tissue
- Time not wasted
- *Slower turnaround*
- *Could be illogical; cases may be missed*

Do We Always Succeed?

- **Diagnostic IHC—rarely insufficient**
 - Occasionally it just doesn't work!
- ***EGFR* mutation**
- ***ALK* rearrangement**
 - IHC screening
 - Confirmation by FISH

EBUS Samples for EGFR Mutation?

Reference	% EBUS INSUFF for <i>EGFR</i> Mutation Test	Comment
Garcia-Olive et al	28%	12% for core biopsy
Schuurbiers et al	23%	
Esterbrook et al	12%	Cell block based
Navani et al	10%	
Rekhtman et al	2%	



INSUFFICIENT FOR TEST


EBUS, endobronchial ultrasound

Garcia-Olivé I, et al. *Eur Respir J*. 2010;35(2):391-395. Schuurbiers OC, et al. *J Thorac Oncol*. 2010;5(10):1664-1667. Esterbrook G, et al. *Lung Cancer*. 2013;80(1):30-34. Navani N, et al. *Am J Respir Crit Care Med*. 2012;185(12):1316-1322. Rekhtman N, et al. *J Thorac Oncol*. 2011;6(3):451-458.


ALK Testing 'Success'??

- In Aberdeen
 - About 4% of cases insufficient for *ALK* IHC
 - About 10% cases insufficient for *ALK* FISH
 - 50-60 assessable cells
 - 4 high power fields to assess
- Up to 20% of samples may be 'insufficient' for *ALK* FISH testing
Lantuéjoul S, et al. In IASLC ALK Atlas.
- Cytology samples less often assessable (69%) versus biopsy (89%) for *ALK* FISH
Vidal J, et al. *J Thorac Oncol.* 2014;9(12):1816-1820.
- Cytology samples suitable for *ALK* IHC
Savic S, et al. *J Thorac Oncol.* 2013;8(8):1004-1011.

A Testing Algorithm



A Testing Algorithm.....



More Complex, Ambitious Testing

- More markers to be tested
- Sequential testing increases risk
 - 30% failure
 - 15% fails in EURTAC trial

Buettner R, et al. *J Clin Oncol*. 2013;31(15):1858-1865.

Benlloch S, et al. *J Clin Oncol*. 2012;30(Suppl): Abstract 10596.

- Trials (Battle, MSKCC SCC)
 - 13% to 17% incomplete test sets

Tam AL, et al. *J Thorac Oncol*. 2013;8(4):436-442.

Paik PK, et al. *J Clin Oncol*. 2012;30(Suppl): Abstract 7505.

NGS for Molecular Testing


- Quoted amounts of DNA required rather variable
 - Technology dependant
 - Size of panel
- Mutation > fusion gene > gene copy number
- Fragmentation of DNA
- Bioinformatic analysis
- 80% samples – Complete panel of mutations
- 95% samples – *EGFR*, *KRAS*, *BRAF*, *HER2* mutations
- ‘Minimum 2000 cells’ - 5 x 10 um thick sections
- Much still to define

Meyerson M, et al. *Nat Rev Genet.* 2010;11(10):685-696.

Next Generation Sequencing: Different Approaches

- Whole Genome Sequencing (WGS): Determines the complete DNA sequence of an organism's genome at a single time
- Whole Exome Sequencing (WES): Selectively sequences only the coding areas of the genome
- “Fully Informative” Sequencing: Sequences a defined subset of genes of interest in their entirety
- Targeted Sequencing (Hot Spot): Sequences only the hot spots of a subset of genes of interest

Lung Cancer Master Protocol: Lung MAP Trial in SCC



Testing Plasma Samples

- Free plasma DNAor CTCs
- Referred to in some guidelines
- Needs sensitive methodology
- Primary analysis
 - cfDNA 58% sensitivity, 86% specificity.
- Monitoring role?
 - 84% sensitivity (*EGFR*) using CTCs
 - Relapse on treatment
- High research priority
- Not currently recommended

Couraud S, et al,
Clin Cancer Res.
2014;20(17):4613-4624.

Marchetti A, et al. *PLoS One.* 2014;9(8):e103883.

CTC, circulating tumour cells





Gridelli C, et al. *Clin Lung Cancer.* 2014;15(3):173-181. Kerr KM, et al, *Ann Oncol.* 2014;25(9):1681-1690.

Gefitinib Treatment in *EGFR*-Mutated Caucasian NSCLC: Circulating-Free Tumour DNA as a Surrogate for Determination of *EGFR* Status

- Mutation status concordance between tumor and matched plasma was 94%, sensitivity 66% and specificity of 100% (n = 652)
- Reproducibility also high: Mutation concordance of 97% for 224 matched plasma specimens
- *Post hoc* analysis of the efficacy of first-line gefitinib revealed that PFS was similar for those with *EGFR* mutation–positive tissue (9.7 months) vs both mutation-positive tissue and plasma (10.2 months)
- Conclusions: Although these results are encouraging and suggest that plasma is a suitable substitute for mutation analysis regardless of mutation subtype, tumor tissue should be considered the preferred sample type when available

26 Sept 2014, CHMP of EMA gave positive opinion to include a label in gefitinib SmPc for the use of circulating tumour DNA (ctDNA) obtained from a blood sample, to be used for the assessment of *EGFR* mutation status in those patients where a tumour sample is not an option.

Biomarkers for Immunotherapy?



PD-L1 Negative

PD-L1 Positive (predictive of response)



1%

5%



10%

50% cell positive

Several therapeutics
Several companion diagnostics.....

Intensity of staining?
Immune cell staining?

The Role of Pathology in the Era of Targeted Therapy

- **Pathologic diagnosis**
- **Pathologic assessment**
- **Tissue handling**
- **Adapt to range of markers required**
- **Multiple test modalities**