

RAISE: A Randomized, Double-Blind, Multicenter Phase III Study of Irinotecan, Folinic Acid, and 5-Fluorouracil (FOLFIRI) Plus Ramucirumab or Placebo in Patients With Metastatic Colorectal Carcinoma Progression During or Following First-Line Combination Therapy With Bevacizumab, Oxaliplatin, and a Fluoropyrimidine


Abstract 512

Tabernero J, Cohn AL, Obermannova R, Garcia-Carbonero R, Ciuleanu T-E, Portnoy DC, Van Cutsem E, Grothey A, Prausová J, Garcia-Alfonso P, Yamazaki K, Clingan PR, Zagonel V, Kim TW, Simms L, Change S-C, Nasrullah F, Yoshino T, The RAISE Study Investigators

Introduction

- **Colorectal carcinoma (CRC) is the fourth most common cause of cancer death worldwide.**
- **VEGF and VEGFR-2 mediated signaling and angiogenesis are important in CRC tumor growth and are established therapeutic targets.**
- **Ramucirumab is a recombinant human IgG1 monoclonal antibody that binds to the extracellular domain of VEGFR-2, preventing ligand binding and receptor activation.**
- **The RAISE study evaluated the efficacy and safety of adding ramucirumab to the standard second-line treatment FOLFIRI following progression after first-line bevacizumab, oxaliplatin, and a fluoropyrimidine.**

Trial Design



*Irinotecan: 180 mg/m²; Folinic acid: 400 mg/m²; 5-Fluorouracil 400 mg/m² bolus, followed by 2400 mg/m² administered intravenously over 46 to 48 hours (continuously)

Stratification factors:

- Geographic regions
- *KRAS* mutation status
- Time to disease progression after beginning first-line therapy

Sample size assumptions:

- Hazard ratio of 0.8
- Median overall survival of 10 months in the control arm vs 12.5 months with ramuciruman with a 2-sided α level of 0.05
- Enrollment of 1050 patients with 756 events for 85% power
- Gatekeeping from OS to PFS to ORR

IG, immunogenicity; PFS, prorgession-free survival; PK, pharmacokinetics; OS, overall survival; ORR, objective response rate

Tabernero J, et al. *J Clin Oncol*. 2015;33(suppl 3): Abstract 512.

Key Inclusion Criteria

- **Diagnosis of metastatic CRC (histological or cytological confirmation)**
- **Known *KRAS* mutation status**
- **Eastern Cooperative Oncology Group performance status score of 0 or 1**
- **Documented progressive disease during or after a first-line combination therapy of bevacizumab, oxaliplatin, and a fluoropyrimidine**
- **At least two doses of bevacizumab in first-line therapy**
- **Disease progression ≤ 6 months after the last dose of first-line therapy**
- **Adequate hematologic and biochemical parameters**

Key Exclusion Criteria


- Previous first-line systemic therapy, other than a combination of bevacizumab, oxaliplatin, and a fluoropyrimidine
- Uncontrolled hypertension
- Arterial thrombotic event within 12 months
- Received bevacizumab ≤ 28 days or chemotherapy ≤ 21 days prior to randomization
- Known brain metastasis
- Grade 3 or higher bleeding event ≤ 3 months prior to randomization

Baseline Characteristics

	Ramucirumab + FOLFIRI (N = 536)	Placebo + FOLFIRI (N = 536)
Median age (range), years	62 (21-83)	62 (33-87)
Age group, n (%) ≥65 years	212 (39.6)	215 (40.1)
Male, n (%)	289 (53.9)	326 (60.8)
Race, n (%)		
Caucasian	405 (75.6)	410 (76.5)
Asian	111 (20.7)	103 (19.2)
ECOG PS, n (%)		
0	263 (49.1)	259 (48.3)
1	268 (50.0)	273 (50.9)
Number of metastatic sites, n (%)		
1	171 (31.9)	157 (29.3)
2	205 (38.2)	194 (36.2)
≥3	157 (29.3)	182 (34.0)
Liver only metastasis, n(%)	92 (17.2)	95 (17.7)
Yes		
Carcinoembryonic antigen, n (%)		
<200 µg/L	389 (72.6)	393 (73.3)
≥200 µg/L	108 (20.1)	107 (20.0)

Some patients may be missing from some categories

Patient Disposition




Treatment discontinuation					
	n	%		n	%
Progressive disease	338	63	Progressive disease	417	78
Subject decision	49	9	Subject decision	26	5
Death	18	3	Death	13	2
Withdrawal of consent	8	1	Withdrawal of consent	4	<1
Lost to follow-up	0	0	Lost to follow-up	0	0
Adverse event	59	11	Adverse event	23	4
Other	4	<1	Other	5	<1
Investigator decision	35	7	Investigator decision	25	5
Total	511	95	Total	513	96

ITT, intent-to-treat; N, number of patients in category; One patient was randomized to the placebo group but received ramucirumab

Tabernero J, et al. *J Clin Oncol*. 2015;33(suppl 3): Abstract 512.

Overall Survival




Patients at Risk

	0	3	6	9	12	15	18	21	24	27	30	33	36	39	42
Ram + FOLFIRI	536	497	421	345	269	195	114	78	53	34	22	12	4	0	0
Placebo + FOLFIRI	536	486	400	329	228	166	108	66	44	22	10	2	2	1	0

CI, confidential interval; HR, hazard ratio; Ram, ramucirumab

Tabernero J, et al. *J Clin Oncol*. 2015;33(suppl 3): Abstract 512.


Overall Survival by Subgroup



CEA, carcinoembryonic antigen; ECOG PS, Eastern Cooperative Oncology Group performance status; N, number of patients; PD, progressive disease

Tabernero J, et al. *J Clin Oncol*. 2015;33(suppl 3): Abstract 512.

Progression-Free Survival



Patients at Risk

Ram + FOLFIRI	536	381	234	142	77	38	20	11	6	5	2	1	1	0	0
Placebo + FOLFIRI	536	345	182	92	52	31	17	10	3	1	0	0	0	0	0

Tumor Response

	Ramicurimab + FOLFIRI N = 536 %	Placebo + FOLFIRI N = 536 %	P-value
Response rate (CR+PR)	13.4	12.5	.6336
Disease control rate (CR+PR+SD)	74.1	68.8	.0587
Complete response	0	0.4	
Partial response (PR)	13.4	12.1	
Stable disease (SD)	60.6	56.3	
Progressive disease (PD)	16.2	25	
Not done or unknown	9.7	6.2	

Tumor assessments based on RECIST 1.1

Post-Treatment Anticancer Therapy

	Ramicurimab + FOLFIRI N = 536 %	Placebo + FOLFIRI N = 536 %
Systemic anticancer therapy	54.1	56.0
Fluoropyrimidines	33.4	34.0
Irinotecan	30.6	28.7
Anti-EGFR antibody (cetuximab or panitumumab)	25.6	28.5
Anti-angiogenic agents (bevacizumab or aflibercept)	12.7	13.1
Regorafenib	10.4	10.8
Platinum	9.0	10.8

Treatment-Emergent Adverse Events

Preferred Term	Any Grade				Grade ≥3			
	Ramucirumab + FOLFIRI N=529		Placebo + FOLFIRI N=528		Ramucirumab + FOLFIRI N=529		Placebo + FOLFIRI N=528	
	n	%	n	%	n	%	n	%
Any TEAE	522	98.7	519	98.3	418	79.0	329	62.3
<i>Neutropenia</i>	311	58.8	241	45.6	203	38.4	123	23.3
<i>Fatigue</i>	305	57.7	275	52.1	61	11.5	41	7.8
Diarrhea	316	59.7	271	51.3	57	10.8	51	9.7
Hypertension	136	25.7	45	8.5	57	10.8	15	2.8
Stomatitis	163	30.8	110	20.8	20	3.8	12	2.3
<i>Abdominal pain</i>	140	26.5	139	26.3	18	3.4	19	3.6
<i>Thrombocytopenia</i>	150	28.4	72	13.6	16	3.0	4	0.8
Vomiting	154	29.1	144	27.3	15	2.8	13	2.5
Nausea	262	49.5	271	51.3	13	2.5	14	2.7
Decreased appetite	198	37.4	144	27.3	13	2.5	10	1.9
<i>Anemia</i>	86	16.3	110	20.8	8	1.5	19	3.6
Constipation	151	28.5	120	22.7	5	0.9	8	1.5
Peripheral edema	108	20.4	48	9.1	1	0.2	0	
Epistaxis	177	33.5	79	15.0	0		0	
Alopecia	155	29.3	165	31.3	0		0	

Italicized terms are consolidated adverse event category comprising synonymous MedDRA preferred terms.

Adverse Events of Special Interest

	Any Grade				Grade ≥3			
	Ramucirumab + FOLFIRI N=529		Placebo + FOLFIRI N=528		Ramucirumab + FOLFIRI N=529		Placebo + FOLFIRI N=528	
	n	%	n	%	n	%	n	%
Bleeding/hemorrhage event	232	43.9	120	22.7	13	2.5	9	1.7
Hypertension	138	26.1	45	8.5	59	11.2	15	2.8
Proteinuria	90	17.0	24	4.5	16	3.0	1	0.2
GI hemorrhage	65	12.3	36	6.8	10	1.9	6	1.1
Venous thromboembolic events	44	8.3	34	6.4	22	4.2	11	2.1
Infusion-related reaction	31	5.9	16	3.0	4	0.8	2	0.4
Renal failure	18	3.4	18	3.4	7	1.3	5	0.9
GI perforation	9	1.7	3	0.6	9	1.7	3	0.6
Arterial thromboembolic event	8	1.5	13	2.5	4	0.8	6	1.1
Pulmonary hemorrhage events	7	1.3	3	0.6	0		0	
Healing complication	6	1.1	1	0.2	1	0.2	0	
Congestive heart failure	4	0.8	3	0.6	4	0.8	3	0.6
Fistula	4	0.8	2	0.4	0		0	
RPLS	1	0.2	1	0.2	0		0	
Thrombotic microangiopathy	1	0.2	0		1	0.2	0	
Hepatic hemorrhage event	0		1	0.2	0		1	0.2

Abbreviations: GI=gastrointestinal; RPLS=reversible posterior leukoencephalopathy syndrome.

Conclusions

- **RAISE met its primary endpoint.**
 - **Demonstrated a statistically significant improvement in overall survival for ramucirumab and FOLFIRI vs placebo and FOLFIRI**
 - **In second-line metastatic CRC patients who progressed after first-line treatment with bevacizumab, oxaliplatin, and a fluoropyrimidine**
- **Consistent survival benefits were observed across subgroups**
- **Ramucirumab in combination with FOLFIRI was well tolerated in patients with mCRC; overall, the adverse events were considered manageable**