



Therapeutic Dual Inhibition of HER2 Pathway for Metastatic Colorectal Cancer (mCRC): The HERACLES Trial

Abstract 565

Siena S, Sartore-Bianchi A, Trusolino L, Martino C, Bencardino K,
Lonardi S, Leone F, Zagonel V, Bertotti A, Valtorta E, Siravegna G,
Amatu A, Vanzulli A, Regge D, Ghezzi S, Ciardiello F, Veronese S,
Comoglio PM, Bardelli A, Marsoni S

Anti-EGFR and Anti-HER2 Therapies in Xenopatient With Cetuximab-Resistant HER2-Amplified mCRC



Growth curves of tumors in xenopatient derived from **cetuximab-resistant quadruple-negative HER2-amplified mCRC** (n = 5 for each treatment arm)

Bertotti A, et al. *Cancer Discov.* 2011;1(6):508-523

Siena S, et al. *J Clin Oncol.* 2015;33(suppl 3): Abstract 565.

Study Flow Chart



Advanced metastatic KRAS WT CRC patients refractory to: Fluoropyrimidines, irinotecan, oxaliplatin, cetuximab/panitumumab, and bevacizumab


Screen for HER2 by IHC and FISH on last available FFPE

HER2 negative

OFF TRIAL

HER2 positive

Informed Consent



IRCC
TARGET
NGS panel

HERACLES

Tumor Assessment (RECIST 1.1) Q8-9 weeks


Trastuzumab IV + Lapatinib OS
(4mg/kg load + 2mg/kg/wk) (1000 mg/die)

PD

Whenever possible



Imaging-guided
biopsy



IRCC TARGET
NGS panel

Liquid Biopsy
Q 2weeks



X4



X4



X4



X4




ctDNA Mutations: KRAS, NRAS, BRAF, PIK3CA; Amplification: HER2, KRAS, MET; Custom Exome Panel in nonresponders and at relapse serum quantification of Extracellular Domain of HER2 (HER2-ECD)

Study Design

Objective	Activity of lapatinib + trastuzumab in HER2 amplified metastatic colorectal cancer
Design	Phase II, open label, multicentric single-stage design according to Fleming and A'Hern
Endpoints	Objective response (primary), safety, and time-to-progression (TTP, secondary)
Population	HER2+, mCRC after failures of previous chemotherapy, antiangiogenesis, and antiEGFR therapies
Main inclusion criteria	<ul style="list-style-type: none"> • Pathology: KRAS exon 2 wild type (WT); HER2-positivity (IHC3+ or IHC2+/SISH+) according to HERACLES diagnostic criteria • Prior therapies: Unless otherwise contraindicated should have received and <u>failed</u> fluoropyrimidine-, oxaliplatin-, irinotecan-, cetuximab- or panitumumab-containing therapy regimens • Performance status ECOG 0-1 • Measurable disease according to RECIST 1.1 • Adequate vital organ function (cardiac, bone marrow, renal, liver)
Treatment	Lapatinib 1000 mg po qd Trastuzumab 4 mg/kg IV load dose, then 2 mg/kg IV weekly
Tumor assessment	RECIST 1,1 imaging every 8 weeks and liquid biopsy every 2 weeks Double-lecture centralized response assessment, Mint Lesion® software assisted
Statistical assumptions	H0 10%, H1 30%, $\alpha = 0.05$ and power =0.85 Primary endpoint: Objective Response according to RECIST 1.1
Sample size	27 patients; 6 responses must be observed for the study to be positive

Study Consort




Patients Characteristics


Characteristics		Patients	
		N	%
Age	Median	63	
	Range	40-86	
Gender	Females	2	9%
	Males	21	91%
Performance status	ECOG 0/1	23	100%
Tumor site	Colon	19	85%
	Rectum	3	15%
Extent of disease	One site only	3	
	Liver	2	
	Lung	1	
	Multiple sites	20	
	Liver	12	
	Lung	11	
	Nodes	9	
	Other sites	12	
Prior treatments	Median number of previous line for metastatic disease (range)		5 (2-8)
		N	%
	More than 3 previous lines	20	87%
	Prior cetuximab or panitumumab	23	100%
	Patients responding to cetuximab or panitumumab	0	0%
	Prior bevacizumab treatment*	10*	100%*

*Only 10/23 patients were eligible for bevacizumab treatment


Patient #121015: Age 62, 5 Prior Chemotherapies + Panitumumab + Bevacizumab + Regorafenib



Dec 31, 2013
Baseline




March 20, 2014
Week 8 **partial response (PR)**




July 10, 2014
Week 24 **PR**

[...]


Patient #121016: Age 86, 4 Prior Chemotherapies + Panitumumab



Jan 27, 2014
Baseline




March 23, 2014
Week 8 **PR**



July 15, 2014
Week 24 **PR**

[...]

Objective Response Rate by HER2 IHC Score or Gene Copy Number (GCN) Variation




*Fisher's exact test

IHC, Immunohistochemistry; SD, stable disease; PD, progressive disease

Siena S, et al. *J Clin Oncol*. 2015;33(suppl 3): Abstract 565.


Response: Spaghetti Plot



*Three patients are not displayed: 122025 (IHC 2+ GCN<20) too early (not yet assessed); 121011 (IHC 3+ GCN>20) and 121013 (IHC 3+ GCN<20) early clinical PDs

Siena S, et al. *J Clin Oncol*. 2015;33(suppl 3): Abstract 565.


Response: Waterfall Plot




*Three patients are not displayed: 122025 (IHC 2+ GCN<20) too early (not yet assessed); 121011 (IHC 3+ GCN>20) and 121013 (IHC 3+ GCN<20) early clinical PDs.

Siena S, et al. *J Clin Oncol*. 2015;33(suppl 3): Abstract 565.


Time-to-Progression: All Patients



Time-to-Progression by HER2 IHC Score



Time-to-Progression by HER2 GCN



Drug Related Toxicities

GI Tract	Total events	% of total AEs	Grade 3	% of total G3 AE
Diarrhea	54	31	0	
Nausea/Vomiting	16	9	0	
Abdominal pain	6	3	0	
Mucositis	2	1	0	
SKIN & ANNEXES				
Paronychia	16	9	0	
Rash/dermatitis	19	11	1	17
Pruritis	4	2	0	
Polliculitis	1	1	0	
Erythema	2	1	0	
Xerosis	4	2	0	
SYSTEMIC				
Fatigue	28	16	3	50
Loss of appetite	2	1	0	
Chills	1	1	0	
Parathesia	1	1	0	
Hand and Foot	5	3	0	
SENSE ORGAN				
Conjunctives	1	1	0	
Dysgeusia	1	1	0	
Dysphonia	1	1	0	
LABORATORY				
Anemia	3	2	0	
Hypomagnesaemia	1	1	0	
CARDIOVASCULAR				
Hypertension	3	2	1	17
LVEF drop	1	1	1	17
Epistaxis	1	1	0	
Total	173	100	6	100

Siena S, et al. *J Clin Oncol*. 2015;33(suppl 3): Abstract 565.

Patient #121001: Liquid Biopsy and HER2-ECD

