

Which of the following (in addition to bone-targeted treatment) would you recommend for treatment of metastatic ER-negative, HER2-positive breast cancer (in lung and bone) in a patient who received 1 year of adjuvant trastuzumab combined with chemotherapy with disease recurrence 18 months later?

- 1. Trastuzumab + single agent chemotherapy**
- 2. Trastuzumab + pertuzumab + taxane**
- 3. Lapatinib + single agent chemotherapy**
- 4. Trastuzumab emtansine (TDM-1)**
- 5. TDM-1 + pertuzumab**
- 6. TDM-1 + pertuzumab + taxane**

Which of the following (in addition to bone-targeted treatment) would you recommend for treatment of metastatic ER-negative, HER2-positive breast cancer (in lung and bone) in a patient who received 1 year of adjuvant trastuzumab combined with chemotherapy with disease recurrence 18 months later?

1. Trastuzumab + single agent chemotherapy



2. Trastuzumab + pertuzumab + taxane



3. Lapatinib + single agent chemotherapy



4. Trastuzumab emtansine (TDM-1)



5. TDM-1 + pertuzumab



6. TDM-1 + pertuzumab + taxane





Clinical Opinion Poll Question #5: Optimal Anti-HER2 Approach in Metastatic HER2+ Breast Cancer

Edith A. Perez, MD

Deputy Director at Large, Mayo Clinic Cancer Center

Serene M. and Frances C. Durling Professor of Medicine and Cancer Biology

Group Vice Chair, Alliance for Clinical Trials in Oncology

Director, Mayo Clinic Breast Cancer Translational Genomics Program

Mayo Clinic



Systemic Therapy for Patients With Advanced Human Epidermal Growth Factor Receptor 2–Positive Breast Cancer: American Society of Clinical Oncology Clinical Practice Guideline

Sharon H. Giordano, Sarah Temin, Jeffrey J. Kirshner, Sarat Chandarlapaty, Jennie R. Crews, Nancy E. Davidson, Francisco J. Esteva, Ana M. Gonzalez-Angulo, Ian Krop, Jennifer Levinson, Nancy U. Lin, Shanu Modi, Debra A. Patt, Edith A. Perez, Jane Perlmutter, Naren Ramakrishna, and Eric P. Winer

Recommendations HER2-targeted therapy is recommended for patients with HER2-positive advanced breast cancer, except for those with clinical congestive heart failure or significantly compromised left ventricular ejection fraction, who should be evaluated on a case-by-case basis. Trastuzumab, pertuzumab, and taxane for first-line treatment and T-DM1 for second-line treatment are recommended. In the third-line setting, clinicians should offer other HER2-targeted therapy combinations or T-DM1 (if not previously administered) and may offer pertuzumab, if the patient has not previously received it. Optimal duration of chemotherapy is at least 4 to 6 months or until maximum response, depending on toxicity and in the absence of progression. HER2-targeted therapy can continue until time of progression or unacceptable toxicities. For patients with HER2-positive and estrogen receptor–positive/progesterone receptor–positive breast cancer, clinicians may recommend either standard first-line therapy or, for selected patients, endocrine therapy plus HER2-targeted therapy or endocrine therapy alone.



First-Line HER2+ Metastatic Breast Cancer (MBC): CLEOPATRA

- Double-blind, placebo-controlled phase III trial
 - Docetaxel 75 mg/m² escalated to 100 as tolerated, about 6 cycles
 - Trastuzumab and pertuzumab (PER) / placebo q3w
- Primary endpoint
 - Independently assessed PFS
- 808 patients centrally confirmed HER2+ MBC
 - Adjuvant therapy
 - 53% no prior chemo
 - 10% prior trastuzumab
 - 49% ER+, 24% received endocrine therapy

Baselga J, et al. *N Engl J Med*. 2012;366(6):520-529. Swain SM, et al. *Lancet Oncol*. 2013;14(6):461-471.



CLEOPATRA: Results and Perspective

- PFS benefit seen in essentially all predefined subsets
- Complete response rare at 4%-5.5%
(Partial response 65%-75%)
- Survival impact is practice changing

	Slamon, et al N = 469*		Marty, et al N = 186		CLEOPATRA N = 808		AVEREL N = 424	
	-TRAS	+TRAS	-TRAS	+TRAS	-PER	+PER	-BEV	+BEV
PFS/TTP, months	4.6	7.3	6.1	11.7	12.4	18.5	13.7	16.5
OS, months	20	26	23	31	37.6	42.6+	38.3	38.5

BEV, bevacizumab


Slamon D, et al. *N Engl J Med*. 2001;344(11):783-792. Mass RD, et al. *Clin Breast Cancer*. 2005;6(3):240-246.

Marty M, et al. *J Clin Oncol*. 2005;23(19):4265-4274. Baselga J, et al. *N Engl J Med*. 2012;366(2):109-119.

Swain SM, et al. *Lancet Oncol*. 2013;14(6):461-471. Gianni L, et al. *J Clin Oncol*. 2013;31(14):1719-1725.



Refractory HER2+ MBC: EMILIA




- **Primary endpoints**
 - OS
 - PFS by independent review
 - Safety

Verma S, et al. *N Engl J Med*. 2012;367(19):1783-1791.


EMILIA: Results



**Progression-Free Survival
by Independent Review**




Overall Survival: Confirmatory Analysis



- Cardiac dysfunction rare
 - All grades: 3.1% vs 1.8%
 - Grade 3: 0.4% vs 0.2%
- Common toxicities
 - Low grade fatigue, nausea, diarrhea
- Unique toxicities
 - Thrombocytopenia
 - 13% grade 3
 - Modest increase in AST and ALT
- PFS in patients with CNS mets at baseline (n = 95)
 - 12.9 months LAP + CAP vs 26.8 months T-DM1
- New standard following progression on trastuzumab based therapy



TH3RESA: Phase III Trial of T-DM1 vs Treatment of Physician's Choice (TPC)




- Stratification factors
 - World region
 - No. of prior regimens for advanced breast cancer
 - Presence of visceral disease
- Primary endpoints
 - PFS by investigator
 - OS
- Secondary endpoints
 - ORR by investigator
 - Safety


Wildiers H, et al. *Eur J Cancer*. 2013;49(Suppl 3): Abstract LBA15. Krop IE, et al. *Lancet Oncol*. 2014;15(7):689-699.

TH3RESA: PFS and Comparison With Trastuzumab Regimen Beyond Progression

PFS by Investigator Assessment



PFS for Patients Treated With Trastuzumab-Containing Regimens



- PFS: 3 months benefit
- OS: Analysis still early
 - 44 events vs 61 events
 - HR 0.552
 - $P = .0034$ (14.9 vs NE)
- Less toxic than “standard” options

Wildiers H, et al. *Eur J Cancer*. 2013;49(Suppl 3): Abstract LBA15. Krop IE, et al. *Lancet Oncol*. 2014;15(7):689-699.



Selected Ongoing First-Line MBC Trials

- VELVET (n = 210)
 - Trastuzumab/pertuzumab/vinorelbine
- MARIANNE (n = 1092)
 - Comparing 3 options
 - (Trastuzumab + taxane) vs (T-DM1 + pertuzumab) vs (T-DM1 + placebo)
- PERTAIN (n = 250)
 - Comparing an aromatase inhibitor vs chemo up front +/- pertuzumab



Summary of Considerations for Clinical Practice (1)

- Pertuzumab approved in first-line setting in combination with trastuzumab/docetaxel
- T-DM1 an effective treatment for HER2+ MBC progressing on trastuzumab
 - Toxicities are unique but generally well tolerated
- Lapatinib + chemotherapy associated with more toxicity, less efficacy than trastuzumab in first-line metastatic setting




Summary of Considerations for Clinical Practice (2)

- Lapatinib/trastuzumab or capecitabine still an option for later line therapy or in special settings (low EF, brain mets)
- No role for bevacizumab (for systemic disease)
- No clear role for everolimus
- More research needed related to brain mets
- Tumor biopsies, plasma DNA and molecular analysis of CTCs important for research

EF, ejection fraction; CTCs, circulating tumor cells

Considerations for Clinical Practice (3)



Multiple agents in clinical trials: inhibitors of mTOR, HSP90, TKs, vaccines, PD-1, PD-L1



Collaborators

NIH

NCI

BCRF

26.2 with Donna Foundation

Feb 13-15, 2015

5K, relay, half and full marathon

breastcancermarathon.com



Feb 13-14, 2015

CME Course: Advances in Systemic
Therapies for Breast Cancer

Mayo Clinic Simulation Center, Jacksonville, FL

RAISING THE BAR IN BREAST CANCER CARE:

Answering Clinically Relevant Questions

