

# Phase I Study of Carfilzomib and Panobinostat for Patients With Relapsed and Refractory Myeloma: A Multicenter MMRC Clinical Trial

Kaufman JL, Zimmerman T, Rosenbaum CA, Mahindra A, Nooka A, Heffner LT, Harvey RD, Gleason C, Lewis C, Sharp C, Barron K, and Lonial S

# Study Rationale

- Multiple myeloma (MM) remains a disease with a need for new treatments for patients with relapsed and refractory disease
- Carfilzomib is a selective proteasome inhibitor that has demonstrated significant activity in patients with relapsed and refractory myeloma
- Panobinostat is a pan-deacetylase inhibitor that can overcome resistance in combination with bortezomib in refractory multiple myeloma patients.
- Based on data showing synergistic cytotoxicity of pan-deacetylase and proteasome inhibitors, we hypothesized that carfilzomib and panobinostat would be safe and effective for the treatment of relapsed/refractory multiple myeloma.

# **Study Design**

## **Objectives and Assessments**

### **Primary endpoint:**

- **Determine MTD and optimal schedule of the combination**

### **Secondary endpoints:**


- **Adverse event incidence, frequency and severity**
- **Response data including response rates, duration, progression free and overall survival**

### **Multi-center, open-label, Phase 1 study:**

- **Dose escalation phase:**
  - **Standard 3 + 3 design to determine MTD**
  - **MTD based on DLT during cycle 1**
  - **Patients assigned to one of 4 cohorts**
- **Expansion phase:**
  - **12 additional patients treated at the MTD to support the secondary objectives**

# Carfilzomib/Panobinostat

## Dosing schedule



■ PAN: Panobinostat 15-20 mg orally

■ CAR: Carfilzomib 20-45 mg/m<sup>2</sup>; all doses of carfilzomib over 27mg/m<sup>2</sup> were infused over 30 minutes

■ Dex: Dexamethasone 4 mg oral for first cycle, optional thereafter

If 3 out of 4 week dosing was not tolerable a change to 2 of 4 week dosing was planned

# Dose Escalation Schedule

Cohort	Panobinostat (oral TIW 3 out of 4 weeks)	Carfilzomib (IV Days 1/2, 8/9, and 15/16 every 4 weeks)	Dexamethasone (oral prior to carfilzomib)
1	15 mg	C1D1/2: 20 mg/m <sup>2</sup> thereafter: 27 mg/m <sup>2</sup>	4 mg (Cycle 1: thereafter as clinically indicated)
2	20 mg	C1D1/2: 20 mg/m <sup>2</sup> thereafter: 27 mg/m <sup>2</sup>	4 mg (Cycle 1: thereafter as clinically indicated)
3	20 mg	C1D1/2: 20 mg/m <sup>2</sup> thereafter: 36 mg/m <sup>2</sup>	4 mg (Cycle 1: thereafter as clinically indicated)
4	20 mg	C1D1/2: 20 mg/m <sup>2</sup> thereafter: 45 mg/m <sup>2</sup>	4 mg (Cycle 1: thereafter as clinically indicated)

# Patient Characteristics

	N = 26*
Female, %	50
Median age, in years (range)	65 (49-75)
ECOG performance status, %	
0-1 / 2	96 / 4
Bortezomib (BTZ) refractory (%)	16 (62)
IMiD refractory (%)	22 (85)
Median lines of therapy (range)	3 (1-7)
Prior Auto (%)	23 (88)
<hr/>	
<ul style="list-style-type: none"><li>• Patients received multiple prior regimens, including multiple prior BTZ combinations</li><li>• *20 patients in the dose escalation phase and 6 patients in the dose expansion phase</li></ul>	

# Results: Cohorts, DLTs and MTD

Cohort	Evaluable	DLT	Non evaluable
1	3	0	1 - progression
2	3	0	1 - missed doses, non toxicity related
3	6	1 - Acute kidney injury, thrombocytopenia	1 - progression
4	5	1 – thrombocytopenia 2 - diarrhea	

One grade 5 cardiac event (sudden cardiac arrest) in cohort 1 was possibly related to study medications

MTD is Cohort 3: Carfilzomib 20/36 mg/m<sup>2</sup> and panobinostat 20 mg TIW 3 of 4 weeks

# Adverse Events

Occurring in  $\geq 20\%$  (All Grades) of Patients (n = 26)

All	26 (100%)	Pan/Bor/Dex <sup>1</sup>
Nausea	17 ( 65%)	60%
Anemia	14 ( 54%)	47%
Diarrhea	13 ( 50%)	71%
Thrombocytopenia	13 ( 50%)	66%
Fatigue	10 ( 38%)	69%
Vomiting	9 ( 35%)	
Hypokalemia	8 ( 31%)	22%
Pyrexia	8 ( 31%)	
Decreased appetite	7 ( 27%)	
Hypocalcemia	7 ( 27%)	
Insomnia	6 ( 23%)	

1. Richardson et al; Panorama 2; Blood: 122 (14), 2013.



# Adverse Events

## Occurring in $\geq 5\%$ (Grade 3/4) of Patients (n = 26)

AE, regardless of relationship to study drugs	Grade 3/4	
Number of Subjects with at Least One Event	20 (77%)	Pan/ Bor/Dex <sup>1</sup>
Anemia	10 (38%)	15%
Thrombocytopenia	10 (38%)	64%
Neutropenia	5 (19%)	15%
Fatigue	3 (12%)	20%
Decreased appetite	2 ( 8%)	
Diarrhea	2 ( 8%)	20%
Elevated creatinine	2 ( 8%)	
Hyperglycemia	2 ( 8%)	
Hypertension	2 ( 8%)	
Hyponatremia	2 ( 8%)	

1. Richardson et al; Panorama 2; Blood: 122 (14), 2013.

# Relative Dose Intensity

Cohort	1	2	3	4
<b>Carfilzomib</b>	<b>20/27</b>	<b>20/27</b>	<b>20/36</b>	<b>20/45</b>
<b>N</b>	<b>4</b>	<b>4</b>	<b>13</b>	<b>5</b>
<b>Mean (%)</b>	<b>100</b>	<b>96</b>	<b>98</b>	<b>89</b>
<b>Panobinostat</b>	<b>15</b>	<b>20</b>	<b>20</b>	<b>20</b>
<b>Mean (%)</b>	<b>94</b>	<b>92</b>	<b>93</b>	<b>90</b>

# Preliminary Efficacy Data

Best confirmed response	N = 26 (%)	BTZ refractory N = 16 (%)
<b>Overall response (CR + VGPR + PR)</b>	<b>12 (46)</b>	<b>7 (44)</b>
Complete response	1 (4)	1 (6)
VGPR	5 (19)	1 (6)
Partial response	6 (23)	5 (31)
MR	3 (12)	1 (6)
SD	4 (15)	3 (19)
PD	6 (23)	4 (25)

- All responses occurred in the first 2 cycles
- Two patients maintained response for 18 months
- Median DOR is 7.5 months and 8 patients remain on treatment
- 1 patient was not evaluable for response


# Patient Disposition

## Prolonged Study Participation

N = 26	
Patients ongoing	8 (31%)
Off treatment	18 (69%)
In follow-up	12 (46%)
Off study	6 (23%)
Death	5 (19%)
Withdrew consent	0 (0%)
Lost due to follow up	1 (4%)

- Patients have been on study an average of 4.6 months (range <1-20)
- 14 (78%) patients discontinued treatment secondary to progression
- 4 (22%) patients discontinued to AE; 1 death; 2 asthenia; 1 GI toxicity

# Progression-Free Survival Entire Cohort



Median PFS is 11.4 months (95% CI is 6.8-16 months)

With a median f/u of 8.7 months over 80% of patients are alive

# Conclusions & Future Directions

- The combination of carfilzomib and panobinostat is safe and effective with manageable adverse events in patients with relapsed and refractory myeloma
- The MTD of the combination in this study is carfilzomib 36mg/m<sup>2</sup> 3 of 4 weeks and panobinostat at 20 mg TIW 3 of 4 weeks
- No unexpected toxicities were seen
- At the MTD patients were able to tolerate more than 90% of the planned doses
- Tolerance, response rate, DOR, PFS and OS are favorable when compared to similar studies using different dosing strategies, Shah et al (ASH 2012) and Berdeja et al (ASH 2013)