
## Tested Triage Strategies

- There is limited evidence regarding the evaluation of specific best-practice mitigation strategies
- Some strategies with limited evidence include a three-space triage protocol [1,2] and a (two-space) COVID-specific psychiatric unit [3]
  - These strategies require space and staff for development, execution, and evaluation

## Three-Space Triage Protocol



## Source of Infection in Patients Testing Positive for SARS-CoV-2 by Week Implementation of Protocol in full -> Implementation of Protocol in full -> Source of Infection in Patients Testing Positive for SARS-CoV-2 by Week Implementation of Protocol in full -> Week Starting Date (2020) Nosocomial Possible Nosocomial Community

The completion of the protocol was associated with a 9-fold decrease in possible nosocomial infections at New York Presbyterian Hospital.

[1] Brody BD, Parish SJ, Kanellopoulos D, Russ MJ. A COVID-19 testing and triage algorithm for psychiatric units: One hospital's response to the New York region's pandemic. *Psychiatry Research*. 2020;291:113244. doi:10.1016/j.psychres.2020.113244 [2] Brody BD, Shi Z, Shaffer C, et al. Universal COVID-19 testing and a three-space triage protocol is associated with a nine-fold decrease in possible nosocomial infections in an inpatient psychiatric facility. *Psychiatry Research*. 2021;302:114036. doi:10.1016/j.psychres.2021.114036