

EECS 16B CSM

Bryan Ngo

UC Berkeley

2020-11-02

SVD

Definition

EECS 16B
CSM

Bryan Ngo

SVD

$$\mathbf{A} = \sum_{i=1}^r \sigma_i \mathbf{u}_i [\mathbf{v}_i]^\top = \mathbf{U} \mathbf{\Sigma} [\mathbf{V}]^\top = \begin{bmatrix} \mathbf{u}_1 & \mathbf{u}_2 & \cdots & \mathbf{u}_r \end{bmatrix} \begin{bmatrix} \sigma_1 & 0 & \cdots & 0 \\ 0 & \sigma_2 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \sigma_n \end{bmatrix} \begin{bmatrix} [\mathbf{v}_1]^\top \\ [\mathbf{v}_2]^\top \\ \vdots \\ [\mathbf{v}_n]^\top \end{bmatrix} \quad (1)$$


- \mathbf{U} : orthonormal matrix
- $\mathbf{\Sigma}$: diagonal matrix
- $[\mathbf{V}]^\top$: orthonormal matrix

Visualization

EECS 16B
CSM

Bryan Ngo

SVD



Gram-Schmidt Process

EECS 16B
CSM

Bryan Ngo

SVD

$$\mathbf{p}_i = \mathbf{v}_i - \sum_{j=1}^{i-1} ([\mathbf{v}_i]^\top \mathbf{w}_j) \mathbf{w}_j \quad (2)$$

$$\mathbf{w}_i = \frac{\mathbf{p}_i}{\|\mathbf{p}_i\|} \quad (3)$$

- turns set of basis vectors to set of *orthogonal* basis vectors
- systematically removing parallel components of our vector