






Free VEGF-A: sensitivity to Free VEGF half-life
Single dose: initial clearance 2/day, degrader half-life 3 days; 25% co-degradation
1 hour
4 hours
8 hours
12 hours
Untreated steady-state baseline
Free VEGF-A (pg/mL)
8.0
7.8
7.6
7.4
7.2
7.0
6.8
6.6
0
2
4
6
8
10
12
14
Time after treatment starts (days)
Production recalibrated to initial untreated steady state for each scenario; all other parameters held at current defaults.
Vertical scales differ across plots. Model predictions; storage pools are exploratory assumptions.



Free PlGF: sensitivity to Free VEGF half-life
Single dose: initial clearance 2/day, degrader half-life 3 days; 25% co-degradation
1 hour
4 hours
8 hours
12 hours
Untreated steady-state baseline
Free PlGF (pg/mL)
38
37
36
35
34
0
2
4
6
8
10
12
14
Time after treatment starts (days)
Production recalibrated to initial untreated steady state for each scenario; all other parameters held at current defaults.
Vertical scales differ across plots. Model predictions; storage pools are exploratory assumptions.



Total sFLT1: sensitivity to Free VEGF half-life
Single dose: initial clearance 2/day, degrader half-life 3 days; 25% co-degradation
Total sFLT1 (ng/mL)
10
9
8
7
6
5
4
3
0
2
4
6
8
10
12
14
Time after treatment starts (days)
1 hour
4 hours
8 hours
12 hours
Untreated steady-state baseline
Scenario curves overlap: total sFLT1 turnover is unchanged by these ligand parameters.
Production recalibrated to initial untreated steady state for each scenario; all other parameters held at current defaults.
Vertical scales differ across plots. Model predictions; storage pools are exploratory assumptions.



Free VEGF-A: sensitivity to Degrader half-life
Single dose: initial clearance 2/day, exponentially decaying; 25% co-degradation
2 hours
12 hours
1 day
3 days
7 days
Untreated steady-state baseline
Free VEGF-A (pg/mL)
7.1
7.0
6.9
6.8
6.7
6.6
6.5
0
2
4
6
8
10
12
14
Time after treatment starts (days)
Production recalibrated to initial untreated steady state for each scenario; all other parameters held at current defaults.
Vertical scales differ across plots. Model predictions; storage pools are exploratory assumptions.


Free PlGF: sensitivity to Degrader half-life
Single dose: initial clearance 2/day, exponentially decaying; 25% co-degradation
2 hours
12 hours
1 day
3 days
7 days
Untreated steady-state baseline
Free PlGF (pg/mL)
34
35
36
37
38
0
2
4
6
8
10
12
14
Time after treatment starts (days)
Production recalibrated to initial untreated steady state for each scenario; all other parameters held at current defaults.
Vertical scales differ across plots. Model predictions; storage pools are exploratory assumptions.


Total sFLT1: sensitivity to Degrader half-life
Single dose: initial clearance 2/day, exponentially decaying; 25% co-degradation
Total sFLT1 (ng/mL)
Time after treatment starts (days)
2 hours
12 hours
1 day
3 days
7 days
Untreated steady-state baseline
Production recalibrated to initial untreated steady state for each scenario; all other parameters held at current defaults.
Vertical scales differ across plots. Model predictions; storage pools are exploratory assumptions.


Free VEGF-A: sensitivity to Ligand co-degradation
Same fraction for VEGF and PlGF; single dose, initial clearance 2/day, half-life 3 days
25%
50%
75%
Untreated steady-state baseline
Free VEGF-A (pg/mL)
7.0
6.9
6.8
6.7
6.6
6.5
0
2
4
6
8
10
12
14
Time after treatment starts (days)
Production recalibrated to initial untreated steady state for each scenario; all other parameters held at current defaults.
Vertical scales differ across plots. Model predictions; storage pools are exploratory assumptions.


Free PlGF: sensitivity to Ligand co-degradation
Same fraction for VEGF and PlGF; single dose, initial clearance 2/day, half-life 3 days
25%
50%
75%
Untreated steady-state baseline
Free PlGF (pg/mL)
38
37
36
35
34
0
2
4
6
8
10
12
14
Time after treatment starts (days)
Production recalibrated to initial untreated steady state for each scenario; all other parameters held at current defaults.
Vertical scales differ across plots. Model predictions; storage pools are exploratory assumptions.


Total sFLT1: sensitivity to Ligand co-degradation
Same fraction for VEGF and PlGF; single dose, initial clearance 2/day, half-life 3 days
Total sFLT1 (ng/mL)
10
9
8
7
6
5
4
3
0
2
4
6
8
10
12
14
Time after treatment starts (days)
25%
50%
75%
Untreated steady-state baseline
Scenario curves overlap: total sFLT1 turnover is unchanged by these ligand parameters.
Production recalibrated to initial untreated steady state for each scenario; all other parameters held at current defaults.
Vertical scales differ across plots. Model predictions; storage pools are exploratory assumptions.