# sFLT1 degrader model

VEGF-A and PlGF distribution, reversible storage, and ligand fate during sFLT1 removal




**LIGAND FATE DURING COMPLEX DEGRADATION**

25% of VEGF and 25% of PlGF are co-degraded; the remaining ligand returns to its free pool. Free sFLT1 removal carries no ligand.

dC(t) = d(t) × relative activity on bound sFLT1 | Other-bound and storage pools have no direct degrader clearance.

| Starting concentration (pg/mL) | VEGF-A | PlGF |
|---|---|---|
| Free plasma | 6.48 | 33.73 |
| sFLT1-bound / other-bound / stored (each) | 31.17 | 22.09 |
| Plasma total / all-pools total | 68.83 / 100.00 | 77.91 / 100.00 |
| Kd: sFLT1 / other / effective storage (pM) | 20 / 20 / 20 | 150 / 150 / 150 |

### NATURAL LOSS & INPUTS

Every state has first-order natural loss.
Complex loss destroys ligand; other-site capacity is maintained.
qS, qL and qT continue throughout treatment.
Storage also releases at rT (default zero).

Production is calibrated to initial untreated steady state.
Single dose day 0: initial 2/day, half-life 1 day.
Storage and affinity are effective assumptions.

Plasma total = L + C + O; all pools = L + C + O + T. Independent O/T sites for each ligand; only sFLT1 is shared.

Exploratory model, not a validated physiological or assay prediction. Source: model.py defaults; regeneration script: generate_model_schematic.py.