

1 Extra material

1.1 Jensens inequality

Finite case: For a real convex function ϕ

$$\phi\left(\sum_{i=1}^N \theta_i x_i\right) \leq \sum_{i=1}^N \theta_i \phi(x_i)$$

Proof. By induction: $n = 2$ follows from convexity and for $N + 1$:

$$\phi\left(\sum_{i=1}^{N+1} \theta_i x_i\right) = \phi\left(\theta_1 x_1 + (1 - \theta_1) \sum_{i=2}^{N+1} \frac{\theta_i}{1 - \theta_1} x_i\right)$$

By convexity of the case $n = 2$ the result follows. \square

Probability formulation: Let X be some random variable, and let $f(x)$ be a convex function (defined at least on a segment containing the range of X). Then:

$$\mathbb{E}[f(x)] \geq f(\mathbb{E}[X])$$

Proof. Let $c = \mathbb{E}[X]$. Since $f(x)$ is convex there exists a supporting line for $f(x)$ at c :

$$\phi(x) = \alpha(x - c) + f(c)$$

for some α , and $\phi(x) \leq f(x)$. Then

$$\mathbb{E}[f(X)] \geq \mathbb{E}[\phi(X)] = \mathbb{E}[\alpha(X - c) + f(c)] = f(c)$$

\square

1.2 Youngs Inequality

Let $a, b, p, q \in \mathbb{R}_{++}$ such that

$$\frac{1}{p} + \frac{1}{q} = 1$$

then:

$$ab \leq \frac{a^p}{p} + \frac{b^q}{q}$$

Proof.

$$ab = \exp\left(\frac{1}{p} \ln(a^p) + \frac{1}{q} \ln(b^q)\right) \leq \frac{1}{p} \exp(\ln(a^p)) + \frac{1}{q} \exp(\ln(b^q)) = \frac{a^p}{p} + \frac{b^q}{q}$$

where the inequality follows from the convexity of the exponential function. \square

1.2.1 Standard version

Let $f : [0, c] \rightarrow R$ be a continuous and strictly increasing function, where $f(0) = 0$.

$$ab \leq \int_0^a f(x)dx + \int_0^b f^{-1}(x)dx$$

Proof. By visual inspection. □

1.3 Hölders inequality

1.3.1 Standard version

Let (S, \sum, μ) be a measure space and let $q \leq p, q \leq \infty$ with $\frac{1}{p} + \frac{1}{q} = 1$ then for all measurable real- or complex-valued functions f and g on S

$$\|fg\|_1 \leq \|f\|_p \|g\|_q,$$

in the case $p = 2$ this becomes the Cauchy inequality.

Proof. If $\|f\|_t = 0$ for $t = p$ or $t = q$ the left side is zero almost everywhere, so we can assume that $\|f\|_t > 0$. For $\|f\|_t = \infty$ the right hand side could be infinite large. If $t = \infty$ then $|fg| \leq \|f\|_\infty |g|$ almost everywhere and Hölders inequality follows from the monotonicity of the Lebesgue integral (?). Normalizing the equation we can expect

$$\|f\|_p = \|g\|_q = 1.$$

Using Youngs inequality

$$|f(s)g(s)| \leq \frac{|f(s)|^p}{p} + \frac{|g(s)|^q}{q} \quad s \in S$$

Integrating both sides:

$$\|fg\|_1 = \int_S |f(s)g(s)|ds \leq \int_S \left(\frac{|f(s)|^p}{p} + \frac{|g(s)|^q}{q} \right) ds = \frac{\|f\|_p^p}{p} + \frac{\|g\|_q^q}{q} = 1 = \|f\|_p \|g\|_q$$

□

1.3.2 With sums

Using summation to define the norm :

$$\|x\|_p = \left(\sum_k |x_k|^p \right)^{\frac{1}{p}}$$

Proof. Then (without normalization)

$$\frac{|\sum_k x_k y_k|}{\|x\|_p \|y\|_q} \leq \sum_k \frac{|x_k| |y_k|}{\|x\|_p \|y\|_q} \leq \frac{1}{p} \sum_k \frac{|x_k|^p}{\|x\|_p^p} + \frac{1}{q} \sum_k \frac{|y_k|^q}{\|y\|_q^q} = \frac{1}{p} + \frac{1}{q} = 1$$

hence

$$|(x, y)| = \left| \sum_k x_k y_k \right| \leq \|x\|_p \|y\|_q$$

□

The equality holds (proof) iff $\arg x_j y_j$ and $|x_j|^p |y_j|^q$ are independent of j . For any $x \in \ell^p$ we can choose $y \in \ell^q$ so we get equality (proof?). Hence we get

$$\|x\|_p = \max_{\|y\|_q=1} \|(x, y)\|_1$$

1.4 Minkowski

For the ℓ^p spaces (same holds for L^p)

$$\|x + y\|_p \leq \|x\|_p + \|y\|_p$$

Proof. (?) Using the second formulation of Hölders inequality and noting that $\|xy\|_1$ could be written as a scalar product and that scalar products are bilinear

$$\|x + y\|_p = \max_{\|u\|_q=1} |(x + y, u)| \leq \max_{\|u\|_q=1} |(x, u)| + |(y, u)| \leq \|x\|_p + \|y\|_q.$$

Where we in the last inequality split the max and use Hölder inequality together with fact that $\|u\|_q = 1$. □

1.5 Cauchy inequality

A scalar product in a linear space X over \mathbb{C} is a real valued function that fulfils

- sesquilinearity: $(ax, y) = a(x, y)$, $(x, ay) = \bar{a}(x, y)$
- Skew symmetry: $(y, x) = \overline{(x, y)}$
- positivity: $(x, x) > 0$, $x \neq 0$

This also lets us define the norm

$$\|x\| = (x, x)^{\frac{1}{2}}$$

1.5.1 Schwarz Inequality in \mathbb{C}

$$|(x, y)| \leq \|x\| \|y\|$$

(This is just a special case of Hölders inequality with $p = 2$)

Proof. For any $t \in \mathbb{R}$ and $y \neq 0$

$$0 \leq \|x + ty\|^2 = \|x\|^2 + 2t \operatorname{Re}(x, y) + t^2 \|y\|^2 \quad (1)$$

By choosing $t = -\operatorname{Re}(x, y)/\|y\|^2$ and multiplying with $\|y\|^2$

$$(\operatorname{Re}(x, y))^2 \leq \|x\|^2 \|y\|^2 \quad (2)$$

Let $x = ax$, $a \in \mathbb{C}$ such that $|a| = 1$ and $\operatorname{Im}(a(x, y)) = 0$ results in

$$|(x, y)| \leq \|x\| \|y\| \quad (3)$$

This also gives as a corollary that every vector in a scalar product space can be written as

$$\|x\| = \max_{\|y\|=1} \|(x, y)\|_1$$

□

1.6 Parallelogram Identity

Set $t \pm 1$ in (1) and we end up with the parallelogram identity, which is true for all scalar product induced norms.

$$\|x + y\|^2 + \|x - y\|^2 = 2(\|x\|^2 + \|y\|^2) \quad (4)$$

Actually every norm that comes from an scalar product must have this equality (von Neumann)

1.7 Polarization Identity

For a real vector space, if the norm fullfills the Parallelogram Identity then it induces a scalar product

$$(x, y) = \frac{1}{4} (\|x + y\|^2 - \|x - y\|^2) \quad (5)$$

And for the complex vector space

$$(x, y) = \frac{1}{4} (\|x + y\|^2 - \|x - y\|^2 + i\|x + iy\|^2 - i\|x - iy\|^2) \quad (6)$$

2 Finite

2.0.1 Lemma x

Let Y be a subset of the metric space X , then $x \in X$ is adherent to Y iff there is a sequence in Y that converges to x .

Proof. (\Rightarrow) For each $n \geq 1$ there exists some point $x_n \in B(x; 1/n) \cap Y$ the sequence (x_n) then converges to x .

(\Leftarrow) If there is a sequence (x_n) converging to x then all balls centered at x contains points from the sequence. \square

2.0.2 Every subspace of a normed space of finite dimension is closed

Let $(V, |\cdot|)$ be a normed vector space of finite dimension, and $S \subset V$ a finite dimensional subspace. Then S is closed.

Proof. Let $a \in \overline{S}$ and choose a sequence (a_n) , with $a_n \in S$ converging to a ???. Then (a_n) is a cauchy sequence in V and also a cauchy sequence in S . Since V is finite dimensional, V is a Banach space, S is complete, so (a_n) converges to an element in S . Since limits in a normed space are unique, that limits must be a , so $a \in S$. \square

2.0.3 Every finite dimensional subspace of a normed space is closed

Proof. Let $(V, |\cdot|)$ be a normed vector space, and $S \subset V$ a finite dimensional subspace. Then S is closed. Let $x \in V$ and let (a_n) be a sequence in S with converges to x . Because S is of finite dimension, we have a basis $\{x_1 \dots x_k$ of S . Also $x \in \text{span}(x_1 \dots x_k, x)$. But as proved in the case when V is finite dimensional, we have that S is closed in $\text{span}(x_1 \dots x_k, x)$ (taken with the normed induced by $(V, |\cdot|)$) with $a_n \rightarrow x$, and $x \in S$. \square

iiiiiii HEAD ===== något att ändra2, jupps något att ändra2, jupps3
123 llllllll test3