

MacLeish: An R package for monitoring environmental conditions in Whately, Massachusetts

Benjamin S. Baumer¹, Dominique Kelly¹, and Albert Y. Kim¹

1 Smith College, Northampton, Massachusetts, USA

DOI:

Software

- Review 🗗
- Repository ☑
- Archive □

Submitted: Published:

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC-BY).

Summary

The Ada & Archibald MacLeish Field Station is a parcel of land in Whately, MA owned and operated by Smith College. The macleish R package provides spatial, meteorological, and photographic data for monitoring environmental conditions at the site. Numerous faculty use the MacLeish Field Station for research purposes.

The macleish package includes multiple layers of spatial data, functions to access historical and live weather readings, data from ongoing forest ecology research, and functions to access photographs taken by a PhenoCam (Phenology Camera) Network digital camera mounted on the 25-meter WhatelyMet tower. The camera captures a photograph of the forest canopy every 30 minutes.

By delivering data from a variety of sources in a common, easy-to-use interface, the macleish package fosters interdisciplinary collaborations within environmental research communities.

About the MacLeish Field Station

From the Smith College website:

The Ada and Archibald MacLeish Field Station, a uniquely liberal arts field station, is a 250-acre patchwork of forest and pasture land in nearby Whately that provides opportunities for all members of the Smith community (students, faculty, staff, and alum) to pursue artistic inquiry, environmental research, outdoor education and low-impact recreation. Students are integral to the field station: they help steward the land, design programs for their peers and participate in making key decisions in the operation of the field station. The Bechtel Environmental classroom provides meeting and dry laboratory space for MacLeish users.

Figure 1 shows a trail map illustrating the MacLeish property. Moise et al. (2021) provides an interactive history of the property.

Statement of Need

The MacLeish Field Station enables scientific inquiry, as the Smith College community explores the diverse ecosystem of plants and animals. While numerous research projects (see Guswa and Spence (2012), Guswa (2012), Zukswert et al. (2014), Visser et al. (2015), Bassar et al. (2016), Wetzel (2017), Ignace, Fassler, and Bellemare (2018), Jackson and Bellemare (2018) for examples) make use of the MacLeish property, the data supporting these efforts is not centralized or easily accessible to the public. The macleish package offers easy, public access to:






Figure 1: Map of MacLeish Field Station, Whately, Massachusetts, USA.

- 1. spatial data on building, landmarks, streams, trails, and other features of the MacLeish Field Station property
- 2. real-time and historical meteorological measurements
- 3. carefully collected data on tree diameters of different species infected by the Hemlock Woolly adelgid
- 4. photographic images of the tree canopy over time

These data are already used in data science textbooks (Benjamin S. Baumer, Kaplan, and Horton 2021) and new research efforts in progress.

Features

To use the CRAN version of the macleish package, first run install.packages(macleish) to install the package, then load it:

library(macleish)

Spatial data

Spatial data from the MacLeish Field Station are available through the macleish_layers data object, which leverages the sf package (Pebesma 2022). The following layers are available:

```
names(macleish_layers)

## [1] "landmarks" "forests" "streams"

## [4] "challenge_courses" "buildings" "wetlands"

## [7] "boundary" "research" "soil"

## [10] "trails" "camp_sites" "elevation"
```

Any sf objects can be placed on an interactive map created by the leaflet package (Cheng, Karambelkar, and Xie 2022). Figure 2 situates the MacLeish property in Whately,





Figure 2: Screenshot from an interactive 'leaflet' map of the MacLeish Field Station.

to the east of the Northampton Reservoir and to the west of the Mt. Esther Wildlife Management Area, pinpoints the OrchardMet and WhatelyMet weather stations, and depicts other geographical features such as streams and trails. Figure 2 displays a static image of the resulting interactive map.

```
library(leaflet)
map <- leaflet() |>
  addTiles() |>
  addPolygons(
    data = macleish_layers[["boundary"]],
    weight = 1, fillOpacity = 0.1
  ) |>
  addPolygons(
    data = macleish_layers[["buildings"]],
    weight = 1, popup = ~name
  ) |>
  addPolylines(
    data = macleish_layers[["trails"]],
    weight = 1, color = "brown",
    popup = ~name
  ) |>
  addPolylines(
    data = macleish_layers[["streams"]],
    weight = 2
  ) |>
  addMarkers(
    data = filter(macleish_layers[["landmarks"]], grepl("Met", Label)),
    popup = ~Label
  setView(-72.67918, 42.44921, zoom = 14)
map
```

Research data

Two tree plot datasets are available: tree_diameter1 and tree_diameter2. The time frame differs between tree_diameter1 (2010–2015) and tree_diameter2 (2009–2016), and tree_diameter2 lacks data on position of individual trees. The datasets present change in tree diameter over a set of years for two plots dominated by the parasitic Hemlock Woolly Adelgid. Tables 1 and 2 show samples from these datasets.



tree diameter1

Table 1: Sample of the first six rows of the first tree diameter dataset.

tag	species	position	semester	year	diameter	notes
2001	Sweet Birch	subcanopy	Fall	2015	NA	Alive in 2009, dead in 2010
2001	Sweet Birch	subcanopy	Fall	2014	NA	Alive in 2009 , dead in 2010
2001	Sweet Birch	subcanopy	Fall	2013	NA	Alive in 2009, dead in 2010
2001	Sweet Birch	subcanopy	Fall	2012	NA	Alive in 2009, dead in 2010
2001	Sweet Birch	subcanopy	Fall	2011	NA	Alive in 2009, dead in 2010
2001	Sweet Birch	subcanopy	Fall	2010	NA	Alive in 2009 , dead in 2010

tree_diameter2

Table 2: Sample of the first six rows of the second tree diameter dataset.

tag	species	semester	year	diameter	notes
2171	Eastern Hemlock	Fall	2016	28.2	NA
2171	Eastern Hemlock	Fall	2012	27.4	NA
2171	Eastern Hemlock	Fall	2011	27.2	NA
2171	Eastern Hemlock	Fall	2010	27.1	NA
2171	Eastern Hemlock	Fall	2009	27.0	NA
2172	Sweet Birch	Fall	2016	24.5	NA

The boxplot in Figure 3 uses the tree_diameter1 data set to demonstrate the distribution of diameter for different species of trees at different positions from 2010–2015.

By viewing an individual species, the distribution of diameter varies based on the position of the species. For instance, the median diameter of emergent Eastern Hemlock trees is the greatest, despite having the smallest overall range. However, the overall range of the diameter of the canopy is the greatest.

Weather data

Weather data from 2015 is available through the whately_2015 and orchard_2015 data objects. These data were recorded every 10 minutes every day from 2015-01-01 to 2015-31-12, and include measurements like temperature, wind speed, and relative humidity.

```
head(whately_2015)
## # A tibble: 6 x 8
##
     when
                          temperature wind speed wind dir rel humidity
##
     <dttm>
                                <dbl>
                                            <dbl>
                                                     <db1>
                                                                   <db1>
## 1 2015-01-01 00:00:00
                                -9.32
                                                      225.
                                                                   54.6
                                            1.40
## 2 2015-01-01 00:10:00
                                -9.46
                                            1.51
                                                      248.
                                                                   55.4
## 3 2015-01-01 00:20:00
                                -9.44
                                            1.62
                                                      258.
                                                                   56.2
## 4 2015-01-01 00:30:00
                                -9.3
                                            1.14
                                                      244.
                                                                   56.4
## 5 2015-01-01 00:40:00
                                -9.32
                                            1.22
                                                      238.
                                                                   56.9
## 6 2015-01-01 00:50:00
                                -9.34
                                             1.09
                                                      242.
                                                                   57.2
## # ... with 3 more variables: pressure <int>, solar_radiation <dbl>,
     rainfall <dbl>
```

Furthermore, live and historical (to present from 2012-01-03 for WhatelyMet and from 2014-06-27 for OrchardMet) weather data from both MacLeish Field Station weather monitors is available for download through functionality provided by the etl package (Benjamin S. Baumer 2021; Benjamin S. Baumer 2019).



Distribution of Diameter Between 2010-2015




Figure 3: Boxplot of tree diameter depending on species and position between 2010–2015.



Figure 4: Images of the MacLeish Field Station retrieved from the PhenoCam Network by the phenocam_image_url() function. The image on the left was taken in June 2022, while the image on the right was taken in December 2021.

Photographic data

The PhenoCam Network is an ecosystem phenology camera network that provides automated, near-surface remote sensing of canopy phenology across the northeastern United States and Canada. One such high-resolution digital camera sits atop the WhatelyMet tower, takes pictures of the canopy every 30 minutes, and then uploads them to the PhenoCam server. The PhenoCam Network permits analysis of archived images, providing an objective means by which canopy phenology can be monitored and quantified at relatively low cost and with minimal expenses.

Figure 4 displays two examples of images retrieved from the PhenoCam Network by the phenocam_image_url() function, which leverages functionality provided by the phenocamr package (Koen 2020).

Acknowledgements

We acknowledge a variety of contributions from Jesse Bellemare, Paul Wetzel, Reid Bertone-Johnson, Jon Caris, Rose Goueth, Wencong Li, Weijia Zhang, and Nicholas Hor-



ton.

References

- Bassar, Ronald D, Benjamin H Letcher, Keith H Nislow, and Andrew R Whiteley. 2016. "Changes in Seasonal Climate Outpace Compensatory Density-Dependence in Eastern Brook Trout." Global Change Biology 22 (2): 577–93. https://doi.org/10.1111/gcb.13135.
- Baumer, Benjamin S. 2019. "A Grammar for Reproducible and Painless Extract-Transform-Load Operations on Medium Data." *Journal of Computational and Graphical Statistics* 28 (2): 256–64. https://doi.org/10.1080/10618600.2018.1512867.
- Baumer, Benjamin S. 2021. Etl: Extract-Transform-Load Framework for Medium Data. https://github.com/beanumber/etl.
- Baumer, Benjamin S., Daniel T. Kaplan, and Nicholas J. Horton. 2021. *Modern Data Science with R.* 2nd ed. Chapman; Hall/CRC Press: Boca Raton. https://doi.org/10.1201/9780429200717.
- Cheng, Joe, Bhaskar Karambelkar, and Yihui Xie. 2022. Leaflet: Create Interactive Web Maps with the JavaScript Leaflet Library. https://rstudio.github.io/leaflet/.
- Guswa, Andrew J. 2012. "Canopy Vs. Roots: Production and Destruction of Variability in Soil Moisture and Hydrologic Fluxes." *Vadose Zone Journal* 11 (3). https://doi.org/10.2136/vzj2011.0159.
- Guswa, Andrew J, and Caitlin M Spence. 2012. "Effect of Throughfall Variability on Recharge: Application to Hemlock and Deciduous Forests in Western Massachusetts." *Ecohydrology* 5 (5): 563–74. https://doi.org/10.1002/eco.281.
- Ignace, Danielle D, Aliza Fassler, and Jesse Bellemare. 2018. "Decline of a Foundation Tree Species Due to Invasive Insects Will Trigger Net Release of Soil Organic Carbon." *Ecosphere* 9 (8): e02391. https://doi.org/10.1002/ecs2.2391.
- Jackson, Michelle R, and Jesse Bellemare. 2018. "The Potential for Indirect Negative Effects of Exotic Insect Species on a Liverwort, Bazzania Trilobata (Lepidoziaceae), Mediated by the Decline of a Foundation Tree Species, Tsuga Canadensis (Pinaceae) 1." The Journal of the Torrey Botanical Society 145 (3): 183–94. https://doi.org/10.3159/TORREY-D-17-00040.1.
- Koen, Hufkens. 2020. Phenocamr: Facilitates PhenoCam Data Access and Time Series Post-Processing. https://CRAN.R-project.org/package=phenocamr.
- Moise, Arris, Arris, Sadie Wiese, and Clara Sorensen. 2021. "Land Use History of MacLeish: A Cultural and Ecological Timeline of the Smith College Field Station." ArcGIS StoryMaps. https://storymaps.arcgis.com/stories/14c7392dbd4e42d4a4ef7775cb56062c.
- Pebesma, Edzer. 2022. Sf: Simple Features for r. https://CRAN.R-project.org/package=sf.
- Visser, Marco D, Sean M McMahon, Cory Merow, Philip M Dixon, Sydne Record, and Eelke Jongejans. 2015. "Speeding up Ecological and Evolutionary Computations in r; Essentials of High Performance Computing for Biologists." *PLoS Computational Biology* 11 (3): e1004140. https://doi.org/10.1371/journal.pcbi.1004140.
- Wetzel, Paul. 2017. "2016 Annual Weather Report: Smith College Ada & Archibald MacLeish Field Station Whately, Massachusetts, USA." Smith College. https://docs.google.com/document/d/1MZc7F7fP5iBrJrjrF6_Ovg7FfCs7WoIk_UIFPB4aAcE/edit?usp=sharing.
- Zukswert, Jenna M, Jesse Bellemare, Amy L Rhodes, Theo Sweezy, Meredith Gallogly, Stephanie Acevedo, and Rebecca S Taylor. 2014. "Forest Community Structure Differs, but Not Ecosystem Processes, 25 Years After Eastern Hemlock Removal in an Accidental Experiment." Southeastern Naturalist 13: 61–87. https://www.jstor.org/stable/26454958.