

OpenCMISS-iron examples and tests used by *OpenCMISS* developers at University of Stuttgart, Germany

Ekin Altan*, Christian Bleiler*, Andreas Hessenthaler*,
Thomas Klotz*, Aaron Krämer[†], Benjamin Maier[‡],
Sergio Morales*, Mylena Mordhorst*, Harry Saini*

June 19, 2017
18:37

CONTENTS

1	Introduction	3
1.1	Cmgui files for cmgui-2.9	3
1.2	Variations to consider	3
1.3	Folder structure	4
2	How to work on this document	4
3	Diffusion equation	7
3.1	Equation in general form	7
3.2	Example-0001	7
3.2.1	Model	7
3.2.2	Results	7
4	Linear elasticity	8
4.1	Equation in general form	8
4.2	Example-0101	8
4.2.1	Model	8
4.2.2	Results	8
5	Finite elasticity	9
6	Navier-Stokes flow	10

LIST OF FIGURES

Figure 1	Results.	7
Figure 2	Results.	8

* Institute of Applied Mechanics (CE), University of Stuttgart, Pfaffenwaldring 7, 70569 Stuttgart, Germany

[†] Institute for Parallel and Distributed Systems, University of Stuttgart, Universitätsstraße 38, 70569 Stuttgart, Germany

[‡] Lehrstuhl Mathematische Methoden für komplexe Simulation der Naturwissenschaft und Technik, University of Stuttgart, Allmandring 5b, 70569 Stuttgart, Germany

LIST OF TABLES

Table 1	Initials of people working on examples, in alphabetical order (surnames).	4
Table 2	Example-0001 to example-0099. Who is doing what? What is finished? See Table 1.	5
Table 3	Example-0101 to example-0199. Who is doing what? What is finished? See Table 1.	6

1 INTRODUCTION

This document contains information about examples used for testing *OpenCMISS-iron*. Read: How-to¹ and [1].

1.1 Cmgui files for cmgui-2.9

1.2 Variations to consider

- Geometry and topology
 - Dimensions
 - Extents
 - Number of elements
 - Interpolation order
 - Generated or user meshes
 - quad/hex or tri/tet meshes
- Initial conditions
- Load cases
 - Dirichlet BC
 - Neumann BC
 - Volume force
 - Mix of previous items
- Sources, sinks
- Time dependence
 - Static
 - Quasi-static
 - Dynamic
- Material laws
 - Linear
 - Nonlinear
- Material parameters, anisotropy
- Solver
 - Direct
 - Iterative
- Test cases
 - Numerical reference data
 - Analytical solution
- A mix of previous items

¹ <https://bitbucket.org/hessenthaler/opencmisshowto>

1.3 Folder structure

TBD..

2 HOW TO WORK ON THIS DOCUMENT

In this section, indicate what you are working on or if a given example was finished

- no mark: to be done
- x: currently working on it
- xx: done

Initials	Full name
EA	Ekin Altan
CB	Christian Bleiler
AH	Andreas Hessenthaler
TK	Thomas Klotz
AK	Aaron Krämer
BM	Benjamin Maier
SM	Sergio Morales
MM	Mylena Mordhorst
HS	Harry Saini

Table 1: Initials of people working on examples, in alphabetical order (surnames).

Progress	Initials	Name of example	Dimensions	Equation	Boundary conditions	Mesh type	Interpolation order
x	AH	Example-0001	3D	Laplace	Dirichlet	generated	linear
		Example-0002					
		...					

Table 2: Example-0001 to example-0099. Who is doing what? What is finished? See Table 1.

Progress	Initials	Name of example	Dimensions	Equation	Boundary conditions	Mesh type	Interpolation order
x	AH, HS	Example-0101	2D	Elasticity	Dirichlet	generated	linear
		Example-0102					
		...					

Table 3: Example-0101 to example-0199. Who is doing what? What is finished? See Table 1.

3 DIFFUSION EQUATION

3.1 Equation in general form

$$\partial_t u + \nabla \cdot \nabla u = f \quad (1)$$

3.2 Example-0001

3.2.1 Model

We solve the following equation,

$$\nabla \cdot \nabla u = 0 \quad \Omega \in [0, 2] \times [0, 1] \times [0, 1], \quad (2)$$

with boundary conditions

$$u = 0 \quad x = y = z = 0, \quad (3)$$

$$u = 0 \quad x = 2, y = z = 1. \quad (4)$$

No material parameters to specify.

3.2.2 Results




Figure 1: Results.

4 LINEAR ELASTICITY

4.1 Equation in general form

$$\partial_{tt}\mathbf{u} + \nabla \cdot \boldsymbol{\sigma}(\mathbf{u},t) = \mathbf{f}(\mathbf{u},t)$$
 (5)

4.2 Example-0101

4.2.1 Model

We solve the following equation,

$$\nabla \cdot \boldsymbol{\sigma}(\mathbf{u},t) = \mathbf{0}(\mathbf{u},t) \qquad \Omega \in [0,160] \times [0,120], t \in [0,5],$$
 (6)

with time step size $\Delta_t = 1$ and boundary conditions

$$\dots \qquad \dots,$$
 (7)

$$\dots \qquad \dots$$
 (8)

2D: specify thickness, Young's modulus and Poisson's ratio.

4.2.2 Results




Figure 2: Results.

5 FINITE ELASTICITY

6 NAVIER-STOKES FLOW

REFERENCES

- [1] Chris Bradley, Andy Bowery, Randall Britten, Vincent Budelmann, Oscar Camara, Richard Christie, Andrew Cookson, Alejandro F Frangi, Thiranjia Babarenda Gamage, Thomas Heidlauf, et al. Openmiss: a multi-physics & multi-scale computational infrastructure for the vph/-physiome project. *Progress in biophysics and molecular biology*, 107(1):32–47, 2011.