



Specimen Label

- May be used on Livestock Animals and Livestock Premises
- May be used in Food Handling Establishments
- Kills Bed Bugs
- Kills Bees and Wasps [Yellow Jackets, Hornets and Paper Wasps]

**ACTIVE INGREDIENTS:**

*Pyrethrins.....	1.00%
OTHER INGREDIENTS .....	99.00%
	100.00%

\*Pyrethrins are a botanical insecticide derived from chrysanthemum flowers.

0.01 lbs. active ingredient per lbs. of Evergreen® Pyrethrum Dust



KEEP OUT OF REACH OF CHILDREN

**CAUTION**

See inside for first aid and precautionary statements.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ ENTIRE LABEL BEFORE EACH USE**

**USE RESTRICTIONS:**

- Apply this product only as specified on the label.
- Aerial applications are prohibited.
- Do not allow contamination of food and feedstuffs.
- In indoor residential areas or other areas where children may come into contact, only apply as a crack-and-crevice or as a spot treatment to areas no greater than 3 feet by 3 feet per room.
- Applications with power duster equipment are prohibited.
- Do not enter or allow others to enter until dusts have settled.
- Do not apply this product in a way that will contact workers or other persons, either directly or indirectly through drift.
- Only protected handlers may enter treated area during application.
- Do not allow adults, children or pets to enter the treated area until dusts have settled.
- For use outdoors, do not apply directly to or near water, storm drains or drainage ditches. Do not apply when windy.
- Do not water the treated area to the point of runoff.
- Do not make applications during rain.
- Do not apply this product in a way that will contact adults, children or pets, either directly or indirectly through drift.
- Remove or cover exposed food and drinking water before application.
- Remove or cover dishes, utensils, food processing equipment and food preparation surfaces, or wash them before use. Do not use in serving areas while food is exposed.

• In Food Areas of Food Handling Establishments: Do not contaminate food or feedstuffs. During applications, cover or remove food. Cover food processing surfaces and equipment or clean after treatment with a suitable detergent and rinse with potable water before use. Do not use in serving areas (dining areas where prepared foods are served) while food is exposed or facility is in operation.

- In Schools, do not apply to classrooms when in use.
- Do not use in aircraft cabins.
- Remove pets, birds and cover fish aquariums before application.
- When used in dairy barns or facilities; close milk bulk tank lids to prevent contamination from spray and from dead or falling insects. Remove or cover milking utensils before application. Wash teats of animals before milking.
- Except when applying to sewers or drains as instructed on this label, do not apply directly into sewers or drains, or to any area like a gutter where drainage to storm sewers, water bodies, or aquatic habitat can occur. Do not allow the product any drain during or after application.

- Not for use on commercial sod farms.
- **USE RESTRICTIONS IN FOOD HANDLING AREAS:**
- Do not apply when food or feed processing facility is in operation or foods are exposed.
- **APPLICATIONS OF THIS PRODUCT IN FOOD/FEED AREAS OF FOOD/FEE**
- DHANDLING ESTABLISHMENTS ARE NOT PERMITTED, OTHER THAN AS A CRACK-AND-CREVICE TREATMENT.**

**Phytotoxicity:**

Plant safety is an important consideration when using insecticides. It is not possible to evaluate the phytotoxicity of Evergreen® Pyrethrum Dust on numerous plant varieties that may react differently to insecticides in different growth stages or under varying environmental conditions. Before making widespread applications of Evergreen® Pyrethrum Dust treat a limited number of plants and observe for phytotoxicity over a 10-day period.

**Greenhouses:**

This product can be used in greenhouses to kill listed insects. Apply to cover upper and lower surfaces of leaves stems and flowers with a thin even film of dust. Apply Evergreen® Pyrethrum Dust to ornamental plants or only to fruit, vegetable and herb crops in the plug stage prior to transplant. Do not apply to fruit, vegetable, or herb crops that are ready for consumption.

**OUTDOOR AREAS for use in and around areas:**

Ant Mounds, Arborets, Backyards, Balconies, Beneath Decks, Beneath Gazebos, Calf Hutches, Compost Sites, Courtyards, Decks, Dog Runs, Drive-In Restaurants, Drive-In Theaters, Dumpsters, Farmers Markets, Flea Markets, Front Yards, Gardens, Gazebos, Home Gardens, Kennels, Lanais, Landscapes, Lath Houses, Loading Docks, Lumber Storage Areas, Nurseries, Outdoor Eating Facilities, Patios, Porches, Sanitary Sewers (Manholes), Side Yards, Swine Yards and Wood Piles.

OUTDOOR APPLICATIONS	
Application Type	Application Rate
Asian Lady Beetles and Box Elder Bugs – Perimeter Building Exterior	16 to 24 oz. / 1,000 sq. ft.
Ornamental Plants	Apply to cover upper and lower surfaces of leaves, stems and flowers, with a thin even film of dust.
Insects Harboring in Lawns, Landscaping and Backyards	Apply 8 to 24 oz. / 1,000 sq. ft.
Building Perimeter Treatments	Apply 16 to 24 oz. / 1,000 sq. ft.

**Building Perimeter Treatments:**

To aid in preventing entry and infestations of buildings by listed insects, apply Evergreen® Pyrethrum Dust to building exterior as a perimeter treatment in a band extending 12"-18" along and adjacent to foundation, and to a height of 2 to 3 feet up the side of the building where these pests are active and may find entrance.

**To Kill Insects Harboring in Lawns, Landscaping, and Backyards:**

To aid in the kill of listed insects, apply Evergreen® Pyrethrum Dust at the specified rate. For best results, mow lawn one to two (1-2) days before applying, and remove clippings. Be sure to cover the entire lawn surface where insects are present, or normally frequent. Apply around the base of ornamental plants, shrubs and trees. Do not water or irrigate immediately following application. Keep pets and children out of the treated area until after the dust has settled.

**LIVESTOCK, DAIRY, SWINE, POULTRY AND OTHER ANIMAL BUILDINGS AND PREMISES**

LIVESTOCK PREMISES	
Application Type	Application Rate
Crack and Crevice	1 oz. / 50 sq. ft.
Spot Treatments and Surface Treatments	16 to 24 oz. / 1,000 sq. ft.
Building Perimeter Treatments	16 to 24 oz. / 1,000 sq. ft.
Spot Treatments for Ticks	16 to 24 oz. / 1,000 sq. ft.

Kills: Bed Bugs, Blacklegged Tick Nymphs, Chicken Mites, Deer Ticks, Face Flies, Fleas, Flies, Northern Fowl Mites, Poultry Lice and Stable Flies

To aid in the kill of listed insects in listed indoor and outdoor animal areas, apply Evergreen® Pyrethrum Dust to cracks, crevices, voids, structural joints, floors, walls, wall joints, support beams, trusses, rafters, ceilings, around scales, chutes, gates, and other areas where insects harbor, alight, hide or travel. For Ticks, apply Evergreen® Pyrethrum Dust in areas where they like to harbor such as cracks and crevices.

Apply Evergreen® Pyrethrum Dust to exterior of livestock buildings in an 18" band along and adjacent to building foundations, and to a height of 2 to 3 feet up the side of the building where insects are active and may find entrance.

**LIVESTOCK ANIMAL USE**

LIVESTOCK ANIMALS	
Application Type	Application Rate
Beef, Dairy Cattle, Horses	2 oz. dust per animal
Swine	1 oz. per head as a uniform coat to head, shoulders and back.
Poultry	1 lb. / 100 birds.

Kills: Bed Bugs, Blacklegged Tick Nymphs, Chicken Mites, Deer Ticks, Face Flies, Fleas, Flies, Northern Fowl Mites, Poultry Lice and Stable Flies

**Beef, Dairy, Cattle, Horses and other Livestock:**

Can be used in dust bags, shaker can and mechanical dust applicator to aid in the kill or repellency of listed livestock insects.

**Direct Application:** Apply 2 oz. of dust per animal by shaker can, or suitable mechanical dust applicator over the head, neck, shoulders, back and legs.

**Application through Dust Bags:** Place contents of this package in any commercially available dust bag, suspend bag in areas frequented by cattle or in gateways or lanes through which the animals must pass daily for water, feed or minerals. Bags may also be placed in loafing sheds or in front of mineral feeders. For dairy cows, bags may be suspended in the exit through which the cows leave the milking barn. Hang the bags 4 to 6 inches below the back line of the cattle. For reduction of Face Flies, bags must be located so animals will be forced to use them daily and hung at a height so that the face is dusted.

**Swine:**

## USE RESTRICTIONS FOR APPLICATION TO SWINE

- Do not apply more often than once every 10 days
- Do not ship animals for slaughter within 5 days of treatment.

**Direct Application:** Apply only 1 oz. per head as a uniform coat to the head, shoulder and back by use of a shaker can or suitable mechanical dust applicator. In severe infestation, both animals and the bedding may be treated.

**Poultry:**

To kill listed livestock insects, apply at a rate of 1 lb. per 100 birds.

**STORAGE AND DISPOSAL**

Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE STORAGE:**

Keep this product in its tightly closed original container, when not in use. Do not store near heat or open flame. Store in a cool, dry, well-ventilated (preferably locked) area that is inaccessible to children and animals, and away from food and pet food.

**PESTICIDE DISPOSAL:**

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

**CONTAINER DISPOSAL:**

Nonrefillable container. Do not reuse or refill this container. Completely empty container by shaking and tapping sides and bottom to loosen clinging particles promptly after emptying. Then offer for recycling if available or reconditioning, if appropriate or dispose of in a sanitary landfill or by other procedures approved by State and Local authorities.

**DIRECTIONS FOR USE CONTINUED****AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval of 12 hours.

PPE required for early entry to treated areas (that is permitted under Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water), is:

- Long-sleeved shirt and long pants;
- Shoes plus socks;
- chemical-resistant gloves made of any waterproof material.

**NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR, Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter until dusts have settled.

**Kills Exposed Listed Insects:**

Adelgids	Brown Spider Beetles	Crickets	Flying Moths	Larger Cabinet Beetles	Pine Engraver	Southern Pine Beetles
Alfalfa Weevil	Brown Spruce Longhorn Beetle	Cross-striped Cabbageworm	Foreign Grain Beetles	Pink Scavenger Caterpillars	Southern Red Mite	(Adult)
Almond Moth	Budworms	Cucumber Beetles	Forest Tent Caterpillar	Larger Grain Borers	Pinworms	Sowbugs
American Black Flour Beetle	Cadelles	Cutworms (exposed)	Fruit Flies	Leafhoppers	Plant Bugs	Spider Beetles
Angoumois Grain Moths	California Oakworm	Dark Mealworms	Fruittree Leafroller	Leafminers	Plum Curculio	Spider Mites
Ants	Cankerworms	Darkling Beetles (lesser meal worm)	Fungus Beetles	Leafrollers	Ponderosa Pine Needle Miner	Spruce Mite (Adult)
Aphids	Carpet Beetles	Deer Tick	Fungus Gnats	Leaftiers	Poultry Lice	Squarenecked Grain Beetles
Apple Maggot	Carrot Weevil	Depressed Flour Beetles	German Cockroaches	Lesser Grain Beetles	Lesser Grain Borers	Powder Post Beetles
Armyworms	Caterpillars	Diamondback Larvae	Giant Root Aphid	Giant Root Aphid	Lesser Meal Worm	Stable Flies
Asian Cerambycid Beetles	Catorama Beetles	Dermestid Beetles	Glassy Winged Sharpshooter	Lesser Peach Tree Borer	Lesser Peach Tree Borer	Stalk Borers
Asian Lady Beetle	Centipedes	Dingy Cutworm Adult	Grain Mites	Longheaded Flour Beetles	Longheaded Flour Beetles	Stink Bugs
Asian Pine Tip Moths	Cereal Beetles	Douglas Fir Midge	Granary Insects	Maize Weevils	Red Flour Beetles	Stored Product Pests
Bagworm	Cheese Mites	Douglas Fir Tussock Moth	Granary Weevils	Loopers	Red Horned Grain Beetles	Subtropical Pine Tip Moth
Barn Flies	Cheese Skippers	Drain Flies	Grape Leaf Skeletonizer	Lugus Bugs	Red Lily Beetles	Sycamore Leaf Beetle
Bat Bugs	Chicken Mites	Dried Fruit Beetles	Grape Leafhopper	Maize Weevils	Red Oak Borer	Tarnished Plant Bug
Bean Beetles	Chocolate Moths	Diamondback Larvae	Grasshoppers	Mealy Bugs	Mediterranean Flour Beetles	Tent Caterpillar
Bean Weevils	Chrysanthemum Leaf Miners	Drugstore Beetles	Green Fruit Worm	Merchant Grain Beetles	Rice Beetles	Thrips
Bed Bugs	Cicadas	Earwigs	Green Peach Aphid	Mexican Bean Beetle	Rice Moths	Tobacco Moths
Bees	Cigarette Beetle	Eastern Tent Caterpillar	Greenbugs	Mexican Grain Beetles	Rice Weevils	Trogoderma Beetles
Beet Armyworms	Clearwing Borer	Elmleaf Beetle	Ground Beetles	Millers	Root Weevils	Two-Banded Beetles
Beetles	Clothes Moths	Emerald Ash Borer	Gypsy Moth (adults & larvae)	Millipedes	Rose Chafer	Vegetable Leaf Miners
Birch Leafminer	Clover Mites	European Grain Moths	Hairy Fungus Beetles	Mimosa Webworm	Royal Palm Bugs	Vinegar Flies
Black Carpet Beetles	Clover Weevil	European Pine Shoot Moth	Hairy Spider Beetles	Mole Crickets	Rusty Grain Beetles	Warehouse Beetles
Black Fungus Beetles	Cluster Flies	Coffee Bean Weevils	Harlequin Bug	Moth Flies	Mimosa Webworm	Wasps [Yellow Jackets, Hornets and Paper Wasps]
Black Turpentine Beetles	Cocoa Bean Moths	Cone Beetles	European Pine Tip Moth	Murmidius Beetles	Mole Crickets	Seedbugs
Black Vine Weevil	Cone Worms	Face Flies	Hide Beetles	Mushroom Flies	Nantucket Pine Tip Beetle	Webworms
Blacklegged Tick Nymphs	Confused Flour Beetles	Fall Cankerworms	Hornworm	Onion Maggot	Onion Maggot	Weevils
Blister Beetles	Conifer Spider Mite	Fall Webworm	Inchworm	Palmetto Bug	Pear Psylla	Western Pine Tip Moth
Blow Flies	Coniferous Beetles	Fannia Flies	Indian Meal Moth	Peach Twig Borer	Small Flying Moths	White Apple Leafroller
Booklice	(Adult)	Field Crickets	Iris Borer	Pine Bark Beetles	Small Fruit Flies	White Marked Spider Beetles
Bottle Flies	Corn Sap Beetles	Firebrats	Japanese Beetle	Pillbugs	Smalleyed Flour Beetles	Whiteflies
Box Elder Bugs	Cottonwood Borer	Fireworms	Katydid	Pine Bark Beetles	Wireworms and Yellow Mealworms	
Broadhorned Flour Beetles	Cottonwood Leaf Beetles	Flat Grain Beetles	Khapra Beetles			
Broadnosed Grain Weevils	Cranberry Girdlers	Flea Beetles	Lace Bugs			
	Crane Flies	Fleas	Large Black Flour Beetles			
		Flies	Beetles			

**Application Methods:**

Apply Evergreen® Pyrethrum Dust using a non-power dust applicator, a shaker can, dust gun, puffer or other bulbous duster, hand pump duster, hand-operated rotary duster, static charge duster, or other appropriate hand-operated duster.

**INDOOR AREAS: For use in and around indoor areas:**

Animal Areas	Coffee Houses	Egg Processing Plants	Goat Houses	Livestock Hauling Equipment	Pet Food Plants	Swine Houses
Animal Quarters	Coffee Shops	Egg Storage Facilities	Grain Elevators	Livestock Holding Areas	Pet Sleeping Areas	Taverns
Apartments	Commercial Buildings	Factories	Grain Handling Equipment	Loafing Sheds	Poultry Houses	Tea Houses
Asian Markets	Condominiums	False Ceilings	Grain Harvesting Equipment	Meat Markets	Poultry Processing Plants	Textile Mills
Attics	Convenience Stores	Farm Structures	Grain Mills	Meat Packaging Plants	Rabbit Houses	Tobacco Factories
Bakeries	Crawl Spaces	Feed Mills	Grain Storage	Meat Packing Plants	Rabbit Processing Plants	Tobacco Warehouses
Barns	Dairies	Flour Mills	Granaries	Meat Packing Plants	Railroad Cars	Tool Sheds
Basements	Dairy and Livestock Barns	Food Handling Establishments	Greenhouses	Mess Halls	Research Animal Areas	Trains
Beef Barns	Dairies	Food Preparation Facilities	Grocery Stores	Milk Houses	Restaurants	Transportation Equipment (except Aircraft)
Beverage Plants	Dairy Food Processing	Food Processing Facilities	Hog Barns	Milk Processing Facilities	Rice Mills	Tree Nuts Storage Areas
Bins	Day Care Centers	Food Quarantine Areas	Homes	Milk Rooms	Schools	Truck Trailers
Bottling Plants	Delicatessens	Food Quarantine Areas	Horse Stables	Milk Sheds	Seed Storage	Vegetable Packing Facilities
Brandy Storage Warehouses	Department Stores	Food Quarantine Areas	Horse Stables and Barns	Milking Rooms	Shade Houses	Vegetable Processing Facilities
Breweries	Diners	Food Service Facilities	Horse Trailers	Mills	Sheep Holding Areas	Vegetable Storage Facilities
Brewery Warehouses	Distilleries	Food Storage Facilities	Hospitals	Motels	Ship Holds	Veterinary Clinics
Buildings Housing Animals	Dog Houses	Food Storage Facilities	Hotels	Mushroom Caves	Ships	Veterinary Hospitals
Buildings Housing Poultry	Dog Sleeping Quarters	Freight Containers	Houses	Mushroom Processing Plants	Spice Markets	Voids in Attics
Cabinets	Drains & Sewers	Fruit Packing Sheds	Indoor Eating Facilities	Office Buildings	Stable Bins	Wall Voids
Cafeterias	Dried Flower Warehouses	Fruit Processing Plants	Industrial Installations	Other Public Buildings	Stables	Warehouses
Calving Barns	Dried Food Products in Processing	Fruit Quarantine Areas	Industrial Plants	Packaged Food Storage	Storage Areas	Wheat Mills
Canneries	Dried Food Products in Storage	Fruit Storage Warehouses	Institution Dining Areas	Pantries	Storage Bins (areas)	Wholesale Stores
Cat Sleeping Quarters	Dried Fruit Processing Plants	Garbage Bins	Institutions	Peanut Processing Plants	Stored Product Areas	Wine Cellars
Cattle Barns	Dried Fruit	Garbage Cans	Kennels	Peanut Warehouses	Stored Products	Wine Storage
Certified Organic Facilities	Dried Fruit Warehouses	Garbage Compactors	Liveries	Pet Bedding	Stores	Warehouses
Cheese Caves	Dumpsters	Garbage Trucks	Livestock Barns	Pet Carriers	Supermarkets	Wine Tasting Rooms
Citrus Packing Houses	Dwellings	Goat Holding Areas				Wineries and Zoos

**APPLICATIONS WITHIN FOOD HANDLING ESTABLISHMENTS**

**Food and/or Feed areas** include areas for receiving, storage (dry, cold, frozen or raw), packaging (canning, bottling, wrapping, boxing and grinding), preparing, edible waste storage and enclosed processing systems (mills, dairies, and systems for processing edible oils and syrups). Serving areas are places where prepared foods are served.

FOOD HANDLING ESTABLISHMENTS	
Application Type	Application Rate
Crack and Crevice Food/Feed Areas of Food/Feed Handling Establishments	1.5 oz. / 50 sq. ft.
Crack and Crevice Stored Product Pests in Non-Food/Non-Feed Areas of Food Handling Establishments	1.0 oz. / 50 sq. ft.
Crack and Crevice Treatments in Storage Sites	1.0 oz/ 50 sq. ft.

Using a non-power duster, apply directly into cracks and crevices, in expansion joints between different elements of construction, under or between equipment bases and the floor, beneath appliances, into wall voids, motor housing, junction boxes or switch boxes, conduits, behind and under cabinets or drawers, in and around drains, around drain pipes, plumbing and utility entrances, hollow equipment legs and other sites where listed insects can enter or hide.

**Stored Product Pests in Non-Food/Non-Feed areas of Food Handling Establishments:**

Non-food/feed areas of food/feed handling establishments are garbage rooms, lavatories, floor drains (to sewers, entries and vestibules), offices, locker rooms, machine rooms, boiler rooms, garages, mop closets and storage (after canning or bottling).

To aid in the kill of stored product pests in listed areas. Apply Evergreen® Pyrethrum Dust at the specified rate. Clean up waste materials, infested materials, dust, dirt, and any other debris. Thoroughly treat floors, walls, shelves, drawers, cabinets and other surfaces in bins, storage and handling areas, around cartons or containers and other areas where these insects tend to congregate. Apply into cracks and crevices and other difficult-to-reach areas where insects may hide. Apply in voids or cracks and crevices in pantries and other infested areas within the structure.

**For Use in Storage Sites:**

This product can be used in warehouse bins, trucks, cargo ships, mills, bin hoppers, elevators and conveying equipment, and other listed sites as a cleanup prior to using them for storage. Remove and dispose of all of the infested stored commodity accumulations and debris, with special attention to material lodged in the cracks and crevices. After sanitation measures have been employed, dust areas prior to use for storage, paying particular attention to cracks and crevices.

For packaged food warehouses, treat potential points of insect entry, as well as harborage sites in, on and around wooden and plastic pallets, pallet-racking systems, loading docks, scales, around cartons.

## INDOOR TREATMENTS

IN INDOOR RESIDENTIAL AREAS OR OTHER AREAS WHERE CHILDREN MAY COME INTO CONTACT, ONLY APPLY AS A CRACK-AND-CREVICE OR AS A SPOT TREATMENT TO AREAS NO GREATER THAN 3 FEET BY 3 FEET PER ROOM.

APPLICATION RATES FOR INDOOR TREATMENTS

Application Type	Application rates for Residential areas such as homes, condominiums, apartments, hotels, motels and pet areas	Application rates for Non-Residential areas such as offices, warehouses, hospitals and other commercial buildings and institutions
Crawling Insects—Crack and Crevice	Apply as a crack-and-crevice treatment to areas no greater than 3 x 3 feet per room.	1.5 oz. / 50 sq. ft.
Bed Bugs—Crack and Crevice	Apply as a crack-and-crevice treatment to areas no greater than 3 x 3 feet per room.	2.0 oz. / 50 sq. ft.
Ants—Crack and Crevice	Apply as a crack and crevice treatment to areas no greater than 3 x 3 feet per room.	1.0 oz. / 50 sq. ft.
Crawling Insects for Severe Infestations—Spot Treatments	Apply as a crack and crevice or as a spot treatment to areas no greater than 3 x 3 feet per room.	16 to 20 oz. / 1,000 sq. ft.
Fleas, Ticks and Mites in Areas where Pets Frequent—Spot Treatments	Apply as a spot treatment to areas no greater than 3 x 3 feet per room.	1.0 oz. / 70 sq. ft.
Powder Post Beetles—Spot Treatments	Apply as a spot treatment to areas no greater than 3 x 3 feet per room.	16 to 20 oz. / 1,000 sq. ft.
Asian Lady Bird Beetles and Box Elder Bugs in Crawl Spaces and Attics—Crack and Crevice	Apply as a spot treatment to areas no greater than 3 x 3 feet per room.	1 oz. / 50 sq. ft.
Asian Lady Bird Beetles and Box Elder Bugs in Crawl Spaces and Attics—Spot Treatments	Apply as a spot treatment to areas no greater than 3 x 3 feet per room.	16 to 24 oz. / 1,000 sq. ft.
Ornamental Plants	Apply to cover upper and lower surfaces of leaves, stems and flowers, with a thin even film of dust.	Apply to cover upper and lower surfaces of leaves, stems and flowers, with a thin even film of dust

### Crawling Insects:

Apply Evergreen® Pyrethrum Dust into voids, cracks and crevices or other places where insects are traveling or harboring. Apply a continuous visible film underneath and behind stairs, behind and beneath sinks, stoves, refrigerators and other appliances, around garbage cans, cabinets. Treat on, around and behind moldings, shelving, baseboards, doorsills, doorframes, floors, around drains, pipes and pipe openings, plumbing, bookcases, storage, utility installations and other areas where insects hide or travel. For severe infestations, apply behind cabinets and walls, in crawl spaces and attics at the specified rate.

### Fleas, Ticks, and Mites:

DO NOT TREAT DOGS OR CATS WITH THIS PRODUCT.

For Fleas, apply Evergreen® Pyrethrum Dust only as a direct treatment. Treat on and around animal bedding, sleeping quarters, kennel, furniture, floor and floor coverings and all areas where these pests inhabit. Treat cracks and crevices behind and around window and door frames, wall cracks, moldings, shelving, baseboards, pipe openings and all other cracks. Treat the entire inside surface of doghouses. For Ticks, apply Evergreen® Pyrethrum Dust in places where the insects are traveling or harboring.

### Bed Bugs:

Thoroughly inspect area of suspected bed bug infestation, and note potential harborage sites or actual infestations. Remove and/or pick up clothing and other obstacles that may prevent thorough inspection or treatment. Remove and sanitize infested bedding (pillows and linens), clothing, or other items.

Take bed apart. Apply Evergreen® Pyrethrum Dust into joints and channels if hollow, such as square or round tubing, and into cracks and crevices on the interior of bed frame, headboards, and footboards. If hollow, ensure that the interior framework is well dusted. Wall-mounted headboards and their hardware should be removed and treated. Mattresses should be dusted, especially seams, edges, and tufts. Mattress treatments must be limited to seams, folds and edges only.

Bed bugs may also harbor in areas of the room away from the bed. Apply Evergreen® Pyrethrum Dust as a crack and crevice application where evidence of bed bugs occurs. All furniture and furniture accessories in the room must be treated, including inside empty drawers and shelving in clothes closets, night stands and other furniture in the room, carpet edges, high and low wall moldings (such as floor molding or window casings or ledges/sills), door jambs and hinges, wall voids, and behind wall hangings and mirrors. Pull carpeting from tack strip and treat under the baseboard, sill plate, and under the carpet tack strips. Remove all clothes and other articles from dressers or closets before application. Treat all cracks and crevices associated with light switches, wall plates, faceplates, outlet covers, mounting boxes and housings. Apply Evergreen® Pyrethrum Dust as a continuous film to other potential harborage sites, such as table and wall-mounted lamps, wall crevices, drapery rods, curtains, blinds/mounting hardware, and other cracks and crevices in the room.

### Ants:

To aid in the kill of listed Ants, apply Evergreen® Pyrethrum Dust at the specified rate. Treat door frames, around window frames, behind, on and around moldings, cabinets, pipe openings, utility openings, shelves, ant trails, hills, wall voids, cracks and crevices and other points of entry into premises. Combine interior treatments with exterior perimeter application of Evergreen® Pyrethrum Dust.

### Powder Post Beetles\*:

To aid in protecting against insect entry. Apply to distribute Evergreen® Pyrethrum Dust through insect entry points, crawl holes into attics and crawl spaces of buildings. Close all vents and blowers so that all target surfaces can be treated. Contact all wood surfaces. In case of localized infestations, probe to locate galleries and drill holes to inject Evergreen® Pyrethrum Dust into the galleries and void areas. If a heavy infestation is suspected, a conventional, industry accepted techniques and treatments should be employed.

\*The purpose of such applications of Evergreen® Pyrethrum Dust is to kill workers or winged reproductive forms that may be present in treated channels at the time of treatment. Such applications are not a substitute for mechanical alteration, soil treatment or foundation treatment, but are merely a supplement. This product is not for use against Termites. For active Termite infestations, use conventional, industry accepted techniques and treatments.

### Asian Lady Beetles and Box Elder Bugs:

To kill or repel these insects, as well as Clover Mites, Cluster Flies and Elmleaf Beetles, inject Evergreen® Pyrethrum Dust into cracks and crevices and/or voids where these insects may be entering the premise, harboring or hibernating. Apply in crawl spaces and attics. On building exterior, treat underneath wood fascia, behind siding, cracks around windows, doors, siding, wall voids, utility pipes and entry points, behind chimneys, around foundations and foundation cracks, patios and deck joints. When treating for Asian Lady Beetles, apply Evergreen® Pyrethrum Dust in, on and around areas with abrupt color contrasts on buildings (shutters, drainpipes, etc.), southwest-facing or sun-exposed sides of buildings, and vertical corners of buildings.

**Ornamental Trees, Shrubs, Flowers and Foliage Plants:** Including: African Violet, Ageratum, Andromeda, Arborvitae, Ash, Aster, Azalea, Beech, Begonia, Birch, Boxwood, Calceolaria, Calendula, Calla, Camellia, Carnation, Ceanothus, Chrysanthemum, Cineraria, Coleum, Cotoneaster, Crabapple, Cyclamen, Cypress, Daffodil, Dahlia, Delphinium, Dogwood, Elm, Eucalyptus, Euonymus, Fern, Ficus, Fir, Firethorn, Foliage Plants, Forsythia, Fuchsia, Gardenia, Geranium, Gladiolus, Gloxinia, Gypsophila, Hawthorn, Hemlock, Hickory, Holly, Honey Locust, Horse Chestnut, Hyacinth, Hydrangea, Iris, Juniper, Larch, Laurel, Lilac, Lily, Linden, Maidenhair Fern, Marigold, Mimosa, Myrtle, Narcissus, Oak, Palm, Pansy, Pelargonium, Peony, Petunia, Philodendron, Phlox, Pine, Privet, Pyracantha, Rhododendron, Roses, Rubber Plant, Snapdragon, Sweet Pea, Tulip Tree, Tulips, Viburnum, Wandering Jew, Willow, Yew, Zinnia.

Evergreen® Pyrethrum Dust must come into contact with insects in order to kill them. Coverage of the target insect is vital to achieving good performance. Apply Evergreen® Pyrethrum Dust to cover upper and lower surfaces of leaves stems and flowers with a thin, even film of dust. Apply when insects or insect damage first appears.

When treating trees, shrubs and similar plants, treat the trunk, the base of the trunk and the critical root zone. Treat the tree crotch, knotholes, wounds and other areas where insects may harbor. Allow dust to settle in treated areas before reentering. Do not water or irrigate immediately following treatment. Do not eat or consume treated plants for 1 year after treatment.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Remove and wash contaminated clothing before reuse.

#### PERSONAL PROTECTIVE EQUIPMENT (PPE)

Mixers, Loaders, Applicators, and other handlers must wear the following:

- long-sleeved shirt
- long pants
- shoes and socks
- chemical-resistant gloves made of any waterproof material

#### USER SAFETY REQUIREMENTS

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

#### USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

#### ENVIRONMENTAL HAZARDS

This product is toxic to aquatic organisms, including fish and invertebrates. To protect the environment, do not allow pesticide to enter or runoff into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are actively visiting the treatment area.

#### PHYSICAL OR CHEMICAL HAZARDS

Do not use or store near heat or open flame.

## KEEP OUT OF REACH OF CHILDREN

#### CAUTION PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

#### FIRST AID

##### IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For additional information on this pesticide product (including health concerns, medical emergencies or pesticide incidents), you may call 1-888-740-8712.



EPA Reg. No. 1021-1871  
EPA Est. No. 1021-MN-2  
Rev. 0214-0315



All other marks are property of their respective holders. Manufactured by MGK®.  
©2015 MGK. All rights reserved.  
8810 Tenth Avenue North, Minneapolis, MN 55427  
2.5M-1111