

SARP: Sosyal Bilimler R Platformu

Burak AYDIN, James ALGINA, Walter LEITE

2017

Contents

1	Kapak	5
1.1	Tanıtım	5
2	Önyüz	7
2.1	Yazarlar	7
2.2	Teşekkür	7
2.3	Data	8
2.4	Finansal Destek	8
3	R'ın popülerliği	9
4	Windows için R kurulumu	11
5	Giriş	13
5.1	Fonksiyonlar	13
5.2	R Data Tipleri	17
5.3	R Paketleri	22
5.4	Çalışma alanı (workspace)	23
6	Veri Setleri	25
6.1	Veri Çekme	25
6.2	Basit Veri İşlemleri	27
6.3	veri Kaydetme	31
7	Betimsleyici İstatistikler ve Hipotez Testi	33
7.1	Betimsleyici İstatistikler	33
7.2	Basit Grafikler	44
7.3	Hipotez Testi Tanıtım	48
8	İki Ortalamanın Karşılaştırılması, t-testi	61
8.1	Bağımsız gruplar t-test (The Independent Groups t-test)	62
8.2	Bağlı gruplar t-testi (Within-subjects t-test)	70
8.3	Yaygın Tasarımlar	75
9	Varyans Analizi (ANOVA)	79
9.1	Terminoloji	79
9.2	Bağlı olmayan gözlemler varyans analizi (Between Subjects ANOVA)	80
9.3	Within Subjects ANOVA	98
9.4	Mixed Design	107
10	Correlation	109
10.1	Pearson correlation coefficient	109
10.2	Spearman's rho and Kendall's tau	118

10.3 Biserial and Point-Biserial Correlation Coefficients with R	119
10.4 Phi Correlation Coefficient with R	120
10.5 Issues in Interpreting Correlation Coefficients	120
11 Multiple Linear Regression, A Short Introduction	123
11.1 Matricies and Least Square Estimation	123
12 Useful R codes	143
12.1 More on the apaStyle package	148
12.2 A useful shiny application	149

Chapter 1

Kapak



Figure 1.1:

Bu platformun hakları korunmuştur CC0 by Burak AYDIN.

1.1 Tanıtım

Bu materyal İngilizce olarak hazırlanıp Türkçeye çevirilmiştir. Bu *platform* sosyal bilimler alanında çalışan ve nicel veri analizlerinin teoriden ziyade uygulama aşamasına ilgi gösteren araştırmacılar için oluşturulmuştur. Bütün istatistiksel prosedürler R (R Core Team, 2016b) ile yürütülmüş, gerçek veri kullanımına özen gösterilmiştir. Bu materyale *platform* denilmesinin üç sebebi vardır, (a) katkıya açıktır, (b) dinamik bir içeriğe sahiptir, (c) bilgisayar anakartı gibi kullanılabilir, R ile oluşturulmuş herhangi bir üst düzey çıktı platforma eklenebilir. Bu materyal Bookdown (Xie, 2016) ile inşaa edilmiştir, Bookdown ise R Markdown (Allaire et al., 2016) üzerine inşa edilmiştir. Materyalin hazırlanma aşamasında R Studio (RStudio Team, 2016) kullanılmıştır.

1.1.1 Neden Bookdown?

Bookdown ile görsel zenginliği mevcut materyaller oluşturulabilir. Shiny uygulamaları, basit olmayan grafikler gibi R'in araştırmacılara sunduğu ve sunacağı teknolojiler bu materyale kolayca eklenebilir. Bookdown ile aynı materyal PDF, HTML veya EBOOK olarak farklı şekilde kolayca okunabilir. Bookdown ile yazılan bir kitap Git Hub içerisinde depolanabilir ve en önemlisi katkı sağlamak isteyen araştırmacıların kolaylık sağlar. Kısacası Bookdown yeni nesil kitap yazma araçlarından biridir.

1.1.2 İçerik

Platformun bu versiyonunda yer alan konular;

- Windows için R kurulumu
- R'a giriş
- Veri Setleri
- Betimsel analizler ve hipotez testi
- t-test

- Varyans analizine giriş
- Korelasyon
- Çoklu regresyona giriş

Kitabın içerięi bir ders kitabından ziyade yardımcı materyal olarak hazırlanmıştır. İçerik az ve öz ele alınmıştır. Eğer açıklamaların yetersiz olduğunu düşünüyorsanız ve katkı sağlamak istiyorsanız isminize Teşekkürler 2.2 kısmında yer verilecektir.

Chapter 2

Önyüz

Bu kitabın yazarları elinde olan notları bir araya getirmek istemişlerdir. Akademik özenle oluşturduğumuz bu içerik her zaman açık kaynak olarak kalacaktır. Katkılarınız, istekleriniz titizlikle dikkate alınacak ve içeriğe dahil edilecektir. Bu içeriğin hangi motivasyonlarla hazırlandığını merak eden okuyucular kitabın sonuna eklediğimiz ¹ proje önerisini okuyabilirler.

2.1 Yazarlar

Akademik çalışmalarımız araştırma tasarıları (research design) ve nicel veri analizine yoğunlaşmıştır. Monte Carlo simulasyonları mutlaka çalışma takvimimizde yer alır. Diğer bir orta noktamız çok düzeyli modellere üzerine yaptığımız çalışmalardır.

2.1.1 Burak Aydın, Ph.D. Eğitimde araştırma ve değerlendirme metotları doktora ve istatistik doktora yandal derecesi mevcuttur. 2010 yılından bu yana R kullanıcısıdır. Yapısal eşitlik modelleri, çok düzeyli modeller ve eğilim puanları üzerine akademik çalışmaları vardır. Detaylı bilgi için **Kişisel or Kurumsal**

2.1.2 James Algina, Ph.D.

Klasik ve Modern Test Teorisi kitabının yazarıdır. Amerikan Eğitim Araştırmaları Birliği ve Amerikan Psikoloji Birliği (APA) üyesidir. 100'den fazla saygın akademik çalışması mevcuttur. Detaylı bilgi için UF Anita Zucker Center

2.1.3 Walter L. Leite, Ph.D.

Florida Üniversitesi Eğitim Fakültesinde Doçenttir. Gerçek deneysel ve yarı deneysel çalışmalardan edinilen verilerin analizinde saygın çalışmaları mevcuttur. Detaylı bilgi için UF College of Education

2.2 Teşekkür

Burada katkı sağlayanların isimlerine yer verilecektir.

¹sadece Türkçe versiyonunda yer alır

2.3 Data

Veri kullanıma açıktır, Dünya Bankası ve İŞKUR tarafından toplanmıştır. Türkiye’de yaşayan, İŞKUR’dan mesleki eğitim talebinde bulunmuş bireyleri temsil eder. 5902 kişi içerir. Materyal içerisinde kullanılan veriye *dataWBT* (data WorldBank Türkiye) ismi verilmiştir ulaşmak için Bölüm 6.1.4 içerisinde yer alan basamaklar takip edilebilir.

dataWBT içerisinde yer alan değişkenler;

1. Id: katılımcı kimliği
2. Program: Mesleki eğitim aldı mı? 1=evet, 2=Hayır
3. Cinsiyet: Erkek, Kadın
4. Kurs: 51 farklı kurs, muhasebeden garsonluğa.
5. Şehir: Katılımcıların yaşadığı şehir
6. Eğitim: en yüksek diploma
7. Babanın eğitim durumu
8. Annenin eğitim durumu
9. Soru 1-6: Toplumsal cinsiyet algısı (4lü likert). Yüksek puan cinsiyet ayrımına işaret eder.
10. Yüksek öğretim durumu: 0=Lise veya altı diplomalı , 1= Yüksek öğretim diplomalı 11: Yaş : 2010 yılında katılımcı yaşları 12: Hane geliri: TL olarak hane geliri 13: Hanede yaşayan kişi sayısı 14: Kişi başı yıllık gelir (hane geliri/hanede yaşayan kişi sayısı) 15: Toplumsal cinsiyet algısı genel puanı: 2,3,4,5 ve 6. soruların ortalaması. 16: Gelir kaynakları (12 farklı kaynak)

Toplumsal cinsiyet algısı puanları ve kullanılışı hakkında detaylı bilgi için Gök ve Aydın (basımda) incelenebilir.

Dünya Bankası Tarafından Katılımcılara Sorulan Sorular

1. Evlendikten sonra hem kadın hem erkek hane gelirine katkıda bulunmalıdır (çift-gelirlilik).
2. Üniversite eğitimi kızlardan ziyade erkekler için önemlidir.
3. MAddi durum zorlamadığı sürece evli bir kadın evinin dışında çalışmamalıdır.
4. Eşinin çalışması bir erkek için onur kırıcıdır.
5. Bir kadın düşüncelerini evde söyleyebilir fakat dışarda asla
6. Bir kadın eşinin sözünden çıkmamalıdır.

2.4 Finansal Destek

Bu materyalin hazırlanması Recep Tayyip Erdoğan Üniversitesi BAP birimi tarafından desteklenmiştir. BAP-53005-601

Öncesinde, bu proje TUBİTAK tarafından Şubat 2016 tarihinde ret edilmiştir.ID 1059B191501734.

Chapter 3

R'ın pop lerliđi

R programının kullanım sıklığı s rekli artmaktadır. Tippmann (2015) Scopus veri tabanında taranan ve 2014 yılında basılmış her 100 makaleden 1 tanesinde R programına veya R paketlerine atıfta bulunulduđunu yazmıştır. 2014 yılında 2925 R paketi mevcuttur. 2016 yılı sonunda bu sayı 10000'e ulaşmıştır.

Data analizi ve istatistiksel modellemenin yanında, işlevsel grafikler çizme, dokümanlar oluşturma, sunum hazırlama ve sim lasyon  retme gibi  eşitli ama lar i in kullanılabilen R, ba ta istatistik iler olmak  zere, m hendisler, ekonometristler ve sosyal bilimlerde modern ve komplike modellerle  alıřan arařtırmacıların dikkatini  ekmiştir.R programının yaygınlığı hakkındaki diđer g stergelere  rnekler;

1. R veri analizikonusunda evrensel bir dil olmuřtur ve yeni metotlar  ođu zaman R ile sunulur (Muenchen, 2011).
2. Elektrik ve elektronik m hendisleri enstit s  (IEEE)  yeleri tarafından en  ok kullanılan 5 programlama dilinden biridir. see
3. R programlama dilini lisans ve y ksek lisans d zeyinde ders olarak sunan  niversiteler ve uzaktan eđitim kurumları mevcuttur.
4.  zel řirketler tarafından kullanılan bir programlama dilidir.

Chapter 4

Windows için R kurulumu

R kurulumu oldukça kolaydır. R-project websitesinde yer alan basamaklar takip edilebilir veya sessiz olarak kaydedilmiş video izlenebilir (Video1 4).

Windows için R yükle

R programını betik dosyası oluşturmada doğrudan kullanmak mümkündür fakat bir betik düzenleyici kullanmak kolaylık sağlar. En basit betik düzenleyicisi R programının içerisinde yer alır. R açık iken *File* ve sonrasında *new script* seçilerek betik düzenleyici açılır. Bu basamaklar Video2 4) ile gösterilmiştir.

R betik

Fakat R içerisinde yer alan betik düzenleyici çok basittir. Kullanışlılığı oldukça yüksek olan ve bu materyalin hazırlanmasında da kullanılmış olan betik düzenleyici R studio'dur. Kurulum basamakları Video3 4 ile gösterilmiştir.

R Studio Yükle

Chapter 5

Giriş

R veri analizi, grafik veya interaktif web uygulaması gibi basit olmayan çıktılar oluşturabilir. Bu bölümün amacı basit olmayan çıktılar oluşturmada önce gereken temel prensipleri göstermektir.

5.1 Fonksiyonlar

R gibi programlanabilir gelişmiş hesap makineleri kullanıcıların fonksiyon yazmasına ve saklamasına izin verir. R kullanıcılarının fonksiyonların nasıl çalıştığını kavraması önemlidir.

5.1.1 R: Basit Hesap makinesi

R hesap yapabilir. Aşağıdaki işlemleri ve ilgili R kodlarını inceleyiniz.

$$1 + 1 = 2 \quad (5.1)$$

```
1+1  
## [1] 2
```

$$1 - 1 = 0 \quad (5.2)$$

```
1-1  
## [1] 0
```

$$1 + (2/3) - (2 * 6.5) = -11.33 \quad (5.3)$$

```
1 + (2 / 3) - (2 * 6.5)  
## [1] -11.3
```

$$\sin(30) + 4^3 + \log(4) + e^3 + \sqrt{7} = 87.13 \quad (5.4)$$

```
sin(30) + 4^3 + log(4) + exp(3) + sqrt(7)  
## [1] 87.1
```

Eşitlik (5.1) 'den (5.4) 'e kadar olan işlemler R tarafından tamamlanır fakat hafızada tutulmaz. Eğer yaptığınız bir işlemin sonucunu tekrar kullanmak istiyorsanız ona isim vermelisiniz. İsim verdiğiniz R çıktıları oturum süresince (session) tekrar erişime açıktır. Çıktılara oturum kapandıktan sonra da ulaşmak istiyorsanız kaydetmelisiniz. Kaydetme işlemleri ilerleyen bölümlerde ele alınmıştır. İsim verme işlemi farklı şekillerde yapılabilir; “=”, “<-” or “<<-”. Bu materyal “=” operatörünü kullanır.

Eşitlik (5.1) 'den (5.4) 'e kadar olan işlemleri oturum süresince saklamak için;

```
a=1 - 1
b=1 + 1
c=1 + (2 / 3) - (2 * 6.5)
d=sin(30) + 4^3 + log(4) + exp(3) + sqrt(7)
```

İsim verdiğiniz çıktılar ile işlem yapabilirsiniz.

```
a+b+c+d
## [1] 77.8
```

İsim verdiğiniz bir çıktıyı değiştirebilirsiniz (overwrite)

```
e=3+2
e
## [1] 5
e=e+10
e
## [1] 15
```

Farklı bir isim vermek için (Not: R büyük harf küçük harf ayrımı yapar)

```
Equation1_output=a
Equation1_output + b + c + d # a+b+c+d ile eşit
## [1] 77.8
```

5.1.2 R: Programlanabilir Hesap makinesi

En basit hali ile fonksiyon 3 parçadan oluşur, girdi, işlem, çıktı. Bu bölümde verilen fonksiyonların test puanlarının analiz basamağında kullanıldığını varsayalım.

5.1.2.1 Tek girdi - Tek Çıktı

Aşağıda verilen fonksiyonun adı *sabit5* olsun. *sabit5* fonksiyonu her öğrencinin puanına 5 puan ekleyecektir. Bir diğer deyişle, son derece basit bir fonksiyon olan *sabit5* verilen bir puana 5 ekleyerek çıktı oluşturur.

```
sabit5=function(girdi){
  cikti=girdi+5
  return(cikti)
}

sabit5(girdi=50)
## [1] 55
sabit5(60)
## [1] 65
sabit5(80)
## [1] 85
```

sabit5 fonksiyonu girdiyi alır , 5 ekler (*input+5*), ve bir çıktı oluşturur (*cikti=girdi+5*), ve çıktıyı rapor eder (*return(cikti)*). Bütün bu işlemler *{ }* içinde verilmelidir.

Diğer basit bir fonksiyon *systematic1* olarak isimlendirilmiştir ve verilen her bir puana %1 ekler.

```
systematic1=function(input){
  output=input+(input/100)
  return(output)
}

systematic1(input=50)
## [1] 50.5
systematic1(100)
## [1] 101
systematic1(120)
## [1] 121
```

5.1.2.2 Çoklu Girdi-Tek Çıktı

Daha önce verilen fonksiyonlar tek girdi alıp tek çıktı oluşturmuştur. Bu örnekte iki farklı girdi ve tek bir çıktı vardır. Fonksiyona *eksipuan* adı verilmiştir. Ham puan ve yanlış sayısı verilen fonksiyon, her yanlış için 0.2 puan düşürür. Örneğin 90 puan ve 6 yanlış girildiğinde çıktı olarak $(90-0.2*6)$ 88.8 verilir.

```
eksipuan=function(puan, yanlis){
  cikti=puan - (0.2 * yanlis)
  return(cikti)
}

eksipuan(puan=90,yanlis=6)
## [1] 88.8
eksipuan(90,17)
## [1] 86.6
```

Bir R fonksiyonunda girdiler *argüman* (arguments) olarak isimlendirilir. *eksipuan* fonksiyonu 2 argümana sahiptir (puan ve yanlış) ve tek bir çıktı verir. Çoklu argüman ve çoklu çıktı içeren fonksiyonlar yazılabilir.

5.1.2.3 Çoklu Girdi ve Çoklu Çıktı

geridonut fonksiyonu doğru yanıt sayısını ve her sorunun kaç puan olduğu argümanlarını alır, çıktı olarak toplam puanı ve 100 almak için eksik kalan soru sayısını hesaplar.

```
geridonut=function(dogruyanit, katsayi){
  total=dogruyanit*katsayi
  kalan=(100-total)/katsayi
  cikti=c(paste("Puan:", total, " eksik:",kalan))
  return(cikti)
}

geridonut(dogruyanit=20,katsayi=2)
## [1] "Puan: 40 eksik: 30"
geridonut(27,2)
## [1] "Puan: 54 eksik: 23"
```

5.1.2.4 Basit Hata

R fonksiyonlarının çalışması için argümanların doğru kullanılması gerekir. Eğer *geridonut* fonksiyonuna *katsayi* parametresini girmezseniz bir hata ile karşılaşsınız.

```
geridonut=function(dogruyanit, katsayi){
  total=dogruyanit*katsayi
  kalan=(100-total)/katsayi
  cikti=c(paste("Puan:", total, " eksik:",kalan))
  return(cikti)
}
geridonut(dogruyanit=20)
## Error in geridonut(dogruyanit = 20): argument "katsayi" is missing, with no default
```

5.1.2.5 Basit uyarı

R fonksiyonları uyarı içerebilir. Daha önce yazdığımız *eksipuan* fonksiyonunu düşünelim

```
eksipuan=function(puan, yanlis){
  cikti=puan - (0.2 * yanlis)
  return(cikti)
}
eksipuan(puan=50,yanlis=10)
## [1] 48
```

Bu fonksiyona bir uyarı ekleyebiliriz. Örneğin hesaplanacak puan sıfırın altında ise bir uyarı verebiliriz.

```
eksipuan2=function(puan, yanlis){
  cikti=puan - (0.2 * yanlis)
  if (cikti<0)
    warning("Yeni puan 0'dan düşük")
  return(cikti)
}
eksipuan2(puan=10,yanlis=60)
## Warning in eksipuan2(puan = 10, yanlis = 60): Yeni puan 0'dan düşük
## [1] -2
```

5.1.2.6 Basit Sekte

Bir R fonksiyonu, yazarın belirlediği durumlarda sekteye uğrayabilir. Örneğin *eksipuan3* fonksiyonunu 20'den düşük puanlar için düzeltme yapmayacak şekilde yazabiliriz.

```
eksipuan3=function(puan, yanlis){

  if ((puan)<(20))
    stop("20den düşük puanlar için bu fonksiyon işlemez")

  cikti=puan - (0.2 * yanlis)
  return(cikti)
}
eksipuan3(10,9)
## Error in eksipuan3(10, 9): 20den düşük puanlar için bu fonksiyon islemez
```

5.1.3 Yardım!

Her R kullanıcısı yeni fonksiyonlar yazmak zorunda değildir, fakat fonksiyonların nasıl çalıştığını bilmek önemlidir. Eğer R bir hata veriyorsa bu genellikle kullanıcı veya datadan kaynaklıdır. Her ne kadar çok

karşılaşılmassa da hatanın fonksiyonun kendisinden kaynaklandığı durumlar da olabilir.

R fonksiyonlar sayesinde çalışır. Dünyanın her yerinden araştırmacılar R fonksiyonları yazmakta, bu fonksiyonları bir R paketi olarak erişime açmaktadırlar. Hali hazırda 10 binden fazla R paketi vardır. R programını indirdiğinizde yaklaşık 30 R paketi bilgisayarınıza otomatik olarak indirilir. Bu 30 R paketinde binlerce fonksiyon bulunur.

R programınızı yüklediğinizde otomatik olarak yüklenen 30 paketten bir tanesi *base* dir. Bu paketin içinde 1200'den fazla fonksiyon bulunur. Örneğin *mean* fonksiyonu aritmetik ortalama hesaplar. Genellikle paketler detaylı açıklamalar ile birlikte sunulur. Kullanıcıların bu açıklamalara ulaşabilmesi için çeşitli yollar mevcuttur; *help*, *?*, *??* veya *example*

```
help("base") #
help(mean)   # aritmetik ortalama fonksiyonu ve argümanları
?mean       # aritmetik ortalama fonksiyonu ve argümanları
??mean      # aritmetik ortalama fonksiyonu ve argümanları
example(mean) # aritmetik ortalama fonksiyonu ve argümanları
```

5.2 R Data Tipleri

Bu bölümde vektörler, matrisler, değişken çeşitleri, kayıp veriler ve data çerçeveleri (data frames) kısaca tanıtılmıştır.

5.2.1 Vektörler

R *c* fonksiyonu ile vektör oluşturabilir. 10 öğrenci için not girelim.

```
notlar=c(40,50,53,65,72,77,79,81,86,90)
notlar
## [1] 40 50 53 65 72 77 79 81 86 90
```

R vektörler üzerinden işlem yapabilir.

```
notlar=c(40,50,53,65,72,77,79,81,86,90)
#her nota 10 ekle
notlar+10
## [1] 50 60 63 75 82 87 89 91 96 100
#her nota yüzde 10 ekle
notlar+(notlar*0.10)
## [1] 44.0 55.0 58.3 71.5 79.2 84.7 86.9 89.1 94.6 99.0
#kendi ile çarp
notlar*notlar
## [1] 1600 2500 2809 4225 5184 5929 6241 6561 7396 8100
# yeni notlar
notlar2=c(30,40,46,58,64,66,69,72,74,81)
# notlar ve notlar2 nin ortalamasını al
(notlar+notlar2)/2
## [1] 35.0 45.0 49.5 61.5 68.0 71.5 74.0 76.5 80.0 85.5

# ilk notların yüzde 40'ı ile ikinci notların yüzde 60'ını topla
notlar*0.4 + notlar2*0.6
## [1] 34.0 44.0 48.8 60.8 67.2 70.4 73.0 75.6 78.8 84.6
```

Vektör oluşturmak için işeyarar birçok fonksiyon vardır. Örneğin *rep* fonksiyonu (bknz: `example(rep)`) aynı değerleri tekrarlamak için kullanışlıdır.

rnorm fonksiyonu normal dağılıma sahip veriler simüle etmek için işe yarar. Eğer *?rnorm* kullanılırsa bu fonksiyonun 3 argümanı olduğu görülür *rnorm*(*n*, *mean* = 0, *sd* = 1). Bu fonksiyon vektör uzunluğu (değişken sayısı) *n* verilmediği sürece çalışmaz. Eğer sadece *n* verilirse, popülasyon ortalaması 0 ve standart sapması 1 olan dağılımdan rasgele seçilen değerler ile bir vektör oluşturulur. Bu parametreler değiştirilebilir. Örneğin *rnorm*(12,*mean*=10,*sd*=2) popülasyon parametreleri 10 ve 2 olan normal bir dağılımdan 12 adet gözlem çeker. Benzer bir fonksiyon *runif*(*n*, *min* = 0, *max* = 1) tekdüzey bir dağılımdan gözlem çeker.

```
a=1:12          # a 1 den 12ye tam sayılar
rep(0,12)        # 0 12 kez tekrarlanır
## [1] 0 0 0 0 0 0 0 0 0 0 0 0
rep(1:5,each=3)  # 1 den 5'e tam sayılar 3er kez tekrarlanır
## [1] 1 1 1 2 2 2 3 3 3 4 4 4 5 5 5
rep(1:5,times=3) # 3 kere 1'den 5'e tekrarlar
## [1] 1 2 3 4 5 1 2 3 4 5 1 2 3 4 5
seq(from=1,to=12) # 1'den 12'ye tam sayılar
## [1] 1 2 3 4 5 6 7 8 9 10 11 12
seq(1,25,by=2)   # 1'den 25'e ikişer atla
## [1] 1 3 5 7 9 11 13 15 17 19 21 23 25
seq(1,6,by=0.5)  # 1'den 6'ya 0.5 atla
## [1] 1.0 1.5 2.0 2.5 3.0 3.5 4.0 4.5 5.0 5.5 6.0
rnorm(12)        # ~N(0,1) 12 gözlem
## [1] 0.718 -2.234 0.191 0.492 0.254 1.952 0.416 -0.655 0.473 -0.153
## [11] -0.618 -1.224
rnorm(12,mean=10,sd=2) #~ N(10,2) 12 gözlem
## [1] 8.37 10.66 8.91 8.04 13.37 8.89 11.21 11.56 8.96 8.83 11.44
## [12] 9.95
runif(12, min = 10, max = 37)
## [1] 25.2 19.0 24.9 13.9 15.9 16.5 25.2 25.2 33.3 12.3 28.3 10.8
```

5.2.2 Matrisler

R matrisler oluşturup işlem yapabilir.

```
A=matrix(1:16,ncol=4,nrow=4) #4x4 matris oluştur
A
##      [,1] [,2] [,3] [,4]
## [1,]    1    5    9   13
## [2,]    2    6   10   14
## [3,]    3    7   11   15
## [4,]    4    8   12   16
B=matrix(runif(16,min=20,max=40),ncol=4) #4x4 matris oluştur

# işlem örnekleri
A+B      # topla
##      [,1] [,2] [,3] [,4]
## [1,] 39.5 33.8 46.0 46.6
## [2,] 23.3 39.9 32.1 34.3
## [3,] 41.9 43.8 37.7 38.6
## [4,] 25.2 47.3 48.3 50.6
A*B      # çarp
##      [,1] [,2] [,3] [,4]
```

```
## [1,] 38.5 144 333 436
## [2,] 42.5 204 221 285
## [3,] 116.6 258 294 354
## [4,] 84.9 315 436 553
A%*%B # matris çarp
##      [,1] [,2] [,3] [,4]
## [1,] 770 1041 860 797
## [2,] 890 1180 982 909
## [3,] 1010 1319 1104 1021
## [4,] 1130 1458 1227 1133
t(B) # çevir
##      [,1] [,2] [,3] [,4]
## [1,] 38.5 21.3 38.9 21.2
## [2,] 28.8 33.9 36.8 39.3
## [3,] 37.0 22.1 26.7 36.3
## [4,] 33.6 20.3 23.6 34.6
```

5.2.3 Değişkenler

Çözümlenecek verisetinin özelliklerini bilmek çok önemlidir. R içerisinde çözümlenecek değişkenler genellikle sınıflama, sıralı, sürekli, kayıp veya tarih tipindedir.

5.2.3.1 Sınıflama

R'da bir sınıflama verisi alfanumerik şekilde girilebilir fakat yorumlanması sayısal değil sınıflama şeklindedir. Örneğin;

```
adres=c("AAX", "BBZ", "CBT", "DBA", "DDC", "XZT")
cinsiyet=c("M", "F", "F", "M", "F", "M")
id=sample(letters,6)
program=rep(c("var", "yok"),each=3)
sehir=as.character(1:6)
```

5.2.3.2 Sıralı

Sıralı bir değişken sınıflama değişkenine göre daha çok bilgi içerir. Sıra ifade eder fakat değerler arasındaki farklılık anlamlı değildir. Örneğin koşucular birinci, ikinci ve üçüncü olarak sıralanabilir fakat bu sıralama verisi birinci ile ikinci arasında kaç dakika farklılık olduğunu belirtmez. Birinci koşucu ikinciden 5 saniye hızlı iken, ikinci koşucu üçüncü koşucudan yarım saat daha hızlı olabilir. R içerisinde *ordered* fonksiyonu ve *level* argümanı ile sıra belirtilebilir. Eğer *level* argümanı boş bırakılırsa R değerleri küçükten büyüğe sıralar.

```
soru1=ordered(c("zayıf", "orta", "iyi", "iyi", "zayıf", "zayıf"),
              levels=c("zayıf", "orta", "iyi"))
ses=ordered(c(1,3,2,2,1,3), levels=c("1", "2", "3"))
```

5.2.3.3 Sürekli

Eşit aralıklı veya eşit oranlı değişkenler sıralı ve sınıflama değişkenlerine göre daha fazla bilgi içerir. Değerler arasındaki farklılık anlamlıdır.

```
notlar=c(52,75,39,62,24,86)
notlar=rnorm(n=6,mean=160,sd=5)
```

5.2.3.4 Tarih

as.Date fonksiyonu ile tarih verisi girilebilir.

```
dt=as.Date(c("1994-06-01","1988-10-20","1990-12-01",
             "1978-03-23","1974-08-22","1994-11-04"))

dt
## [1] "1994-06-01" "1988-10-20" "1990-12-01" "1978-03-23" "1974-08-22"
## [6] "1994-11-04"

tatil=as.Date(c("01/01/2016","04/23/2016","05/19/2016","08/30/2016","09/29/2016"),
              format="%m/%d/%Y")

tatil
## [1] "2020-01-01" "2020-04-23" "2020-05-19" "2020-08-30" "2020-09-29"

Sys.Date( )
## [1] "2017-03-23"
Sys.Date( )-dt
## Time differences in days
## [1] 8331 10381 9609 14245 15554 8175
```

5.2.3.5 Doğru-Yanlış (logical)

Bu değişken TRUE veya FALSE değerlerini alır. Eğer sayısal veri olmaya zorlanırsa 1 ve 0 değerlerini alır. Aşağıda verilen kod girilen notların ortalamadan düşük olup olmadığını gösterir.

```
notlar=c(52,75,39,62,24,86)    # notlar
notlar>mean(notlar)
## [1] FALSE TRUE FALSE TRUE FALSE TRUE
as.numeric(notlar>mean(notlar)) # 1 ve 0.
## [1] 0 1 0 1 0 1
```

5.2.4 Faktörler

R içerisinde yer alan *factor* veri tipi sıralı ve sınıflama verileri için kullanılan bir çatıdır.

```
kurs=factor(c("muhassebe","garson","temizlik","garson","muhassebe","garson"))
ga1=factor(c(1,1,3,4,2,3),levels = 1:4,
           labels=c("tamamenkatilmiyorum","katilmiyorum","katililiyorum","tamamenkatililiyorum"))
ga2=factor(c(1,3,4,4,2,3),ordered = T)
ga3=gl(n=3,k=2,labels=c("A","B","C"),ordered=F)
```

Faktörler önemlidir. Faktörlerin alt sınıfları (levels) dikkatli bir şekilde incelenmelidir. Çözümleme aşamasında kullanılmayan alt sınıflar silinmelidir. Örneğin veri seti bölünmeden önce *Renk* faktörü girilmiş olsun, “mavi”, “yeşil”, “sarı” alt sınıflar olsun. Daha sonra veri seti bölündüğünde alt sınıf “sarı” kullanılmamış olsun. R *Renk* değişkenini hala 3 alt sınıflı olarak düşünecektir ve ona göre işlem yapacaktır. Bu hatalara sebep olur. *Droplevel* fonksiyonu kullanılarak faktör değişkeni düzeltilmelidir.

```

renk=factor(c(1,1,1,2,2,3),levels = 1:3,labels=c("mavi","yesil","sari"))
renk
## [1] mavi mavi mavi yesil yesil sari
## Levels: mavi yesil sari
renk2=renk[1:5] # renk2 değişkeni son degeri almadı
renk2 #fakat hala 3 level mevcut
## [1] mavi mavi mavi yesil yesil
## Levels: mavi yesil sari
droplevels(renk2) #kullanılmayan level silindi
## [1] mavi mavi mavi yesil yesil
## Levels: mavi yesil

```

5.2.5 Kayıp Veriler

Veri seti kayıp veriler içerebilir. R kayıp verileri *NA* (not available) ile belirtir.

```

gelir=c("maas","maas","destek",NA,NA,"maas")
hanekisi=c(3,2,3,NA,NA,4)

```

NOT: Kayıp veri belirleyiciler çetrefilli olabilir. *NA*, , " " (boşluk) veya önceden belirlenmiş bir sayı,örneğin -99 kayıp verileri temsil edebilir.

```

ornek = factor(c('maas','destek', NA, 'NA'," ",-99,"-99"))

# faktör içerisinde kayıp veriler <NA> olarak verilir.
# < > içinde yer almayan NA faktör sınıfını gösterir.
# " " bu da faktör alt sınıfını gösterir
#-99 and "-99" aynı faktör sınıfını gösterir

# is.na fonksiyonu kayıp verileri gösterir.
# ornek için bakıldığında sadece 3. eleman kayıp veri olarak görünür
is.na(ornek)
## [1] FALSE FALSE TRUE FALSE FALSE FALSE FALSE

#çözüm 'NA', " ", -99 ve "-99" ları NA'ye çevirelim
ornek[ornek=='NA' | ornek==" " | ornek== -99 | ornek== "-99"]=NA

#kontrol
is.na(ornek)
## [1] FALSE FALSE TRUE TRUE TRUE TRUE TRUE

#droplevel kullanalım
ornek=droplevels(ornek)

```

5.2.6 Veri Çerçeveleri (Data Frames)

Bir veri çerçevesi değişkenlerden oluşur. Sosyal bilimcilerin genellikle değişkenler arası ilişkileri araştırdığını düşünürsek, veri çerçeveleri kullanıcılarının temel R ögesidir. Daha önce oluşturduğumuz değişkenleri bir veri çerçevesine alabiliriz;

```
# hatırlatma
# id=sample(letters,6)

# program=rep(c("var", "yok"), each=3)

# cinsiyet=c("M", "F", "F", "M", "F", "M")

# soru1=ordered(c("zayif", "orta", "iyi", "iyi", "zayif", "zayif"),
#               levels=c("zayif", "orta", "iyi"))

# ses=ordered(c(1,3,2,2,1,3), levels=c("1", "2", "3"))

# notlar=c(52,75,39,62,24,86)

# gelir=c("maas", "maas", "destek", NA, NA, "maas")

# dt=as.Date(c("1994-06-01", "1988-10-20", "1990-12-01",
#              "1978-03-23", "1974-08-22", "1994-11-04"))

# kurs=factor(c("muhassebe", "garson", "temizlik", "garson", "muhassebe", "garson"))

basit_data=data.frame(id,program,cinsiyet,soru1,ses,
                      notlar,gelir,dt,kurs)

basit_data
##   id program cinsiyet soru1 ses notlar gelir      dt      kurs
## 1 y      var      M zayif   1    52  maas 1994-06-01 muhassebe
## 2 v      var      F  orta   3    75  maas 1988-10-20   garson
## 3 e      var      F   iyi   2    39 destek 1990-12-01 temizlik
## 4 x      yok      M   iyi   2    62   <NA> 1978-03-23   garson
## 5 k      yok      F zayif   1    24   <NA> 1974-08-22 muhassebe
## 6 u      yok      M zayif   3    86  maas 1994-11-04   garson
```

Veri setleri el yordamı ile girildiğinde veya hazır olarak R a aktarıldığında (örneğin excel dosyasından) veri setinin yapısını incelemek önemlidir. *str* (structure) fonksiyonu kullanılabilir.

```
str(basit_data)
## 'data.frame':   6 obs. of  9 variables:
##  $ id      : Factor w/ 6 levels "e","k","u","v",...: 6 4 1 5 2 3
##  $ program : Factor w/ 2 levels "var","yok": 1 1 1 2 2 2
##  $ cinsiyet: Factor w/ 2 levels "F","M": 2 1 1 2 1 2
##  $ soru1   : Ord.factor w/ 3 levels "zayif"<"orta"<...: 1 2 3 3 1 1
##  $ ses     : Ord.factor w/ 3 levels "1"<"2"<"3": 1 3 2 2 1 3
##  $ notlar  : num  52 75 39 62 24 86
##  $ gelir   : Factor w/ 2 levels "destek","maas": 2 2 1 NA NA 2
##  $ dt      : Date, format: "1994-06-01" "1988-10-20" ...
##  $ kurs    : Factor w/ 3 levels "garson","muhassebe",...: 2 1 3 1 2 1
```

5.3 R Paketleri

R bilgisayarımıza kurulurken 30'dan fazla paket yükler. Bu paketler *sistem kütüphanesinde* saklanır. R paketleri otomatik olarak yüklenen bu 30 paketle sınırlı değildir, örneğin doğrusal karma etkiler modellerini

(linear mixed models) çözümlmek için *lme*(Bates et al., 2015) paketi kullanılabilir. Bu paket 60000'den fazla bilgisayara yüklenmiş ve 1500'den fazla akademik yayında kullanılmıştır. R paketleri genellikle CRAN (comprehensive R archive network) içerisinde bulunur. Paketler yazarlar tarafından güncellendiği sürece CRAN'da bulunur. R paketlerini CRAN'dan çekerek kendi bilgisayarınızda saklayabilirsiniz. Yüklediğiniz paketler *kullanıcı kütüphanesinde* tutulur. Paketleri bir R oturumunda kullanabilmek için onları aktif hale getirmeniz gerekir.

R ve R Studio'yu kurma aşamasında R-RStudio-R paketleri arasında bilgisayar tarafından sağlanan otomatik bir bağ olduğunu farketmiş olabilirsiniz. R studio R'dan sonra yüklendiğinde bilgisayarınızı tarayacak, R programının yerini bulacak ve ona bağlanacaktır. Hem R hem de R Studio R paketlerinizin yerini bulabilir (eğer siz yerlerini değiştirmediyse). Eğer R paketlerinizin nerede olduğunu öğrenmek isterseniz *.libPaths()* fonksiyonunu kullanabilirsiniz.

CRAN'da yer alan R kütüphaneleri bilgisayarınıza kolayca yüklenebilir. R studio'da yer alan *Packages* ve *install* sekmesinden veya *install.packages("paketismi")* fonksiyonu ile paketleri indirebilirsiniz. Paketlerin oturum esnasında aktif hale getirilmesi gerekir. Bu işlem R studio *Paketler* sekmesinde yer alan paket isimlerinin yanındaki kutucuğa tıklayarak veya *library("paketismi")* fonksiyonu ile tamamlanabilir. Bu basamaklar Video 4 ile gösterilmiştir.5.3.

R Paketi Yükle

5.4 Çalışma alanı (workspace)

Bir R oturumu açtığınızda ve R işlemleri yaptığınızda bu işlemler çalışma alanında yürütülür. Her adımınız R Studio sağ üst köşede yer alan *History* sekmesinde görülür. Çalışma alanınızı oturum sonunda kaydedebilirsiniz. Oturum esnasında oluşturduğunuz R çıktıları çalışma alanında tutulur. *ls()* fonksiyonu ile bu çıktıları görebilirsiniz.

R çıktıları çalışma alanına getirilebilir veya çalışma alanı dışına uzun süreliğine kaydedilebilir. Dosyaların yerlerini bulmak ile uğraşmak istemiyorsanız bütün işlemlerinizi aynı klasörde tamamlamayı tercih edebilirsiniz. *getwd()* fonksiyonu size hangi klasör içinde (working directory) olduğunuzu gösterir. Uzun süreliğine kaydetmek istediğiniz bir R çıktısı bu klasöre kolayca kaydedilebilir. Aktif olan klasörünüzü *setwd()* fonksiyonu ile değiştirebilirsiniz. Tabiki bilgisayarınızda her hangi bir klasörde, hatta internette sakladığınız bir nesneyi R çalışma alanınıza getirebilir veya çalışma alanınızda oluşturduğunuz bir dosyayı bilgisayarınızda her hangi bir klasöre kaydedebilirsiniz. Fakat bu durumlarda adresi (location) hatasız bir şekilde R'a bildirmeniz gerekir. Girdi ve Çıktı konuları bir sonraki bölümde ele alınmıştır.

Chapter 6

Veri Setleri

Verileri teker teker kaydetme 5.2.6 bölümünde verilmiştir. Fakat veri setleri genellikle çözümleme yapacak araştırmacıya hazır şekilde gelir. Bu bölüm (a) veri çekme , (b) basit veri işleme etme yöntemleri ve (c) veri kaydetme konularını içerir.

6.1 Veri Çekme

Bir veri seti farklı formatlarda bulunabilir. R kullanıcılarının çok karşılaştığı veri formatları arasında .csv, .sav, .Rdata, .txt sayılabilir. Çözümleme işleminden önce verilerin düzgün bir şekilde çalışma alanına getirilmesi önemlidir. Eğer çalışacağınız veri seti ve R betiği aynı klasör içerisinde ise, bir diğer deyişle veri setinizin çalışma klasörünün (working directory) içerisinde ise adres belirtmeden veriyi çalışma alanınıza çağırabilirsiniz.

6.1.1 CSV

CSV (comma separated values) virgülle ayrılmış değerler içeren dosyalardır. Microsoft Excel kullanıcıları excel formatında yer alan verileri kolayca csv olarak kaydedebilirler. Diğer excel formatları ile kıyaslandığında (xls,xlsx,xlsb, vd.) işlemesi daha kolay veri formatıdır. *read.csv* fonksiyonu ile veri çalışma alanına çağırılabilir. En basit hali ile;

```
data1=read.csv("dataismi.csv") # eğer çalışma klasöründe dataisim.csv dosyası
                                # mevcut ise çalışır

#Windows için
data1=read.csv("C:\\Users\\Desktop\\folderX\\dataismi.csv") # adres (path)
data1=read.csv("C:/Users/Desktop/folderX/dataismi.csv") # adres (path)
#NOTE: \ karakteri hata verir / veya \\ kullanılmalıdır.
```

?*read.csv* komutu ile fonksiyonun argümanlarını görebilirsiniz. Önemli olan argümanlara örnek;

- veya değişken isimleri mevcut ise *header=TRUE* aksi halde *header=FALSE* .
- Kayıp veri belirticiler için *na.strings* . Örneğin *na.strings = "-99"* bütün -99 değerlerinin kayıp veriyi belirttiğini R'a iletir. Benzer şekilde *na.strings = c("-99", "-9")* hem -99 hem de -9 değerlerinin kayıp veri olduğunu belirtir.
- Eğer karakter verilerini faktör olarak kullanmak istiyorsanız *stringsAsFactors=TRUE* aksi halde *stringsAsFactors=FALSE*.

- d) Veriyi çağırma esnasında değişkenlere yeni isim vermek istiyorsanız *col.names* argümanı kullanılabilir. Örneğin 3 değişkeniniz varsa *col.names=c("A1","B2","C3")* argümanı ile sütunlara isim verebilirsiniz.

Eğer csv dosyanızda ondalık sayılar nokta yerine virgül ile ayrılmış ise (Avrupa ve Türkiye) bu *read.csv* fonksiyonu için problem oluşturur. Çözüm olarak *read.csv2* fonksiyonunu kullanabilir, veya *read.csv* içerisinde *sep=";"* ve *dec=","* argümanlarını kullanabilirsiniz. CSV dosyasından veri çağırma basamakları Video 5 ile gösterilmiştir.

CSV Oku

6.1.2 SPSS

SAV sosyal bilimciler tarafından kullanılan bir veri formatıdır. *foreign* (R Core Team, 2016a) paketinde yer alan *read.spss* fonksiyonu sav dosyalarını okumak için kullanılabilir.

```
require(foreign)
?read.spss
data=read.spss("dataismi.sav",to.data.frame=TRUE)
# eğer dataisim.sav çalışma klasöründe ise çalışılır
```

6.1.3 Rdata

Rdata formatı genelde daha az bilgisayar hafızası işgal eder. Rdata olarak kaydedilecek her R çıktısı ismi ile birlikte kaydedilir.

```
load("dataisim.Rdata") #eğer dataisim.Rdata çalışma klasöründe ise çalışılır
```

6.1.4 Sanal Depolardan Veri Çekmek

R ile sanal dünyadan veri çekilebilir. Süreci basite indirgersek, (a) öncelikle verinin nerede yer aldığı doğru şekilde belirlenmelidir, (b) verinin formatı doğru şekilde belirlenmelidir, (c) veri indirilip R' çağrılır veya doğrudan R'a çağrılır. Aşağıda verilen komutlar 2.3 bölümünde tanıtılan *dataWBT*'nin çalışma alanınıza getirilmesini sağlar.

```
#sanal depodan CSV oku
urldosyasi='https://raw.githubusercontent.com/burakaydin/materyaller/gh-pages/ARPASS/dataWBT.csv'
dataismi=read.csv(urldosyasi)
str(dataismi)

#sanal depodan Rdata oku
urldosyasi2='https://github.com/burakaydin/materyaller/blob/gh-pages/ARPASS/dataWBT.Rdata?raw=true'
load(url(urldosyasi2))
str(dataWBT)
```

Bu veri setleri Github Depo veya , excel dosyası olarak buradan indirilebilir.

6.1.5 R Stuido Aracılığı ile Veri Çağırma

Veri dosyanız bilgisayarınızda farklı bir klasörde (çalışma klasörü dışında) ise veya tıkla-bırak yöntemini (point-click) tercih ederseniz R Studio'nun sağ üst köşesinde *Environment* sekmesi altında yer alan *import dataset* ile veri çağırabilirsiniz. Bu basamaklar Video 6 ile gösterilmiştir.

R Studio ile CSV Oku

6.2 Basit Veri İşlemleri

Genellikle çözümleme basamağına geçilmeden önce verinin işlenmesi gerekir. Bu bölüm (a) değişkenleri yeniden kodlama, (b) alt küme seçme, (c) yeni değişken oluşturma, (d) veri çerçevesini değiştirme, (e) değişken türünü değiştirme ve (f) veri silme işlemlerini kısaca özetler.

6.2.1 Değişkenleri yeniden kodlama

Bir satırı, bir sütünü veya bir satır/sütun keşiminde yer alan tek bir elemanı değiştirmek mümkündür. Değişkenlerin yeni isimler vermek mümkündür. *plyr* (Wickham, 2011) paketi değişkenleri yeniden kodlamada yardımcı olabilir.

```
# sanal depodan CSV oku
urldosya='https://raw.githubusercontent.com/burakaydin/materyaller/gh-pages/ARPASS/dataWBT.csv'
veriseti1=read.csv(urldosya)

#URL adresini sil
rm(urldosya)

# 151. satır 16. sütunu değiştir ve 30 yap
veriseti1[151,16]=30

# aynı işlem satır ismi ve sütun ismi verilerek yapılabilir.
# id numarası 67034022 olan satırın yaş değerini 32 yap.
veriseti1[veriseti1$id==67034022,"age"]=32

# tekrar kodlama
# Veri setinde yer alan treatment değişkeni 0 ve 1 olarak girilmiştir.
# 1leri "trt" ve 2leri "cnt" yapmak için
veriseti1[veriseti1$treatment==1,"treatment"]="trt"
veriseti1[veriseti1$treatment==2,"treatment"]="cnt"

# ifelse fonksiyonu benzer şekilde çalışır
# "wage01" değişkeninde "wage01" "Yes" ise 0.5, değil ise -0.5 olarak kodlayalım
veriseti1$wage01=ifelse(veriseti1$wage01=="Yes",0.5,-0.5)

# plyr paketi kullanarak
require(plyr)
# pension01yeni değişkeni pension01 değişkeni üzerinden tanımlanmıştır
# eski değerler olan Yes ve No yerine 1 ve 0 kodlayalım
veriseti1$pension01yeni <- mapvalues(veriseti1$pension01,
                                     from=c("Yes","No"),to=c("1","0"))

#bir değişkene yeni isim verelim
#4. ve 5. değişkenlere yeni isim verelim
colnames(veriseti1)[4]="kurs"
colnames(veriseti1)[5]="bolge"

#isim verme işlemini tek sıra kod ile yapalım
colnames(veriseti1)[c(17,21)]=c("Tgelir","maas1")
```

```
#plyr paketini kullanalım gen_att değişkenine toplumsalCinsiyet ismi verelim
veriseti1 <- rename(veriseti1,c('gen_att'='toplumsalCinsiyet'))

#kontrol etmek için head(veriseti) veya summary(veriseti) kullanılabilir

# veriseti1'i çalışma alanından sil
rm(veriseti1)
```

6.2.2 Alt Küme Seçme (Subsetting)

R ile alt küme oluşturmak oldukça kolaydır.

```
# CSV yükle
urldosya='https://raw.githubusercontent.com/burakaydin/materyaller/gh-pages/ARPASS/dataWBT.csv'
veriseti1=read.csv(urldosya)

# URL adresi sil
rm(urldosya)

# sadece İstanbul'u seç
istDAT=veriseti1[veriseti1$city=="İSTANBUL",]

# sadece İstanbul'dan ilk sekiz katılımcıyı seç
istDAT18=veriseti1[veriseti1$city=="İSTANBUL",1:8]

# sadece İstanbul'dan gen_att puanı 2den yüksek olanları seç
istDATGAT2=veriseti1[veriseti1$city=="İSTANBUL" | veriseti1$gen_att >2 ,]

# subset fonksiyonu
# sadece İstanbul'dan gen_att puanı 2den yüksek olan ilk sekiz katılımcıyı seç
istDATGAT2B=subset(veriseti1, city=="İSTANBUL" | veriseti1$gen_att >2, select=1:8)

#item 1 değeri 1,2 ve 3 olan katılımcıları seç
item1_123 <- veriseti1[veriseti1$item1 %in% c(1,2,3), ]

#çalışma alanını temizle
rm(list=ls())
```

6.2.3 Yeni Değişken Oluştur

Daha önce 5 bölümünde değişken oluşturma yöntemlerine değinilmiştir. Bu bölüm hatırlatma olarak görülebilir.

```
# CSV yükle
urldosya='https://raw.githubusercontent.com/burakaydin/materyaller/gh-pages/ARPASS/dataWBT.csv'
veriseti1=read.csv(urldosya)

#URL dosyası yükle
rm(urldosya)

# item2'den 6'ya kadar olan sütunları topla
```

```

veriseti1$itemTOPLAM=with(veriseti1,item2+item3+item4+item5+item6)

# item2'den 6'ya kadar olan sütunların ortalamasını al (na.rm =T önemli)
veriseti1$itemAVE=with(veriseti1,
                        rowMeans(cbind(item2,item3,item4,item5,item6),na.rm=T))

#veya rowMeans fonksiyonu
veriseti1$itemAVE=rowMeans(veriseti1[,10:14],na.rm = T)

# Şehirler için ortalama hesaplama
veriseti1$CityAVEScore =with(veriseti1, ave(itemAVE,city,FUN=function(x) mean(x, na.rm=T)))

#veya
veriseti1=merge(veriseti1, aggregate(itemAVE ~ city, data = veriseti1, FUN=mean, na.rm=TRUE),
                by = "city", suffixes = c("", "citymean"),all=T)

#veya her bir soru için şehir ortalaması hesaplama
veriseti1=merge(veriseti1, aggregate(cbind(item2,item3,item4,item5,item6) ~ city,
                                    data = veriseti1, FUN=mean, na.rm=TRUE),
                by = "city", suffixes = c("", "Citymean"),all=T)

# değişkenlerin kategorize edilmesi. Eğer item1AVE 2'den küçük ise 0 aksi halde 1
veriseti1$itemAVE01=ifelse(veriseti1$itemAVE<2,0,1)

# 0 ile 1.8 arasına 1 ver
# 1.8 ve 2.5 arasına 2 ver
# 2.5 ile 5 arasına 3 ver
veriseti1$itemAVE123=with(veriseti1,cut(itemAVE, breaks=c(0,1.8,2.5,5), labels = FALSE))
# cut fonksiyonu içerisinde yer alan right=T argümanına göz gezdirin
# örneğin right=T ise değeri tam olarak 1.8 olan değişkenler 1 olur
#               right=F ise değeri tam olarak 1.8 olan değişkenler 2 olur

```

6.2.4 Veri Çerçevesini Değiştirme (Reshaping data)

Veri çerçevesini değiştirmek, uzun formattan geniş formata geçmek veya tam tersi gerekli olabilir. *tidyr* (Wickham, 2016) yardımcı olabilir.

```

# CSV yükle
urldosya='https://raw.githubusercontent.com/burakaydin/materyaller/gh-pages/ARPASS/dataWBT.csv'
veriseti1=read.csv(urldosya)

#adresi sil
rm(urldosya)

# genişten uzuna item 1den 6 ya kadar olan sütunları item adı altında birleştir
library(tidyr)
data_long = gather(veriseti1, item, score, item1:item6, factor_key=TRUE)

#id'ye göre diz
data_long=data_long[order(data_long$id),]

```

```
# uzundan geniş.
data_wide = spread(data_long, item, score)

## belirlediğiniz nesneler dışında çalışma alanını temizle
rm(list=setdiff(ls(),c("veriseti1")))
```

6.2.5 Değişken Türünü Değiştirme

Sayısal girilen verileri faktöre çevirme gibi işlemler çözümleme basamağından önce gerekli olabilir.

```
# CSV yükle
urldosya='https://raw.githubusercontent.com/burakaydin/materyaller/gh-pages/ARPASS/dataWBT.csv'
veriseti1=read.csv(urldosya,stringsAsFactors = F)

#URL sil
rm(urldosya)

#treatment değişkenini incele
str(veriseti1$treatment)

#sayısal veriyi faktöre çevir
veriseti1$treatmentFactor=factor(veriseti1$treatment,labels=c("treatment","control"))

#karakter olarak girildiğinde faktörleri sayıya çevirme

veriseti1$iv1=factor(rep(c("1","2","3"),length=nrow(veriseti1)))
veriseti1$iv1numeric=as.numeric(levels(veriseti1$iv1))[veriseti1$iv1]
#veya
veriseti1$iv1numeric=as.numeric(as.character(veriseti1$iv1))

#NAleri -99'a çevir
veriseti1[is.na(veriseti1)]= (-99)

#çalışma alanını temizle
rm(list=ls())
```

6.2.6 Veri Silme

Bir tek hücreyi, bir satırı veya bir sütunu silmek gerekebilir.

```
# CSV yükle
urldosya='https://raw.githubusercontent.com/burakaydin/materyaller/gh-pages/ARPASS/dataWBT.csv'
veriseti1=read.csv(urldosya,stringsAsFactors = F)

#URL sil
rm(urldosya)

#3. satır 5. sütunda yer alan hücreyi sil
veriseti1[3,5]=NA

#3. satırı sil
veriseti1[3,]= NA
```

```

#veya
veriseti1=veriseti1[-3,]

#course taken isimli sütunu sil
veriseti1$course_taken=NULL

#gösterim amaçlı veri oluştur
temp=veriseti1[,1:10]

# kayıp verili satırı silme (listwise)
temp=na.omit(temp)

#çalışma alanını temizle
rm(list=ls())

```

6.3 veri Kaydetme

Adres belirtmediğiniz sürece kaydetme işlemi mevcut çalışma klasörünüz (working directory) içerisinde tamamlanır.

```

# CSV yükle
urldosya='https://raw.githubusercontent.com/burakaydin/materyaller/gh-pages/ARPASS/dataWBT.csv'
veriseti1=read.csv(urldosya,stringsAsFactors = F)

#URL sil
rm(urldosya)

#nesne oluştur.
subset1=veriseti1[1:20,1:5]
object2=mean(veriseti1$item1,na.rm = T)

#çalışma klasörünüzü kontrol edin
getwd()

# Rdata olarak sakla
save(subset1,file="subset1Rfile.Rdata")
# adres vererek sakla
save(object2,file="C:/Users/Desktop/object2Rfile.Rdata")

# csv olarak sakla
write.csv(subset1,file="subset1CSVfile.csv",row.names = F)

#sps dosyası olarak sakla
library(foreign)
write.foreign(subset1, "subset1SPSfile.txt", "subset1SPSfile.sps", package="SPSS")

#çalışma alanını temizle
rm(list=ls())

```


Chapter 7

Betimleyici İstatistikler ve Hipotez Testi

Betimleyici istatistikler örnekleme tanımlamayı amaçlar. Bu bölüm içerisinde daha önce tanıtılan dataWBT (2.3) kullanılarak (a) betimleyici istatistikler hesaplanmış (b) basit grafikler çizilmiş ve (c) hipotez testi açıklanmıştır.

Bu bölümde yer alan R kodlarını kullanmak isteyen araştırmacıların bir önceki bölümü inceledikleri varsayılmıştır. Bu bölümde yer alan basamakların atlanmadan takip edildiği varsayımı da yapılmıştır. dataWBT çalışma alanınıza çağırarak için;

```
# CSV yükle
urldosya='https://raw.githubusercontent.com/burakaydin/materyaller/gh-pages/ARPASS/dataWBT.csv'
dataWBT=read.csv(urldosya)

#remove URL
rm(urldosya)
```

7.1 Betimleyici İstatistikler

Bu alt bölümde ortalama, ortanca, varyans, standart sapma, çarpıklık ve basıklık hesaplanmıştır. Örneklerde toplumsal cinsiyet algısı (gen_att) değişkeni kullanılmıştır.

7.1.1 Ortalama

Eşitlik (7.1) 'de verildiği gibi, ortalama, bir değişkeni oluşturan değerlerin toplamının toplam değer sayısına bölünmesi ile hesaplanır.

$$\bar{Y} = \frac{1}{n} \sum_{i=1}^n Y_i \quad (7.1)$$

```
# gen_att değişkeninin ortalamasını hesapla
mean(dataWBT$gen_att, na.rm = T)
## [1] 1.94
```

```
# birden fazla değişkenin ortalamasını hesapla
# ?colMeans
colMeans(dataWBT[,c("gen_att","item1")],na.rm = T)
## gen_att item1
## 1.94 3.45
```

7.1.2 Ortanca

Büyükten küçüğe veya küçükten büyüğe dizilmiş bir değişkenin orta noktasına ortanca denir. Eğer değişkenin eleman sayısı (n) tek sayı ise $((n+1)/2)$. sırada yer alan, eğer çift sayı ise $(n/2)$. ve $((n+1)/2)$. değerlerin ortalaması ortancayı verir.

```
# Ortanca hesapla
median(dataWBT$gen_att,na.rm = T)
## [1] 2
```

7.1.3 Varyans

Varyans değişkenin ne kadar yayıldığını anlamada çok kullanılan bir ölçüdür. Eşitlik (7.2) ile hesaplanır.

$$s_Y^2 = \frac{1}{n-1} \sum_{i=1}^n (Y_i - \bar{Y})^2 \quad (7.2)$$

```
#varyans hesapla
var(dataWBT$gen_att,na.rm = T)
## [1] 0.364
```

7.1.4 Standart Sapma

Varyansın kareköküdür ve Eşitlik (7.3) ile hesaplanır.

$$s_Y = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (Y_i - \bar{Y})^2} \quad (7.3)$$

```
#SS hesapla
sd(dataWBT$gen_att,na.rm = T)
## [1] 0.603
```

7.1.5 Çarpıklık (Skewness)

Çarpıklık değeri dağılımın şekli hakkında bilgi verir. Tamamen simetrik olan bir dağılımın çarpıklık değeri 0'dır.

Dağılımın sol kuyruğu sağ kuyruğuna nazaran uzun olduğunda çarpıklık değerinin sıfırdan küçük çıkması tipiktir. Bu tür dağılımlar sola çarpık veya negatif çarpık olarak isimlendirilir. Bu tür dağılımlarda medyan ortalama ortalamadan yüksektir.

Dağılımın sağ kuyruğu sol kuyruğuna nazaran uzun olduğunda çarpıklık değerinin sıfırdan büyük hesaplanması tipiktir. Bu tür dağılımlar sağa çarpık veya pozitif çarpık olarak isimlendirilir. Bu tür dağılımlarda medyan ortalama ortalamadan küçüktür.

Örneklem için çarpıklık formülü ¹

$$\sqrt{n} \frac{\sum_i^n (X_i - \bar{X})^3}{\left(\sum_i^n (X_i - \bar{X})^2\right)^{3/2}} \quad (7.4)$$

Örneklem için çarpıklık değeri *moments* (Komsta and Novomestky, 2015) paketinde yer alan *skewness* fonksiyonu ile hesaplanabilir.

```
#çarpıklık hesapla
library(moments)
skewness(dataWBT$gen_att,na.rm = T)
## [1] 0.377
```

NOT: Çarpıklık ve basıklık değerleri için standart hata ve sonrasında z-puanı hesaplanabilir. Hesaplanan bu z-puanı seçilen bir kritik değer ile (ör. 1.96) kıyaslanarak çarpıklık veya basıklığın istatistiksel olarak anlamlı olup olmadığı sınanabilir. Benzer şekilde normallik testleri de (ör. Shapiro-Wilk) yapılabilir. Fakat bu testler örneklem büyüklüğüne hassastır. Bir diğer deyişle örneklem büyüdükçe çok küçük farklılıklar istatistiksel olarak anlamlı bulunabilir. Çarpıklık, basıklık veya normallik testlerinin varsayım ihlallerini tespit etmek üzere kullanılışı nispeten eskimiş yöntemlerdir. Bu testleri kullanmak yerine normallik grafik üzerinden incelenip, dirençli tahminleyicilerin (robust estimators) veya Monte Carlo simulasyon tekniklerinin çıktıları incelenebilir.

7.1.5.1 Çarpıklık örnekleri

Normal bir dağılım ve çarpıklık istatistiği;

Sola çarpık sürekli değişken;

Sağa çarpık sürekli değişken;

7.1.6 Basıklık (Kurtosis)

Basıklık değeri dağılımın şekli hakkında bilgi verir. Normal bir dağılımın Pearson basıklık değeri 3'tür. Eşitlik (7.5) basıklık değerinin hesaplanışını gösterir.

$$n \frac{\sum_i^n (X_i - \bar{X})^4}{\left(\sum_i^n (X_i - \bar{X})^2\right)^2} \quad (7.5)$$

Eşitlik (7.5) sıfırdan küçük değerler vermez. 0 ile 3 arasında yer alan değerler genellikle düz dağılımlarda hesaplanır, örneğin tekdüzey dağılımlar. Uzun kuyruklu dağılımlarda 3'ten büyük değerler görülebilir. Alan yazında yorumu kolaylaştırmak için Eşitlik (7.5) 'ten 3 çıkarıldığı durumlar mevcuttur.

Örneklem için Pearson basıklık değeri *moments* (Komsta and Novomestky, 2015) paketinde yer alan *kurtosis* fonksiyonu ile hesaplanabilir.

```
#basıklık hesapla
library(moments)
kurtosis(dataWBT$gen_att,na.rm = T)
## [1] 2.9
```

¹Bu formül popülasyon için yanlı (biased) bir tahminleyicidir. R yanlı olmayan çarpıklık ve basıklık istatistikleri hesaplayabilir, *describe* fonksiyonu *type* argümanı incelenebilir.




Figure 7.1: Normal dagilim gösteren bir sürekli degisken




Figure 7.2: Sola çarpık sürekli degisken




Figure 7.3: Saga çarpık sürekli degisken

7.1.6.1 Basıklık Örnekleri

Normal bir dağılım ve basıklık ölçüsü




Figure 7.4: Normal dagilim gösteren bir sürekli degisken

Tekdüzey bir dağılım ve basıklık değeri

Beta dağılımı gösteren bir sürekli değişken

7.1.7 Betimleyici İstatistiklerin Raporlanması

psych (Revelle, 2016), *doBy* (Højsgaard and Halekoh, 2016) ve *apaStyle* (de Vreeze, 2016) paketleri betimsel analizleri rapor etmede yardımcı olabilir.

```
# psych paketi describe fonksiyonu sırasıyla;
# n: gözlem sayısı (kayıp veriler hariç)
# ortalama, ss, ortanca, budanmış ortalama (trim=0.05 5% budanmış)
# ortanca mutlak dağılımı (median absolute deviation),
# minimum, maksimum, ranj
# çarpıklık ve basıklık-3 (type=2 popülasyon basıklık ve çarpıklık)
# standart hata
library(psych)
```




Figure 7.5: Tekdüzey dagilim gösteren sürekli degisken




Figure 7.6: Beta dagilimi gösteren bir sürekli degisken

```

desc1=describe(dataWBT[,c("gen_att","age")],trim = 0.05,type=3)
desc1
##          vars      n mean   sd median trimmed  mad min max range skew
## gen_att    1 5302  1.94 0.60      2    1.92 0.59   1  4    3 0.38
## age        2 5308 27.08 7.21     25    26.62 5.93  15 60   45 0.96
##          kurtosis    se
## gen_att    -0.10 0.01
## age         0.63 0.10

# kaydet
write.csv(desc1,file="pscyhbetimsel.csv")

#doBy
# program değişkenine göre betimleyici istatistikler
library(doBy)
library(moments)
desc2=as.matrix(summaryBy(gen_att+age~treatment, data = dataWBT,
  FUN = function(x) { c(n = sum(!is.na(x)), nmis=sum(is.na(x)),
    m = mean(x,na.rm=T), s = sd(x,na.rm=T),
    skw=moments::skewness(x,na.rm=T),
    krt=moments::kurtosis(x,na.rm=T)) } ))

#yuvarlama
round(desc2,2)
##      treatment gen_att.n gen_att.nmis gen_att.m gen_att.s gen_att.skw
## 1           1      2736          265     1.93      0.6      0.38
## 2           2      2566          335     1.95      0.6      0.38
##      gen_att.krt age.n age.nmis age.m age.s age.skw age.krt
## 1           2.90 2739      262 26.9 7.17 0.99 3.69
## 2           2.91 2569      332 27.3 7.24 0.93 3.57
write.csv(round(desc2,2),file="doBydesc.csv")

#apaStyle
# APA formatında tablo
library(apaStyle)
apa.descriptives(data = dataWBT[,c("gen_att","age")],
  variables = c("Gender Attitude","Age"), report = c("M", "SD"),
  title = "APAtableGenderAge", filename = "APAtableGenderAge.docx",
  note = NULL, position = "lower", merge = FALSE,
  landscape = FALSE, save = TRUE)
##
## Word document succesfully generated in: C:/Users/Burak/Desktop/github/SARP

#apaStyle paketi hata veriyorsa;
#https://www.r-statistics.com/2012/08/how-to-load-the-rjava-package-after-the-error-java_home-cannot-be
#Sys.setenv(JAVA_HOME='C:\\Program Files\\Java\\jre1.8.0_111')

```

7.1.7.1 Betimsel İstatistik Rapor Örneği

Toplumsal cinsiyet algısı puanları 5302 katılımcı için 1 ve 4 arasında değişmiştir, ortanca 2, ortalama 1.94 ve standart sapma 0.6 olarak hesaplanmıştır. Puanların dağılımı örneklem bazında 0.38 çarpıklık ve -0.1




Figure 7.7: APA Tablo.docx

basıklık değerine sahiptir.

7.2 Basit Grafikler

R programı ile basit olmayan grafikler çizilebilir. Popüler olan grafik oluşturma yöntemlerinden dört tanesi, base(R Core Team, 2016b), lattice(Sarkar, 2016), ggplot2(Wickham and Chang, 2016) ve plotrix(Lemon et al., 2016). Bu materyal ggplot2 kullanmıştır. Bir ggplot fonksiyonunda argüman sayısı oldukça fazladır, bu sayede kullanıcılar grafiğin her noktasında değişiklik yapabilirler.

7.2.1 Histogram

Dikdörtgenlerden oluşan histogram grafikleri değişken içerisinde yer alan değerlerin frekanslarına göre oluşturulur.

7.2.1.1 Tek değişken için histogram

Dağılım hakkında bilgi sahibi olmak için kullanışlıdır.

```
library(ggplot2)
ggplot(dataWBT, aes(x = gen_att)) +
  geom_histogram(binwidth = 0.2)+ theme_bw()+labs(x = "Toplumsal Cinsiyet Algısı ") +
  theme(axis.text=element_text(size=15),
        axis.title=element_text(size=14,face="bold"))
```

7.2.1.2 Tek değişken tek faktör histogram

Gruplara dayalı farklılıkları görmek için kullanışlı

```
dataWBT$HEF=droplevels(factor(dataWBT$higher_ed,
                              levels = c(0,1),
                              labels = c("Lise ve altı", "Üniversite"))))

ggplot(dataWBT, aes(x = gen_att, fill=HEF,drop=T)) +
  geom_histogram(breaks=seq(1, 4, by =0.2),alpha=.5,col="black")+
  theme_bw()+labs(x = "Toplumsal Cinsiyet Algısı",fill='Yüksek Öğretim Durumu')+
  theme(axis.text=element_text(size=15),
        axis.title=element_text(size=14,face="bold"))

dataWBT2=na.omit(dataWBT[,c("gen_att", "HEF")])
ggplot(dataWBT2, aes(x = gen_att)) +
  geom_histogram(breaks=seq(1, 4, by =0.2),alpha=.5,col="black")+
  theme_bw()+labs(x = "Toplumsal Cinsiyet Algısı")+ facet_wrap(~ HEF)+
  theme(axis.text=element_text(size=15),
        axis.title=element_text(size=14,face="bold"))

library(ggplot2)
ggplot(dataWBT, aes(x = gen_att)) +
  geom_histogram(binwidth = 0.2)+ theme_bw()+
  facet_wrap(~city, ncol = 8)
```




Figure 7.8: Toplumsal Cinsiyet Algisi Puan Dagilimi





Figure 7.9: Eğitime Göre Toplumsal Cinsiyet Algisi




7.2.1.3 Tek değişken iki faktör histogram

Useful for two way interactions

```
dataWBT2=na.omit(dataWBT[,c("gen_att","HEF","gender")])

ggplot(dataWBT2, aes(x = gen_att,fill=gender)) +labs(fill='Gender')+
  geom_histogram(binwidth = 0.2,alpha=.5)+ theme_bw()+
  facet_grid(~HEF)
```




7.2.1.4 Tek değişken üç faktör histogram

Etkileşim (interaction) açıklamada kullanışlı

```
dataWBT$Condition=droplevels(factor(dataWBT$treatment,
                                     levels = c(1,2),
                                     labels = c("treatment", "control")))

dataWBT2=na.omit(dataWBT[,c("gen_att", "HEF", "gender", "Condition")])

ggplot(dataWBT2, aes(x = gen_att, fill=gender)) +labs(fill='Gender')+
  geom_histogram(binwidth = 0.2, alpha=.5)+ theme_bw()+
  facet_grid(~HEF+Condition)
```



7.3 Hipotez Testi Tanıtım

Cambridge sözlüğü evren (popülasyon) tanımı olarak “aynı ülke, aynı alan veya aynı yerde yaşayan insan veya canlı grubu” cümlesini kullanır. Sosyal bilimlerde evren genellikle “belli bir gruba ait bütün insanlar” olarak belirlenir. Örneğin *sekiz yaşındaki tüm öğrenciler, belli bir ülkede bulunan sekiz yaşındaki tüm öğrenciler, 8 yaşında disleksi teşhisi konulan öğrenciler*.

Sosyal bilimciler araştırma soruları doğrultusunda ilgili evreni tanımlar. Evrende yer alan bireylerin (unit) gözlenlenebilen karakteristik özellikleri değişkenleri oluşturur. Diğer bir ifade ile değişkene ait popülasyon tanımlanabilir. Bölüm 5.2.3 'de değişken türleri açıklanmıştır. Evren değişkene ait bütün değerleri kapsar, bu değerlere ait bir ranj ve görülme olasılığı (probability of occurrence) vardır. Yoğunluk (probability, sürekli değişken için) ve çoğunluk (mass, sürekli olmayan değişken için) fonksiyonları görülme olasılıklarını formüle etmek için kullanışlıdır. Dağılım hakkında yapılacak geçerli bir varsayım ile örneklemden evrene genellemeler yapılabilir.

Seçkisiz seçilen bir örneklem ile ulaşılan değişken evrende sahip olduğu bütün değerleri içermeyebilir. Fakat, özellikle gözlem sayısı küçük değil ise, seçkisiz seçme işleminde sistematik bir yanlılık görülmeceği düşünüldüğünde, örneklemin evrene benzer özellikler göstermesi beklenir. Bu bilgi kullanılarak evrene ait bir parametre örneklemden yola çıkarak tahmin edilebilir. Bu işlem genellikle bir model sayesinde olur. Modelin veriye gösterdiği uyum araştırmacı tarafından değerlendirilir. Hipotez testleri kullanılan bir model sonrasında araştırma soruları ile ilgili karara varma sürecidir.

7.3.1 Örneklem Dağılım (Sampling Distribution)

Seçkisiz bir örneklem için hesaplanmış bir istatistik aslında bir değişkendir ve belli bir dağılıma sahiptir. Örneklem dağılım konusunu açıklamak için en çok kullanılan istatistik aritmetik ortalamadır. Merkezi limit teoremine göre basit seçkisiz örneklem kullanıldığında ² değişkenin evrende gösterdiği dağılımdan bağımsız olarak, o değişkene ait örneklem ortalamalarının dağılımı yaklaşık olarak normaldir. Örneklem büyüdükçe ;

$$\bar{X}_n \sim N(\mu, \frac{\sigma^2}{n}). \quad (7.6)$$

Eğer evren bazında dağılım normal ise (7.6) küçük örneklem için de doğrudur. Ortalamanın örneklem dağılımına ait standart sapma *ortalamanın standart hatası* olarak isimlendirilir ve istatistiksel çıkarımlarda kullanılır. Eşitlik (7.6) içinde yer alan μ ve σ^2 bilinmezdir. Fakat bu eşitlik örneklem ait ortalamanın evrene ait ortalamayı ne derecede kestirebileceğini anlamada önemlidir. Örneğin bir araştırmacının basit seçkisiz yöntemle 10 kişilik bir örneklem seçtiğini düşünelim. Araştırmacının bilemediği parametrelerin ise $\mu = 100$ ve $\sigma = 15$ olduğunu düşünelim. Bu durumda örneklem dağılım için standart sapma $(15/\sqrt{10}) = 4.74$ olarak bulunur. %95 olasılıkla araştırmacının 10 kişilik örneklem ile ulaşacağı aritmetik ortalama 90.7 ve 109.3 arasında olacaktır. Bu oldukça geniş bir aralıktır. Fakat araştırmacı 10 kişi yerine 100 kişiyi aynı örneklem ile seçseydi ulaşacağı örneklem ortalaması %95 olasılıkla 97.1 ve 102.9 olacaktır.

Bu noktada önem taşıyan konu, örneklemden gelen bilgilerle evrene ait parametrelerin hangi tahminleme yöntemleri (estimator) ile yansız, tutarlı ve keskin (unbiased, consistent and efficient) olarak kestirebileceğidir. Örneğin μ , Eşitlik (7.1) ile, σ^2 ise Eşitlik (7.2) ile yansız olarak kestirilebilir.

7.3.1.1 Yansız tahminleme ve örneklem seçimi

Eklenecek

7.3.2 Güven Aralıkları (The Confidence Intervals (CI))

Dağılım hakkında yapılacak bir varsayım, örneklemden gelen bilgi ve uygun bir tahminleyici (estimator to produce a point estimate) kullanılarak güven aralıkları oluşturulabilir. Bir güven aralığı evren parametresinin muhtemelen hangi aralıkta olduğunu gösterir. Fakat bu evren parametresinin bu aralıkta kesinlikle yer aldığı anlamına gelmez. Örneklem ait aritmetik ortalamadan yola çıkarak evren parametresi için güven aralığı hesaplamak oldukça basittir. Değişkene ait dağılımın normal olduğu varsayıldığında, ortalamanın örneklem dağılımı da normaldir ve örneklem ait ortalama yansız bir tahmindir. Normal dağılım bilindik özellikleri vardır, yoğunluk fonksiyonu değerlerin %95'inin ortalamadan 1.96 standart sapma aşağıda ve yukarıda olduğunu gösterir. Normal dağılımın bu özelliği grafik 7.10 ile gösterilmiştir. Mavi ile işaretlenen bölgeden bir gözlem yapma olasılığı %5'tir. Benzer şekilde, mavi veya sarı ile işaretlenen bölgeden gözlem yapma olasılığı da %10'dur. Gri bölge (± 1) yoğunluğun yaklaşık olarak %68'ini kapsar. Bu bilgi kullanışlıdır. Örneklem için hesaplanan ortalama ve varyans μ için güven aralığı hesaplamada kullanılabilir.

7.3.2.1 Güven Aralığı Örneği

Toplumsal Cinsiyet Algısına ait ortalama ve güven aralığı hesaplamaları

```
# gözlem sayısı, n
GA_n=sum(!is.na(data$BT$gen_att))

#ortalama
```

²evrende yer alan her üyenin eşit seçilme olasılığı olduğunda ve seçilen bir üyenin diğer bir üyenin seçilme olasılığını etkilemediği durumlarda




Figure 7.10: The z distribution

```
GA_m=mean(dataWBT$gen_att,na.rm = T)

#ss
GA_s=sd(dataWBT$gen_att,na.rm = T)

#95% güven aralığı
alt=GA_m - 1.96 * (GA_s/sqrt(GA_n))
alt
## [1] 1.92
ust=GA_m + 1.96 * (GA_s/sqrt(GA_n))
ust
## [1] 1.96

#veya
GA_m +c(-1,1)*1.96 * (GA_s/sqrt(GA_n))
## [1] 1.92 1.96

#1.96 için qnorm(0.975)
```

7.3.2.2 Ortalama için güven aralığı raporlama

5302 katılımcı için Toplumsal Cinsiyet Algısı puanlarına ait ortalama 1.94 (%95 GA [1.92-1.96]), standart sapma 0.60 bulunmuştur.

7.3.3 Boş Hipotez

Hipotez testinin amacı evren hakkında öne sürülen iki hipotezden hangisinin örneklem tarafından desteklen-
diği karar vermektir. Bir hipotez testi beş basamaktan oluşur;

- 1) Boş hipotezin³ belirlenmesi (örneğin $\mu = 0$)
- 2) Alternatif hipotezin belirlenmesi. (örneğin $\mu \neq 0$)
- 3) Test istatistiğinin seçilmesi
- 4) Test istatistiği ve belirlenen kritik değeri karşılaştırarak karar verilmesi. Eğer hesaplanan test istatistiği kritik değerden daha yüksekse boş hipotezin ret edilmesi (kritik değer alternatif hipoteze göre değişir).
- 5) Sonucun açıklanması. Araştırma sorusuna yanıt vermek üzere kararın ifade edilmesi.

Boş hipotez (H_0) ve alternatif hipotez (H_1) araştırma sorusunu cevaplamak üzere belirlenir. İstatistiksel kanıtlar boş hipotezini kabul veya terketmek için kullanılır. Boş hipotezi kabul etmek veya terketmek bir karardır, teorik istatistik açısından bu karara yönelik oluşabilecek durumlar şunlardır;

Gerçekte Durum	Karar	Sonuç
H_0	Kabul H_0	Doğru Karar
H_0	Red H_0	Yanlış Karar (<i>Tip I hata, α</i>)
H_1	Red H_0	Doğru Karar
H_1	Kabul H_0	Yanlış Karar (<i>Tip II hata, β</i>)

Tip-I hata: Gerçekte doğru olduğu halde boş hipotezin terkedilmesi durumudur. Alfa , α , boş hipotezin

³yokluk veya sıfır hipotezi olarak da bilinir

gerçekte doğru olduğu halde red edilme olasılığıdır. Hipotez testi sürecinde önemli olan noktalardan biri α 'nın yeterince küçük olması gerektiğidir. Sosyal bilimlerde sıklıkla $\alpha = .05$ kullanılır.

Tip-II hata: Gerçekte yanlış olan bir boş hipotezin terkedilememesidir. Beta, β , gerçekte yanlış olan bir boş hipotezi kabul etme olasılığıdır.

7.3.4 z-puanı ve z-testi

z-puanı için genel formül;

$$z_X = \frac{X - \bar{X}}{s_X}$$

Hesaplanan bu z değişkeni 0 ortalamaya ve 1 standart sapmaya sahiptir. Eğer X normal dağılım gösteriyorsa z de normal dağılım gösterir.

z puanı hesapla

```
GA_m=mean(dataWBT$gen_att,na.rm = T)
GA_s=sd(dataWBT$gen_att,na.rm = T)
z_GA=(dataWBT$gen_att-GA_m)/GA_s
```

#veya

```
z_GA=scale(dataWBT$gen_att, center=T, scale=T)
```

scale fonksiyonu ile birden fazla değişken için z hesaplanabilir.

center=T her X puanından ortalamayı çıkarır

scale=T farkı standart sapmaya böler

scale(dataWBT\$gen_att, center=3, scale=2)her değerden 3 çıkarıp 2'ye böler.

Ortalama için z-istatistiği hesaplamak kolaydır;

$$z = \frac{\bar{X} - \mu_{hipotez}}{\text{StandartHata}} = \frac{\bar{X} - \mu_{hipotez}}{\sigma_X / \sqrt{n}}$$

Hesaplanan bu z istatistiği bir z dağılımı kullanılarak yorumlanabilir (Figür 7.10);

- Eğer alternatif hipotez, gözlemlenen ortalamanın, hipotez değerinden küçük olacağını belirtiyorsa, hesaplanan z istatistiği z_{α} veya $-z_{(1-\alpha)}$ ile kıyaslanır. Eğer hesaplanan z , z_{α} 'ya eşit veya küçük ise boş hipotez terkedilir.
- Eğer alternatif hipotez, gözlemlenen ortalamanın, hipotez değerinden farklı olacağını belirtiyorsa, hesaplanan z istatistiğinin mutlak değeri $z_{1-(\alpha/2)}$ ile kıyaslanır. Eğer mutlak z , $z_{1-(\alpha/2)}$ 'ya eşit veya büyük ise boş hipotez terkedilir.
- Eğer alternatif hipotez, gözlemlenen ortalamanın, hipotez değerinden büyük olacağını belirtiyorsa, hesaplanan z istatistiği $z_{1-\alpha}$ ile kıyaslanır. Eğer hesaplanan z , $z_{1-(\alpha)}$ 'ya eşit veya büyük ise boş hipotez terkedilir.

Burada dikkat edilmesi gereken nokta a ve c senaryolarında (yönlü alternatif) kullanılan kritik değerin b senaryosunda (yönsüz alternatif) kullanılan kritik değerden farklı oluşudur. Araştırmacılar alternatif hipotezlerinin yönlü veya yönsüz oluşunu savunabilmelidir.

7.3.4.1 z testi örnek-1 (yönsüz)

Boş hipotez $H_0 : \mu_{CinsiyetAlgisi} = 2$ ve alternatif hipotez $H_1 : \mu_{CinsiyetAlgisi} \neq 2$ ve $\alpha = 0.05$;

```
# n
GA_n=sum(!is.na(data$WT$gen_att))

#ortalama
GA_m=mean(data$WT$gen_att,na.rm = T)

#ss
GA_s=sd(data$WT$gen_att,na.rm = T)

# boş hipotez
mu_hyp=2

# z istatistiği
(GA_m-mu_hyp)/(GA_s/sqrt(GA_n))
## [1] -7.17

#alpha=0.05 ve yönsüz alternatif için kritik değer
qnorm(1-(0.05/2))
## [1] 1.96
```

5302 katılımcıya ait Toplumsal Cinsiyet Algısı puanları için ortalama 1.94 ve standart sapma 0.6 olarak hesaplanmıştır. Tek örneklem için hesaplanan z testi, gözlemlenen ortalamanın, hipotez ile öne sürülen 2'den 7.17 standart hata daha düşük olduğunu göstermiştir. Kritik değer olarak 1.96 ($z_{1-(0.05/2)}$) seçildiğinde ve gözlemlenen ortalama ve hipotez ile öne sürülen ortalama arasındaki farkın istatistiksel olarak anlamlı olduğu kararı verilmiştir.

7.3.4.2 z testi örnek-2 (yönlü)

Bu örnekte Toplumsal Cinsiyet Algısı değişkeninin evren bazında ortalaması 1.9 ve standart sapması 0.75 olarak varsayılmıştır. Eğer evren bazında standart sapma biliniyorsa istatistik hesaplarken kullanılmalıdır. $H_0 : \mu_{CinsiyetAlgisi} = 1.9$ ve $H_1 : \mu_{CinsiyetAlgisi} > 1.9$ ve $\alpha = 0.01$;

```
# boş hipotez
mu_hyp=1.9

# z istatistik
(GA_m-mu_hyp)/(0.75/sqrt(GA_n))
## [1] 3.94

#yönlü alternatif ve alfa=.01
qnorm(1-(0.01))
## [1] 2.33
```

Kritik değer olarak 2.33 kullanıldığında ($z_{0.99}$) gözlemlenen ortalamanın hipotez ile öne sürülen ortalamadan büyük olduğu ve bu farkın istatistiksel olarak anlamlı olduğu kararına varılmıştır $z = 3.94$.

7.3.5 Tek örneklem t-testi

Evrene ait dağılım normal olsa dahi küçük örneklem için z dağılımı yerine t dağılımı kullanmak daha geçerlidir. bu t dağılımının serbestlik derecesi n-1'dir.

7.3.5.1 t test örnek-1 (yönsüz)

Örnek için Düzce ilinde yaşadığını belirten katılımcılara ait Toplumsal Cinsiyet Algısı puanları kullanılmıştır.

$H_0 : \mu_{CinsiyetAlgisi} = 1.94$ ve alternatif $H_1 : \mu_{CinsiyetAlgisi} \neq 1.94$ ve $\alpha = 0.05$;

```
dataWBT_DUZCE=dataWBT[dataWBT$city=="DUZCE",]
#betimleyici
describe(dataWBT_DUZCE[, "gen_att"], type=3)
##      vars  n mean   sd median trimmed  mad min max range skew kurtosis   se
## X1      1 47 2.18 0.55      2    2.14 0.59   1 3.8   2.8 0.56      0.28 0.08

#t test
t.test(dataWBT_DUZCE$gen_att,
       alternative="two.sided",
       mu=1.94,
       conf.level = 0.95)

##
## One Sample t-test
##
## data:  dataWBT_DUZCE$gen_att
## t = 3, df = 50, p-value = 0.005
## alternative hypothesis: true mean is not equal to 1.94
## 95 percent confidence interval:
##  2.01 2.34
## sample estimates:
## mean of x
##      2.18

#kritik değer
qt(.975,df=46)
## [1] 2.01
```

Düzce ilinde yaşayan katılımcıların Cinsiyet Algısı puanları 1 ve 3.8 arasında değişmiştir, ortanca 2, ortalama 2.18, standart sapma 0.55, örnekleme ait dağılımın çarpıklığı 0.56 ve basıklığı 0.28 olarak hesaplanmıştır. Düzce şehrinde yaşayan katılımcılara ait Toplumsal Cinsiyet Algısı ortalaması, hipotez ile öne sürülen 1.94 değerinden farklıdır ve bu farklılık istatistiksel olarak anlamlıdır, $t(46)=2.94$ ve $t_{.975,46} = 2.01$

7.3.5.2 t test örnek-2

$H_0 : \mu_{CinsiyetAlgisi} = 1.94$ ve alternatif $H_1 : \mu_{CinsiyetAlgisi} \leq 1.94$ ve $\alpha = 0.05$;

```
#t test
t.test(dataWBT_DUZCE$gen_att,
       alternative="less",
       mu=1.94,
       conf.level = 0.95)

##
## One Sample t-test
##
## data:  dataWBT_DUZCE$gen_att
## t = 3, df = 50, p-value = 1
## alternative hypothesis: true mean is less than 1.94
## 95 percent confidence interval:
## -Inf 2.31
```

```
## sample estimates:
## mean of x
##      2.18

#kritik deęer
qt(.05,df=46)
## [1] -1.68
```

Test istatistięi $t(46)=2.94$ ve kritik deęer $t_{.05,46} = -1.68$ kullanılarak rneklemenin, evrene ait ortalamannın 1.94'ten kk olduęu hipotezini destekleyecek kanıtı iermedięi kararı verilmiřtir.

7.3.6 p-deęeri

Tek rneklem ile t testi iin kullanılan t.test fonksiyonu bir p deęeri rapor etmiřtir. Bu p deęeri hesabı, boř hipotezin ve daęılım iin yapılan varsayımın doęru olduęu kabul zerine yapılır. Bu p-deęerinin amacı arařtıracıyı hesaplanan istatistięin sıradan olup olmadıęı ynnde bilgilendirmektir. Uzun yıllardır arařtırmacılar hesapladıkları bu p-deęerini daha nceden belirledikleri bir alfa kriteri ile kıyaslayıp, buldukları sonuların istatistiksel olarak anlamlı olup olmadıęına karar vermiřlerdir.

7.3.7 p deęeri rnek-1

Bir z-daęılımın geerli olduęu ve z deęerinin 1.80 hesaplandıęı durum iin grafik;




Figure 7.11: z daęilimi ve $z=1.8$

Mavi alan yoğunluğun %3.6'sını gösterir, $p=0.0359$

```
1-pnorm(1.8)
```

```
## [1] 0.0359
```

Bu p değeri yönlü bir alternatif hipotez için hesaplanmıştır. Yönsüz alternatif için geçerli değildir. Yönsüz alternatif için ;




Figure 7.12: z dağılımı ve $|z|=1.8$

Mavi alan yoğunluğun %7.2'sini gösterir, $p=0.0719$;

```
2*(1-pnorm(1.8))
```

```
## [1] 0.0719
```

7.3.8 İstatiksel Güç

İstatistiksel güç, gerçekte yanlış olan bir boş hipotezi terketme olasılığıdır ve $(1-\beta)$ 'ya eşittir. Bu olasılık istatistiksel sınamalar yapıldıktan önce (a-priori) veya sonra (post-hoc) hesaplanabilir, fakat sınama sonrasında yapılan güç analizi genellikle işlevsizdir. Sınama yapılmadan önce, daha doğrusu veriler toplanmadan önce yapılacak bir güç hesabı ile çalışma tasarısı gözden geçirebilir, yeniden düzenlenebilir ve örneklem sayısı belirlenebilir. Amerika'da bir çok proje başvurusu istatistiksel güç analizlerini mecbur tutar.

İstatistiksel gücü açıklamak için R ile çizilen grafik ve kod⁴;

⁴partially based on <http://multithreaded.stitchfix.com/blog/2015/05/26/significant-sample/>


```

x <- seq(-4, 8, 0.02)
zdat <- data.frame(x = x, y1 = dnorm(x, 0, 1), y2 = dnorm(x, 2.5, 1))
ggplot(zdat, aes(x = x)) +
  geom_line(aes(y = y1), size=2) +
  geom_line(aes(y = y2), color='red', size=2) +
  geom_vline(xintercept = c(0,2.5), color="black", linetype = "longdash")+
  geom_vline(xintercept = qnorm(1 - 0.05))+
  scale_x_continuous(breaks = c(-4,0,1.65,2.5,4))+
  annotate("text", label="beta", x=1.1, y=0.05, parse=T, fontface =2, size=6)+
  annotate("text", label="alpha", x=2, y=0.02, parse=T, fontface =2, size=6)+
  annotate("text", label="1-~beta", x=3.3, y=0.1, parse=T, fontface =2, size=6)+
  geom_area(aes(y=y1, x = ifelse(x > qnorm(.95), x, NA)), fill = 'blue', alpha=0.25) +
  geom_area(aes(y=y2, x = ifelse(x > qnorm(.95), x, NA)), fill = 'green', alpha=0.25) +
  geom_area(aes(y=y2, x = ifelse(x < qnorm(.95), x, NA)), fill = 'yellow', alpha=0.25) +
  xlab("z") + ylab("dnorm(z)") + theme_bw()

```




Figure 7.13: z dağılımı ile istatistiksel güç

$H_0 : \mu = 0$ boş hipotezini doğru kabul eden bir z dağılımı siyah çizgiler ile gösterilmiştir, bu dağılımın ortalaması sıfırdır ve kesik çizgiler ile gösterilmiştir. Kırmızı çizgiler ile gösterilen dağılım (a) boş hipotezin yanlış olduğu ve (b) evrene ait ortalama ve standard sapma ile hesaplanan z -istatistiğinin $((\mu - \mu_{\text{hypothesis}})(\sigma\sqrt{n}) = 2.5)$ olduğu varsayımı ile çizilmiştir. Bu görsel yönlü bir alternatif hipotez ve $\alpha=0.05$, dolayısıyla kritik değer $z_{0.95} = 1.65$ için geçerlidir. Mavi alan α 'yı, sarı alan β 'yı ve yeşil alan istatistiksel gücü yansıtır. Bu görselde istatistiksel güç .804'tür.

```
1-pnorm(qnorm(0.95),mean=2.5)
```

```
## [1] 0.804
```

Figure⁵ 7.13 , istatistiksel güç hesaplamak için iki farklı dağılımın, bir alfa değerinin ve bir test istatistiğinin olması gerektiğini gösterir. Bu bilinenler ile istatistiksel güç hesaplanabilir. Burda önemli olan detay bir test istatistiğinin kendi bileşenleri olduğudur, genellikle bu bileşenler bir bölünen ve bir bölendir. z testi için bölünen , hipotez ile öne sürülen değer ile gözlemlenen değer arasındaki fark, bölen ise ortalamanın standart hatasıdır (σ/\sqrt{n}). Eğer istatistiksel güç sabit tutulursa (örneğin .8) eşitlik seçilen bir bilinmeyene göre çözülebilir. Genellikle de seçilen bilinmeyen örneklem sayısıdır, n .

İstatistiksel güç ilerleyen bölümlerde tekrar değinilmiştir. Test istatistikleri, parametre kestirimleri, standart hatalar, dağılımsal varsayımlar tasarlanan araştırmaya göre değişecektir. Tek örneklem için t testi düşünüldüğünde *power.t.test* fonksiyonu işeyarardır.

```
#power.t.test
power.t.test(delta=.1, sd=.6,sig.level=0.05, power=0.9,
              type="one.sample", alternative="one.sided")

##
##      One-sample t test power calculation
##
##              n = 310
##            delta = 0.1
##              sd = 0.6
##          sig.level = 0.05
##            power = 0.9
##      alternative = one.sided
```

Bu örnek, belirlenmiş bir ortalama fark 0.1, standart sapma 0.6, alfa 0.05, yönlü alternatif ve istenilen güç 0.9 için örneklem 310 olması gerektiğini gösterir. Bir diğer ifade ile, araştırmacı 310 kişiden gelen bir veride, ortalama fark 0.1 , standart sapma 0.6 tespit eder ve alfa 0.05 ile yönlü bir test kullanırsa boş hipotezi ($H_0 : \mu = 0$) terketme olasılığı 0.90'dır.

7.3.9 z ve t dağılımları geçerli değil ise

Bilinenlerden (örneklem) bilinmeyenlere (evren) genelleme varsayımların yapılmasını gerektirir. Bir test istatistiğine ait örneklem dağılımın, belli bir örneklem sayısında, belirli bir varsayımın ihlal edilmesi ile büyük ölçüde değişmemesi durumu *direnç* (robustness) olarak isimlendirilir (Verzani (2014)). Burda dikkat edilmesi gereken, bir test istatistiğinin bir varsayım ihlaline dirençli iken başka bir varsayım ihlaline dirençsiz olabileceğidir. Ayrıca, bir varsayım ihlaline dirençli olan bir test istatistiği, ikinci bir varsayım ihlalinin de yaşanması durumunda direncini yitirebilir. Bir test istatistiği dirençli olduğu için kullanılması şart değildir çünkü aynı şartlar altında daha iyi çalışan bir başka test istatistiği olabilir.

z istatistiği, örneklem 30'dan büyük ise normallik varsayımının ihlallerine karşı dirençlidir(Field et al. (2012), page 198). Örneklem dağılımın z dağılımına yakınlığını etkileyen bir faktör diğer faktörde örneklem nasıl bir dağılım gösterdiğidir. Ayrıca, evrene ait dağılımın normal olduğu varsayımı ile küçük örneklem için t dağılımı geçerlidir.

Tek örneklem ile aritmetik ortalama için yapılacak dirençli istatistikler detaylı olarak Wilcox (2012) tarafından verilmiştir. Verilen R kodu bootstrap-t metodu için %95 güven aralığı hesaplar (Wilcox (2012), page 117).

```
#ikinci tür bootstrap t metodu
# Düzce katılımcılarını seç
dataWBT_DUZCE=na.omit(dataWBT[dataWBT$city=="DUZCE",c("id","gen_att")])
```

⁵accurate only for post hoc power

```

# normallik varsayımı ve t test kullanarak
# evren ortalamasının 1.94 olup olmadığını sına
t.test(dataWBT_DUZCE$gen_att,mu=1.94,conf.level = 0.95)
##
## One Sample t-test
##
## data: dataWBT_DUZCE$gen_att
## t = 3, df = 50, p-value = 0.005
## alternative hypothesis: true mean is not equal to 1.94
## 95 percent confidence interval:
## 2.01 2.34
## sample estimates:
## mean of x
## 2.18

# bootstrap ile 95% GA (normallik varsayımı yok)
set.seed(04012017)
B=5000 # bootstrap sayısı
alpha=0.05 # alfa

#x değişken
# xBAR gözlemlenen ortalama
tstar=function(x,xBAR) sqrt(length(x))*abs(mean(x)-xBAR)/sd(x)

output=c()
for (i in 1:B){
  output[i]=tstar(sample(dataWBT_DUZCE$gen_att,
                        replace=T,
                        size=length(dataWBT_DUZCE$gen_att)),
                  xBAR=mean(dataWBT_DUZCE$gen_att))
}
output=sort(output)
Tc=output[as.integer(B*(1-alpha))]

#bootstrap GA
mean(dataWBT_DUZCE$gen_att)+c(-1,1)*(Tc*sd(dataWBT_DUZCE$gen_att)/sqrt(length(dataWBT_DUZCE$gen_att)))
## [1] 2.01 2.34

```

7.3.9.1 Raporlama

Düzce ilinden 47 katılımcının verdiği yanıtlar ile hesaplanan Toplumsal Cinsiyet Algısı puanları 1 ve 3.8 arasında değişmiş, ortancası 2, ortalaması 2.18, standart sapması 0.55 bulunmuştur. Puanların dağılımına ait çarpıklık değeri 0.56, basıklık değeri 0.28 olarak hesaplanmıştır. Kritik değer olarak 2.01 ($t_{.975,46}$) kullanıldığında, tek örneklem t testi anlamlı bir farklılığa işaret etmiştir, $t(46)=2.94$, bu şehirdeki katılımcıların puanları hipotez ile öne sürülen 1.94 değerinden farklıdır. Normallik varsayımı yapıldığında %95 güven aralığı [2.01,2.34] olarak bulunmuştur. Normallik varsayımı yapılmadığında 5000 tekrarlı bootstrap metodu ile hesaplanan güven aralığı [2.01,2.34] olarak bulunmuştur.

Chapter 8

İki Ortalamamanın Karşılaştırılması, t-testi

Bölüm 7.3.1 örnekleme dağılım (sampling distribution) konusunun ana hatlarını tek bir aritmetik ortalama üzerinden ele almıştır. Eğer iki farklı aritmetik ortalama kıyaslanmak isteniyorsa t testi kullanılabilir, bu prosedür aritmetik ortalamaların örnekleme dağılımı üzerine inşaa edilmiştir.

$\bar{Y}_1 - \bar{Y}_2$ ($\mu_{\bar{Y}_1 - \bar{Y}_2}$) sonucunun örnekleme dağılım ortalaması daima $\mu_1 - \mu_2$ 'dir. Fakat örnekleme dağılım standart sapması ($\sigma_{\bar{Y}_1 - \bar{Y}_2}$) araştırmanın tasarısına göre değişir.

Örnek Bir cerrahın yaraların iyileşmesi konusunda araştırma yaptığını düşünelim. Cerrahın kapanan bir yaradan sonra gerilme direncini araştırdığını, yara bandı ve dikiş atma tedavisinin arasında bir fark olup olmadığını araştırdığını varsayalım. Bu çalışmada tek bir faktör vardır, yara kapatma yöntemi ve bu faktöre ait iki alt sınıf vardır, yara bandı ve dikiş. Bu araştırmayı iki farklı şekilde tasarlamak mümkündür.

Bağlı gözlemler (within-subjects) 10 tavşanın her birinin sırtına (omurganın sağına ve soluna) 2 kesik oluşturulur. 2 kesikten bir tanesi yeni geliştirilen bir yara bandı ile diğeri dikiş ile kapatılır, hangi kapatma yönteminin hangi yarayı kapatacağı rassal (random) seçilmelidir. Bu tasarı *bağlı-gözlemler* olarak isimlendirilmiştir çünkü faktöre ait iki alt sınıf aynı tavşan üzerinde gözlemlenmiştir.

Bağlı olmayan gözlemler (between-subjects) 20 tavşan rassal olarak 2 gruba ayrılır, birinci grupta yer alan tavşanların yaraları bant ile, ikinci grupta yer alan tavşanların yaraları ise dikiş ile kapatılır. Yaralar omurganın sağ veya sol tarafında rassal olarak açılmalıdır. Bu tasarı *bağlı olmayan gözlemler* olarak isimlendirilmiştir çünkü faktöre ait alt sınıflar farklı tavşanlar üzerinde gözlemlenmiştir ve bu tavşanların herhangi bir şekilde eşlenmiş değillerdir. Örneğin aynı anneden gelen iki tavşan rassal olarak gruplara atansa idi gözlemler bağlı olurdu.

Her iki yara kapatma yönteminden sonra yapılacak gerilme direnci ölçümlerinin evren bazında bir ortalaması ve standart sapması olduğu aşikardır.

Konuyu açıklama amaçlı, yara bandı yönteminden sonra yapılan gerilme direnci ölçümlerine ait ortalamamanın ve standart sapmanın bağlı gözlem veya bağlı olmayan gözlem tasarılarında aynı olduğunu düşünelim.

Parametres	Bant	Dikiş
Ortalama	μ_B	μ_D
Standart Sapma	σ_B	σ_D
Örneklem	n_B	n_D

Buradan itibaren μ_B yerine μ_1 , μ_D yerine μ_2 , σ_B yerine σ_1 , σ_D yerine σ_2 kullanılmıştır.

Örnekleme dağılım parametresi	Bağlı olmayan gözlem	Bağlı gözlem
Ortalama ($\mu_{\bar{Y}_1 - \bar{Y}_2}$)	$\mu_1 - \mu_2$	$\mu_1 - \mu_2$
Standart sapma ($\sigma_{\bar{Y}_1 - \bar{Y}_2}$)	$\sqrt{\frac{\sigma_1^2 + \sigma_2^2}{n}}$	$\sqrt{\frac{\sigma_1^2 + \sigma_2^2 - 2\sigma_1\sigma_2\rho_{12}}{n}}$

1. ρ_{12} bağlı gözlemlerde, bant ve dikiş sonrası yapılan ölçümlerin arasındaki korelasyondur.
2. Standart sapmadaki değişiklik ρ_{12} 'den kaynaklanır. Eğer bu korelasyon 0 ise iki tasarımın da örneklem dağılımına ait standart sapma aynıdır.

Bir araştırma tasarımında hedeflerden biri standart hatayı mümkün olduğunca küçük tutmaktır. Standart hatanın küçük olması elde edilen test istatistiğinin tahmin edilen parametreye yakın olduğu anlamına gelir.

Veri çözümleme sürecinde hata varyansı (error variance) hesaplamak için bir formül seçilir. Yanlış formülün kullanılması büyük bir hatadır.

Uygulamada, standart hatanın hesaplanması tasarımın bağlı mı bağımsız mı olduğuna göre değişir. Tasarımın yanlış sınıflandırılması, çözümleme sürecinde büyük bir hatadır.

8.1 Bağımsız gruplar t-test (The Independent Groups t-test)

Uşak ilinde yaşayan katılımcıların Toplumsal Cinsiyet Algısı (TCA) puanları yüksek öğretim durumuna göre karşılaştırılmıtır. Her grup için yoğunluk grafikleri;

```
# csv yükle
urlfile='https://raw.githubusercontent.com/burakaydin/materyaller/gh-pages/ARPASS/dataWBT.csv'
dataWBT=read.csv(urlfile)

#URL sil
rm(urlfile)
dataWBT_USAK=dataWBT[dataWBT$city=="USAK",]

# factor ve droplevels fonksiyonları bölüm 5.2.4 ile verilmiştir.
# yeni oluşturulan HEF (Higher Education Factor)
# katılımcı lise veya altı diplomaya sahipse 0, non-college
# katılımcı lise üstü diplomaya sahip ise 1, college
dataWBT_USAK$HEF=droplevels(factor(dataWBT_USAK$higher_ed,
                                     levels = c(0,1),
                                     labels = c("non-college", "college")))

require(ggplot2)
plotdata=na.omit(dataWBT_USAK[,c("gen_att", "HEF")])
ggplot(plotdata, aes(x = gen_att)) +
  geom_histogram(aes(y = ..density..),col="black",binwidth = 0.2,alpha=0.7) +
  geom_density(size=2) +
  theme_bw()+labs(x = "Gender Attitude by HEF in USAK")+ facet_wrap(~ HEF)+
  theme(axis.text=element_text(size=15),
        axis.title=element_text(size=14,face="bold"))
```

8.1.1 Bağımsız gruplar t testi için R betiği

Takip edilen basamaklar;




Figure 8.1: Yüksek Öğretim Durumuna göre TCA puanları

1. Betimsel istatistikler
2. Test istatistiğinin hesabı

$$t = \frac{\bar{Y}_1 - \bar{Y}_2}{S_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}}$$

$$S_p = \sqrt{\frac{(n_1 - 1)S_1^2 + (n_2 - 1)S_2^2}{n_1 + n_2 - 2}}$$

3. Kritik değerin hesabı $\pm t_{\alpha/2, n_1+n_2-2}$

$$H_0 : \mu_1 - \mu_2 = 0$$

$$H_1 : \mu_1 - \mu_2 \neq 0$$

```
library(psych)
descIDT=with(dataWBT_USAK,describeBy(gen_att, HEF,mat=T,digits = 2))
descIDT
##      item      group1 vars  n mean  sd median trimmed  mad min max range
## X11      1 non-college   1 86 1.83 0.54   1.8   1.80 0.59   1 3.8   2.8
## X12      2 college     1 51 1.64 0.61   1.6   1.54 0.59   1 3.4   2.4
##      skew kurtosis  se
## X11 0.72      0.90 0.06
## X12 1.19      1.09 0.09
# rapor etmek için
#write.csv(descIDT,file="independent_t_test_desc.csv")

# ss
sp=sqrt((85*.543^2 + 50*.608^2)/(86+51-2))

# t-istatistik
tstatistic=(1.832-1.635)/(sp*sqrt(1/86+1/51))

# alfa=0.05 kritik değer
qt(.975,df=135)
## [1] 1.98
```

1.963 kritik değer $t_{.975,135} = 1.978$ 'den küçük olduğu için, H_0 kabul edilir.

$H_1 : \mu_1 - \mu_2 > 0$ alternatif hipotezi kurulsa idi, kritik değer $t_{.95,135} = 1.66$, 1.963'ten küçük olduğu için H_0 terkedilirdi.

$H_1 : \mu_1 - \mu_2 < 0$ alternatif hipotezi kurulsa idi, 1.963 kritik değer olan $t_{.05,135} = -1.66$ ' ten küçük olmadığı için H_0 kabul edilirdi (Burada alternatif hipotez ile örneklem arası farklılık zıt yönde).

Daha kullanışlı bir R betiği;

```
# dataWBT HEF faktörünü içermez, yukarıda HEF faktörü oluşturulmuştur.

t.test(gen_att~HEF,data=dataWBT_USAK,var.equal=T,
       alternative="two.sided",
       conf.level=0.95)

##
## Two Sample t-test
##
## data:  gen_att by HEF
```



```
## t = 2, df = 100, p-value = 0.05
## alternative hypothesis: true difference in means is not equal to 0
## 95 percent confidence interval:
##  -0.0019  0.3949
## sample estimates:
## mean in group non-college      mean in group college
##                               1.83                1.64

# büyüktür
t.test(gen_att~HEF,data=dataWBT_USAK,var.equal=T,
       alternative="greater",
       conf.level=0.95)

##
## Two Sample t-test
##
## data:  gen_att by HEF
## t = 2, df = 100, p-value = 0.03
## alternative hypothesis: true difference in means is greater than 0
## 95 percent confidence interval:
##  0.0303      Inf
## sample estimates:
## mean in group non-college      mean in group college
##                               1.83                1.64

# küçüktür
t.test(gen_att~HEF,data=dataWBT_USAK,var.equal=T,
       alternative="less",
       conf.level=0.95)

##
## Two Sample t-test
##
## data:  gen_att by HEF
## t = 2, df = 100, p-value = 1
## alternative hypothesis: true difference in means is less than 0
## 95 percent confidence interval:
##  -Inf 0.363
## sample estimates:
## mean in group non-college      mean in group college
##                               1.83                1.64
```

8.1.1.1 Yönsüz alternatif için rapor örneği

Uşak ilinde yaşayan katılımcılardan yüksek öğretim diplomasına sahip olan 51 kişi için TCA puanları ortalaması 1.64, standart sapması 0.61, puanlara ait dağılım çarpıklık değeri 1.19 ve basıklık değeri 1.09 olarak bulunmuştur. Aynı ilde yüksek öğretim diploması olmayan 86 katılımcının ise TCA puanları ortalaması 1.83, standart sapması 0.54, puanlara ait dağılımın çarpıklığı 0.72 ve basıklığı 0.90 olarak hesaplanmıştır. Bağımsız gruplar t-testi sonuçları, Uşak ilinde yaşayan katılımcıların Toplumsal Cinsiyet Algısı puanlarının yüksek öğretim durumuna göre değişeceği tezini desteklememiştir, $t(135)=1.96$, $p=0.052$. Puanlar arasındaki fark için %95 güven aralığı $[-0.002, 0.395]$ olarak hesaplanmıştır.¹

¹Çözümler R programlama dili kullanılarak tamamlanmıştır, betimsel istatistikler *psych* paketi (Revelle, 2016), t test istatistiği ise *stats* paketi (R Core Team, 2016b) ile hesaplanmıştır.

8.1.1.2 Yönlü alternatif için rapor örneği

Uşak ilinde yaşayan katılımcılardan yüksek öğretim diplomasına sahip olan 51 kişi için TCA puanları ortalaması 1.64, standart sapması 0.61, puanlara ait dağılım çarpıklık değeri 1.19 ve basıklık değeri 1.09 olarak bulunmuştur. Aynı ilde yüksek öğretim diploması olmayan 86 katılımcının ise TCA puanları ortalaması 1.83, standart sapması 0.54, puanlara ait dağılımın çarpıklığı 0.72 ve basıklığı 0.90 olarak hesaplanmıştır. Bağımsız gruplar t testi sonuçları, yüksek öğretim diploması olmayanların TCA puanları yüksek öğretilere nazaran daha yüksektir tezini desteklemiştir, $t(135)=1.96$, $p=0.026$. Puanlar arasındaki fark için %95 güven aralığı $[0.030, \infty]$ olarak hesaplanmıştır.

8.1.2 Bağımsız gruplar t testi varsayımları

Geleneksel t-testi sonuçlarının geçerliği 3 varsayımın ihlal edilmemesi ile mümkündür.

1. Yanıtların bağımsızlığı (independence). Her gruba ait puanların dağılımı birbirinden bağımsız olmalıdır. Yanıtların bağımsızlığını tehdit eden durumlardan biri aynı grup içerisinde yer alan bireylerin birbirlerinin yanıtlarını etkilemesidir (Yantların bağımsızlığı 9.2.1.4 bölümünde daha detaylı ele alınmıştır.)
2. Normallik. Her gruba ait puanlar normal bir dağılımdan çekilmiştir. Myers et al. (2013) grupların örneklem sayısı (n) eşit olduğunda ve toplam örneklem 40 veya daha fazla olduğu durumlarda t test istatistiğinin normallik varsayımı ihlallerine dirençli olduğunu belirtmiştir. Fakat bu direnç normal dağılımdan büyük çaplı sapmalar (extreme) için geçerli değildir. Bu kitabın yazarları normallik testlerinin bu varsayımı kontrol etmek için kullanılmasına sıcak bakmamaktadır. Görsel bir değerlendirmeden sonra özellikle küçük örneklerde, normallik varsayımının ihlal edildiği kaygısı varsa araştırmacılar dirençli tahminleme yöntemlerini kullanmalıdırlar.
3. Eş varyanslılık. Varyans homojenliği olarak da bilinen bu varsayım, iki grupta yer alan puanların evren bazında eşit varyanslı dağılımlardan çekildiğini kabul eder. Myers et al. (2013) grup örneklem sayılarının **eşit** ve en az 5 olduğu durumlarda varyans eşitliği sağlanamasa dahi 1. tip hata oranlarının önemli ölçüde değişmeyeceğini belirtmiştir. Fakat bu direnç büyük çaplı heterojenlikler için geçerli değildir, örneğin $s_1^2/s_2^2 > 100$. Field et al. (2012) Levene testi gibi eşvaryanslılık testlerinin örneklem sayılarının eşit olmadığı ve küçük örneklerde sağlıklı sonuçlar vermediğini belirtmiştir, halbuki bu testlere en çok ihtiyaç durumlar küçük örneklem ve eşit olmayan örneklem sayısı durumlarıdır. Eş varyanslılık testleri ile ilgili bir diğer kaygı, bu test sonucunda varyansların eşit sayılabileceği kararı verilmiş olsa da varyanslar arası farklılığın t test istatistiğini etkileyebileceğidir. *t.test* fonksiyonu, aksi istenmediği sürece, varyans eşitliği varsayımı yapmaz ve Welch t testini hesaplar.

Varsayımlar hakkındaki bu kısa tanıtımın ele almadığı durumlar vardır. Örneğin hem normallüğün hem eş varyanslılığın ihlal edildiği durumlar tartışılmamıştır. Ayrıca direnç konusu tartışmaya açıktır. Örneğin $n_1=n_2=10$ örneklem sayısı ve eş olmayan varyans durumu için 100000 tekrarlı yaptığımız bir simülasyon, $\alpha=.01$ ve yönsüz bir t testi için 1. tip hata oranını .018 bulmuştur. Bu oranın kabul edilebilir olup olmadığı tartışmaya açıktır.

Sonuç olarak, eğer yanıtların bağımsızlığı kabul ediliyorsa, örneklem sayısı eşitse ve her grupta en az 20 ise bağımsız gruplar t istatistiğinin büyük ölçüde dirençli olduğu kabulü makul bir kabuldur. Diğer durumlarda varsayım ihlalleri tespit edildi ise araştırmacılar alternatif çözümleme yöntemlerini kullanabilirler.

8.1.3 Welch t test

Normallüğün ciddi ölçüde zedelenmediği ve örneklem sayılarının her grup için en az 20 olduğu durumlarda Welch testi geçerli sonuçlar üretir. Bu test varyans eşdeğerli varsayımı yapmaz.

```
t.test(gen_att~HEF,data=dataWBT_USAK,var.equal=F,
      alternative="two.sided",
```

```

                                conf.level=0.95)
##
## Welch Two Sample t-test
##
## data:  gen_att by HEF
## t = 2, df = 100, p-value = 0.06
## alternative hypothesis: true difference in means is not equal to 0
## 95 percent confidence interval:
##  -0.00848  0.40146
## sample estimates:
## mean in group non-college      mean in group college
##                1.83                1.64

```

8.1.3.1 Welch t testi için raporlama örneği

Uşak ilinde yaşayan katılımcılardan yüksek öğretim diplomasına sahip olan 51 kişi için TCA puanları ortalaması 1.64, standart sapması 0.61, puanlara ait dağılım çarpıklık değeri 1.19 ve basıklık değeri 1.09 olarak bulunmuştur. Aynı ilde yüksek öğretim diploması olmayan 86 katılımcının ise TCA puanları ortalaması 1.83, standart sapması 0.54, puanlara ait dağılımın çarpıklığı 0.72 ve basıklığı 0.90 olarak hesaplanmıştır. Bağımsız gruplar Welch t-testi sonuçları, Uşak ilinde yaşayan katılımcıların Toplumsal Cinsiyet Algısı puanlarının yüksek öğretim durumuna göre değişeceği tezini desteklememiştir, $t(95.89)=1.9$, $p=0.06$. Puanlar arasındaki fark için %95 güven aralığı $[-0.082, 0.402]$ olarak hesaplanmıştır.

Eğer normallik varsayımı ciddi ölçüde şüpheli ise ve özellikle iki grup farklı şekillerde dağılım gösteriyor ise yüzdeli bootstrap prosedürü (percentile bootstrap) kullanılabilir (Wilcox (2012), page 171).

```

#bootstrap ile %95 güven aralığı (normallik varsayımı yok)
set.seed(04012017)
B=5000          # bootstrap tekrar sayısı
alpha=0.05      # alfa

# grupları tanımla
GroupCollege=na.omit(dataWBT_USAK[dataWBT_USAK$HEF=="college", "gen_att"])
GroupNONcollege=na.omit(dataWBT_USAK[dataWBT_USAK$HEF=="non-college", "gen_att"])

output=c()
for (i in 1:B){

  x1=mean(sample(GroupCollege,replace=T,size=length(GroupCollege)))
  x2=mean(sample(GroupNONcollege,replace=T,size=length(GroupNONcollege)))
  output[i]=x2-x1
}
output=sort(output)

## yönsüz
# D yıldız alt
output[as.integer(B*alpha/2)+1]
## [1] -0.0134

# D yıldız üst
output[B-as.integer(B*alpha/2)]
## [1] 0.39

##Yönlü x2>x1

```

```
# D yıldız alt
output[as.integer(B*alpha)+1]
## [1] 0.022

#hatalı yön x2<x1
# D yıldız üst
output[as.integer(B*(1-alpha))]
```

8.1.3.2 Yüzdeli bootstrap yöntemi için raporlama örneği

Uşak ilinde yaşayan katılımcılardan yüksek öğretim diplomasına sahip olan 51 kişi için TCA puanları ortalaması 1.64, standart sapması 0.61, puanlara ait dağılım çarpıklık değeri 1.19 ve basıklık değeri 1.09 olarak bulunmuştur. Aynı ilde yüksek öğretim diploması olmayan 86 katılımcının ise TCA puanları ortalaması 1.83, standart sapması 0.54, puanlara ait dağılımın çarpıklığı 0.72 ve basıklığı 0.90 olarak hesaplanmıştır. Puanlar arasındaki fark için %95 bootstrap güven aralığı $[-0.013, 0.390]$ olarak hesaplanmıştır. Güven aralığı 0 değerini içerdiği için puanlar arasındaki farkın istatistiksel olarak anlamlı olduğu savunulamaz.

Yönlü test Alternatif hipotez, yüksek öğretim mezunu olmayan katılımcıların puanlarının yüksek olacağını belirtmiş ise; Puanlar arasındaki fark için %95 bootstrap güven aralığı $[0.022, \infty]$ olarak hesaplanmıştır. Eldeki veri yüksek öğretim mezunu olmayanların puanlarının daha yüksek olacağı tezini destekleyecek kanıt sunmuştur, $H_0 : \mu_{non-college} = \mu_{college}$ in favor of $H_1 : \mu_{non-college} - \mu_{college} > 0$.

Yönlü test Alternatif hipotez, yüksek öğretim mezunu olmayan katılımcıların puanlarının düşük olacağını belirtmiş ise; Puanlar arasındaki fark için %95 bootstrap güven aralığı $[-\infty, 0.358]$ olarak hesaplanmıştır. Eldeki veri yüksek öğretim mezunu olmayanların puanlarının daha düşük olacağı tezini desteklememektedir. $H_0 : \mu_{non-college} = \mu_{college}$ ve $H_1 : \mu_{non-college} - \mu_{college} < 0$.

Yönlü ve yönsüz alternatif testler farklı kriterler kullanır. Yönlü testlerde yönün nasıl belirlendiği savunulmalıdır. Kullanılan örnekte yönsüz alternatif kullanıldığında Welch t testi p değeri 0.06 bulunmuştur. Bu sonuç marjinal anlamlılık olarak yorumlanabilir. Ülkemizde TCA puanları ve yüksek öğretim ilişkisi hakkında alanyazın sınırlı olduğu için yönlü bir alternatif savunulması zordur.

8.1.4 Bağımsız gruplar t testi için etki büyüklüğü

Bir t testi istatistiği ortalamaların birbirinden farklı olup olmadığı hakkında karar vermeye yardımcı olsa da, bu farklılığın büyüklüğünü yorumlamak için etki büyüklüğü hesaplamaları geliştirilmiştir. Bağımsız gruplar t testi için Cohen etki büyüklüğü, ortalamaların farkını bileşik standart sapmaya (pooled) bölünmesi ile hesaplanır.

$$EB = \frac{t}{\sqrt{\frac{n_1 n_2}{n_1 + n_2}}}$$

Cohen (1962) etki büyüklüğü sınıflaması

Etki büyüklüğü	Tanım
.2	Küçük
.5	Orta
.8	Büyük

```
## normallik ve eş varyanslılık varsayımı yapıldığında
## (dirençli yöntem benzer sonuç verdiği için varsayımların kabulü makuldür.)
```

```

n1=51
n2=86
tval=1.96

EB=tval/sqrt((n1*n2)/(n1+n2))
EB
## [1] 0.346

#veya effsize paketi ile
t.test(gen_att~HEF,data=dataWBT_USAK,var.equal=F,
       alternative="two.sided",
       conf.level=0.95)

##
## Welch Two Sample t-test
##
## data:  gen_att by HEF
## t = 2, df = 100, p-value = 0.06
## alternative hypothesis: true difference in means is not equal to 0
## 95 percent confidence interval:
##  -0.00848  0.40146
## sample estimates:
## mean in group non-college      mean in group college
##                1.83                1.64
library(effsize)
cohen.d(gen_att~HEF,data=dataWBT_USAK, paired=F, conf.level=0.95,noncentral=F)
##
## Cohen's d
##
## d estimate: 0.346 (small)
## 95 percent confidence interval:
##      inf      sup
## -0.00843  0.70078
# noncentral=T argümanınızı araştırabilirsiniz.

```

effsize paketi (Torchiano, 2016) etki büyüklüğünü 0.35 ve ilgili %95 güven aralığını [-0.008, 0.701] olarak hesaplamıştır.

8.1.5 Extra: Pratikte anlamlılık ve istatistiksel anlamlılık

Bir çalışma sonucunda elde edilen sonuçların istatistiksel olarak anlamlı olmadığı fakat pratikte anlamlı olduğu öne sürülebilir. Bu sakıncalı bir durumdur ve sadece küçük örneklerde görülür. Küçük bir örneklem ile yapılan çalışmadan sonra pratikte anlamlılıktan bahsetmek tezat oluşturur.

Bir çalışma sonucunda elde edilen sonuçların istatistiksel olarak anlamlı olduğu fakat pratikte anlamlı olmadığı öne sürülebilir. Bu doğru olabilir. 400 kişi ile tamamlanan bir çalışmada istatistiksel anlamlı farklılık .05 etki büyüklüğüne sahip olabilir. Eğer .05 etki büyüklüğü çalışmanın yapıldığı alanda küçük adlediliyorsa istatistiksel anlamlılık pratikte anlamlılığı desteklemez.

8.1.6 Kayıp veriler ile bağımsız gruplar t testi

Eklenecek

8.1.7 Destekleyici grafikler

Eklenecek

8.1.8 İstatistksel güç

İstatistksel güç konusunun ana hatlarına bölüm 7.3.8’de değinilmiştir.

```
#power.t.test
power.t.test(delta=.35, sd=.6, sig.level=0.05, power=0.95,
              type="two.sample", alternative="two.sided")

##
##      Two-sample t test power calculation
##
##              n = 77.4
##              delta = 0.35
##              sd = 0.6
##              sig.level = 0.05
##              power = 0.95
##              alternative = two.sided
##
## NOTE: n is number in *each* group
```

Bu örnekte belirlenmiş değerler ortalamalar farkı 0.35, standart sapma 0.6, alfa 0.05, yönsüz test ve hedeflenen güç 0.95 seçildiğinde gereken örneklem sayısı her grup için 78’dir. Diğer bir ifade ile, 156 katılımcı ile belirlenen değerlere (ortalamalar farkı 0.35, standart sapma 0.6, alfa 0.05, yönsüz test) ulaşılması durumunda $H_0 : \mu_1 - \mu_2 = 0$ boş hipotezinin terkedilme olasılığı %95’tir.

8.2 Bağlı gruplar t-testi (Within-subjects t-test)

20 tavşan ile gerçekleştirilen deneyde, yara bandı ve dikiş yöntemlerinin yara kapandıktan 10 gün sonra ölçülen germe mukavemeti değerleri üzerinde etkisi araştırılmıştır.

```
gerMUK=data.frame(tavid=1:20,
                  bant=c(6.59,9.84 ,3.97,5.74,4.47,4.79,6.76,7.61,6.47,5.77,
                        7.36,10.45,4.98,5.85,5.65,5.88,7.77,8.84,7.68,6.89),
                  dikis=c(4.52,5.87,4.60,7.87,3.51,2.77,2.34,5.16,5.77,5.13,
                        5.55,6.99,5.78,7.41,4.51,3.96,3.56,6.22,6.72,5.17))

# Grafik verisi
library(tidyr)
plotdata=gather(gerMUK, metot, mukavemet, bant:dikis, factor_key=TRUE)

require(ggplot2)
ggplot(plotdata, aes(x = mukavemet)) +
  geom_histogram(aes(y = ..density..), col="black", alpha=0.7) +
  geom_density(size=2) +
  theme_bw()+labs(x = "mukavemet")+ facet_wrap(~ metot)+
  theme(axis.text=element_text(size=15),
        axis.title=element_text(size=14, face="bold"))
```




Figure 8.2: Bağlı gruplar örneği

8.2.1 Bağlı gruplar için t testi

Takip edilen basamaklar;

1. Betimsel istatistikler
2. Test istatistiğinin hesabı

$$t = \frac{\bar{Y}_1 - \bar{Y}_2}{\sqrt{\frac{S_1^2 + S_2^2 - 2S_1 S_2 r_{12}}{n}}}$$

3. Kritik değerin hesabı $\pm t_{\alpha/2, n-1}$,

$$H_0 : \mu_1 - \mu_2 = 0$$

$$H_1 : \mu_1 - \mu_2 \neq 0$$

```
library(psych)
descDT=with(gerMUK,describe(cbind(bant,dikis)))
descDT
##          vars  n mean   sd median trimmed  mad  min   max range  skew
## bant         1 20 6.67 1.71   6.53   6.54 1.45 3.97 10.45  6.48  0.55
## dikis        2 20 5.17 1.49   5.17   5.19 1.30 2.34  7.87  5.53 -0.08
##          kurtosis   se
## bant        -0.45 0.38
## dikis        -0.87 0.33

corDT=with(gerMUK,cor(bant,dikis,use="complete.obs"))
corDT
## [1] 0.354

# tahmin edilen standart hata (tsh)
tsh=sqrt(((1.71^2+1.49^2)-(2*1.71*1.49*corDT))/(20))

# t-istatistik
tstatistic=(6.67-5.17)/tsh

# alfa=0.05 kritik değer
qt(.975,df=19)
## [1] 2.09
```

Hesaplanan 3.67, kritik değerdan ($t_{.975,19} = 2.09$) büyük olduğu için boş hipotez terkedilebilir.

Daha kolay bir R satırı;

```
library(psych)
with(gerMUK, t.test(bant,dikis,paired=T,
                    alternative="two.sided",
                    conf.level=0.95))

##
## Paired t-test
##
## data:  bant and dikis
## t = 4, df = 20, p-value = 0.002
## alternative hypothesis: true difference in means is not equal to 0
```



```
## 95 percent confidence interval:
##  0.643 2.352
## sample estimates:
## mean of the differences
##                1.5
```

8.2.1.1 Raporlama örneği:

Bağlı gruplar t testi hesaplarına dayanarak, gerilim mukavemetinin bant tedavisi (ort=6.67,SS=1.71,çarpıklık=0.55,basıklık=-0.45) ve dikiş tedavisi (ort=5.17,SS=1.49,çarpıklık=-0.08,basıklık=-0.87) arasında farklılık gösterdiği sonucuna varılmıştır, $t(19)=3.67$, $p=0.002$, $r=0.35$. Aritmetik ortalamaların farkı için 95% güven aralığı [0.64,2.35] olarak hesaplanmıştır.

8.2.2 Varsayımlar: bağlı gruplar için t testi

Puanların farkı ($Y_{1i} - Y_{2i}$) normal bir dağılımdan çekilmiş olmalıdır. Puanların farkları her birey için bağımsız olmalıdır. Bağlı gruplar t testi , örneklem küçük değil ise, normallik varsayımı ihlallerine karşı genellikle dirençlidir.

8.2.3 Dirençli yöntem: bağlı gruplar için t testi

Eğer dağılım normallikten büyük ölçüde ayrılıyor ise, yüzdeli bootstrap prosedürü kullanılabilir (Wilcox (2012),page 201).

```
#bootstrap ile %95 güven aralığı (normallik varsayımı yok)
set.seed(04012017)
B=5000          # bootstrap tekrar sayısı
alpha=0.05      # alfa

gerMUK=data.frame(ratid=1:20,
                  bant=c(6.59,9.84 ,3.97,5.74,4.47,4.79,6.76,7.61,6.47,5.77,
                        7.36,10.45,4.98,5.85,5.65,5.88,7.77,8.84,7.68,6.89),
                  dikis=c(4.52,5.87,4.60,7.87,3.51,2.77,2.34,5.16,5.77,5.13,
                        5.55,6.99,5.78,7.41,4.51,3.96,3.56,6.22,6.72,5.17))

output=c()
for (i in 1:B){
  #satırları örnekle
  bs_rows=sample(gerMUK$ratid,replace=T,size=nrow(gerMUK))
  bs_sample=gerMUK[bs_rows,]
  mean1=mean(bs_sample$bant)
  mean2=mean(bs_sample$dikis)
  output[i]=mean1-mean2
}
output=sort(output)

## yönsüz
# d yıldız alt
output[as.integer(B*alpha/2)+1]
## [1] 0.686
```

```
# d yıldız üst
output[B-as.integer(B*alpha/2)]
## [1] 2.24

##Yönlü x2>x1
# d yıldız alt
output[as.integer(B*alpha)+1]
## [1] 0.837

#yanlış yön x2<x1
# d yıldız üst
output[as.integer(B*(1-alpha))]
## [1] 2.14
```

8.2.3.1 Yönsüz yüzdeli bootstrap için örnek rapor:

Yaraların tedavisinden 10 gün sonra gerilim mukavemeti ölçümleri yapılmıştır. Yara bandı ile (ort=6.67,SS=1.71,çarpıklık=0.55,basıklık=-0.45) dikiş tedavisi (ort=5.17,SS=1.49,çarpıklık=-0.08,basıklık=-0.87) ölçümleri arasındaki fark için 5000 tekrarlı bootstrap prosedürü %95 güven aralığı [0.667,2.2555] olarak hesaplanmıştır. Yeni geliştirilen yara bandı sonrası gerilim mukavemetinin daha yüksek olduğu ve bu farklılığın istatistiksel olarak anlamlı olduğu sonucuna varılmıştır.

8.2.4 Effect size for the dependent groups t-test

En basit etki büyüklüğü hesaplama yöntemlerinden biri; ²

$$ES = \frac{t}{\sqrt{n}}$$

```
## dirençli prosedürler farklı sonuç vermediği için
## normallik ve varyans eşitliği varsayımları yapılmıştır.
n=20
tval=3.6678

EB=tval/sqrt(n)
EB
## [1] 0.82

library(effsize)
cohen.d(gerMUK$bant,gerMUK$dikis,
        paired=T, conf.level=0.95,noncentral=F)
##
## Cohen's d
##
## d estimate: 0.82 (large)
## 95 percent confidence interval:
##   inf   sup
## 0.135 1.505
```

²Lakens (2013) Eşitlik 7, fakat bu değer korelasyon 1'e yaklaştıkça sonsuza yaklaşır. Lakens (2013) Eşitlik 10 daha uygun bir tercih olabilir. $\frac{meandifference}{(SD_1+SD_2)/2}$

effsize paketi (Torchiano, 2016) etki büyüklüğünü 0.820 ve ilgili %95 güven aralığını [0.135, 1.505] olarak hesaplamıştır.

8.2.5 Kayıp veriler ile bağlı gruplar t testi

Eklenecek

8.2.6 Destekleyici grafikler

Eklenecek

8.2.7 İstatistiksel güç: bağlı gruplar t testi

```
#power.t.test
power.t.test(delta=.35, sd=.6, sig.level=0.05, power=0.95,
              type="paired", alternative="two.sided")
##
##      Paired t test power calculation
##
##              n = 40.2
##            delta = 0.35
##              sd = 0.6
##      sig.level = 0.05
##            power = 0.95
##      alternative = two.sided
##
## NOTE: n is number of *pairs*, sd is std.dev. of *differences* within pairs
```

Bu örnekte belirlenmiş değerler ortalamalar farkı 0.35, standart sapma 0.6, alfa 0.05, yönsüz test ve hedeflenen güç 0.95 seçildiğinde gereken örneklem sayısı 41'dir. Diğer bir ifade ile, 41 katılımcı ile belirlenen değerlere (ortalamalar farkı 0.35, standart sapma 0.6, alfa 0.05, yönsüz test) ulaşılması durumunda $H_0 : \mu_1 - \mu_2 = 0$ boş hipotezinin terkedilme olasılığı %95'tir.

8.3 Yaygın Tasarılar

Sosyal bilimlerde iki ortalamayı kıyaslamak üzere kurulan tasarılardan yaygın olanları özetlenmiştir.

8.3.1 Grupların ilişkili olduğu durumlar

Ortalamaların hesaplandığı puanların farklı gruplar altında ilişkili olup olmadığı önemlidir.

8.3.1.1 Tekrarlanan ölçümler

Aynı katılımcı için aynı değişkenin birden fazla ölçülmesi durumu.

1. Bireylerin kendi kontrol grubunu oluşturması:

Semantik hafızanın aktivasyon hızını ölçmek üzere 100 üniversite öğrencisi ile araştırma yapılmıştır. Her öğrenci çiftler halinde verilen kelimeleri okumuştur. Okudukları ilk kelime ya bir silah (örneğin mermi, hançer) ya da silah olmayan bir kelimedir. Okudukları ikinci kelime mutlaka agresif bir kelimedir (örneğin yarala, imha et). Her bir öğrenci 192 çift kelimeyi bilgisayarda görüp sesli olarak okumuştur. Bilgisayar ilk kelimeyi 1.25 saniye ekranda tutmuş, .5 saniye siyah ekran göstermiş ve ikinci kelimeyi ekranda göstermiş, her ikinci kelimeden önce reaksiyon zamanını ölçmüştür. Her bireyin reaksiyon zamanı ortalamaları alınmış ve analiz edilmiştir.

İlk kelime		
Öğrenci	Silah	Silah Değil
1		
2		
...		
100		

Her öğrencinin kendi okuma hızı olduğundan, Silah ve Silah olmayan kelimeleri okumak için gereken reaksiyon zamanları birbiri ile ilişkilidir.

2. **Boylamsal tasarılar:** 6. sınıf öğrencilerinin matematik başarıları dönem başında ve dönem sonunda ölçülmüştür. Amaç puanların değişip değişmediğini görmektir.

Zaman		
Öğrenci	Dönem başı	Dönem sonu
1		
2		
...		
48		

Ölçümler aynı öğrenciler ile tekrarlandığı için puanlar ilişkilidir.

8.3.1.2 Blok tasarıları

Katılımcıların blok olarak ikili eşleştirilmesidir. Her çiftin (blok içindeki) benzer davranması beklenir.

1. **Rassal blok tasarısı:** Okuma hızını artırmak için bir program geliştirilmiştir, etkililiğini araştırmak üzere, 30 ikinci sınıf öğrencisi bir okuma testini cevaplamış, ve puanlarına göre çiftler oluşturmuştur.

Çift	Okuma testi puan sırası
1	1,2
2	3,4
...	...
15	29,30

Görüldüğü gibi en hızlı okuyan 2 öğrenci ilk çifti, en yavaş okuyan iki öğrenci son çifti oluşturmuştur. Çiftlerden her biri rassal olarak yeni program grubuna veya kontrol grubuna atanmıştır. As shown, the students with the two highest scores were in the first pair, the students with the

Yeni program son bulduktan sonra öğrencilerin okuma hızları ölçülmüştür

[:——:]::————-:[:——-: | Çift | Yeni Program | Kontrol | | 1 | | | | 2 | | | | ... | | | | ... | | | | 15 | | |

2. **Rassal olmayan blok tasarı:** Şiddete maruz kalmış grup çocuğun, şiddet görmemiş daha kalabalık bir gruptan çocuklar ile genel kaygı düzeyi üzerinden eşleştğini düşünelim. Sonrasında stres anında gösterdikleri endişe durumlarını ölçelim

Çift	Şiddete maruz kalan	Kontrol
1		
2		
...		
20		

3. **Kalıtsal tasarılar (Famialial):** 25 anne ve erişkin kızlarının politik görüşleri alınmıştır.

Çift
Anne
Kız
1
2
...
25

4. **Dyadik tasarıları:** Afrikalı-Amerikalı ve Avrupalı-Amerikalı çocuklardan oluşturulan çiftler işbirliği gerektiren küçük oyunlar oynamışlardır. Her bir çocuk eşinin işbirlikçiliğini puanlamıştır.

Etnisite
Çift
Afrikalı-Amerikalı
Avrupalı-Amerikalı
1
2
...
25

8.3.2 Bağlı olmayan grup tasarılarına örnekler

1. **Tamamen rassal tasarı**

İki farklı manyetik ağı kesici makinesinin performansı karşılaştırılacaktır. 50 hasta rassal olarak, 25-25, iki makineye alınmış, tedaviden sonra ağı düzeylerini rapor etmişlerdir.

Makine
1
2
.
.
.

2. **Rassal olmayan tasarı:** Sekizinci sınıf öğrencilerinden 50 kız ve 50 erkek seçilmiş 2 basamaklı toplama işlemi yapma hızları ölçülmüştür.

Chapter 9

Varyans Analizi (ANOVA)

Varyans analizi (ANOVA) bölümünde kısaca tanıtılan konular; (a) terminoloji, (b) gruplar-arası ANOVA, ve (b) gruplar-ıçi ANOVA olarak üç başlıktır. Kitabın bir sonraki versiyonunda karma ANOVA (mixed ANOVA) eklenecektir.

9.1 Terminoloji

Varyans analizi kapsamında kullanılan terimler kısaca açıklanmıştır. Fakat bu bölüm bir önceki bölümde tanıtılan *yaygın tasarılar* (8.3) paragraflarının devamıdır.

Faktör Kategorik bağımsız değişkenler ANOVA çerçevesinde faktör olarak isimlendirilir. Örneğin bir önceki bölümde (8.3.2) rassal blok tasarısı içerisinde tanımlanan *okuma hızını artırmaya yönelik yeni program* ve *kontrol* grubu üyeliğini belirten iki kategorili (iki alt sınıflı) değişken bir faktör oluşturur. Toplumsal Cinsiyet Algısının (TCA) yüksek öğretim durumuna göre (yüksek öğretim mezunları ve yüksek öğretim mezunu olmayanlar) değişip değişmediğini test eden bir modelde TCA bağımlı değişken, yüksek öğretim durumu ise bir faktördür.

Alt sınıf Faktörü oluşturan kategorilere alt sınıf denir. TCA örneğinde faktör yüksek öğretim durumudur. Bu faktöre ait alt sınıflar *yüksek öğretim mezunu* ve *yüksek öğretim mezunu değil* olmak üzere iki tanedir.

Kesişen faktörler (Crossed factors) Bir faktöre ait alt sınıfların diğer bir faktörün bütün alt sınıfları ile kesişmesi durumudur. Tekarlanan ölçümler için bir önceki bölümde verilen örneği düşünelim,

İlk kelime		
Öğrenci	Silah	Silah Değil
1		
2		
...		
100		

Öğrenciler de bir faktör olarak düşünüldüğünde, her öğrencinin reaksiyon süreleri ilk kelime faktörünün her iki alt sınıfında da ölçüldüğü için öğrenci ve ilk kelime faktörü kesişmiştir.

Kesişmeyen faktörler (Nesting) Bir faktöre ait bir alt sınıfın, ikinci faktörün sadece bir alt sınıfında gözlemlenmesi durumudur. Tamamen rassal tasarı için bir önceki bölümde verilen örneği düşünelim,

Makine	
1	2
H_1	H_{n+1}
H_2	H_{n+2}
H_3	H_{n+3}
...	
H_n	H_{2n}

Her bir hasta 1. veya 2. makine ile tedavi edildiği için, katılımcı faktörü makine faktörünün içinde düşünülür ve faktörler kesişmemiştir.

Bağlı gözlemler faktörü (Within-subjects factor) katılımcı faktörü (öğrenci, çalışan, hasta vb.) ile kesişen faktörlerdir. Tekrarlanan ölçümler örneğinde olduğu gibi her bir katılımcı diğer bir faktörün bütün alt sınıflarında gözlemlenir. Boylamsal çalışmalarda zaman (örneğin öntest, sontest) bağlı gözlemler faktörüne örnektir.

Bağlı bloklar faktörü bloklar ile kesişen faktördür. Rassal blok tasarımı için bir önceki bölümde verilen örneği düşünelim,

|:—:|:—:|:—:| | Çift | Yeni Program | Kontrol | | 1 | | | 2 | | | ... | | | ... | | | 15 | | |

Rassal olmayan blok, kalıtsal ve diyardik tasarımlar da bağlı bloklar faktörüne örnektir.

Bağlı gözlemler faktörü ve bağlı bloklar faktörü aynı şekilde analiz edildiği için çoğu zaman aynı isimle kullanılır, bu bölümde de her iki faktör *bağlı gözlemler* olarak kullanılmıştır.

Gözlemler arası faktör veya Bağlı olmayan gözlemler faktörü (Between-subjects factor) katılımcı faktörü ile kesişmeyen faktörlerdir. Toplumsal Cinsiyet Algısının (TCA) yüksek öğretim durumuna göre (yüksek öğretim mezunları ve yüksek öğretim mezunu olmayanlar) değişip değişmediğini test eden bir modelde TCA bağımlı değişken, yüksek öğretim durumu ise gözlemler arası faktördür.

9.2 Bağlı olmayan gözlemler varyans analizi (Between Subjects ANOVA)

Gözlemlerin (katılımcıların) tek bir alt sınıf kombinasyonuna ait olduğu durumlarda yapılan çözümlemelerdir.

9.2.1 Tek-yönlü bağlı olmayan gözlemler varyans analizi

Tek faktörlü bağlı olmayan gözlemler varyans çözümlemesi $Y_{ij} = \mu + \alpha_j + \epsilon_{ij}$ eşitliğinde yer alan parametrelerin tahmini ile tamamlanabilir. Bu eşitlikte Y_{ij} j alt sınıfında yer alan i katılımcısına ait puanı, μ bütün puanların ortalaması, α_j j alt sınıfının etkisi ve ϵ_{ij} hata terimidir. $\mu_j = \mu + \alpha_j$ 'dir, μ_j j alt sınıfında yer alan katılımcıların aritmetik ortalamasıdır.

Genellikle ilgi α_j üstünedir, çünkü $\mu_j - \mu$ 'yi temsil eder. Bu ilgi $H_0 : \mu_1 = \mu_2 = \dots = \mu_j$ boş hipotezinin test edilmesini gerektirir. Alternatif hipotez en az bir alt sınıfa ait ortalamanın farklı olduğunu belirtir. Bu boş hipotez varyansın ayrıştırılması ile test edilebilir. Myers et al. (2013) tarafından verilen notasyon kullanırsak;

VK	sd	KT	KTO	F
Toplam	$N - 1$	$\sum_{j=1}^J \sum_{i=1}^{n_j} (Y_{ij} - \bar{Y}_{..})^2$		
A	$J - 1$	$\sum_{j=1}^J n_j (\bar{Y}_{.j} - \bar{Y}_{..})^2$	SS_A/df_A	$MS_A/MS_{S/A}$

VK	sd	KT	KTO	F
S/A	$N - J$	$\sum_{j=1}^J \sum_{i=1}^{n_j} (Y_{ij} - \bar{Y}_{.j})^2$	$SS_{S/A}$	$SS_{S/A} / df_{S/A}$

VK	BKTO
Toplam	
A	$\sigma_{S/A}^2 + \frac{1}{J-1} \sum_j n_j (\mu_j - \mu)^2$
S/A	$\sigma_{S/A}^2$

VK = varyans kaynağı, sd = serbestlik derecesi, KT = kareler toplamı, KTO = kareler toplamı ortalaması, BKTO = beklenen kareler toplamı ortalaması.

A , J adet alt sınıfa sahip gruplar arası faktörü, S/A A faktörü içerisindeki katılımcıları, N toplam örneklem sayısını, j=1,...,J faktör alt sınıflarını, i=1,...,n_j katılımcıları, Y_{ij} katılımcı puanlarını, $\bar{Y}_{.}$ genel ortalamayı, $\bar{Y}_{.j}$ j alt sınıfı katılımcı ortalamasını temsil eder.

$MS_A/MS_{S/A}$ oranı, boş hipotez doğru olduğunda ve varsayımlar ihlal edilmediğinde, J-1 ve N-J serbestlik derecesine sahip bir F dağılımını takip eder. Dolayısı ile $MS_A/MS_{S/A}$, $F_{\alpha, J-1, N-J}$ kritik değerinden büyük ise boş hipotez terkedilir.

9.2.1.1 Etki büyüklüğü tek-yönlü bağlı olmayan gözlemler varyans analizi

Tanıtmı kolaylaştırmak amaçlı her alt sınıfın eşit sayıda gözlem içerdiğini düşünelim, $n_1 = n_2 = \dots = n_j = n$. Bu durumda A faktörü için BKTO $\sigma^2 + n\theta_A^2$ 'dır ve ;

$$\theta_A^2 = \sum_{j=1}^J \frac{(\mu - \mu_j)^2}{J-1}$$

θ_A^2 parametresinin tahmini olan $\hat{\theta}_A^2$, $\frac{MS_A - MS_{S/A}}{n}$ ile ve $\sigma_{S/A}^2$ parametresinin tahmini olan $\hat{\sigma}_{S/A}^2$, $MS_{S/A}$ ile hesaplanır.

8.1.4 bölümünde değinildiği gibi, aritmetik ortalamalar arasındaki farkın büyüklüğünü yorumlamak için etki büyüklüğü hesaplaması yapılabilir. Günümüzde varyans çözümlemesi kullanmış çoğu bilimsel makalede etki büyüklüğü de raporlanmıştır. Bu etki büyüklükleri arasında omega-kare ($\hat{\omega}^2$), eta-kare ($\hat{\eta}^2$) ve f en çok raporlananlar arasındadır.

9.2.1.1.1 Omega-kare

Omega-kare faktör tarafından açıklanan varyansın toplam varyansa oranını tahmin etmek için türetilmiştir.

$$\hat{\omega}^2 = \frac{(J-1)\hat{\theta}_A^2/J}{((J-1)\hat{\theta}_A^2/J) + \hat{\sigma}_{S/A}^2}$$

Omega-kare 0.01 ise küçük, 0.06 ise orta ve 0.14 ise büyük etki olarak yorumlanır Myers et al. (2013).

9.2.1.1.2 Eta-kare

Eta-kare de , $\hat{\eta}^2 = \frac{SS_A}{SS_{Total}}$, faktör tarafından açıklanan varyansın toplam varyansa oranını tahmin etmek için türetilmiştir.

Aynı çözümleme için, $\hat{\eta}^2$, $\hat{\omega}^2$ 'den büyüktür, çünkü $\hat{\eta}^2$, özellikle n küçük ise, pozitif yönde yanlış bir tahmindir. Buna rağmen $\hat{\eta}^2$ bildiğimiz kadarı ile en çok raporlanan etki büyüklüğüdür. $\hat{\eta}^2$, regresyon çözümlemesi çerçevesinde R^2 olarak da raporlanır.

9.2.1.1.3 Etki büyüklüğü f

Cohen'in f katsayısı, $f = \frac{\hat{\theta}_A}{\hat{\sigma}_{S/A}}$ ile hesaplanır. Bir f değeri 0.10 ise küçük, 0.25 ise orta, 0.40 ise büyük etki olarak yorumlanır.

9.2.1.1.4 A general note on the Effect Size Measures

For illustrative purposes, we briefly summarized effect size measures for equal sample size in each group. In practice it is generally not common to have equal sample sizes. It is also not common to have a single factor design. In addition, factors in a design are either measured or manipulated, which which affect the effect size computation. The *ezANOVA* function (Lawrence (2016)) reports generalized eta-squares based on Bakeman (2005). The work by Bakeman (2005) encourages researchers to use generalized eta-squared defined by Olejnik and Algina (2003). Hence, a convenient choice for a researcher is to use the *ezANOVA* function, while paying attention to the *observed* argument to declare the measured factors. On the other hand, if it is not desired to be dependent on an R package, the researcher can examine and apply the formulae by Olejnik and Algina (2003).

9.2.1.2 Testing specific contrasts of means

Either in addition to or in place of the ANOVA, specific contrasts (comparisons) of means may be tested. A contrast is a weighted sum of means in which the weights sum to zero. There are two classes of contrasts: pairwise contrasts and complex contrasts. To illustrate these classes consider a one-way design in which the factor has three levels, a control treatment and two active treatments. Let the population means for these levels be μ_1 , μ_2 , and μ_3 , respectively. In a pairwise contrast two means are compared and the weights are 1 for one mean, -1 for another and zero for all others. A pairwise contrast of the means for the active treatments is $(0)\mu_1 + (1)\mu_2 + (-1)\mu_3$. The complex contrast $(-1)\mu_1 + (-.5)\mu_2 + (-.5)\mu_3$ is a comparison of the mean for the control group to the average of the means for the two active treatments. Under the assumptions of a one-way between-subjects ANOVA, the null hypothesis that a contrast is equal to zero can be tested using

$$t = \frac{\sum_{j=1}^J (w_j \bar{Y})}{\sqrt{MS_{S/A} \sum_{j=1}^J \left(\frac{w_j^2}{n_j}\right)}}$$

9.2.1.3 Testing all possible pairwise comparisons

There are several procedures for testing all possible pairwise contrasts. An important issue in such testing is the error rate to control. Controlling an error rate means keeping it at or below some conventional level (e.g., .05). Two of the most common error rates are the per comparison error rate and the familywise error rate. The per comparison error rate is the probability of making a Type I error when one of the contrasts is tested. To control the per comparison error rate the critical value for a pairwise comparison is $\pm t_{(1-\alpha/2), N-J}$. When this critical value is used, the per comparison error rate is α . The family wise error rate is the probability of falsely rejecting one of more of the contrasts. If all pairwise contrasts are equal to zero, the family wise error rate is between α and $[J(J-1)/2]\alpha$. The upper limit can be quite high even when the number of levels of the factor is small. For example if there are $J=3$ levels, the upper limit is 3α . There are several procedures for controlling the familywise error rate.

9.2.1.3.1 Trend analyses following one-way between-subjects ANOVA

To be added.

9.2.1.4 Assumptions of the one-way between-subjects ANOVA

The assumptions of the one-way between-subjects ANOVA are the same as the assumptions of the independent samples t test.

1. Independence. The scores in each group should be independently distributed and the scores in different groups should also be independent. The validity of this assumption in regard to independence within groups is questionable when (a) scores for participants within a group are collected over time or (b) the participants within a group work together in a manner such that a participant's response could have been influenced by another participant in the study. The validity of this assumption in regard to independence between groups is questionable when the factor is a within-subjects factor rather than a between-subjects factor. Violating the independence assumption is a critical violation that usually can be addressed by adopting an analysis appropriate for the lack of independence. For example, if there are different participants in each group, but within each group there are subgroups of participants who work together then according to (b) above independence is likely to have been violated. This violation can be addressed by using multilevel analysis. If there are different participants in each group, but the participants in the groups have been matched, using a randomized block ANOVA can address the violation of independence.
2. Normality. The scores with each group are drawn from a normal distribution. Statistical power is likely to be compromised if the distributions of scores have long tails. When the sample sizes are equal violating normality is not likely to affect the type I error rate, unless the non-normality is severe and the sample sizes are small.
3. Equal variance. This assumption is also called the homogeneity of variance assumption and means it is assumed that samples in the J groups are drawn from J populations with equal variances. Violation of the equal variance assumption is likely to affect the Type I error rate except when the sample sizes are equal and fairly large.

Even though we briefly summarized the assumptions of the one-way between subjects ANOVA above, they were only introductory. If independence does not appear to be violated, then when the sample sizes are equal and at least 20 in each group and the scores are approximately normally distributed the one-way between subjects ANOVA can be used. In other situations alternatives should be used. When the robust analyses (e.g. Wilcox (2012)) and conventional analyses yield the same decisions about **all hypothesis tests**, results of the conventional analyses can be reported due to their greater familiarity to most readers.

9.2.1.5 R codes for a one-way between-subjects ANOVA

For illustrative purposes, the city of KOCAELI is subsetting from the DataWBT (Section 2.3). The gender attitudes scores is the dependent variable and the highest degree completed is the between subjects factor. This factor had seven levels; no-degree, primary school, middle school, high school, vocational high school, 2 year college and bachelors. However, there is only one participant in the *no-degree* group. We combined the no-degree and primary school groups. The gender attitude score for this participant is 1.6.¹

For illustrative purposes, the city of KOCAELI is subsetting from the DataWBT (Section 2.3). The gender attitudes scores is the dependent variable and the highest degree completed is the between subjects factor. This factor had seven levels; no-degree, primary school, middle school, high school, vocational high school, 2 year college and bachelors. However, there is only one participant in the *no-degree* group. We relocated this individual into primary school level. The gender attitude score for this participant is 1.6. Removing this participant from the ANOVA has no substantial effect on the results.

¹Removing this participant from the ANOVA would have had no substantial effect on the results.

Step 1: Set up data and report descriptive

```
# load csv from an online repository
urlfile='https://raw.githubusercontent.com/burakaydin/materyaller/gh-pages/ARPASS/dataWBT.csv'
dataWBT=read.csv(urlfile)

#remove URL
rm(urlfile)

#select the city of KOCAELI
# listwise deletion for gen_att and education variables
dataWBT_KOCAELI=na.omit(dataWBT[dataWBT$city=="KOCAELI",
                                c("id","gen_att","education")])

#There is only 1 participant in the level "None", merge it into Primary school
# the gender attitude score for this participant is 1.6
library(car)
dataWBT_KOCAELI$eduNEW <- recode(dataWBT_KOCAELI$education,
                                "'None'='Primary School (5 years)'" )

dataWBT_KOCAELI$eduNEW <- recode(dataWBT_KOCAELI$eduNEW,
                                "'High School (Lycee)'='
                                'High School (Lycee) (4 years)'" )

dataWBT_KOCAELI$eduNEW <- recode(dataWBT_KOCAELI$eduNEW,
                                "'Vocational School'='
                                'Vocational High School (4 years)'" )

#table(dataWBT_KOCAELI$eduNEW)

##optional re-order levels (cosmetic)
#levels(dataWBT_KOCAELI$eduNEW)
dataWBT_KOCAELI$eduNEW = factor(dataWBT_KOCAELI$eduNEW,
                                levels(dataWBT_KOCAELI$eduNEW)[c(4,3,1,6,2,5)])

#which(dataWBT_KOCAELI$education=="None")

#drop empty levels
dataWBT_KOCAELI$eduNEW=droplevels(dataWBT_KOCAELI$eduNEW)

#get descriptives
library(psych)
desc1BW=data.frame(with(dataWBT_KOCAELI,
                        describeBy(gen_att, eduNEW,mat=T,digits = 2)),
                    row.names=NULL)

#select relevant descriptives
# Table 1
desc1BW[,c(2,4,5,6,7,13,14)]
##
## 1 Primary School (5 years) 70 2.11 0.41 2.2 -0.19 0.81
```

```
## 2 Junior High/ Middle School (8 years) 94 2.08 0.52 2.1 -0.35 -0.37
## 3 High School (Lycee) (4 years) 158 1.84 0.58 2.0 0.29 0.64
## 4 Vocational High School (4 years) 74 2.04 0.50 2.0 -0.14 0.41
## 5 Higher education of 2 years 112 1.80 0.53 1.8 0.28 -0.36
## 6 University - Undergraduate degree 62 1.78 0.53 1.8 0.06 -0.63
#write.csv(desc1BW,file="onewayB_ANOVA_desc.csv")
```

Step 2: Check assumptions

```
require(ggplot2)
ggplot(dataWBT_KOCAELI, aes(x = gen_att)) +
  geom_histogram(aes(y = ..density..),col="black",binwidth = 0.2,alpha=0.7) +
  geom_density(size=1.5) +
  theme_bw()+labs(x = "Gender Attitude by Degree in Kocaeli")+ facet_wrap(~ eduNEW)+
  theme(axis.text=element_text(size=14),
        axis.title=element_text(size=14,face="bold"))
```




Figure 9.1: Gender Attitudes by Degree

Departures from the normality do not seem to be severe.

```
require(ggplot2)
ggplot(dataWBT_KOCAELI, aes(eduNEW,gen_att)) +
  geom_boxplot() +
  labs(x = "Education",y="Gender Attitude by degree in Kocaeli")+coord_flip()
```




Figure 9.2: Gender Attitudes by Degree

Homogeneity of variance is questionable but not severely violated.

Step 3: Run ANOVA

For illustrative purposes, let us ignore the violations first. The *ezANOVA* function (Lawrence (2016)) reports the F test, the Levene Test and an effect size. Type of the effect size depends on the model. For further details, please carefully study the Table 1 in Bakeman (2005), an open access article, or Olejnik and Algina (2003). The Levene test rejects the null hypothesis of equal variances across factor levels.

```
library(ez)
#the ezANOVA function throws a warning if id is not a factor

dataWBT_KOCAELI$id=as.factor(dataWBT_KOCAELI$id)

# set the number of decimals (cosmetic)
options(digits = 3)

#alternative 1 the ezANOVA function

alternative1 = ezANOVA(
  data = dataWBT_KOCAELI,
  wid=id, dv = gen_att, between = eduNEW,observed=eduNEW)
## Warning: Data is unbalanced (unequal N per group). Make sure you specified
## a well-considered value for the type argument to ezANOVA().

alternative1
## $ANOVA
##   Effect DFn DFd    F      p p<.05    ges
## 1 eduNEW   5 564 7.27 1.31e-06 * 0.0605
##
## $`Levene's Test for Homogeneity of Variance`
##   DFn DFd SSn  SSd    F      p p<.05
## 1   5 564 1.35 63.5 2.4 0.0361 *

# critical F value
qf(.95,5,564)
## [1] 2.23
```

ABOUT the warning of ez function;
*#Warning: Data is unbalanced (unequal N per group). Make sure you specified
 #a well-considered value for the type argument to ezANOVA().*

ezANOVA can calculate three different types of sums of squares for main effects and interactions.
 For a one-way between-subjects design the F test is the same for all three types and this warning can be ignored.

The same results can be obtained with the *lm* (linear model) function in R Core Team (2016b).

```
# alternative 2 the lm function
alternative2=lm(gen_att~eduNEW,data=dataWBT_KOCAELI)
```

```
#Table 2
anova(alternative2)
## Analysis of Variance Table
##
## Response: gen_att
##          Df Sum Sq Mean Sq F value    Pr(>F)
## eduNEW      5   10.1    2.026     7.27 1.3e-06 ***
## Residuals 564  157.2    0.279
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

The *aov* function in R Core Team (2016b) is the third alternative.

```
#alternative 3 the aov function
alternative3=aov(gen_att~eduNEW,data=dataWBT_KOCAELI)
summary(alternative3)
##          Df Sum Sq Mean Sq F value    Pr(>F)
## eduNEW      5   10.1    2.026     7.27 1.3e-06 ***
## Residuals 564  157.2    0.279
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

The *pairwise.t.test* function in the *stats* package (R Core Team (2016b)) is convenient. Provide the preferred procedure by using *p.adjust.method* argument, for example *p.adjust.method = "Holm"* to use the adjustment given by Holm (1979). Five other procedures are available with this function, please see *?p.adjust*.

```
# pairwise comparisons
# Table 3
with(dataWBT_KOCAELI, pairwise.t.test(gen_att,eduNEW,p.adjust.method ="holm"))
##
## Pairwise comparisons using t tests with pooled SD
##
## data:  gen_att and eduNEW
##
##                                     Primary School (5 years)
## Junior High/ Middle School (8 years) 1.000
## High School (Lycee) (4 years)         0.004
## Vocational High School (4 years)      1.000
## Higher education of 2 years           0.001
## University - Undergraduate degree     0.004
##
##                                     Junior High/ Middle School (8 years)
## Junior High/ Middle School (8 years) -
## High School (Lycee) (4 years)         0.005
## Vocational High School (4 years)      1.000
## Higher education of 2 years           0.002
## University - Undergraduate degree     0.006
##
##                                     High School (Lycee) (4 years)
## Junior High/ Middle School (8 years) -
## High School (Lycee) (4 years)         -
## Vocational High School (4 years)      0.044
## Higher education of 2 years           1.000
## University - Undergraduate degree     1.000
##
##                                     Vocational High School (4 years)
## Junior High/ Middle School (8 years) -
```



```
## High School (Lycee) (4 years)      -
## Vocational High School (4 years)  -
## Higher education of 2 years       0.018
## University - Undergraduate degree 0.036
##                                   Higher education of 2 years
## Junior High/ Middle School (8 years) -
## High School (Lycee) (4 years)      -
## Vocational High School (4 years)  -
## Higher education of 2 years       -
## University - Undergraduate degree  1.000
##
## P value adjustment method: holm
```

9.2.1.6 Robust estimation and hypothesis testing for a one-way between-subjects design

Several approaches to conducting a robust one-way between subjects ANOVA, have been presented by Wilcox (2012). One of the convenient robust procedure, a heteroscedastic one-way ANOVA for trimmed means, has been compressed into the *t1way* function, available via WRS-2 (Mair and Wilcox (2016)). Please use *?t1way* for the current details, this promising package is being improved frequently.

```
library(WRS2)

#t1way
# 20% trimmed
t1way(gen_att~eduNEW,data=dataWBT_KOCAELI,tr=.2,nboot=5000)
## Call:
## t1way(formula = gen_att ~ eduNEW, data = dataWBT_KOCAELI, tr = 0.2,
##       nboot = 5000)
##
## Test statistic: 7.57
## Degrees of Freedom 1: 5
## Degrees of Freedom 2: 144
## p-value: 0
##
## Explanatory measure of effect size: 0.29

# 10% trimmed
t1way(gen_att~eduNEW,data=dataWBT_KOCAELI,tr=.1,nboot=5000)
## Call:
## t1way(formula = gen_att ~ eduNEW, data = dataWBT_KOCAELI, tr = 0.1,
##       nboot = 5000)
##
## Test statistic: 9.54
## Degrees of Freedom 1: 5
## Degrees of Freedom 2: 188
## p-value: 0
##
## Explanatory measure of effect size: 0.3

# 5% trimmed
t1way(gen_att~eduNEW,data=dataWBT_KOCAELI,tr=.05,nboot=5000)
## Call:
## t1way(formula = gen_att ~ eduNEW, data = dataWBT_KOCAELI, tr = 0.05,
```

```

##      nboot = 5000)
##
## Test statistic: 9.41
## Degrees of Freedom 1: 5
## Degrees of Freedom 2: 212
## p-value: 0
##
## Explanatory measure of effect size: 0.31

## heteroscedastic pairwise comparisons

#level order
lincon(gen_att~eduNEW,data=dataWBT_KOCAELI,tr=.1)[[2]]
## [1] "Higher education of 2 years"
## [2] "Junior High/ Middle School (8 years)"
## [3] "University - Undergraduate degree"
## [4] "Vocational High School (4 years)"
## [5] "High School (Lycee) (4 years)"
## [6] "Primary School (5 years)"
round(lincon(gen_att~eduNEW,data=dataWBT_KOCAELI,tr=.1)[[1]][,c(1,2,6)],3)
##      Group Group p.value
## [1,]      1      2  0.701
## [2,]      1      3  0.000
## [3,]      1      4  0.360
## [4,]      1      5  0.000
## [5,]      1      6  0.000
## [6,]      2      3  0.000
## [7,]      2      4  0.597
## [8,]      2      5  0.000
## [9,]      2      6  0.000
## [10,]     3      4  0.004
## [11,]     3      5  0.460
## [12,]     3      6  0.467
## [13,]     4      5  0.001
## [14,]     4      6  0.003
## [15,]     5      6  0.911

```

9.2.1.7 Example writeup for one-way between-subjects ANOVA

For our illustrative example, results of hypothesis tests conducted using robust procedures did not disagree with the results of the ANOVA and pairwise comparisons of means. This was expected given the assumptions were not severely violated. When the robust analyses and conventional analyses yield the same decisions about *all hypothesis tests*, results of the conventional analyses can be reported due to their greater familiarity to most readers. A possible write up for our illustrative example would be:

An ANOVA was performed to investigate whether the gender attitudes scores differ across education level. The means, standard deviations, skewness and kurtosis values of the gender scores, grouped by the highest-degree obtained, are presented in Table 1. The analysis of variance indicated a significant difference in the gender attitudes scores, $F(5,564) = 7.27$, $p < .001$, $\eta_G^2 = .06$. Table 2 is the ANOVA table for this analysis. Pairwise comparisons were planned a priori. The familywise error rate was selected for control and the Holm procedure (Holm (1979)) was used. The results of the pairwise comparisons are presented in the Table 3. Nine out of fifteen comparisons yielded statistically significant results; (primary school vs lycee, primary school vs 2-year-collage, primary school vs undergraduate,... (provide details). Robust statistical procedures

yielded the same conclusions.

9.2.1.8 Missing data techniques for one-way between-subjects ANOVA

To be added

9.2.1.9 Power calculations for one-way between-subjects ANOVA

To be added

9.2.2 Two-Factor Between Subjects ANOVA

This topic concerns designs in which there are two between-subjects factors: factor A with J levels and factor B with K levels for a total of JK combinations of levels; each combination is called a cell. The factors are between-subjects if (a) different subjects appear in each cell and (b) subjects are not matched in any way. In the simplest version of this design, each factor has two levels. For example, consider if a researcher is interested in the effect of the gender and college education on the gender attitudes scores. The following is a depiction of a study designed to investigate these two factors.

	non-college	college	
Female	μ_{11}	μ_{12}	$\mu_{1\cdot}$
Male	μ_{21}	μ_{22}	$\mu_{2\cdot}$
	$\mu_{\cdot 1}$	$\mu_{\cdot 2}$	

Also shown are the parameters about which hypotheses will be tested: the population cell means ($\mu_{11}, \mu_{12}, \mu_{21}, \mu_{22}$), row means ($\mu_{1\cdot}, \mu_{2\cdot}$) and column means ($\mu_{\cdot 1}, \mu_{\cdot 2}$). The general term for a row or column mean is a marginal mean.

Symbolically the hypothesis of no interaction can be written as $H_0 : \mu_{11} - \mu_{12} = \mu_{21} - \mu_{22}$. The interaction is also a comparison of the two simple effects of gender ($\mu_{21} - \mu_{11}$ and $\mu_{22} - \mu_{12}$) leading to the null hypothesis $H_0 : \mu_{21} - \mu_{11} = \mu_{22} - \mu_{12}$. If one of the null hypotheses is true the other must also be true and if one is false the other must also be false.

Interaction The first null hypothesis of interest is the hypothesis of no interaction between the two factors. Before defining an interaction, we first define a simple main effect. A simple main effect refers to differences among the cell means in a particular row or in a particular column. In the current example, there are two types of simple main effects: simple main effects of gender and simple main effects of college education.

For each level of graduation from college type there are two simple main effects gender: the simple main effect of gender at college graduates (μ_{12} versus μ_{22}) and the simple main effect of gender at non-college graduates (μ_{11} versus μ_{21}). The interaction is a comparison of these two effects.

For each type there are two simple main effects. There is a simple main effect of gender at college graduates (μ_{12} versus μ_{22}) and a simple main effect of gender at non-college graduates (μ_{11} versus μ_{21}).

There is a simple main effect of education for Female (μ_{11} versus μ_{12}) and a simple main effect of education for Male (μ_{21} versus μ_{22}).

The main effects Effects defined in terms of marginal (row and column) means are called main effects. Symbolically, the main effect of gender is $\mu_{1\cdot} - \mu_{2\cdot}$, and the hypothesis of no main effect due to gender is $H_0 : \mu_{1\cdot} - \mu_{2\cdot} = 0$. Similarly, the hypothesis of no main effect due to college education is $H_0 : \mu_{\cdot 1} - \mu_{\cdot 2} = 0$.

When there is an interaction:

1. Inspection of the main effect for a factor is misleading when the directions of the simple main effects of the factor are not the same at all levels of the second factor.
2. It is a matter of opinion as to whether it is misleading to inspect the main effect for a factor the directions of the simple main effects of the factor are the same at all levels of the second factor.

When the main effect is misleading about the effect of a factor, the cell means are the proper basis for studying the effects of the factor.

The structural model for a two-factor between subjects ANOVA is $Y_{ijk} = \mu + \alpha_j + \beta_k + \alpha\beta_{jk} + \epsilon_{ij}$, in which Y_{ijk} is the score for the participant i in first factor level j , and the second factor level k ; μ is the grand mean of the scores, α_j is the effect of the level j of the first factor, β_k is the effect of the level k of the second factor, $\alpha\beta_{jk}$ is the interaction effect and ϵ_{ij} is the error term (nuisance).

SV	df	F
A	$J - 1$	$\frac{MS_A}{MS_{S/AB}}$
B	$K - 1$	$\frac{MS_B}{MS_{S/AB}}$
AB	$(J - 1)(K - 1)$	$\frac{MS_{AB}}{MS_{S/AB}}$
S/AB	$N - JK$	
Total	$N - 1$	

9.2.2.1 Type I, II and III sum of squares

As we pointed out in the section on one-way between-subjects designs, the F test of the main effect is the same for all three types of sums of squares. This is not true in designs with two or more between-subjects factors. In designs with three or more between-subjects of effects F tests for interaction other than the highest order interaction can vary across the types of sums of squares. Selecting among the three types can be an important decision and we refer the reader to Carlson and Timm (1974) for a discussion of the issues in selecting among the three types of sums of squares in experimental studies and to Appelbaum and Cramer (1976) for a discussion of the issues in survey studies.

9.2.2.2 R codes for a two-way between-subjects ANOVA

For illustrative purposes, the city of Kayseri is subsetting from the DataWBT (Section 2.3). The gender attitudes scores is the dependent variable, gender and higher education indicator are the between subjects factors.

Step 1: Set up data and report descriptive

```
# load csv from an online repository
urlfile='https://raw.githubusercontent.com/burakaydin/materyaller/gh-pages/ARPASS/dataWBT.csv'
dataWBT=read.csv(urlfile)

#remove URL
rm(urlfile)

#select the city of Izmir
# listwise deletion for gen_att and education variables
dataWBT_Kayseri=na.omit(dataWBT[dataWBT$city=="KAYSERI",c("id","gen_att","higher_ed","gender")])

# Higher education is coded as 0 and 1, change it to non-college, college
dataWBT_Kayseri$HEF=droplevels(factor(dataWBT_Kayseri$higher_ed,
                                     levels = c(0,1),
```

```

labels = c("non-college", "college"))

#table(dataWBT_Kayseri$gender)
#table(dataWBT_Kayseri$HEF)

#drop empty levels
dataWBT_Kayseri$gender=droplevels(dataWBT_Kayseri$gender)

with(dataWBT_Kayseri,
      table(gender,HEF))
##           HEF
## gender    non-college college
## Female           99      50
## Male             67      36

# set the number of decimals (cosmetic)
options(digits = 3)

#get descriptives
library(doby)
library(moments)
desc2BW=as.matrix(summaryBy(gen_att~HEF+gender, data = dataWBT_Kayseri,
                             FUN = function(x) { c(n = sum(!is.na(x)),
                                                     mean = mean(x,na.rm=T), sdv = sd(x,na.rm=T),
                                                     skw=moments::skewness(x,na.rm=T),
                                                     krt=moments::kurtosis(x,na.rm=T)) } ))

# Table 4
desc2BW
##      HEF      gender  gen_att.n gen_att.mean gen_att.sdv gen_att.skw
## 1 "non-college" "Female" "99"      "1.93"      "0.424"      "-0.548"
## 2 "non-college" "Male"   "67"      "2.32"      "0.419"      "-0.191"
## 3 "college"     "Female" "50"      "1.80"      "0.346"      " 0.263"
## 4 "college"     "Male"   "36"      "2.13"      "0.543"      " 0.159"
##      gen_att.krt
## 1 "2.51"
## 2 "3.18"
## 3 "1.94"
## 4 "2.25"
#write.csv(desc2BW,file="twowayB_ANOVA_desc.csv")

```

Step 2: Inspect assumptions

```

require(ggplot2)
ggplot(dataWBT_Kayseri, aes(x = gen_att)) +
  geom_histogram(aes(y = ..density..),col="black",binwidth = 0.2,alpha=0.7) +
  geom_density(size=1.5) +
  theme_bw()+labs(x = "Gender Attitudes by HEF and Gender in Kayseri")+ facet_wrap(~ HEF+gender)+
  theme(axis.text=element_text(size=14),
        axis.title=element_text(size=14,face="bold"))

```

Departures from the normality do not seem to be severe.




Figure 9.3: Gender Attitudes by HEF and Gender

```
require(ggplot2)
ggplot(dataWBT_Kayseri, aes(x=gender, y=gen_att))+
  geom_boxplot()+
  facet_grid(.~HEF)+
  labs(x = "Gender",y="Gender Attitude by Gender and HEF in Kayseri")
```




Figure 9.4: Gender Attitudes by Degree

Variances look similar.

Step 3: Run ANOVA

The *ezANOVA* function (Lawrence (2016)) reports the F test, the Levene Test and an effect size. Type of the effect size depends on the model and indirectly depends on the type of sum of squares used. The *type* argument (1,2 or 3) transmits the choice.

```
library(ez)
#the ezANOVA function throws a warning if id is not a factor

dataWBT_Kayseri$id=as.factor(dataWBT_Kayseri$id)

#alternative 1 the ezANOVA function
alternative1 = ezANOVA(
  data = dataWBT_Kayseri,
  wid=id, dv = gen_att, between = .(HEF,gender),observed=.(HEF,gender),type=2)
## Warning: Data is unbalanced (unequal N per group). Make sure you specified
```

```
## a well-considered value for the type argument to ezANOVA().
```

```
alternative1
## $ANOVA
##      Effect DFn DFd      F      p p<.05      ges
## 1      HEF    1 248  6.739 9.99e-03    * 0.022436
## 2    gender    1 248 45.389 1.12e-10    * 0.151106
## 3 HEF:gender    1 248  0.251 6.17e-01    0.000837
##
## $`Levene's Test for Homogeneity of Variance`
##      DFn DFd  SSn  SSd      F      p p<.05
## 1      3 248 0.469 17.5 2.22 0.0867
```

```
# Type III SS
# alternative1b = ezANOVA(
#   data = dataWBT_Kayseri,
#   wid=id, dv = gen_att, between = HEF+gender,type=3)
#
# alternative1b
# critical F value
qf(.95,1,248)
## [1] 3.88
```

9.2.2.3 Robust estimation and hypothesis testing for a two-way between-subjects design

Several approaches to conducting a robust two-way between subjects ANOVA, have been presented by Wilcox (2012). One of the convenient robust procedure, a heteroscedastic two-way ANOVA for trimmed means, has been compressed into the *t2way* function, available via WRS-2 (Mair and Wilcox (2016)). Please use *?t2way* for the current details, this promising package is being improved frequently.

```
library(WRS2)

#t2way
# 20% trimmed
t2way(gen_att~HEF*gender,data=dataWBT_Kayseri,tr=.2)
## Call:
## t2way(formula = gen_att ~ HEF * gender, data = dataWBT_Kayseri,
##       tr = 0.2)
##
##              value p.value
## HEF           7.1310  0.011
## gender        20.2039  0.001
## HEF:gender     0.0855  0.772

# 10% trimmed
t2way(gen_att~HEF*gender,data=dataWBT_Kayseri,tr=.1)
## Call:
## t2way(formula = gen_att ~ HEF * gender, data = dataWBT_Kayseri,
##       tr = 0.1)
##
##              value p.value
```



```
## HEF          8.4235    0.005
## gender       33.1599    0.001
## HEF:gender   0.0361    0.850

# 5% trimmed
t2way(gen_att~HEF*gender,data=dataWBT_Kayseri,tr=.05)
## Call:
## t2way(formula = gen_att ~ HEF * gender, data = dataWBT_Kayseri,
##       tr = 0.05)
##
##              value p.value
## HEF          6.169   0.015
## gender       29.838   0.001
## HEF:gender    0.164   0.687
```

9.2.2.4 Example writeup two-way between-subjects ANOVA

For our illustrative example, robust procedures did not disagree with our initial analyses. This was expected given the assumptions were not severely violated. When the robust analyses yield very similar results, we prefer to report initial results to ease communication. A possible write up for our illustrative example would be:

Descriptive statistics for the gender attitudes scores as a function of gender and higher education in the city of Kayseri are presented in Table 4. A 2x2 ANOVA was reported. F tests were conducted at $\alpha = .05$. There was a significant difference for gender $F(1, 248) = 45.39, p < .001$. There was also a significant difference for the college effect $F(1, 248) = 6.24, p = .013$. However, there was no significant interaction between the gender and higher education status, $F(1, 248) = 0.25, p = .617$. The *ezANOVA* (Lawrence (2016)) function reported generalized eta hat squared ($\hat{\eta}_G^2$) of 0.15 for the gender effect and 0.02 for the college-effect. Table 5 is the ANOVA table for these analyses.

9.2.2.5 Follow ups for two-way between-subjects ANOVA

To be added.

9.2.2.5.1 Pairwise comparisons following two-way between-subjects ANOVA

To be added.

9.2.2.5.2 Contrasts comparisons following two-way between-subjects ANOVA

To be added.

9.2.2.6 Missing data techniques for two-way between-subjects ANOVA

To be added

9.2.2.7 Power calculations for two-way between-subjects ANOVA

To be added

9.3 Within Subjects ANOVA

This procedure can be used when there are score on the same participants under several treatments or at several time points and is then called repeated measures ANOVA. It can also be used in blocking designs and is then called randomized block ANOVA. Compared to between-subjects designs, this procedure is expected to eliminate influence of individual differences, in other words, to reduced variability, and thus, to reduce error. This development results in more power than independent-samples ANOVA with the same sample size. Of course there are issues other than power that must be considered in selecting a design. For example, if the goal is to compare reading achievement under three instructional methods, using a design in which each participant is exposed to the three methods will be problematic because the effect of exposure to one treatment will continue to influence reading ability during the other treatments.

9.3.1 One-way Within-Subjects ANOVA

The structural model following Myers et al. (2013) notation for a non-additive model;

$$Y_{ij} = \mu + \eta_i + \alpha_j + (\eta\alpha)_{ij} + \epsilon_{ij} \quad (9.1)$$

where i represents the individual, $i=1,\dots,n$; j represents the levels of the within-subjects factor (i.e, the repeated measurement or the treatment factor), $j=1,\dots,P$. Y is the score; μ is the grand mean; η_i represents the difference between individual's average score over the levels and the grand mean; α_j represents the difference between the average score under level j of the within-subjects factor and the grand mean; $(\eta\alpha)_{ij}$ represents the interaction, and ϵ_{ij} represent the error component. Because $(\eta\alpha)_{ij}$ and ϵ_{ij} have the same subscripts, they are confounded.

Generally, the interest is on α_j . This interest leads to hypothesis testing:

$$H_0 : \mu_1 = \mu_2 = \dots = \mu_P$$

The alternative hypothesis states that at least one population mean is different. The ANOVA table for a one-way with-subjects ANOVA is;

SV	df	F
Subjects (S)	$n - 1$	
Waves (A)	$P - 1$	$\frac{MS_A}{MS_{SA}}$
SA	$(n - 1)(P - 1)$	
Total	$nP - 1$	

Note on additivity The simplest explanation of additivity is the situation that would justify $(\eta\alpha)_{ij} = 0$ in Equation (9.1). This unrealistic restriction implies that the effect of levels of the within-subject factor waves is the same for all individuals.

Shown below in Table 9.8 are data for an experiment in which each person participates in four treatments defined by the amount of alcohol consumed. The dependent variable is a reaction time measure.

Treatment means, treatment standard deviations, and subject means are shown below. A subject mean is the average of the four scores for that subject.

```
# set the number of decimals (cosmetic)
options(digits = 2)

#participants mean
```

Table 9.8: Original Alcohol Data

id	noALC	twoOZ	fouroz	sixoz
1	1	2	5	7
2	2	3	5	8
3	2	3	6	8
4	2	3	6	9
5	3	4	6	9
6	3	4	7	10
7	3	4	7	10
8	6	5	8	11

Table 9.9: Correlation Coefficients for Reaction Time Data

	noALC	twoOZ	fouroz	sixoz
noALC	1.00	0.93	0.88	0.88
twoOZ	0.93	1.00	0.89	0.94
fouroz	0.88	0.89	1.00	0.95
sixoz	0.88	0.94	0.95	1.00

```

apply(owadata,1, mean)
## [1] 3.2 4.0 4.4 4.8 5.4 6.0 6.2 7.6

#treatment mean
apply(owadata[, -1], 2, mean)
## noALC twoOZ fouroz sixoz
## 2.8 3.5 6.2 9.0

#treatment sd
apply(owadata[, -1], 2, sd)
## noALC twoOZ fouroz sixoz
## 1.49 0.93 1.04 1.31

```

Because each participant has a score for each treatment, amount of alcohol is a within-subjects factor and it is possible to calculate a correlation for each pair of treatments. These correlations, which are presented in Table 9.2, indicate that reaction time is highly correlated for each pair of treatments. Recall that corresponding to each correlation there is a covariance;

$$Cov_{pp'} = S_p S_{p'} r_{pp'}$$

where p and p' are two levels of the alcohol consumption factor. For example the correlation between the scores in the first two levels of the alcohol consumption factor is $r_{02} = 0.93$. And the corresponding covariance is $Cov_{02} = 1.5 * 0.9 * 0.93 = 1.26$

The alcohol consumption factor is a within-subjects factor. Consequently the F statistic for comparing the four treatment means should be appropriate for a design with a within-subjects factor. Let

P = the number of levels of the within-subjects factor, in the example $P=4$;

\bar{C} = the average covariance; in the example $\bar{C} = 1.26$.

The appropriate F statistic is

$$F_W = \frac{MS_A}{MS_{SA}} = \frac{MS_A}{MS_{S/A} - \bar{C}}$$

where MS_A and $MS_{S/A}$ are calculated as they are for a between-subjects factor and the W emphasizes the F statistic is for a within-subjects factor. The critical value is $F_{\alpha, P-1, (P-1)(n-1)}$. The denominator mean square, MS_{SA} , is read mean square Subjects x A where A is the generic label for the treatment factor. Recall that the F statistic for a between-subjects factor is $F_B = MS_A / MS_{S/A}$. Comparison of F_W and F_B shows that F_W incorporates the correlations between the pairs of treatments and F_B does not. (Recall that, in like fashion, the dependent samples t statistic incorporates the correlation whereas the independent samples t statistic does not.) As a result, when applied to a design with a within-subjects factor $F_W \geq F_B$. Therefore, incorrectly using will usually result in a loss of power.

9.3.1.1 Assumptions of one-way within-subjects ANOVA

Sphericity is an assumption about the pattern of variances and covariances. If the data are spherical, the difference between each pair of repeated measures has the same variance for all pairs.

Example covariance matrix;

	Y_1	Y_2	Y_3
Y_1	10	7.5	10
Y_2	7.5	15	12.5
Y_3	10	12.5	20

Sphericity holds;

$Y_p - Y_{p'}$	$\sigma_p^2 + \sigma_{p'}^2 - 2\sigma_{pp'}$
$Y_1 - Y_2$	$10 + 15 - 2(7.5) = 10$
$Y_1 - Y_3$	$10 + 20 - 2(10) = 10$
$Y_2 - Y_3$	$15 + 20 - 2(12.5) = 10$

Box's epsilon —measures how severely sphericity is violated.

$$\frac{1}{P-1} \leq \epsilon \leq 1$$

Estimates of ϵ are Greenhouse-Geisser ($\hat{\epsilon}$) and Huynh-Feldt ($\tilde{\epsilon}$)

$\tilde{\epsilon}$ can be larger than 1.0; if it is $\tilde{\epsilon}$ is set equal to 1.0.

Critical value assuming sphericity is $F_{\alpha, (P-1), (n-1)(P-1)}$.

Approximately correct critical value when sphericity is violated $F_{\alpha, \epsilon(P-1), \epsilon(n-1)(P-1)}$.

normality of errors in Equation (9.1), ϵ_{ij} is assumed to be normally and independently distributed with a mean value of zero.

normality of η_i in Equation (9.1), η_i is assumed to be normally and independently distributed with a mean value of zero.

Together the assumptions listed immediately above imply that the repeated measures are drawn from a multivariate normal distribution.

9.3.1.1.1 The relationship between additivity and sphericity

Although assumptions can be stated in terms of η_i and ϵ_{ij} , a simpler approach is that the repeated measures are drawn from a multivariate normal distribution with covariance matrix that meets the sphericity assumption. If the data meet the sphericity assumption, the difference between each pair of repeated measures has the same variance for all pairs.

Assuming that the data are drawn from a multivariate normal distribution and within each level of the within-subjects factor the scores are independent, having equal variance and covariances (compound symmetry) is a sufficient condition for the F test on the within-subjects factor to be valid.

If additivity holds and the equal variance assumption holds then compound symmetry holds. But compound symmetry is a stricter assumption than the sphericity. Considering that sphericity is a necessary and sufficient requirement for the F test on the within-subjects factor to be valid (assuming that data are drawn from a multivariate normal distribution and scores are independent within each level of the within-subjects factor the) checking for sphericity is more important than checking for additivity. In addition because there are adjusted degrees of freedom procedures that adjust the F test on the within-subjects factor for violation of sphericity, it is not necessary to test for sphericity.

9.3.1.2 R codes for a one-way within-subjects ANOVA

For illustrative purposes, a subsample from an original cluster randomized trial study (for details see Daunic et al. (2012)) was taken. The subsample included 1 control-classroom and 17 students. The variable of interest is the problem solving knowledge. Each wave was approximately one year apart. Higher scores mean higher knowledge.

Step 1: Set up data

```
#enter data
PSdata=data.frame(id=factor(1:17),
                  wave1=c(20,19,13,10,16,12,16,11,11,14,13,17,16,12,12,16,16),
                  wave2=c(28,27,18,17,29,18,26,21,15,26,28,23,29,18,26,21,22),
                  wave3=c(21,24,14,8,23,15,21,15,12,21,23,17,26,18,14,18,19))
```

Report descriptive

```
# set the number of decimals (cosmetic)
options(digits = 3)

##the long format will be needed
#head(PSdata)
library(tidyr)
data_long = gather(PSdata, wave, PrbSol, wave1:wave3, factor_key=TRUE)

#get descriptives
library(doby)
library(moments)
desc1W=as.matrix(summaryBy(PrbSol~wave, data = data_long,
                          FUN = function(x) { c(n = sum(!is.na(x)),
                                                mean = mean(x,na.rm=T), sdv = sd(x,na.rm=T),
                                                skw=moments::skewness(x,na.rm=T),
                                                krt=moments::kurtosis(x,na.rm=T)) } ))

# Table 6
desc1W
##   wave   PrbSol.n PrbSol.mean PrbSol.sdv PrbSol.skw PrbSol.krt
## 1 "wave1"  "17"      "14.4"      "2.91"      " 0.311"    "2.10"
```

```
## 2 "wave2" "17"      "23.1"      "4.67"      "-0.224"     "1.64"
## 3 "wave3" "17"      "18.2"      "4.77"      "-0.315"     "2.45"
#write.csv(desc1W,file="onewayW_ANOVA_desc.csv")
```

The covariance matrix might be helpful.

```
# check covariance Table 7
cov(PSdata[,-1])
##      wave1 wave2 wave3
## wave1  8.49  8.85  9.87
## wave2  8.85 21.81 18.49
## wave3  9.87 18.49 22.78
```

Step 2: Check assumptions

```
ggplot(data_long, aes(x=wave, y=PrbSol))+
  geom_boxplot()+
  labs(x = "Wave",y="Problem Solving Knowledge scores")
```




Figure 9.5: Problem Solving Knowledge score by wave

We will test for the sphericity assumption using Mauchly's test integrated in the *ezANOVA* function, but this graph implies that it might be violated.

```
require(ggplot2)
ggplot(data_long, aes(x=wave, y=PrbSol, group=id))+
  geom_line() + labs(x = "Wave",y="Problem Solving Knowledge scores")
```




Figure 9.6: Problem Solving Knowledge score by wave, line graph

This graph, which plots the problem solving scores by wave, suggests that the $\eta\beta_{ij}$ interaction terms are not likely to all be zero; therefore assuming sphericity while testing the hypothesis of equal wave means is not likely to be justified. The `tukey.add.test` function in *asbio* (Aho (2016)) investigates H_0 : *main effect and blocks are additive*.

```
library(asbio)
with(data_long,tukey.add.test(PrbSol,wave,id))
##
## Tukey's one df test for additivity
## F = 5.943   Denom df = 31   p-value = 0.021

# if additivity exists a randomized block approach might be appropriate
#additive=with(data_long,lm(PrbSol~id+wave))
#anova(additive)
```

The Tukey additive test rejects the null hypothesis of additivity agrees with the line graph. In other words, a non-additive model is more appropriate for the problem solving knowledge data.

Step 3: Run ANOVA (including checks for the sphericity and normality of residuals assumptions)

The *ezANOVA* function (Lawrence (2016)) reports the F test, the Levene Test and an effect size. Type of the effect size depends on the model and indirectly depends on the type of sum of squares used, the *type* argument (1,2 or 3) transmits the choice.

```

library(ez)
#alternative 1 the ezANOVA function

alternative1 = ezANOVA(
  data = data_long,
  wid=id, dv = PrbSol, within = wave,
  detailed = T,return_aov=T)

alternative1
## $ANOVA
##      Effect DFn DFd  SSn SSd    F      p p<.05  ges
## 1 (Intercept)   1  16 17510 680 412.0 7.62e-13 * 0.954
## 2      wave    2  32   647 169  61.2 1.16e-11 * 0.433
##
## $`Mauchly's Test for Sphericity`
##      Effect      W      p p<.05
## 2      wave 0.918 0.526
##
## $`Sphericity Corrections`
##      Effect  GGe      p[GG] p[GG]<.05  HFe      p[HF] p[HF]<.05
## 2      wave 0.924 6.17e-11          * 1.04 1.16e-11          *
##
## $aov
##
## Call:
## aov(formula = formula(aov_formula), data = data)
##
## Grand Mean: 18.5
##
## Stratum 1: id
##
## Terms:
##              Residuals
## Sum of Squares      680
## Deg. of Freedom      16
##
## Residual standard error: 6.52
##
## Stratum 2: id:wave
##
## Terms:
##              wave Residuals
## Sum of Squares   647      169
## Deg. of Freedom    2      32
##
## Residual standard error: 2.3
## Estimated effects may be unbalanced

PrbSolres=sort(alternative1$aov$id$residuals)
qqnorm(PrbSolres);qqline(PrbSolres)

```

The distribution of the residuals is not severely non-normal.

The *aov* function is the second alternative.




Figure 9.7: Inhibit model residuals

```
# alternative 2 the aov function
summary(aov(PrbSol ~ wave + Error(id/wave), data=data_long))
##
## Error: id
##           Df Sum Sq Mean Sq F value Pr(>F)
## Residuals 16   680    42.5
##
## Error: id:wave
##           Df Sum Sq Mean Sq F value Pr(>F)
## wave       2   647    324   61.2 1.2e-11 ***
## Residuals 32   169      5
## ---
## Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

9.3.1.3 Robust estimation and hypothesis testing for a one-way within-subjects design

One of the convenient robust procedures, a heteroscedastic one-way repeated measures ANOVA for trimmed means, has been compressed into the *rmanova* function, available via WRS-2 (Mair and Wilcox (2016)).

```
library(WRS2)

#rmanova
# 20% trimmed
with(data_long, rmanova(PrbSol, wave, id, tr=.20))
## Call:
## rmanova(y = PrbSol, groups = wave, blocks = id, tr = 0.2)
##
## Test statistic: 34.9
## Degrees of Freedom 1: 1.9
## Degrees of Freedom 2: 19
## p-value: 0
```

9.3.1.4 Example writeup one-way within-subjects ANOVA

Descriptive statistics for the problem solving knowledge scores at each wave are presented in Table 6. The covariance matrix is presented in Table 7. A one-way within ANOVA was reported. F test was conducted at $\alpha = .05$. The assumptions of one-way within subjects ANOVA are satisfied. There was a significant difference between waves $F(2, 32) = 61.2, p < .001$. The *ezANOVA* function reported a generalized eta hat squared ($\hat{\eta}_G^2$) of 0.43.

9.3.1.5 Follow ups for one-way within-subjects ANOVA

To be added.

9.3.1.6 Missing data techniques for one-way within-subjects ANOVA

To be added.

9.3.1.7 Power calculations for one-way within-subjects ANOVA

To be added.

9.4 Mixed Design

To be added.

Chapter 10

Correlation

In our chapter 7, we introduced descriptive statistics; mean, variance, median, kurtosis, etc. These descriptive statistics aimed to ease the communication for a single variable. In other words, instead of transferring the entire raw data set to a colleague (or to a machine), providing these descriptives is generally satisfying and easier. However when the interest is in the association between variables, other measures are needed.

The sum of cross products, $S_{XY} = \sum(X - \bar{X})(Y - \bar{Y})$, can provide some information about the association. For example Figure 10.1 depicts an X and a Y variable. The sum of cross products for these two variables is zero.

##	x	y	deviationX	deviationY	crossPRODUCT
## 1	1.00	0.00	0.93	0.00	0.00
## 2	0.90	0.43	0.83	0.43	0.36
## 3	0.62	0.78	0.56	0.78	0.44
## 4	0.22	0.97	0.16	0.97	0.15
## 5	-0.22	0.97	-0.29	0.97	-0.28
## 6	-0.62	0.78	-0.69	0.78	-0.54
## 7	-0.90	0.43	-0.97	0.43	-0.42
## 8	-1.00	0.00	-1.07	0.00	0.00
## 9	-0.90	-0.43	-0.97	-0.43	0.42
## 10	-0.62	-0.78	-0.69	-0.78	0.54
## 11	-0.22	-0.97	-0.29	-0.97	0.28
## 12	0.22	-0.97	0.16	-0.97	-0.15
## 13	0.62	-0.78	0.56	-0.78	-0.44
## 14	0.90	-0.43	0.83	-0.43	-0.36
## 15	1.00	0.00	0.93	0.00	0.00

The covariance between two variable is simply $Cov_{XY} = S_{XY}/n - 1$, but its a scale dependent measure, the correlation coefficient on the other hand generally has its bounds.

10.1 Pearson correlation coefficient

Pearson introduced a correlation coefficient in 1896. This coefficient ranges between -1 and +1, can be calculated as $Cov_{XY}/S_X S_Y$. This coefficient measures the linear relationship between two variables. Figure 10.1 depicts a correlation of zero. Even though X and Y in this figure are related to form a 14-sided polygon, the relation is not linear. Hence the correlation is zero. Figure 10.2 depicts several other associations; (A) is a perfect positive linear relationship, (B) is a positive correlation of .7, (C) substantially no linear relation, (D) is a correlation of -.4 and (E) is a correlation of -1.




Figure 10.1: Sum of cross products=0




Figure 10.2: Correlation examples

10.1.1 Inference on a Pearson correlation coefficient

Information from the sample (r) can be utilized to make judgement about the population (ρ).

The z transformation, assuming a bivariate normality and a sample size of at least 10 (Myers et al. (2013)), is a helpful procedure to reach a judgement. The transformation equation is;

$$z_r = \frac{1}{2} \ln \left(\frac{1+r}{1-r} \right)$$

The standard error is;

$$\sigma_r = \frac{1}{\sqrt{n-3}}$$

Hence the confidence intervals are $z_r \pm z_{\alpha/2} \sigma_r$. Back transformation is needed to make interpretation about the correlation coefficient; $r = \frac{e^{2z_r} - 1}{e^{2z_r} + 1}$.

Utilizing a normal distribution, a null hypothesis can be tested;

$$z = \frac{z_r - z_{\rho_{null}}}{\frac{1}{\sqrt{n-3}}}$$

The t distribution can also be utilized to test $H_0 : \rho = 0$.

$$t = r \sqrt{\frac{n-2}{1-r^2}}$$

The distribution for this statistic follows a t distribution with a degrees of freedom of $n - 2$.

10.1.2 R codes for Pearson Correlation coefficient

For illustrative purposes we selected the city of Bayburt. The Pearson correlation is computed for the association between the Gender Attitudes scores and the annual income per person. The income per person is calculated as “total household income” divided by the “total number of residents in the house”.

```
# load csv from an online repository
urlfile='https://raw.githubusercontent.com/burakaydin/materyaller/gh-pages/ARPASS/dataWBT.csv'
dataWBT=read.csv(urlfile)

#remove URL
rm(urlfile)

#select the city of Elazig
# listwise deletion for gen_att and education variables
dataWBT_Bayburt=dataWBT[dataWBT$city=="BAYBURT",]
#hist(dataWBT_Bayburt$income_per_member)
```

The bivariate distribution can be seen in 10.1.2. This is an interactive graph, please use your mouse to inspect it, created with the *rgl* package (Adler and Murdoch (2017)).

```
## wgl
## 3
```

Gender Attitudes and Income




Figure 10.3: Bivariate Normal Distribution




Figure 10.4: Bayburt: Gender attitudes vs income scatterplot

Bivariate normality seems to be violated. For comparison, below graph 10.3 depicts a bivariate normal distribution with $r=0.7$. Nevertheless, for illustrative purposes we use these data to test the null hypothesis $H_0 : \rho = 0$ against the non-directional alternative hypothesis $H_1 : \rho \neq 0$. The scatter plot is provided in 10.4.

The correlation between these two variables is computed by the `cor` function in the `stats` package (R Core Team (2016b)). The `cor.test` function in the same package performs the t-test and provides a confidence interval based on Fisher's z transformation.

```
#use ?cor to see use="complete.obs" is doing casewise deletion

with(dataWBT_Bayburt,cor(gen_att,income_per_member,
                          use = "complete.obs",method="pearson"))

## [1] 0.0664

with(dataWBT_Bayburt,cor.test(gen_att,income_per_member,
                              alternative = "two.sided",
                              method="pearson",
                              conf.level = 0.95,
                              na.action="na.omit"))

##
## Pearson's product-moment correlation
##
## data: gen_att and income_per_member
## t = 0.8, df = 100, p-value = 0.4
## alternative hypothesis: true correlation is not equal to 0
## 95 percent confidence interval:
## -0.102 0.232
## sample estimates:
## cor
## 0.0664
```

These procedures can easily be hard coded. Stating $H_0 : \rho = 0$ and $H_0 : \rho \neq 0$

```
sample_r=0.06641641
r0=0          #the null
sample_n=137  # the number of complete.cases
zr=(0.5)*log((1+sample_r)/(1-sample_r)) # Z transformasyonu
z0=(0.5)*log((1+r0)/(1-r0)) # Z transformasyonu
sigmar=1/(sqrt(sample_n-3))

#the z test statistic
(zr-z0)/sigmar
## [1] 0.77

ll=zr-(qnorm(0.975)*sigmar) # lower limit

ul=zr+(qnorm(0.975)*sigmar) # upper limit

(exp(2*ll)-1)/(exp(2*ll)+1) #transformback
## [1] -0.102
(exp(2*ul)-1)/(exp(2*ul)+1) #transformback
## [1] 0.232
```

```

t=sample_r*(sqrt((sample_n-2)/(1-sample_r^2)))
qt(c(.025, .975), df=(sample_n-2))
## [1] -1.98  1.98
p.value = 2*pt(-abs(t), df=sample_n-2)
p.value
## [1] 0.441

```

A percentile bootstrap method might perform satisfactorily as a robust approach (Myers et al. (2013))

```

#Calculate 95% CI using bootstrap (normality is not assumed)
set.seed(31012017)
B=5000          # number of bootstraps
alpha=0.05      # alpha

#gender attitudes and income
originaldata=dataWBT_Bayburt2

#add id
originaldata$id=1:nrow(originaldata)

output=c()
for (i in 1:B){
  #sample rows
  bs_rows=sample(originaldata$id,replace=T,size=nrow(originaldata))
  bs_sample=originaldata[bs_rows,]
  output[i]=cor(bs_sample$gen_att,bs_sample$income_per_member)
}
output=sort(output)

## Non-directional
# lower limit
output[as.integer(B*alpha/2)]
## [1] -0.138

# d star upper
output[B-as.integer(B*alpha/2)+1]
## [1] 0.252

# investigate the WRS package
# library(WRS2)
# pball(originaldata[,-3])

```

There are alternatives to percentile bootstrapping for a correlation coefficient, extensively discussed by Wilcox (2012). The WRS 2 package offers two alternatives, the percentage bend correlation and the Winsorized correlation. Only for illustrative purposes below is an R code;

```

# investigate the WRS package
library(WRS2)
pbcor(dataWBT_Bayburt2$gen_att,dataWBT_Bayburt2$income_per_member,beta=.2)
## Call:
## pbcor(x = dataWBT_Bayburt2$gen_att, y = dataWBT_Bayburt2$income_per_member,
##      beta = 0.2)
##
## Robust correlation coefficient: -0.0351
## Test statistic: -0.407

```

```
## p-value: 0.684

wincor(dataWBT_Bayburt2$gen_att,dataWBT_Bayburt2$income_per_member,tr=.2)
## Call:
## wincor(x = dataWBT_Bayburt2$gen_att, y = dataWBT_Bayburt2$income_per_member,
##       tr = 0.2)
##
## Robust correlation coefficient: -0.0197
## Test statistic: -0.229
## p-value: 0.82
```

Write up: We tested a null hypothesis stating the gender attitudes scores and income variable are correlated against an non-directional alternative hypothesis. The Pearson correlation coefficient was $r = .066$, $p = .44$, the confidence interval with a .05 probability of a type I error using the z transformation is -.10 to .23. The null hypothesis is retained. This conclusion is consistent with the bootstrap results, using 5000 iterations, the 95% CI is -.138 to .252.

Sign difference note The Pearson correlation coefficient is .066 but not significantly different than zero. The WRS package functions also agreed to retain the null but the coefficient was negative. The income variable was slightly skewed due to a small number of relatively large income values. In fact, when the World Bank team analyzed the data using a regression, they top-coded and transformed the income variable (for details Hirshleifer et al. (2016)). Let us top-code and transform the income variable, inspect bivariate normality and calculate the Pearson correlation;




Figure 10.5: Top-coded and transformed income variable

```

with(dataWBT_Bayburt2, cor.test(gen_att, incomeTC,
                                alternative = "two.sided",
                                method="pearson",
                                conf.level = 0.95,
                                na.action="na.omit"))

##
## Pearson's product-moment correlation
##
## data:  gen_att and incomeTC
## t = -0.009, df = 100, p-value = 1
## alternative hypothesis: true correlation is not equal to 0
## 95 percent confidence interval:
##  -0.169  0.167
## sample estimates:
##      cor
## -0.00081

```

Top-coding and transforming the income variable produced a distribution relatively closer to normal. The sign of the Pearson correlation coefficient is negative.

10.2 Spearman's rho and Kendall's tau

When the data is in the rank format, or there is a need for protection against outliers¹ when working with continuous data the Spearman correlation coefficient is used. If the number of ties in the ranks is not large, procedures provided for the Pearson correlation coefficient can be utilized. Setting the method argument to "spearman", the *cor.test* function first transforms the data into ranks and performs the procedures introduced for the Pearson coefficient.

10.2.1 The R code for Spearman's rho and Kendall's tau

We calculated the Pearson correlation coefficient to assess the association between the gender attitudes scores and the income for the participants in Bayburt. The Spearman correlation coefficient can conveniently be calculated by R;

```

#use ?cor to see use="complete.obs" is doing casewise deletion
with(dataWBT_Bayburt, cor.test(gen_att, income_per_member,
                                alternative = "two.sided",
                                method="spearman",
                                conf.level = 0.95,
                                na.action="na.omit",
                                exact=FALSE))

##
## Spearman's rank correlation rho
##
## data:  gen_att and income_per_member
## S = 5e+05, p-value = 0.6
## alternative hypothesis: true rho is not equal to 0
## sample estimates:

```

¹Here protection refers to being less sensitive to outliers compared to Pearson coefficient. However Spearman's rho and Kendall's tau might be more sensitive to outliers compared to robust procedures, see Wilcox (2012).

```
##      rho
## -0.0508
```

When there are ties, the `cor.test` function corrects the Spearman coefficient but the exact p value can not be calculated. Instead `exact=FALSE` argument yields a p value based on a t distribution. Field et al. (2012) suggests using Kendall's tau with large number of ties;

```
#use ?cor to see use="complete.obs" is doing casewise deletion
with(dataWBT_Bayburt,cor.test(gen_att,income_per_member,
  alternative = "two.sided",
  method="kendall",
  conf.level = 0.95,
  na.action="na.omit",
  exact=FALSE))

##
## Kendall's rank correlation tau
##
## data:  gen_att and income_per_member
## z = -0.6, p-value = 0.5
## alternative hypothesis: true tau is not equal to 0
## sample estimates:
##      tau
## -0.0373
```

The `exact=FALSE` argument with `method="kendall"` uses normal approximation.

The Spearman correlation between the gender attitudes scores and income was $r_S = -.051, p = .56$, and the Kendall's tau was $\tau = -.037, p = .54$

10.3 Biserial and Point-Biserial Correlation Coefficients with R

The association between a continuous variable and a dichotomously reflected latent continuous variable can be examined with a biserial correlation. In psychometrics, for example, biserial correlation is used for calculating the correlation between a total test score (continuous) and a dichotomous item score (assumed to underlie a latent variable).

For illustrative purposes let us use dichotomized item1² and the gender attitudes score. The `biserial` function in the *psych* (Revelle (2016)) package can calculate the bi-serial correlation;

```
dataWBT_Bayburt$binitem1=ifelse(dataWBT_Bayburt$item1==4,1,0)
require(psych)
with(dataWBT_Bayburt,biserial(gen_att,binitem1))
##      [,1]
## [1,] 0.317
```

The point-biserial correlation is calculated for an association between a dichotomous variable and a continuous variable. The `cor.test` function with `method="pearson"` can be used to calculate a point-biserial correlation. The association between the gender and the gender attitudes scores is examined below;

```
dataWBT_Kayseri=dataWBT[dataWBT$city=="KAYSERI",]
dataWBT_Kayseri$genderNUM=ifelse(dataWBT_Kayseri$gender=="Female",1,0)
with(dataWBT_Kayseri,cor.test(gen_att,genderNUM,
  alternative = "two.sided",
  method="pearson",
```

²This item is indeed dichotomized by the Worldbank team in their analyses

```

    conf.level = 0.95,
    na.action="na.omit"))
##
## Pearson's product-moment correlation
##
## data:  gen_att and genderNUM
## t = -7, df = 200, p-value = 2e-10
## alternative hypothesis: true correlation is not equal to 0
## 95 percent confidence interval:
##  -0.487 -0.277
## sample estimates:
##      cor
## -0.387

```

10.4 Phi Correlation Coefficient with R

When the two variables are dichotomous, a phi (ϕ) correlation coefficient is calculated. For illustrative purposes we calculated the phi coefficient between the gender and the wage variable. This variable equals to “yes” if one of the house members receives wage in the past 12 months. The *phi* function in the *psych* package requires the 2 x 2 matrix of frequencies to calculate the phi coefficient.

```

dataWBT_Kayseri=dataWBT[dataWBT$city=="KAYSERI",]
table(dataWBT_Kayseri$gender,dataWBT_Kayseri$wage01)
##
##           No Yes
## Female   52  97
## Male     49  54
## Unknown   0   0

genderWAGE=matrix(c(52,49,97,54),ncol=2)
library(psych)
phi(genderWAGE)
## [1] -0.13

phi(genderWAGE)
## [1] -0.13

```

10.5 Issues in Interpreting Correlation Coefficients

Several issues arise in interpreting correlation coefficients.

Causation A correlation coefficient does not imply causation. For any correlation there are at least four possible interpretations involving causation: (a) X causes Y, (b) Y causes X, (c) both X and Y share one or more common causes, and (d) X and Y have different causes, but these causes are correlated.

The magnitude Whether a correlation of .6 is large or not depends on the context. For example suppose the .6 is the correlation between scores on two forms of a standardized achievement tests. This correlation is called an alternate forms reliability coefficient. Alternate forms reliability coefficients for standardized tests are expected to be at least .70 and preferably higher, so the .6 correlation would be regarded as small. Now suppose the correlation is between GRE scores and GPA. The correlation between GRE scores and GPA is typically somewhere between .10 and .30, so a .60 correlation would be a very large correlation coefficient.

Outliers Correlation coefficients can be misleading when the data set contains outliers.

Reliability If either X or Y contains measurement error, the effect of the measurement error is to attenuate the correlation coefficient. Attenuate means to make the correlation coefficient closer to zero than it would have been if there had been no measurement error.

Its possible to correct for attenuation using

$$r_{T_x T_y} = \frac{r_{xy}}{\sqrt{(r_{xx} r_{yy})}}$$

where r_{xx} and r_{yy} is the reliability coefficients.

- *When NOT to correct for attenuation:* When a variable is used for practical decision making and we are interested in the validity of those decisions, we should NOT correct for attenuation because the decisions are made on the basis of an observed variable, not a true variable.
- *When to correct for attenuation:* We can correct for attenuation when our motivation is to examine theory.
- *Comparison of Correlation Coefficients:* A comparison of correlation coefficients for two variables with a third variable can be affected by differences in reliability for the first two variables. If we are interested in theoretical relationships between variables and we want to compare the strength of relationship of two constructs (call these A and B and let them be measured by X1 and X2) with a third (call this C and let it be measured by Y), the comparison of the strength of relationship between A and C to the strength of relationship between B and C is compromised if X1 and X2 have different reliability coefficients. To compare strength of relationship we want the reliability of X1 and X2 to be the similar. Of course, it is best if both reliability coefficients are high, but it is critical that that are quite similar.

Unit of analysis A correlation calculated for one unit of analysis (e.g., individuals without regard to school) should not be applied to other units of analysis (i.e., individuals within schools or school means).

Variance in the two variables being correlated The correlation coefficient for two variables can be strongly affected by the amount of variance for the variables being correlated. Other things being equal when the variance of either or both variables is small, the correlation will tend to be small. If the variance for either or both variables is artificially small, misleading small correlation coefficients can occur. Variance can be artificially small due to

- Categorizing Based on Quantitative Variables
- Limited Range Scales
- Restriction of range
- Floor and Ceiling Effects

Chapter 11

Multiple Linear Regression, A Short Introduction

Scientific development requires that knowledge be transferred reliably from one study to another and, as Galileo showed 350 years ago, such transference requires the precision and computational benefits of a formal language. Pearl (2009)

The *formal language* in the quote refers to *mathematical equations*. Galton for example, in late 1800s, used equations to describe the relationship between the weights of mother and daughter pea seeds. Galton's work followed by Pearson's contributions led to initial idea of regression ¹.

In the year 2016, the Web of Science reported that 60000+ abstracts of academic articles included the term "regression". The literature is vast, oftentimes the regression is mentioned as the *workhorse*. It is extensively used by frequentist and Bayesian statisticians, and more generally by data scientists in hundreds of different disciplines. The explanation of the popularity of regression analysis is simple, unless they are simulated by a machine, connections between variables, whether observed or latent variables, in a data set requires more complex statistical solutions than are provided by correlation coefficients.

It is not feasible to cover regression in a book chapter. We briefly introduce basics of a relatively simple multiple regression model.

11.1 Matrices and Least Square Estimation

In a multiple regression framework, demonstrating process of model fitting based on matrices and least squares estimation should have at least two benefits; (a) a simple demystification of the procedure, (b) a workable and sensible foundation for readers with a desire to move further in the advanced topics. The following sections use two different data sets. The first data set includes only 12-cases to show calculations and is named the synthetic data. The second data is simulated with a larger sample size for illustrative purposes and is named the simulated data.

Consider a case in which data on three variables are collected and the researcher is interested in the relationship of one of the variables (i.e., the dependent variable) to the other two variables (i.e., the independent variables). Further, these three variables are continuous. The regression model in this scenario is

$$Y_i = \beta_0 + \beta_1 X_{i1} + \beta_2 X_{i2} + \epsilon_i$$

¹<http://ww2.amstat.org/publications/jse/v9n3/stanton.html>

where i represents individuals $i=1,\dots,n$, Y is the dependent variable, X_1 and X_2 are independent variables, β s are the regression coefficients and ϵ is the random error term (residuals). This model can be presented in a matrix equations

$$Y = X\beta + \epsilon$$

In this more general form, all the independent variables are represented in the X matrix and the regression coefficients are represented by the β matrix. Let us assume the researcher has the following data

id	Y	X1	X2
ind 1	8	0	3
ind 2	4	-2	1
ind 3	6	6	3
ind 4	6	-2	0
ind 5	5	5	0
ind 6	9	4	2
ind 7	7	3	3
ind 8	-6	-4	-5
ind 9	-8	-4	-6
ind 10	-1	-3	0
ind 11	0	-2	-2
ind 12	5	-1	1

This synthetic data set has only 12 cases. The researcher can form 2 matrices and use these to calculate $\hat{\beta}$, the estimate of β .

$$Y = \begin{bmatrix} 8 \\ 4 \\ 6 \\ 6 \\ 5 \\ 9 \\ 7 \\ -6 \\ -8 \\ -1 \\ 0 \\ 5 \end{bmatrix}, X = \begin{bmatrix} 1 & 0 & 3 \\ 1 & -2 & 1 \\ 1 & 6 & 3 \\ 1 & -3 & 0 \\ 1 & 5 & 0 \\ 1 & 4 & 2 \\ 1 & 3 & 3 \\ 1 & -4 & -5 \\ 1 & -4 & -6 \\ 1 & -3 & 0 \\ 1 & -2 & -2 \\ 1 & -1 & 1 \end{bmatrix}$$

Using the least square procedure the β coefficients can easily be estimated;

$$\hat{\beta} = (X'X)^{-1}X'Y \quad (11.1)$$

Let's calculate this with R for the synthetic-data;

```
Y=matrix(c(8,4,6,6,5,9,7,-6,-8,-1,0,5),ncol=1)
X=matrix(cbind(rep(1,12),
               c(0,-2,6,-2,5,4,3,-4,-4,-3,-2,-1),
               c(3,1,3,0,0,2,3,-5,-6,0,-2,1)),ncol=3)

solve(t(X)%*%X)%*%t(X)%*%Y
##      [,1]
## [1,] 2.917
```

```
## [2,] 0.199
## [3,] 1.552
```

The regression equation is

$$\hat{Y}_i = 2.9167 + 0.1989X_{i1} + 1.5519X_{i2}$$

where \hat{Y}_i is the predicted value for the i^{th} individual. Equation (11.1) was derived to minimize the error sum of squares: $\sum_{i=1}^n (Y_i - \hat{Y}_i)^2 = Y'Y - \beta'X'X\beta$. These estimates are Best Linear Unbiased Estimates.

Each independent variable has a mean of zero because they are mean-centered. Therefore, zero represents a score at the center of the distribution for both X_1 and X_2 and is therefore an interpretable score for both X_1 and X_2 . When both predictors are zero (at their mean), the (\hat{Y}_i) is 2.92. That is, for participants with independent variables scores equal to the mean on both independent variables the expected dependent variable score is 2.92. An increase in X_1 of 1 unit is predicted to correspond to an increase of 0.20 units in Y when the X_2 variable is held constant. Similarly, an increase in X_2 of 1 unit is predicted to correspond to an increase of 1.55 units in Y while controlling for X_1 . The term “controlling for” (“*ceteris paribus*”) is necessary to describe the effect of an independent variable in a multiple regression. The coefficients .20 and 1.55 would provide information about the association of the dependent and independent variables, if the researcher had substantial understanding of the unit of measurement for the independent variables, that is, the importance of a “1 unit” change in each variable.

11.1.1 a) “Essentially, all models are wrong, but some are useful.”

This aphorism belongs to Box and Draper (1987). The researcher should provide a convincing discussion about the relevance of the variables included in the regression model to the research questions addressed by the research. If there are important omitted variables, the beta coefficients are probably not valid. Hence the researcher is obligated to provide justifications on variable selections to claim usefulness of the results.

Consider the case below ;

```
#omit X2 from the synthetic-data
X2omitted=matrix(cbind(rep(1,12),c(0,-2,6,-2,5,4,3,-4,-4,-3,-2,-1)),ncol=2)
solve(t(X2omitted)%*%X2omitted)%*%t(X2omitted)%*%Y
##      [,1]
## [1,] 2.92
## [2,] 1.09
```

For our synthetic data, X_1 and X_2 had a correlation of .68. If the researcher fails to include X_2 in the model, the coefficient for X_1 is estimated to be 1.09. This is a dramatic change from 0.20. Omitting predictors that are related to both the other predictors in the model and the dependent variable will cause the coefficients for the variables that have not been omitted to be misleading. Therefore an important part of the theoretical justification of a regression model is a discussion of variables that may have been omitted.²

In addition to omitted variable issue, the validity of the results from a regression model (the usefulness) is also directly related to the sampling process and appropriate reflection of this process in the model. For example, if sampling weights exist they should not be ignored in the analyses. *Sampling and regression* is beyond the scope of this chapter.

11.1.2 b) Strength of relationship between the dependent and independent variables

The sum of squares for Y , which is also known as the total sum of squares, can be decomposed into two parts, *the model sum of squares*, which is also the sum of squares for the predicted values, and *the error*

²This might lead to a clue on popularity of controlled randomized trials.

sum of squares. The ratio of the model sum of squares to the *total sum of squares*, is called the sample squared multiple correlation coefficient and symbolized as R^2 . The coefficient R^2 measures the strength of association between the dependent variable and the independent variables. Examine the R code below given for the synthetic data;

```
# SS total
n=length(Y)
TotalSS=t(Y)%*%Y-(n*mean(Y)^2)

# SS Model
betahat=solve(t(X)%*%X)%*%t(X)%*%Y
ModelSS=t(betahat)%*%t(X)%*%Y-(n*mean(Y)^2)

ModelSS/TotalSS
##          [,1]
## [1,] 0.879
```

Also known as *coefficient of determination*, R^2 is a biased estimator of the population squared multiple correlation coefficient. A more nearly unbiased estimate is the adjusted squared multiple correlation coefficient. One benefit of adjusted R^2 is computational simplicity. Examine the R code below given for the synthetic data

```
Rsquared=ModelSS/TotalSS
#sample size
n=12

#the number of predictors
p=2

# include intercept? 1 for yes, 0 for no
int_inc=1

AdjustedRsquared=1-(1-Rsquared)*((n-int_inc)/(n-int_inc-p))
AdjustedRsquared
##          [,1]
## [1,] 0.852
```


R^2 and R^2_{Adj} are useful coefficients; they provide information on how much of the variance is explained. Note that in this example $R^2 = .879$ and $R^2_{Adj} = .852$ are very similar. However if were $R^2 = .25$ then R^2_{Adj} would be .08. When R^2 is 1, the model successfully explains 100% of the variance in Y and when R^2 is 0 the model does not explain any of the variance in Y. The coefficients R^2 and R^2_{Adj} are also useful for comparing the strength of relationship for different set of predictors to predict a specific outcome. The interpretation of the R^2 is similar to the interpretation of a correlation coefficient. Depending on the context a small R^2 value might be regarded as substantial, or an R^2 value of .7 might be regarded as low.

11.1.3 c) Residuals and influential data points

Residuals provide information for assessing potential problems with the model. Inspecting residuals can provide information about deviations from the assumed linearity of the relationships of the dependent variables to the independent variable. Inspecting the distributional properties of residuals is needed to provide evidence for the validity of statistical inference. For example, because the normality assumption is made when conducting significance tests and calculating confidence interval, residuals should follow a straight line

on a Quantile-Quantile (QQ) plot. Examine the R code below given for the synthetic data:

```
#Predicted values
Yhat=X%*%betahat
residuals=Y-Yhat
residuals
##           [,1]
## [1,]  0.4276
## [2,] -0.0708
## [3,] -2.7658
## [4,]  3.4811
## [5,]  1.0888
## [6,]  2.1839
## [7,] -1.1691
## [8,] -0.3615
## [9,] -0.8096
## [10,] -3.3199
## [11,]  0.5850
## [12,]  0.7303
qqnorm(residuals)
```



There are three common types of residuals;

- Unstandardized residuals, that is, $Y_i - \hat{Y}_i$. Unstandardized residuals are on the same scale as Y .
- Standardized residuals: The residuals divided by the overall standard deviation of residuals; Standardized residuals are on a z-score scale ($M = 0$, $SD = 1$). When residuals are assumed to be normally distributed, it is common practice to identify outliers as Y values for which the absolute value of the standardized residual

is larger than 2. However, it should be noted that this practice can be misleading because outliers can cause the regression coefficients to be poorly estimated and/or can increase the standard deviation of the residuals and both effects can cause poor outlier detection. In addition, if the residuals are in fact normally distributed approximately 5% of the participants will have residuals beyond ± 2.00 . See Wilcox (2012) for more information about outlier detection.


- Studentized residual: A studentized residual is ratio of the unstandardized residual to the estimated standard error of the residual.

When investigating residuals, these three types of residuals generally lead to same conclusions. The standardized residuals are forced to have a z-scale, and thus, -2 and +2 are commonly pronounced cut offs. The studentized residuals are connected to the t distribution; $t_{n-p'-1}$ where n is the sample size p' is the number of coefficients in the model (i.e. intercept + two predictors = 3). It is argued that when detecting outliers in residuals, investigating the studentized residuals is more convenient (Rawlings et al. (1998)).

Scatter plots of residuals vs. predicted values can provide information about whether the assumed linear relationships between the independent variables and the dependent variable are adequate. Ideally the scatter plot should not show a detectable pattern. Here is a plot of studentized residuals vs fitted values, from a regression model fitted to simulated data in which the linearity assumption is adequate. The simulated data have a sample size of 500 and two independent variables.

```
#simulate data
library(mvtnorm)
sigma <- matrix(c(4,2,2,3), ncol=2)
xx <- rmvnorm(n=500, mean=c(0,0), sigma=sigma)
yy=5+xx[,1]*2+xx[,2]*-3+rnorm(500,0,1.5)
model=lm(yy~xx[,1]+xx[,2])
errors=rstudent(model)
predicted=predict(model)

#Standardized Residuals vs Yhat
library(ggplot2)
plotdata=data.frame(errors,predicted)
ggplot(plotdata, aes(x = predicted, y = errors)) +
  geom_point() + geom_hline(yintercept=0) + ylab("Studentized residuals")+
  theme_bw()+stat_smooth()
```





The blue line above, which is determined independently from the regression model, should be compared to the horizontal line at 0. The more similar the two lines, the less likely the linearity assumption is violated.

Here is a plot, studentized residuals vs fitted values, from a mis-specified regression model on a simulated data. The sample size is 100, there are two independent variables, and the relationship of Y and X2 is quadratic.

```
#simulate data
library(mvtnorm)
sigma <- matrix(c(4,2,2,3), ncol=2)
xx <- rmvnorm(n=100, mean=c(10,10), sigma=sigma)
yy=150+(xx[,1]*4)+(xx[,2]*-3)+(xx[,2]^2*1.2)+rnorm(100,0,3)
model=lm(yy~xx[,1]+xx[,2])
errors=rstudent(model)
predicted=predict(model)

#Studentized Residuals vs Yhat
library(ggplot2)
plotdata=data.frame(errors,predicted)
ggplot(plotdata, aes(x = predicted, y = errors)) +
  geom_point() + geom_hline(yintercept=0)+ylab("Studentized residuals")+
  theme_bw()+stat_smooth()
```



There is a pattern indicating that the model is omitting a quadratic association. However, this graph does not inform about the source of the quadratic association, see non-linearity section below.

Unusual residuals should be inspected. Even when the residuals are substantially normally distributed and there is substantially no-pattern for the residual vs predicted value plot, there might be unusual residuals. Deciding whether a residual is unusual or not (e.g 3,4 or 5 standard deviation above), and more importantly whether to keep the observation in the data set or not requires justifications. Examine the code below to simulate data and examine the studentized residuals:

```
#simulate data
set.seed(04022017)
library(mvtnorm)
sigma <- matrix(c(4,2,2,3), ncol=2)
xx <- rmvnorm(n=100, mean=c(10,10), sigma=sigma)
yy=(xx[,1]*4)+(xx[,2]*-3)+rnorm(100,0,3)
tempdata=data.frame(yy,xx,id=1:100)
model=lm(yy~X1+X2,data=tempdata)
tempdata$SUTresiduals=rstudent(model)
# how many of the residuals are larger than a critical value?
# lets use alpha=.05
sum(abs(tempdata$SUTresiduals)>qt(c(.975), df=100-3-1))
## [1] 8

#which observations?
tempdata[which(abs(tempdata$SUTresiduals)>qt(c(.975), df=100-3-1)),]
##      yy    X1    X2 id SUTresiduals
## 13 21.39 11.49 10.29 13          2.02
```

```
## 32  8.85 11.96 10.65 32      -2.20
## 43 15.80 11.14  7.56 43      -1.99
## 50  9.21  8.00 10.21 50       2.53
## 51 19.96 10.11  8.97 51       2.02
## 68 25.33 10.96  8.33 68       2.04
## 84  2.03  7.94  7.84 84      -2.03
## 91  5.51 10.74 10.25 91      -2.10
```

Assume we justified the use of $t_{.975,96}$ as the critical value, in which $\alpha=.05$. We should expect approximately $n * .05$ (in our case $100*.05=5$) cases larger than the critical value. In this particular case, even though 8 cases were identified, none of them seems unusual.

If the researcher detects an abnormality and further, if the researcher decides to remove the observation from the data, it should be done one observation at a time. The justification of removing a data point should be given clearly. A better alternative, on the other hand, may be to use an estimation method that is robust to outlying data points.

R program is convenient for investigating influential data points. Examine *?influence.measures* below for the simulated data set;

```
summary(influence.measures(model))
## Potentially influential observations of
##   lm(formula = yy ~ X1 + X2, data = tempdata) :
##
##      dfb.1_ dfb.X1 dfb.X2 dffit cov.r   cook.d hat
## 12   0.08  -0.02  -0.08  -0.10  1.12_*  0.00  0.08
## 33   0.09  -0.03  -0.07  -0.11  1.11_*  0.00  0.07
## 41  -0.01  -0.03   0.03  -0.04  1.10_*  0.00  0.06
## 42   0.05  -0.12   0.07   0.13  1.11_*  0.01  0.07
## 50   0.20  -0.40   0.21   0.47  0.88_*  0.07  0.03
## 64  -0.03   0.03   0.00   0.04  1.10_*  0.00  0.06
## 100  0.01   0.13  -0.15  -0.18  1.10_*  0.01  0.07
```

This output reports 5 different measures.

In this example, cases 12, 33, 41, 42, 50, 64 and 100 are reported to be *potentially* influential. As they highlighted by an asterisk, they labeled as potential using the covariance ratio criteria (cov.r). This value reports the impact of an observation on the sampling variances of the regression coefficients. Values larger than $1 + (3p'/n)$ and lower than $1 - (3p'/n)$ are labeled as influential, in our case, $n=100$ and $p'=3$, hence the cut offs are 1.09 and .91.

The Dfb (DFBETAS) for each predictor reports how much the coefficient for the predictor changes when the case is removed. It is the difference between the two coefficients divided by an estimate of the standard error of the new coefficient and therefore is on the scale of a t statistic. R places an asterisk if the value is larger than $2/\sqrt{n}$. For this specific illustration the cut off value is $2/\sqrt{(100)} = .2$.

The dffit reports the change in the predicted value for the i^{th} case when the i^{th} case is removed from the data. The criterion for identifying potentially influential data points is $2 * \sqrt{\frac{p'}{n}}$.

Cook's distance (cook.d) measures the influence of a particular case on all of the estimated coefficients and values larger than $F_{5,p',n-p'}$ are highlighted. Cook's distance also measures influence of omitting a particular case of the predicted values for all of the remaining cases.

Leverage Values (Hat Diag) measure the distance of an observation compared to other independent variables. Values larger than $2p/n$ are considered to identify potentially influential data points.

It is researcher's responsibility to examine any potentially influential data points.

11.1.4 d) *Equal variance assumption*

The standard errors of the coefficients are calculated as the square roots of the diagonal elements of $\hat{\sigma}^2(X'X)^{-1}$, where $\hat{\sigma}^2$ is the variance of the residuals. Examine the code below given for the synthetic data set:


When using the OLS with an assumption of normally distributed Y variable, the distribution of β can be obtained. Examine the code below given for the synthetic-data set;

```
#Residuals
s2 <- (t(residuals) %*% residuals)/(nrow(Y)-nrow(betahat))
Var_betahat <- s2[1,1]*solve(t(X)%*%X)
```


The equation $\sigma^2(X'X)^{-1}$ is valid under the assumption of homogeneity, that is, observations on the Y variable have a common variance controlling for the independent variables. In other words, every observation of Y has the same amount of information (Rawlings et al. (1998)). With this assumption, regression coefficients are selected to minimize $\sum_{i=1}^n (Y_i - \hat{Y}_i)^2$. In this expression equal weights are given to the residuals for every case.. If homogeneity is questionable the estimator can be modified to allow for unequal weights or replaced. Alternatively the Y variable can be transformed or the estimator of the standard error can be modified (see package ‘sandwich’ Lumley and Zeileis (2015)). Otherwise, the standard error of $\hat{\beta}$ could be underestimated or overestimated. Underestimation results in Type I error rates that are larger than the alpha level used in hypothesis tests and confidence intervals and over estimation results in reduced statistical power. It is common practice to plot residuals against the predicted values to study heterogeneity. Examine the code below to simulate data with unequal variance and examine the studentized residuals:

```
#simulate data
set.seed(03032017)
library(mvtnorm)
sigma <- matrix(c(1,.7,.7,1), ncol=2)
xx <- rmvnorm(n=100, mean=c(1,1), sigma=sigma)
#heteroscedasticity function
hts=function(v1,v2){2+.5*v1+.5*v2}
yy=5+xx[,1]*5+xx[,2]*5+rnorm(100,0,hts(xx[,1],xx[,2]))
model=lm(yy~xx[,1]+xx[,2])
#summary(model)
errors=rstudent(model)
predicted=predict(model)

#Studentized Residuals vs Yhat
library(ggplot2)
plotdata=data.frame(errors,predicted)
ggplot(plotdata, aes(x = predicted, y = errors)) +
  geom_point() + geom_hline(yintercept=0)+ylab("Studentized residuals")+
  geom_segment(mapping=aes(xend = predicted, yend = 0)) +
  theme_bw()
```



The variance with smaller \hat{Y} values are smaller. Below is a graph for a regression model on a simulated data with equal variance.



11.1.5 e) Hypothesis testing

The F test is used within a multiple regression framework to test $H_0 : \beta_1 = \dots = \beta_p = 0$, a hypothesis stating that the p regression coefficients are all equal to zero in the population. The alternative hypothesis states that at least one coefficient is not zero. The null hypothesis can be tested using the statistic $MS_{\text{regression}}/MS_{\text{residual}}$. This statistic follows an F distribution with p and $n - p'$ degrees of freedom. As mentioned earlier, p is the number of predictors and p' is the number of coefficients ($p' = p$ if there is no intercept). Examine the code below given for the synthetic data, setting Type I error rate = .05;

```
# Model SS and Total SS calculated before
```

```
dfREG=2 # (p=2, predictors X1 and X2)
```

```
dfRES=9 # (n-p', 12-3)
```

```
MSreg=ModelSS/dfREG
```

```
MSres=(TotalSS-ModelSS)/dfRES
```

```
MSreg/MSres
```

```
##      [,1]
```

```
## [1,] 32.8
```

```
#critical F
```

```
qf(.95,dfREG,dfRES)
```

```
## [1] 4.26
```

```
1-pf(MSreg/MSres,dfREG,dfRES)
```

```
##      [,1]
```

```
## [1,] 7.39e-05
```

The t-test is used for investigating $H_0 : \beta_X = \beta_{hyp}$ vs $H_1 : \beta_X \neq \beta_{hyp}$. Most commonly $\beta_{hyp} = 0$

The statistic $(b_X - \beta_{hyp})/SE(b_X)$ follows a t-distribution with N-p' degrees of freedom. Examine the code below given for the synthetic-data;

```
# test if the coefficient for X2 is different than 0
Bhyp=0 #hypothesized value

# estimated coefficient for X2 (see betahat calculated before)
bx2=betahat[3]

# estimated SE for X2 (see var_betahat calculated before)
se_bx2=sqrt(Var_betahat[3,3])

#t statistic
(bx2-Bhyp)/se_bx2
## [1] 5.33

# t critic
qt(.975,9)
## [1] 2.26

#p value
2*(pt(-abs((bx2-Bhyp)/se_bx2),9))
## [1] 0.000478
```

11.1.6 f) Variable Selection

Broadly speaking there are two situations in which multiple regression is used to analyze data.

The first is illustrated by the following example. A social science researcher conducts an extensive literature review, identifies all independent variables relevant to the research questions, collects the data, estimates a model in which all independent variables are included and reports results for this model.

The second is illustrated by an example in which the researcher has data on a very large set of variables and does not know prior to analyzing the data which variables will be included in the final model that will be reported. This might happen because the researcher is working in a relatively new research area and collects data on a wide variety of variables or is conducting a secondary data analysis of a data set with a wide variety of predictors. In either case the researcher may want to begin by conducting variable selection that is using statistical results to select the best subset of many independent variables. There are several approaches to select the best subset of predictors. For example, stepwise regression, backward selection or forward selection is covered in many sources. However, in our experience, when applied to the same data set these three approaches are likely to give different answers.

A convenient approach with R is to study all possible regressions. For introductory purposes, examine the code below given for a simulated data set;

```
#simulate data
set.seed(02082017)
library(mvtnorm)
sigma=matrix(c(5.899559,4.277045,3.906341,
               4.277045,5.817412,3.654419,
               3.906341,3.654419,5.642258),ncol=3)
xx <- rmvnorm(n=200, mean=c(0,0,0), sigma=sigma)
yy=5+xx[,1]+xx[,2]*1.5+xx[,3]*2+rnorm(200,0,3)
simdata=data.frame(yy,xx,id=1:200)
```

```
library(leaps)
formula <- formula(paste("yy ~ ",
  paste(names(simdata[2:4]), collapse=" + ")))
allpossreg <- regsubsets(formula,nbest=3,data=simdata)
aprout <- summary(allpossreg)

#this functions reports more than R-squared and adjusted R-squared
#examine str(aprout)

APRtable=with(aprout,round(cbind(which,rsq,adjr2),3))
APRtable=data.frame(APRtable,check.rows = F,row.names = NULL)
APRtable$ppri=rowSums(APRtable[,1:4])
kable(APRtable)
```

X.Intercept.	X1	X2	X3	rsq	adjr2	ppri
1	0	1	0	0.753	0.751	2
1	0	0	1	0.696	0.695	2
1	1	0	0	0.630	0.629	2
1	0	1	1	0.871	0.870	3
1	1	0	1	0.811	0.809	3
1	1	1	0	0.808	0.806	3
1	1	1	1	0.890	0.888	4

This table reports that intercept and X_2 only model results in an R^2 value of .753. When all predictors included, the R^2 reaches to .890; however, excluding the X_1 from the full model reduced the R^2 only by .019. Below is a graphical depiction.

```
require(ggplot2)
ggplot(APRtable, aes(x=ppri-1, y=rsq)) +
  geom_point(shape=1,size=3)+
  scale_y_continuous(breaks = seq(0.5, 1, by = 0.05)) +
  scale_x_continuous(breaks = seq(0, 3, by = 1))+
  theme_bw()+labs(x = "R-squared")+
  theme(axis.text=element_text(size=15),
    axis.title=element_text(size=14,face="bold"))

ggplot(APRtable, aes(x=ppri-1, y=adjr2)) +
  geom_point(shape=1,size=3)+
  scale_y_continuous(breaks = seq(0.5, 1, by = 0.05)) +
  scale_x_continuous(breaks = seq(0, 3, by = 1))+
  theme_bw()+labs(x = "Adjusted R-squared")+
  theme(axis.text=element_text(size=15),
    axis.title=element_text(size=14,face="bold"))
```

11.1.7 g) Collinearity

Collinearity is the degree to which the predictors are correlated among themselves. The correlation between predictors is a concern in regression because the standard errors of the coefficients increase as collinearity increase and therefore collinearity hides the individual contribution of each predictor in the regression equation.

As an illustration suppose there are two independent variables with $r = .9$. You MIGHT have two types of problems: The regression coefficients become unstable (i.e. they would vary a great deal across different




Figure 11.1: All Possible Regressions

samples obtained from the same population).

You may obtain a statistically significant R^2 but not statistically significant regression coefficients.

Variance inflation factor (VIF) is helpful to detect collinearity in regard to a particular independent variables and can be applied in models two or more independent variables. The formula is $VIF_x = \frac{1}{1-R_x^2}$ where R_x^2 is the R^2 when the predictor is regressed on the remaining independent variables. Large VIF values are indicator of possible multicollinearity. Commonly pronounced cut off values are 4 and 10, however VIF values are indirectly affected by sample size and variance (Obrien (2007)). When large VIF values are detected, the researcher should examine the problem. It might be justifiable to (a) leave out one of the highly correlated predictor, (b) combine the two highly correlated variable. The decision should be made cautiously given that the possible solution might be more problematic than a large VIF value, see Obrien (2007). Examine the code below given for a simulated data set;

```
#check correlations among predictors
cor(simdata[,2:4])
##      X1      X2      X3
## X1  1.00  0.730  0.640
## X2  0.73  1.000  0.666
## X3  0.64  0.666  1.000

#the largest correlation is.73
#no multicollinearity expected

library(car)
vif(lm(yy~X1+X2+X3,data=simdata))
##      X1      X2      X3
## 2.36  2.50  1.98
# no problematic VIF values
```

11.1.8 h) Non-linearity

In the presence of a non-linear relation between the dependent variable and any given independent variable, ignoring non-linearity is simply a validity concern due to the omitted variable issue. Examining the residuals is helpful to detect non-linearity. Residuals should be plotted against predicted and independent variables. A common practice is to include higher order variables in the model, for example, if the plot indicates a non linear pattern for X_k against residuals, X_k^2 might be needed in the model. The type of the non-linearity ,

such as quadratic, cubic or quartic should be treated accordingly. Gelman and Hill (2007) , commenting on age variable when the age and dependent variable are not linearly associated, prefers treating the variable as a categorical predictor. Alternatively transformations of the dependent or independent variable may be employed.

11.1.9 i) Correlated errors and nonindependent errors

Errors should not be correlated or more broadly should be independent. When such dependency is not addressed, regression results are invalid. This topic ,however, is well beyond the scope of this introductory material. Correlated errors are likely to distort the standard errors for the beta coefficients. This is not desired. In social sciences, correlated errors might be present when measurements are repeated. Multilevel models and latent growth models has been developed to address appropriate modeling of repeated measure designs. Nesting of participants in subgroups is another common source of non-independent errors in social sciences. Multilevel models are one popular solution to model clustered data.

11.1.10 j) Centering and Scaling

Consider an example in which mother's age at the date of her child's birth (maternal age) is used to predict IQ at age 10. The intercept estimates average IQ for children whose mother's maternal age was zero and cannot be meaningfully interpreted. Centering maternal age around its mean results in an intercept which estimates average IQ for children whose mother's maternal age was at the mean of the sample and can be meaningfully interpreted. Or consider predicting absences from work from an anxiety measure. A score of zero is possible score on the anxiety measure, but does not occur in the sample. The intercept estimates average absences for employees whose anxiety level is outside the range of the data and therefore represents an extrapolation for the data. Centering around the mean for the sample solves this problem. Another approach would be to center around an anxiety score that is in the range of the data and considered high. Or consider a study of income and an index of health. Income is on a scale in which a change of 1 represents a change of 1 dollar in income. The regression coefficient is .001, which represents a trivial change in the index. Dividing X by 1000 so that a change of 1 represents a change of 1000 dollars in income results in a regression coefficient of 1, which is a small but not trivial change in the index may make the results easier to think about.

11.1.11 k) Standardized coefficients

A related topic to linear transformations is to use a z-score for the continuous predictors by subtracting the mean and dividing by the standard deviation. Depending on the nature of the variable, using the z scores might ease the communication between researchers. Here are interpretation examples; Raw scores: An increase in anxiety of 1 unit is predicted to correspond to an increase of 3 units in achievement, holding the remaining predictors constant. z-scores: An increase in motivation of 1 standard deviation is predicted to correspond to an increase of 0.25 standard deviations in achievement, holding the the remaining predictors constant.

11.1.12 l) Interactions

We covered the basic idea of interaction in our ANOVA section. Ignoring an interaction is an omitted variable problem because an interaction affects the interpretation of main effects. For example, suppose a researcher investigates the relationship between mathematics achievement at the end of the school year (Y), effort measured by voluntary homework completed and submitted during the year (X_1), and mathematics achievement at the end of the preceding year (X_2). Using the model $Y_i = \beta_0 + \beta_1 X_{i1} + \beta_2 X_{i2} + \epsilon_i$ assumes that the relationship between Y and X_1 does not depend on X_2 and will be misleading if the assumption is false. A common model used to investigate interactions is ;


$$Y_i = \beta_0 + \beta_1 X_{i1} + \beta_2 X_{i2} + \beta_3 X_{i1} X_{i2} + \epsilon_i \quad (11.2)$$

The slope of the relationship between Y and X_1 , for example, is $\beta_1 + \beta_3 X_{i2}$ indicating that the relationship depends on X_2 . Similarly the slope relationship between Y and X_2 is $\beta_2 + \beta_3 X_{i1}$. Consideration of $\beta_1 + \beta_3 X_{i2}$ shows that β_1 is the slope of the relationship between Y and X_1 when $X_2 = 0$ and therefore β_1 cannot be meaningfully interpreted if $X_2 = 0$ is not a meaningful score or is outside the range of the data. This problem can be addressed by centering X_1 and X_2 around their respective means. It should be noted that the model in Equation (11.2) assumes that the interaction can be accurately modeled by including $\beta_3 X_{i1} X_{i2}$ in the model. This assumes the relationship between Y and X_1 is linear when X_2 is controlled. Violations of assumption of the model in Equation (11.2) should be investigated. R can be helpful in interpreting interactions by 3-dimension graphs. Examine the code below given for a simulated data set to highlight the use of R package *visreg* (Breheny and Burchett (2016))

```
## manipulate simdata
## Yvar: dependent variable on no interaction
simdata$Yvar=3+simdata$X1*2+simdata$X2*3+rnorm(nrow(simdata),0,5)

library(visreg)

model=lm(Yvar~X1+X2+X1*X2,data=simdata)
visreg2d(model, "X1", "X2", plot.type="persp")
```




The surface is flat indicating no interaction.

```
## manipulate simdata
## Yvarint: dependent variable on a interaction
simdata$Yvarint=3+simdata$X1*1+simdata$X2*2+simdata$X1*simdata$X2*1.5+rnorm(nrow(simdata),0,5)
```

```
library(visreg)

model2=lm(Yvarint~X1+X2+X1*X2,data=simdata)
visreg2d(model2, "X1", "X2", plot.type="persp")
```



The surface is no longer flat in the presence of an interaction.

11.1.13 m) Estimators

To be added

11.1.14 n) Robust Regression

To be added

11.1.15 o) Sample size and statistical power

To be added

11.1.16 p) Reliability of variables

To be added

11.1.17 q) The nature of the variables

To be added

11.1.18 r) Multiple dependent variables

To be added

11.1.19 s) Missing variables

To be added

Chapter 12

Useful R codes

```
# Convert numeric to factor

temdata[,2:9] <- lapply(temdata[,2:9], as.factor)

# Convert factor to numeric
as.numeric.factor <- function(x) {as.numeric(levels(x))[x]}
temdata[,2:5] <- lapply(temdata[,2:5], as.numeric.factor)

# Have frequencies table for multiple columns

dems=apply(temdata[,5:11], 2, function(x){table(x,temdata$grp)})
library (plyr)
mydems <- ldply (mydems, data.frame)

# Aggregate variables by grp

uncagg=aggregate(. ~ grp, data = temdata, FUN=mean, na.rm=TRUE)

uncaggfaster=temdata[, lapply(.SD, mean,na.rm=T), by = grp]

# Find max in a table
which.max(x)

# Update R
if(!require(installr)) {
  install.packages("installr"); require(installr)} #load / install+load installr
updateR()

# Create dummy variable from a factor
head(temdata)
for(level in unique(temdata$zp)){
  temdata[paste("dummy", level, sep = "_")] <- ifelse(temdata$zp == level, 1, 0)
}
```

```

# Using semi colon to send multiple input
x=rnorm(10000,5,10)
mean(x);var(x);sqrt(var(x))

# Remove an object
y=rnorm(10)
rm(y)

# Empty the working space
rm(list=ls())

# Remove all but some
rm(list=setdiff(ls(),c("temdata", "temdata2")))

# Integer division
7%/%2

# Modulo = remainder
5%%2

# Define and print
(count=c(25,12,7,4,6,2,1,0,2))

# Read csv by clicking
data=read.csv(file.choose(),header=TRUE,)

#Combine more than 1 csv files
filenames <- list.files()
temdata=do.call("rbind", lapply(filenames, read.csv, header = F))
write.table(temdata, file ="temdata.binded.csv" , sep = ",",col.names = F, row.names = F)

#Multiple QQ plot
#split screen
layout(matrix(1:9, nc = 3))
sapply(names(temdata)[1:9], function(x) {
  qqnorm(temdata[[x]], main = x)
  qqline(temdata[[x]])
})

#Split for more plots
par(mfrow=c(3,3))

#Double for loop
x=matrix(1:15,3,5)
for(i in seq_len(nrow(x)))
{

```



```

for(j in seq_len(ncol(x)))
{
  print(x[i,j])
}
}

#While loop
count=0
while(count<10){
  print(count)
  count=count+1
}

#Missing data
convert -999s to NAs

read.csv("x.csv", na.strings="-999")
temdata[is.na(temdata)] <- 0

#convert NAs to -99s

vector[which(vector== NA)]= (-99)
temdata[is.na(temdata)]= (-99)

#if you are having trouble converting <NA> (but not NA)
temdata=read.csv("temdata.csv",stringsAsFactors=FALSE)

# add group mean

temdata2=merge(temdata, aggregate(X ~ grp, data = temdata, FUN=mean, na.rm=TRUE),
  by = "grp", suffixes = c("", ".mean"),all=T)

temdata2=merge(temdata, aggregate(cbind(X1 ,X2 ,X3 , X4) ~ grp, data = temdata, FUN=mean,
  by = "grp", suffixes = c("", ".mean"),all=T))

temdata2=merge(temdata,
  ddply(temdata, c("grp"), function(x) colMeans(x[c("X1" ,"X2","X3" , "X4")])),
  by = "grp", suffixes = c("", ".mean"),all=T)

#ifelse
y=c(1,2,3,4,5,5,5)
y2=ifelse(y==5,NA,y)
y2

```

```

temdata <- data.frame (ID=c(2,3,4,5), Hunger =c(415,452,550,318 ))

temdata$newcol<-ifelse(temdata[,2]>=300 & temdata[,2]<400,350,
                      ifelse(temdata[,2]>=400 &temdata[,2]<500,450,
                              ifelse(temdata[,2]>=500 & temdata[,2]<600,550,NA)))

#if
x=5
y=if(x>6){1}else{0}
y=if(x>6){1} else if(x==5) {99} else {0}

#sort a dataframe by the order of the elements in B
temdata[order(temdata$B),]

#sort the dataframe in reverse order
temdata[rev(order(temdata$B)),]

#create combinations
m=c(54,38,51,62,18,31,58,74,35,34)
f=c(41,18,19,39,44,18,58,21,38)

mean(m)
mean(f)

combn(m,8,FUN=mean)
combn(f,8)

min(combn(m,8,FUN=mean))
max(combn(f,8,mean))

#setting contrasts
options('contrasts')
options(contrasts=c('contr.sum','contr.poly'))
options(contrasts=c('contr.treatment','contr.poly'))

# delete if all NA
temdata=temdata[apply(temdata,1,function(x)any(!is.na(x))),]

# add group frequency
temdata=ddply(temdata, "grp", transform, cellsize = count(grp)[2])

#create new folder
dir.create("testdir")

#split data frame

```

```
library(datasets)
head(airquality)
splitdata=split(airquality,airquality$Month)
splitdata
str(splitdata)
splitdata[[2]]
```

```
x=list(a=1:5, b=rnorm(10))
x
lapply(x,mean)
```

output is always a list

```
x=1:4
lapply(x,runif)
lapply(x,runif,min=10, max=20)
```

```
x=list(a=matrix(1:4,2,2),b=matrix(1:6,3,2))
lapply(x,function(elt) elt[,1])
```

sapply

```
x=list(a=1:5, b=rnorm(10),c=runif(10))
x
lapply(x,mean)
sapply(x,mean)
```

#apply generally used for rows or columns

```
x=matrix(rnorm(200),20,10)
x
apply(x,2,mean)
apply(x,1,sum)
```

#tapply

```
x=c(1:10,rnorm(10),runif(10,3,5))
f=gl(3,10)
?gl
h=factor(rep(1:3,each=10))
tapply(x,f,mean)
tapply(x,h,mean)
tapply(x,h,mean,simplify=F)
tapply(x,h,range)
```

```

#missing data proportion percentage
propmiss <- function(temdata) lapply(temdata,function(x) data.frame(nmiss=sum(is.na(x)), n=length(x), p
propmiss(temdata)

#upper case
temdata$childid=toupper(temdata$childid)

# plot graph individual all variables

plotpdf="C:/Users/Desktop/work/multiplePLOTS.pdf"
pdf(file=plotpdf)
for (i in 7:55){
  muis=round(mean(temdata[,i],na.rm=T),3)
  sdis=round(sd(temdata[,i],na.rm=T),3)
  meansc=c("mean",muis)
  hist(temdata[,i],freq=F,main=names(temdata)[i],xlab=meansc)
  #lines(density(temdata[,i],na.rm=T))
  curve(dnorm(x, mean=muis, sd=sdis), add=TRUE)
  lines(density(temdata[,i],na.rm=T, adjust=2), lty="dotted", col="darkgreen", lwd=2)
  abline(v=muis,col="blue")
  abline(v=muis+3*sdis,col="red")
  abline(v=muis-3*sdis,col="red")
}

dev.off()

# read in upper directory
dd=read.csv("../temdata.csv")

```

12.1 More on the apaStyle package

Here is more details on the apaStyle package;

```

require(pasteecs)
require(apaStyle)
library(rJava)
#if this throws an error
Sys.setenv(JAVA_HOME='C:\\Program Files\\Java\\jre1.8.0_111') # for 64-bit version

#define a data set

apa.descriptives(data = temdataet[,1:5], variables = names(temdataet[,1:5]), report = "", title = "test

example <- data.frame(c("Column 1", "Column 2", "Column 3"), c(3.45, 5.21, 2.64), c(1.23, 1.06, 1.12) )
apa.table(data = example, level1.header = c("Variable", "M", "SD"))

example <- data.frame( c("Column 1", "Column 2", "Column 3"),

```

```
      c(3.45, 5.21, 2.64),  
      c(1.23, 1.06, 1.12),  
      c(8.22, 25.12, 30.27),  
      c("+", "**", "***") )  
  
apa.table( data = example, level1.header = c("", "Descriptives", "Inferential"),  
           level1.colspan = c(1, 2, 1),  
           level2.header = c("Variable", "M", "SD", "t-value", "*") )$table
```

12.2 A useful shiny application

Below is a Shiny app example (Figure 12.2) to calculate sample size for an analyses of covariance design;

```
knitr::include_app('https://burakaydin.shinyapps.io/ancovaN/', height = '800px')
```

ANCOVA sample size calculator

Bibliography

- Adler, D. and Murdoch, D. (2017). *rgl: 3D Visualization Using OpenGL*. R package version 0.97.0.
- Aho, K. (2016). *asbio: A Collection of Statistical Tools for Biologists*. R package version 1.3-4.
- Allaire, J., Cheng, J., Xie, Y., McPherson, J., Chang, W., Allen, J., Wickham, H., Atkins, A., and Hyndman, R. (2016). *rmarkdown: Dynamic Documents for R*. R package version 1.0.9014.
- Appelbaum, M. I. and Cramer, E. M. (1976). Balancing - analysis of variance by another name. *Journal of Educational Statistics*, 1(3):233–252.
- Bakeman, R. (2005). Recommended effect size statistics for repeated measures designs. *Behavior Research Methods*, 37(3):379–384.
- Bates, D., Mächler, M., Bolker, B., and Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67(1):1–48.
- Box, G. E. P. and Draper, N. R. (1987). *Empirical model-building and response surfaces*. Wiley, New York.
- Breheny, P. and Burchett, W. (2016). *visreg: Visualization of Regression Models*. R package version 2.3-0.
- Carlson, J. E. and Timm, N. H. (1974). Analysis of nonorthogonal fixed-effects designs. *Psychological Bulletin*, 81(9):563–570.
- Cohen, J. (1962). The statistical power of abnormal-social psychological research: A review. *The Journal of Abnormal and Social Psychology*, 65(3):145–153.
- Daunic, A. P., Smith, S. W., Garvan, C. W., Barber, B. R., Becker, M. K., Peters, C. D., Taylor, G. G., Van Loan, C. L., Li, W., and Naranjo, A. H. (2012). Reducing developmental risk for emotional/behavioral problems: A randomized controlled trial examining the tools for getting along curriculum. *Journal of School Psychology*, 50(2):149–166.
- de Vreeze, J. (2016). *apaStyle: Generate APA Tables for MS Word*. R package version 0.4.
- Field, A. P., Miles, J., and Field, Z. (2012). *Discovering statistics using R*. Sage, Thousand Oaks, Calif;London;.
- Gelman, A. and Hill, J. (2007). *Data analysis using regression and multilevel/hierarchical models*. Cambridge University Press, Cambridge;New York;.
- Hirshleifer, S., McKenzie, D., Almeida, R., and Ridao-Cano, C. (2016). The impact of vocational training for the unemployed: Experimental evidence from turkey. *The Economic Journal*, 126(597):2115–2146.
- Højsgaard, S. and Halekoh, U. (2016). *doBy: Groupwise Statistics, LSmeans, Linear Contrasts, Utilities*. R package version 4.5-15.
- Holm, S. (1979). A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics*, 6(2):65–70.

- Komsta, L. and Novomestky, F. (2015). *moments: Moments, cumulants, skewness, kurtosis and related tests*. R package version 0.14.
- Lakens, D. (2013). Calculating and reporting effect sizes to facilitate cumulative science: a practical primer for t-tests and anovas. *Frontiers in Psychology*, 4:863.
- Lawrence, M. A. (2016). *ez: Easy Analysis and Visualization of Factorial Experiments*. R package version 4.4-0.
- Lemon, J., Bolker, B., Oom, S., Klein, E., Rowlingson, B., Wickham, H., Tyagi, A., Eterradosi, O., Grothen-dieck, G., Toews, M., Kane, J., Turner, R., Witthoft, C., Stander, J., Petzoldt, T., Duursma, R., Biancotto, E., Levy, O., Dutang, C., Solymos, P., Engelmann, R., Hecker, M., Steinbeck, F., Borchers, H., Singmann, H., Toal, T., and Ogle, D. (2016). *plotrix: Various Plotting Functions*. R package version 3.6-3.
- Lumley, T. and Zeileis, A. (2015). *sandwich: Robust Covariance Matrix Estimators*. R package version 2.3-4.
- Mair, P. and Wilcox, R. (2016). *WRS2: A Collection of Robust Statistical Methods*. R package version 0.9-1.
- Muenchen, R. A. (2011). *R for SAS and SPSS users*. Springer, New York, 2nd edition.
- Myers, J. L., Well, A., Lorch, R. F., and Corporation, E. (2013). *Research design and statistical analysis*. Routledge, New York, 3rd edition.
- Obrien, R. M. (2007). A caution regarding rules of thumb for variance inflation factors. *Quality and Quantity*, 41(5).
- Olejnik, S. and Algina, J. (2003). Generalized eta and omega squared statistics: Measures of effect size for some common research designs. *Psychological Methods*, 8(4):434–447.
- Pearl, J. (2009). *Causality: models, reasoning, and inference*. Cambridge University Press, Cambridge; New York, 2nd edition.
- R Core Team (2016a). *foreign: Read Data Stored by Minitab, S, SAS, SPSS, Stata, Systat, Weka, dBase, ...*. R package version 0.8-67.
- R Core Team (2016b). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.
- Rawlings, J. O., Pantula, S. G., and Dickey, D. A. (1998). *Applied regression analysis: a research tool*. Springer, New York, 2nd edition.
- Revelle, W. (2016). *psych: Procedures for Psychological, Psychometric, and Personality Research*. R package version 1.6.9.
- RStudio Team (2016). *RStudio: Integrated Development Environment for R*. RStudio, Inc., Boston, MA.
- Sarkar, D. (2016). *lattice: Trellis Graphics for R*. R package version 0.20-34.
- Tippmann, S. (2015). Programming tools: adventures with r: a guide to the popular, free statistics and visualization software that gives scientists control of their own data analysis. *Nature*, (7532):109.
- Torchiano, M. (2016). *effsize: Efficient Effect Size Computation*. R package version 0.7.0.
- Verzani, J. (2014). *Using R for introductory statistics*. CRC Press Taylor and Francis Group, Boca Raton, second edition.
- Wickham, H. (2011). The split-apply-combine strategy for data analysis. *Journal of Statistical Software*, 40(1):1–29.
- Wickham, H. (2016). *tidyr: Easily Tidy Data with ‘spread()’ and ‘gather()’ Functions*. R package version 0.6.0.

- Wickham, H. and Chang, W. (2016). *ggplot2: Create Elegant Data Visualisations Using the Grammar of Graphics*. R package version 2.2.0.
- Wilcox, R. R. (2012). *Introduction to Robust Estimation and Hypothesis Testing*. Academic Press, US, 3rd;3; edition.
- Xie, Y. (2016). *bookdown: Authoring Books with R Markdown*. R package version 0.1.