

BEE 4750/5750

Homework 3 Solutions

2026-09-24

Overview

Load Environment

The following code loads the environment and makes sure all needed packages are installed. This should be at the start of most Julia scripts.

```
import Pkg
Pkg.activate(@__DIR__)
Pkg.instantiate()
```

```
using Plots
using LaTeXStrings
```

Problems (Total: 50 Points)

Problem 1 (15)

Problem 1.1 (4)

$$\frac{dC}{dt} = 5 - 0.25C$$

(a) Denoting the right-hand side $f(C) = 5 - 0.25C$, the forward Euler update is

$$C(t + \Delta t) = C(t) + \Delta t(5 - 0.25 C(t)).$$

(b) This is a linear system with a single rate constant, so its characteristic timescale is

$$1 / |-0.25| = 4$$

. Since $\Delta t = 1$ is four times smaller than the timescale, it is a reasonable starting choice, though it might not be small enough to account for discretization error.

(c) Assuming first-order convergence (as expected from the Taylor approximation underlying Forward Euler discretization), error scales linearly with Δt . So halving Δt from 2 to 1 should halve the error:

$$\text{error}(\Delta t = 1) \approx 0.6 \times \frac{1}{2} = 0.3.$$

Problem 1.2 (5)

$$\frac{dC}{dt} = \frac{Q}{V}(C_{\text{in}} - C) - kC$$

(a) With $Q/V = 50/200 = 0.25 \text{ yr}^{-1}$, the forward Euler update is

$$\begin{aligned}
C(t + \Delta t) &= C(t) + \Delta t \left[\frac{Q}{V} (C_{\text{in}} - C(t)) - kC(t) \right] \\
&= C(t) + \Delta t [0.25(2 - C(t)) - 0.3C(t)] \\
&= C(t) + \Delta t (0.5 - 0.55C(t)).
\end{aligned}$$

(b) $\frac{dC}{dt} = 0.5 - 0.55C$: the two loss terms combine into a single effective rate -0.55 yr^{-1} . The characteristic timescale is $1/|-0.55| \approx 1.82 \text{ yr}$. $\Delta t = 1 \text{ yr}$ could be a reasonable starting choice, though as with Problem 1.1, there could still be discretization error that would need to be reduced by refining Δt .

(c) Here Δt shrinks by a factor of 4 (from 1.6 to 0.4), so under first-order convergence the error should also shrink by a factor of 4:

$$\text{error } (\Delta t = 0.4) \approx 0.8 \times \frac{1}{4} = 0.2.$$

Problem 1.3 (6)

$$\frac{dx}{dt} = 3 - 0.5x^2, \quad x \geq 0$$

(a) Denoting $f(x) = 3 - 0.5x^2$, the forward Euler update is

$$x(t + \Delta t) = x(t) + \Delta t (3 - 0.5x(t)^2).$$

(b) The equilibrium solves $3 - 0.5(x^*)^2 = 0 \Rightarrow x^* = \sqrt{6} \approx 2.449$ (taking the positive root, per the problem's domain). $f'(x) = -x$, so $g = f'(x^*) = -\sqrt{6} \approx -2.449$. The local characteristic timescale there is $1/|\sqrt{6}| \approx 0.408$.

$\Delta t = 0.5$ is *larger* than this local timescale so this is not a reasonable starting value near this equilibrium.

(c) Here Δt shrinks by a factor of 3, so

$$\text{error } (\Delta t = 0.1) \approx 1.5 \times \frac{1}{3} = 0.5.$$

Problem 2 (35)

Problem 2.1 (5)

Denoting the right-hand sides $f(P, M) = L - s_{\text{out}}P - hP + \kappa M \frac{P^q}{m^q + P^q}$ and $g(P, M) = hP - bM - \kappa M \frac{P^q}{m^q + P^q}$, the coupled forward Euler update is

$$\begin{aligned}
P(t + \Delta t) &= P(t) + \Delta t \left[L - s_{\text{out}}P(t) - hP(t) + \kappa M(t) \frac{P(t)^q}{m^q + P(t)^q} \right] \\
M(t + \Delta t) &= M(t) + \Delta t \left[hP(t) - bM(t) - \kappa M(t) \frac{P(t)^q}{m^q + P(t)^q} \right]
\end{aligned}$$

i Note

Note: **both** updates must use the recycling term evaluated at the *current* $P(t)$ and $M(t)$. A common mistake is to update P first and then use the *new* $P(t + \Delta t)$ when updating M — this is not forward Euler (it's closer to a Gauss-Seidel-style update), which requires joint linearization (recall the multivariate Taylor expansion we used to derive the condition for higher-dimensional stability). As a result, it does not have the same error properties we derived in class. You won't lose points for this — we didn't talk about it in class, but it's good to be aware of.

Problem 2.2 (13)

Choosing T . The slow variable is the sediment pool M , whose timescale is $1/b = 1/0.02 = 50$ yr. $T = 300$ yr is six of those — comfortably long enough for M to fill.

Choosing the quantity of interest. $P(30)$, while the lake is still changing. The value at the *end* of the run would be useless here: by $t = 300$ the system has relaxed, and every stable step size returns the same number to six or more digits.

```
outflow_rate, sediment_transfer = 0.3, 0.3
burial_rate, recycling_coefficient = 0.02, 0.15
half_saturation, steepness = 25.0, 8.0

function recycling_flux(phosphorus, sediment)
    saturation = phosphorus^steepness /
        (half_saturation^steepness + phosphorus^steepness)
    recycling_coefficient * sediment * saturation
end

function simulate_lake(loading, horizon, step)
    steps = Int(round(horizon / step))
    trajectory = zeros(steps)
    phosphorus, sediment = 0.0, 0.0
    for i in 1:steps
        recycled = recycling_flux(phosphorus, sediment)
        change_in_phosphorus = loading - outflow_rate*phosphorus -
            sediment_transfer*phosphorus + recycled
        change_in_sediment = sediment_transfer*phosphorus -
            burial_rate*sediment - recycled
        phosphorus += step * change_in_phosphorus
        sediment += step * change_in_sediment
        trajectory[i] = phosphorus
    end
    trajectory
end

phosphorus_at_30(step) = simulate_lake(10.0, 300.0, step)[Int(30 / step)]

step_reference = 0.001
reference_value = phosphorus_at_30(step_reference)

step_sizes = [2.0, 1.0, 0.5, 0.25, 0.125, 0.0625]
values = [phosphorus_at_30(step) for step in step_sizes]
errors = [abs(value - reference_value) for value in values]
orders = [log2(errors[i] / errors[i+1]) for i in 1:length(errors)-1]
```

Reference solution at $\Delta t = 0.001$ yr gives $P(30) = 18.58479 \mu\text{g} / \text{L}$.

Δt [yr]	$P(30)$	error	empirical order
2.0000	18.6349	5.01e-02	—
1.0000	18.6092	2.44e-02	1.04

Δt [yr]	$P(30)$	error	empirical order
0.5000	18.5970	1.22e-02	1.00
0.2500	18.5909	6.09e-03	1.00
0.1250	18.5878	3.04e-03	1.00
0.0625	18.5863	1.51e-03	1.01




Figure 1: Error in $P(30)$ against step size. The slope is the order of convergence.

The error is first order, as expected for forward Euler — and the errors span a factor of 33 from coarsest to finest. $\Delta t = 0.125$ yr is used for the rest of this problem: the error there is $0.003 \mu\text{g} / \text{L}$, far below anything that would change the interpretation in Problem 2.3, and the run is still fast.

i The horizon and the step size are set by different processes

T was chosen based on the *slowest* process in the system, the sediment pool at $1/b = 50$ yr. Δt is constrained by the *fastest*, which is P itself at $1/(s_{\text{out}} + h) \approx 1.7$ yr. A step chosen by looking only at M would be wildly too large.

Steps well above the fast timescale do not merely lose accuracy — they can destabilise. At $\Delta t = 4$ yr this model blows up entirely. Note the transition is not a perfectly abrupt: $\Delta t = 3.5$ still returns a finite (but badly wrong) answer.

Problem 2.3 (10)

Simulating at $L = 10 \mu\text{g} / (\text{L} \cdot \text{yr})$ with the step size chosen in Problem 2.2 gives the trajectory below.




Figure 2: $P(t)$ for the two-stock lake model at $L = 10 \mu\text{g}/(\text{L}\cdot\text{yr})$.

From Problem 2.2, $P(30) = 18.6 \mu\text{g} / \text{L}$ — a modest, plausibly-stable oligotrophic reading, a little above the naive one-stock estimate but not alarmingly so. Once the simulation has fully run, P settles to a **much higher** equilibrium of about $29.0 \mu\text{g} / \text{L}$ — solidly eutrophic, well above $m = 25$.

Physically: at $t = 30$, the sediment pool M hasn't yet accumulated enough phosphorus for recycling to matter much, so the lake behaves close to a simple outflow-dominated system. As M slowly builds (over the sediment timescale, $1/b = 50 \text{ yr}$), the recycling term $\kappa M \cdot P^q / (m^q + P^q)$ becomes large enough to meaningfully add phosphorus back to the water column. This triggers a rapid transition — visible as a sharp rise and an overshoot, peaking around $P = 38.2 \mu\text{g} / \text{L}$ near $t = 45 \text{ yr}$ — before the system settles into its new, eutrophic equilibrium by roughly $t = 80\text{--}90 \text{ yr}$.

Problem 2.4 (7)

The naive one-stock model, $\frac{dP}{dt} = L - s_{\text{out}}P - hP$, has equilibrium

$$P_{\text{naive}}^* = \frac{L}{s_{\text{out}} + h} = \frac{10}{0.6} \approx 16.7 \mu\text{g} / \text{L} .$$

Since this is well below $m = 25 \mu\text{g} / \text{L}$, a manager using this simplified model (or, equivalently, only running a short simulation of the full model — recall $P \approx 18.6$ at $t = 30$ in Problem 2.3) would conclude this loading is **safely oligotrophic**.

But the true long-run behavior of the full two-stock model shows the lake actually settles into a **eutrophic** state ($P \approx 29 \mu\text{g} / \text{L}$) at this same loading, once the sediment pool has had time to fill. The naive model isn't just quantitatively a bit off — it gets the **qualitative outcome wrong**.

References

List any external references consulted, including classmates.