

# 1 Timeline of HIV in the US

1. **June 1981:** MMWR report
2. **September 1982:** AIDS defined

“a disease, at least moderately predictive of a defect in cell-mediated immunity, occurring in a person with no known cause for diminished resistance to that disease”
3. **December 1982:** Case of AIDS after blood transfusion reported

1985: FDA instituted lifetime ban on blood donation for MSM
4. **March 1983:** CDC addresses transmission risks

*‘AIDS may be caused by an infectious agent that is transmitted sexually or through exposure to blood products’*
5. **January 1983:** ‘Ward86’ opens at UCSF as the first outpatient AIDS clinic
6. **June 1983:** AIDS activism begins
7. **May 1983:** HIV is discovered as the virus which causes AIDS (Nobel Prize)

**January 1985:** Conclusively determined that AIDS caused by HIV

1985 First public mention of AIDS by Reagan
8. **March 1987:** ACT UP (AIDS Coalition to Unleash Power) forms grassroots political group to end AIDS pandemic
9. **March 1987:** FDA approves first antiretroviral drug (Zidovudine = ‘AZT’)
10. **1992:** AIDS becomes #1 cause of death for US men aged 25-44

# 2 HIV Viral Lifecycle

## 2.1 Reverse Transcription and Integration

1. gp120 on the virus attaches to CD4 receptor on T cells
2. gp41 mediates membrane fusion: viral capsid and RNA genome are delivered
3. RNA genome is **reverse transcribed** into DNA and the DNA is then **integrated** into the T cell genome by integrase
4. The ‘Provirus’ is the integrated DNA copy of the RNA genome

Integrated virus can remain silent for a long time (**latent**)

Retroviruses stably integrate DNA into host cell chromosomes

1. Persistence: Infection is not easily cleared
2. Latency: Integrated, but silent proviruses are not seen by the immune system
3. Normal cell division increases number of infected cells (clonal expansion)

→ Reason why HIV is chronic and has no cure

## 2.2 Mutation Rate

Our DNA polymerases that replicate our DNA: 1 in 1 billion base pairs mutate per cell division (3 mutations per cell division)

HIV reverse transcriptase: 1 in 10,000 base pairs mutate per replication (HIV RNA genome is around 10,000 base pairs)

Approximately one mutation every time HIV replicates

Low estimate of release into blood of virions is 10 billion per day

On average every possible mutation and every possible position in the HIV genome is predicted to occur each day

Since the start of the HIV-1 group M pandemic (early 1900s), HIV sequences have massively diverged




Figure 1: HIV mutation in a single individual is equivalent to that of an entire flu season

## 2.3 Assembly and Release

1. The integrated provirus directs the creation of new RNA viral genomes (by transcription) and new viral proteins (by translation)
2. HIV Protease cuts up a precursor protein (e.g. Pol) into smaller functional proteins (reverse transcriptase, protease, integrase)
3. New viral particles are then assembled and released from the cell to infect new cells ('budding')




Figure 2: Protease in the HIV life Cycle (Maturation)




Figure 3: Mature and infectious virions

### 3 HIV Pathogenesis

HIV infection has 2 stages: Acute and Chronic

Acute infection:

1. flu-like symptoms (50-90% of people)
2. 2-4 weeks after infection, lasting a few days or weeks

Chronic infection:

1. Severe immunodeficiency → AIDS ]item CD4 T cells are lost by direct killing by virus and immune mechanisms

AIDS is a disease of helper CD4 T cells

Allows recruitment and activation of immune cells, helps CD8 T cells, and helps B cells make antibodies

The immune system is damaged during all infections, but in untreated people, the virus is chronic and the disease is progressive

Destruction of the immune system leads to opportunistic infections

### 3.1 Acute Phase

Initial peak of viremia (virus in plasma) with flu-like symptoms

### 3.2 Chronic Phase

Viremia is controlled and reaches a low level

Number of CD4+ T cells slowly declines as virus / immune responses kill them

CD4+ T cells reach low enough level that the immune system begins to fail

Virus replicates out of control → Further loss of immune function → opportunistic infection




Figure 4: HIV Pathogenesis