

## European Organisation for Astronomical Research in the Southern Hemisphere

OBSERVING PROGRAMMES OFFICE • Karl-Schwarzschild-Straße 2 • D-85748 Garching bei München • e-mail: opo@eso.org • Tel.: +49 89 320 06473

## APPLICATION FOR OBSERVING TIME

PERIOD: 96Z

Category:

Important Notice:

Title

DDT

X-0

By submitting this proposal, the PI takes full responsibility for the content of the proposal, in particular with regard to the names of CoIs and the agreement to act according to the ESO policy and regulations, should observing time be granted.

First exopianet image							
2. Abstract / Total Time Requested Total Amount of Time:							
The TW Hydrae association is a group of approximately thirty very young stars located 50 parsecs from Earth that are 5-10 million years old. The best studied members of this stellar association are TW Hydrae (nearest known accreting T Tauri star to the Earth), HR 4796 (an A-type star with resolved dusty debris disk; the most massive known group member), HD 98800 (a quadruple star system with debris disk), and 2M1207 (accreting brown dwarf with remarkable planetary-mass companion 2M1207b). This 25 MJup brown dwarf(2M1207), located 70 pc from Earth, has been recently identified as a member of the TW Hydrae Association (age 8 Myr). Astrometry and photometry in the H, Ks and L' bands and upper limit in J-band reveal a very red object(2M1207b) at a close separation of 780 mas (55 AU) with a spectral type L5-L9.5. Different evolutionary models predict an object within the planetary regime with a mass of M=5±2 MJup and an effective temperature							
3. Run Period Instrument Time A 96 NACO 1h	Month any	Moon Seeing Sky Mo	ode Type				
<ul><li>4. Number of nights/hours</li><li>a) already awarded to this project:</li></ul>	Telescope(s)	Amount of time					
b) still required to complete this project:	VLT VLT	4h 3h					
5. Special remarks:  Take advantage of this box to provide any	special remark using up	to three lines					
6. Principal Investigator: beevageeva							
6a. Co-investigators:							
E. F. Van Dishoeck 1096 E. Dartois 1150							

7.	Description of the proposed programme
	A – Scientific Rationale: We already have a few photometric and astrometric results from Two Micron All Sky Survey (2MASS) in bands: J , H , Ks (The transformations between the two filters Ks and K were found smaller than the measuring errors.) and from Keck telescope in L' band The parent dwarf appears to be of spectral type M8 with magnitudes: $13.00(J)$ , $12.39(H)$ , $11.95(K)$ , $11.38(L')$ and the giant planet candidate: spectral type L5 - L9.5 with magnitudes: $\geq 18.5(J)$ , $18.09(K)$ , $16.93(K)$ , $15.28(L')$
	B – Immediate Objective: It is not established yet the nature of the companion of the brown dwarf that we believe is a giant planet. We need to repeat the photometric measurements using adaptive optics infrared wavefront sensing to acquire sharp images and to obtain spectroscopic results

## 7. Description of the proposed programme and attachments




Fig. 1: image obtained from 2 mass centered on the companion of the parent dwarf: RA: 12h:07m:34.01s, dec: -39d:32m:58.9s, 30 arcsec

8.	Justification of requested observing time and observing conditions
	Lunar Phase Justification: The object we want to observe is very faint
	Time Justification: (including seeing overhead) We need images in band J, H, Ks and L' and specter of the object with upper limit estimated magnitude $\approx 20$ in band J; ETC gave us NDIT = 9 (J), NDIT = 1 (H, Ks and L') and NDIT = 120(specter) for DIT = 20s making a total time 2640s (we estimated to 1 h with overheads)
8a	VLT is the world's most advanced optical instrument, consisting of four Unit Telescopes with main mirrors of 8.2m diameter and four movable 1.8m diameter Auxiliary Telescopes. We want to use UT1 beacuse we want to use intrument NACO.  NACO = Nasmyth Adaptive Optics System (NAOS) + Near-Infrared Imager and Spectrograph (CONICA)  NACO has been contributing to major discoveries in different fields of astronomy: exoplanets, the Galactic centre of the Milky Way, young stellar objects, Solar System bodies, etc.
8b	The only option was for cycle 96z(DDT cycle)
8c	Calibration Request: Standard Calibration

9. Report on the use of ESO facilities during the last 2 years  September 2003, VLT(UT3) telescope with instrument ISAAC at Nasmyth A focus: Spectroscopic surveys of large numbers of young low-mass stars which are still deeply embedded in their parental clouds. Our infrared spectra show a rich variety of features due to ices and gas-phase molecules, each of which trace different aspects of the physical and chemical state of the objects.  We already published the data: Origin and Evolution of Ices in star-forming regions a VLT-ISAAC 35 μm spectroscopic survey
9a. ESO Archive - Are the data requested by this proposal in the ESO Archive (http://archive.eso.org)? If so, explain the need for new data.  No
9b. GTO/Public Survey Duplications:  There are some results from 2 mass survey that our research is based on. We want to repeat the observations in J, H, Ks and L' band using adaptative optics and a bigger integration time and to obtain its specter in order to determine the nature of the companion of the brown dwarf that appears faintly in the images from 2 mass.
10. Applicant's publications related to the subject of this application during the last 2 years E. F. Van Dishoeck , E. Dartois, K. M.Pontoppidan, W. F. Thi, L. D'Hendecourt, A. C. A. Boogert, H. J.Fraser, W. A. Schutte, A. G. G. M Tielen: Origin and Evolution of Ices in star-forming regions a VLT-ISAAC 35 μm spectroscopic survey

Run	f targets propose Target/Field	$\alpha$ (J2000)	$\delta$ (J2000)	ToT M	lag. D	iam.	Additional info	Reference star
A	2M1207b	12:07:34.01	-39:32:58.9	1 18	3			2M1207
star for		he parent brov	vn dwarf with	coordina	ates: R	A: 12	h:07m:33.47s	nat. We use as references and dec: -39d:32m:54.0 stars.

This proposal involves time-critical observations, or observations to be performed at specific time intervals.

13. Instrument configuration						
Period	Instrument	Run ID	Parameter	Value or list		
96	NACO	A	IMG filters: J, H, Ks	N90C10 dichroic, FowlerN-samp(HighSensitivity) detector mode, detector FOV: 27.6x27.6 arcsec		
96	NACO	A	IMG filter: L'	VIS dichroic, Un- corr(HighWellDepth) detector mode, detector FOV: 27.6x27.6 arcsec		
96	NACO	A	Spectroscopy	VIS dichroic, S54 4 SJ spectroscopic mode(camera S54), FowlerNsamp(HighSensitivity) detector mode, 86 mas slit width		