

# Vectors

MATH 3512, BCIT

Matrix Methods and Statistics for Geomatics

October 8, 2018

The **projection**  $u_H$  of a vector  $u$  onto a hyperplane  $H$  is the vector in the hyperplane that is “most similar” to  $u$ . The formal definition for  $u_H$  requires that

- 1  $u$  is in  $H$
- 2  $(u - u_H)$  is orthogonal to all basis vectors of  $H$

**Example 1: Finding a Projection.** Let  $H$  be the line spanned by  $\vec{v} = (-1, 1)^\top$  in  $\mathbb{R}^2$ . What is the projection  $\vec{w}$  of  $\vec{u} = (3, -2)^\top$ ?



Let  $\vec{w} = (w_1, w_2)^\top$ . Then (1)  $\vec{u} - \vec{w}$  is orthogonal to  $\vec{v}$  and (2)  $\vec{w} = \alpha \vec{v}$  for some  $\alpha \in \mathbb{R}$ .

$$\begin{aligned} w_1 - w_2 &= 5 \\ w_1 + w_2 &= 0 \end{aligned} \tag{1}$$

Cramer's rule tells us that  $\vec{w} = (2.5, -2.5)^\top$ .

Let  $u = (u_1, \dots, u_n)^T$  be a vector and  $H$  be a  $k$ -dimensional hyperplane in the vector space  $\mathbb{R}^n$ . Let  $x_1, \dots, x_k$  be a basis for  $H$ . Then it is true for all vectors  $v$  in the hyperplane that

$$\|u - v\| \geq \|u - u_H\| \quad (2)$$

Proof: use the theorem of Pythagoras for

$$\|u - v\|^2 = \|u - u_H\|^2 + \|u_H - v\|^2 \geq \|u - u_H\|^2 \quad (3)$$

The claim follows. It illustrates what I mean when I say that  $u_H$  is the vector in  $H$  that is most similar to  $u$ .

**Example 2: Finding Another Projection.** What is the projection of  $\vec{u} = (5, 2, 10)^\top$  onto the plane  $T$  characterized by  $2x + y + 3z = 0$ ?

First we find two linearly independent vectors in  $H$  to form a basis of  $H$ , for example  $\vec{v}_1 = (1, 1, -1)^\top$  and  $\vec{v}_2 = (0, -3, 1)^\top$ . The conditions

- ①  $u_H \in T$
- ②  $(u - u_H) \perp v_1$
- ③  $(u - u_H) \perp v_2$

give us the system of linear equations

$$\begin{bmatrix} 2 & 1 & 3 \\ -1 & 1 & 1 \\ 0 & 3 & -1 \end{bmatrix} \cdot \begin{bmatrix} \hat{x} \\ \hat{y} \\ \hat{z} \end{bmatrix} = \begin{bmatrix} 0 \\ 3 \\ -4 \end{bmatrix} \quad (4)$$

for which the solution is  $u_H = (\hat{x}, \hat{y}, \hat{z})^\top = (-1, -1, 1)^\top$ .

# Least Squares Method

Let there be two linearly independent vectors  $u$  and  $v$  in  $\mathbb{R}^n$ . Then the formula for the projection  $u_v$  of  $u$  onto the line spanned by  $v$  is

$$u_v = \left( \frac{u \cdot v}{v \cdot v} \right) \cdot v \quad (5)$$

To verify the formula, note that  $u_v = av$  for some real number  $a$ . Therefore

$$(u - av) \perp v \quad (6)$$

Isolate  $a$  in the equation  $(u - av) \cdot v = 0$  to yield the formula.

# Least Squares Method

Formula (5) only works when the hyperplane is a line. You can scale up the idea in terms of dimensions by the following theorem.

## Formula for Projection Onto Plane with Orthogonal Basis

Let  $\{u, v\}$  be an orthogonal basis for  $H$ . Then the projection of  $w$  onto  $H$  is the sum of  $w_u$  and  $w_v$ , the projections of  $w$  onto the lines spanned by  $u$  and  $v$ , respectively.

Proof: check the following

- 1  $(w_u + w_v) \in H$  (trivial)
- 2  $(w - (w_u + w_v)) \perp u$  (use the fact that  $u \perp v$ )
- 3  $(w - (w_u + w_v)) \perp v$  (same idea)

# Least Squares Method


Consider the following table of measurements for the length of shoe prints and the height of the person wearing the shoes.

Shoe Print (cm)	Height (cm)
29.7	175.3
29.9	177.8
31.4	185.4
31.8	175.3
27.6	172.7

In the statistics portion of this course, we will learn whether the paired data provide evidence of a linear relationship. In the linear algebra portion, we will learn how to find the line which is closest to the data points in the least squares sense.



# Least Squares Method



# Least Squares Method

## Least Squares Method

If  $L$  is a given line, the **error** for each data point is the vertical distance from that point to the line. The **squared error** is the sum of the squares of the errors. The line that best fits the data in the least squares sense is the line that minimizes the squared error.

You can find the **regression line** using calculus optimization. However, there is also an elegant method using linear algebra.

# Least Squares Method

Let  $L$  be a line with slope  $m$  and  $y$ -intercept  $b$ . Let  $(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)$  be a set of paired data. Then the following equations hold:

$$\begin{aligned} y_1 &= mx_1 + b + \epsilon_1 \\ y_2 &= mx_2 + b + \epsilon_2 \\ &\vdots \\ y_n &= mx_n + b + \epsilon_n \end{aligned} \tag{7}$$

where the  $\epsilon_i$  are the errors ( $i = 1, \dots, n$ ). This system is equivalent to the following vector equation,

$$Y = AV + E \tag{8}$$

where  $Y, A, V, E$  are defined on the next slide.

# Least Squares Method

$$Y = \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix}, A = \begin{bmatrix} x_1 & 1 \\ x_2 & 1 \\ \vdots & \vdots \\ x_n & 1 \end{bmatrix}, V = \begin{bmatrix} m \\ b \end{bmatrix}, E = \begin{bmatrix} \epsilon_1 \\ \epsilon_2 \\ \vdots \\ \epsilon_n \end{bmatrix}$$

$E$  is called the error vector. According to (8), it is

$$E = Y - AV \quad (9)$$

We are trying to choose  $m, b$  so that

$$\|E\|^2 = \|Y - AV\|^2 \quad (10)$$

is minimal.

Let

$$X = \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} \text{ and } B = \begin{bmatrix} 1 \\ \vdots \\ 1 \end{bmatrix} \quad (11)$$

Then  $AV = mX + bB$ . The set  $S = \{AV | m, b \in \mathbb{R}\}$  is a plane in  $n$ -dimensional space. The ordered pair  $(m, b)$  that minimizes the squared error corresponds to the projection of  $Y$  onto  $S$ .

# Least Squares Method

Let  $C$  be  $B - B_X$ , where  $B_X$  is the projection of  $B$  onto the line spanned by  $X$ . Then

$$C = \frac{B \cdot X}{X \cdot X} \cdot X \quad (12)$$

Note that  $X \perp B_X$ .

# End of Lesson

Next Lesson: Eigenvalues and Eigenvectors