
Part 5: Moment Generating Functions

Suppose random variable X has some univariate distribution, discrete or continuous.

We define $E(X)$, $E(X^2)$, $E(X^3)$, $\dots$ to be the **moments of the distribution of X** . As we've already seen, $E(X^k)$ can be finite, infinite, or not exist.

The **moment generating function (MGF)** of X is

$$M_X(t) = E(e^{tX}).$$

Note that it is a function of t .

The MGF is simply the expected value of some function of X , and hence can be found using the Rule for the Lazy Statistician.

Exercise: Derive the MGF for the Poisson(λ) distribution.

Exercise: Derive the MGF for the Exponential(λ) distribution.

Name	Density/Mass Function	MGF	Mean	Variance
Normal	$\frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right),$ $-\infty < x < \infty, \sigma > 0$	$\exp\left(\mu t + \frac{\sigma^2 t^2}{2}\right)$	μ	σ^2
Log-Normal	$\frac{1}{x\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(\log(x)-\mu)^2}{2\sigma^2}\right),$ $0 < x < \infty, \sigma > 0$	—	$\exp\left(\mu + \frac{\sigma^2}{2}\right)$	$e^{2(\mu+\sigma^2)} - e^{2\mu+\sigma^2}$
Gamma	$\frac{\beta^\alpha}{\Gamma(\alpha)} x^{\alpha-1} e^{-\beta x},$ $0 < x < \infty, \alpha, \beta > 0$	$\left(\frac{\beta}{1-t}\right)^\alpha, t < \beta$	$\frac{\alpha}{\beta}$	$\frac{\alpha}{\beta^2}$
Exponential	$\beta e^{-\beta x},$ $0 < x < \infty, \beta > 0$	$\frac{\beta}{\beta-t}, t < \beta$	$\frac{1}{\beta}$	$\frac{1}{\beta^2}$
Beta	$\frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} x^{\alpha-1} (1-x)^{\beta-1},$ $0 < x < 1, \alpha, \beta > 0$	$1 + \sum_{k=1}^{\infty} \left(\prod_{r=0}^{k-1} \frac{\alpha+r}{\alpha+\beta+r} \right) \frac{t^k}{k!}$	$\frac{\alpha}{\alpha+\beta}$	$\frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$
Uniform (continuous)	$\frac{1}{b-a},$ $a \leq x \leq b$	$\frac{e^{bt}-e^{at}}{t(b-a)}, t \neq 0$ $1, t = 0$	$\frac{b+a}{2}$	$\frac{(b-a)^2}{12}$
Bernoulli	$p^x (1-p)^{1-x},$ $x = 0, 1, 0 \leq p \leq 1$	$pe^t + (1-p)$	p	$p(1-p)$
Binomial	$\binom{n}{x} p^x (1-p)^{n-x},$ $x = 0, 1, \dots, n, 0 \leq p \leq 1$	$[pe^t + (1-p)]^n$	np	$np(1-p)$
Negative Binomial	$\binom{r+x-1}{x} p^r (1-p)^x,$ $x = 0, 1, \dots, r \in \{1, 2, \dots\}, 0 \leq p \leq 1$	$\left(\frac{p}{1-(1-p)e^t} \right)^r, t < \log(1/(1-p))$	$\frac{r(1-p)}{p}$	$\frac{r(1-p)}{p^2}$
Geometric	$(1-p)^x p,$ $x = 0, 1, \dots, 0 \leq p \leq 1$	$\frac{p}{1-(1-p)e^t}, t < \log(1/(1-p))$	$\frac{1-p}{p}$	$\frac{(1-p)}{p^2}$
Poisson	$\frac{e^{-\lambda} \lambda^x}{x!},$ $x = 0, 1, \dots, \lambda > 0$	$e^{\lambda(e^t-1)}$	λ	λ
Hypergeometric	$\frac{\binom{A}{x} \binom{B}{n-x}}{\binom{A+B}{n}},$ $x = 0, 1, \dots, n$	—	$\frac{nA}{A+B}$	$\frac{nAB(A+B-n)}{(A+B)^2(A+B+1)}$

Using the MGF

1. Calculating Moments

Successive derivatives of the MGF, evaluated at $t = 0$, return the moments of the random variable:

$$E(X^k) = \frac{\partial^k}{\partial t^k} M_X(t) \Big|_{t=0}$$

Exercise: Why does this work? Sketch it out in the continuous case.

1 **Exercise:** Find the moments of the $\text{Poisson}(\lambda)$ distribution.

2 _____

3 _____

4 _____

5 _____

6 _____

7 _____

8 _____

9 _____

10 _____

11 _____

1 **2. Uniqueness of the MGF**

2 **The uniqueness result is the following:** If $M_X(t) = M_Y(t)$ for all t ,

3 then X and Y have the same distribution.

4 It is interesting to note that two random variables can have matching

5 moments, i.e., $E(X^k) = E(Y^k)$ for $k = 1, 2, \dots$, but X and Y have

6 different distributions.

7 **Exercise:** Suppose that X has the $\text{Exponential}(\lambda)$ distribution, and

8 $k > 0$ is some constant. Does kY have an Exponential distribution?

9 _____

10 _____

11 _____

12 _____

13 _____

14 _____

15 _____

16 _____

17 _____

18 _____

19 _____

1 **Exercise:** Suppose that the number of events during an interval of
2 length one time unit is a random variable X which has the Poisson(λ)
3 distribution. For each of these events, there is probability p that it is
4 of “Type I.” The determination of whether or not an event is of “Type
5 I” is independent from event to event. Let Y denote the number
6 of “Type I” events during the time interval. Show that Y has the
7 Poisson distribution.

1 3. Sums of Independent Random Variables

2 Suppose that $X_1, X_2, \dots, X_n$ are independent random variables, and
3 let $Y = \sum_i X_i$. Then,

$$M_Y(t) = \prod_{i=1}^n M_{X_i}(t)$$

5 for all t such that all $M_{X_i}(t)$ is finite.

6 **Exercise:** Prove the above result.

1 **Exercise:** Show that the sum of independent Poisson random vari-
2 ables also has the Poisson distribution.

3 _____
4 _____
5 _____
6 _____
7 _____
8 _____

9 **Exercise:** Under what condition(s) does the sum of Gamma random
10 variables still have a Gamma distribution?

11 _____
12 _____
13 _____
14 _____
15 _____
16 _____
17 _____
18 _____
19 _____
20 _____
21 _____

1 **Exercise:** Prove the following properties of the Gamma distribution.

- 2 1. If $X_1, X_2, \dots, X_n$ are iid $\text{Exponential}(\lambda)$ random variables, the
3 $\sum_i X_i$ is $\text{Gamma}(n, \lambda)$.
4 2. If X has the $\text{Gamma}(\alpha, \beta)$ distribution and $k > 0$, then kX is
5 $\text{Gamma}(\alpha, \beta/k)$.

6 _____
7 _____
8 _____
9 _____
10 _____
11 _____
12 _____
13 _____
14 _____
15 _____
16 _____
17 _____
18 _____
19 _____
20 _____

1 4. Establishing Convergence in Distribution

2 In some situations it will be important to formalize the notion of “the
3 distribution of $X_1, X_2, \dots$ converges to the distribution of X .”

4 In such cases, we would write $X_n \xrightarrow{D} X$, and say that “ X_n converges
5 in distribution to X .”

6 The formal definition is as follows: X_n **converges in distribution** to
7 X if $F_{X_n}(x) \rightarrow F_X(x)$ as $n \rightarrow \infty$ at all x where F_X is continuous.

8 In practical terms, the value of knowing $X_n \xrightarrow{D} X$ is that (for large n),
9 the distribution of X can be taken as an approximation to the distri-
10 bution of X_n . The **central limit theorem** is an important example.

11 **The useful result:** $X_n \xrightarrow{D} X$ if $M_{X_n}(t) \rightarrow M_X(t)$ as $n \rightarrow \infty$ for all
12 $-\epsilon < t < \epsilon$ for some $\epsilon > 0$.

1 **Exercise:** Suppose that X equals zero with probability one and X_n
2 equals $1/n$ with probability one. Does $X_n \xrightarrow{D} X$?

3 _____

4 _____

5 _____

6 _____

7 _____

8 _____

9 _____

10 _____

11 _____

12 _____

13 _____

14 _____

15 _____

16 _____

17 _____

18 _____

19 _____

20 _____

Exercise: Suppose that X_n has the Binomial($n, \lambda/n$) distribution for $\lambda > 0$. Let X have the Poisson(λ) distribution. Show that $X_n \xrightarrow{D} X$.

The Normal Distribution and the MGF

If X has the normal(μ, σ^2) distribution (with pdf given previously in the notes), then

$$M_X(t) = e^{t\mu + t^2\sigma^2/2}, \quad -\infty < t < \infty$$

The following properties of the normal(μ, σ^2) distribution are easy to prove once we know the MGF and its properties:

1. $E(X) = \mu$

2. $E(X^2) = \mu^2 + \sigma^2$, and hence $V(X) = \sigma^2$

3. If $Y = aX + b$, with $a \neq 0$, then $Y \sim N(a\mu + b, a^2\sigma^2)$.

4. The linear combination of independent normal random variables also has the normal distribution. In particular,

- (a) If $X_1, X_2, \dots, X_n$ are i.i.d. $N(\mu, \sigma^2)$, then $\sum_i X_i \sim N(n\mu, n\sigma^2)$.

- (b) If $X_1, X_2, \dots, X_n$ are i.i.d. $N(\mu, \sigma^2)$, then $\bar{X} \sim N(\mu, \sigma^2/n)$.

5. **The Central Limit Theorem:** Suppose $X_1, X_2, \dots$ are i.i.d. and $E(X_i) = \mu$ and $V(X_i) = \sigma^2$ both exist and are finite, then

$$\frac{\bar{X}_n - \mu}{\sigma/\sqrt{n}} \xrightarrow{D} Z$$

where $\bar{X}_n = \sum_{i=1}^n X_i/n$ and $Z \sim N(0, 1)$.

and

$$\sum_{i=1}^{99} p_i(1-p_i) = \frac{1}{100} \sum_{i=1}^{99} i - \frac{1}{(100)^2} \sum_{i=1}^{99} i^2$$

$$= 49.5 - \frac{1}{(100)^2} \cdot \frac{(99)(100)(199)}{6} = 16.665.$$

It follows from the central limit theorem that the distribution of the total number of questions that are answered correctly, which is $\sum_{i=1}^{99} X_i$, will be approximately the normal distribution with mean 49.5 and standard deviation $(16.665)^{1/2} = 4.08$. Therefore, the distribution of the variable

$$Z = \frac{\sum_{i=1}^n X_i - 49.5}{4.08}$$

will be approximately the standard normal distribution. It follows that

$$\Pr\left(\sum_{i=1}^n X_i \geq 60\right) = \Pr(Z \geq 2.5735) \simeq 1 - \Phi(2.5735) = 0.0050.$$

Outline of Proof of Central Limit Theorem

Convergence of the Moment Generating Functions Moment generating functions are important in the study of convergence in distribution because of the following theorem, the proof of which is too advanced to be presented here.

Theorem 6.3.5

Let $X_1, X_2, \dots$ be a sequence of random variables. For $n = 1, 2, \dots$, let F_n denote the c.d.f. of X_n , and let ψ_n denote the m.g.f. of X_n .

Also, let X^* denote another random variable with c.d.f. F^* and m.g.f. ψ^* . Suppose that the m.g.f.'s ψ_n and ψ^* exist ($n = 1, 2, \dots$). If $\lim_{n \rightarrow \infty} \psi_n(t) = \psi^*(t)$ for all values of t in some interval around the point $t = 0$, then the sequence $X_1, X_2, \dots$ converges in distribution to X^* .

In other words, the sequence of c.d.f.'s $F_1, F_2, \dots$ must converge to the c.d.f. F^* if the corresponding sequence of m.g.f.'s $\psi_1, \psi_2, \dots$ converges to the m.g.f. ψ^* .

Outline of the Proof of Theorem 5.7.1 We are now ready to outline a proof of Theorem 6.3.1, which is the central limit theorem of Lindeberg and Lévy. We shall assume that the variables $X_1, \dots, X_n$ form a random sample of size n from a distribution with mean μ and variance σ^2 . We shall also assume, for convenience, that the m.g.f. of this distribution exists, although the central limit theorem is true even without this assumption.

For $i = 1, \dots, n$, let $Y_i = (X_i - \mu)/\sigma$. Then the random variables $Y_1, \dots, Y_n$ are i.i.d., and each has mean 0 and variance 1. Furthermore, let

$$Z_n = \frac{n^{1/2}(\bar{X}_n - \mu)}{\sigma} = \frac{1}{n^{1/2}} \sum_{i=1}^n Y_i.$$

From DeGroot and Schervish,
Fourth Ed.

We shall show that Z_n converges in distribution to a random variable having the standard normal distribution, as indicated in Eq. (6.3.1), by showing that the m.g.f. of Z_n converges to the m.g.f. of the standard normal distribution.

If $\psi(t)$ denotes the m.g.f. of each random variable Y_i ($i = 1, \dots, n$), then it follows from Theorem 4.4.4 that the m.g.f. of the sum $\sum_{i=1}^n Y_i$ will be $[\psi(t)]^n$. Also, it follows from Theorem 4.4.3 that the m.g.f. $\zeta_n(t)$ of Z_n will be

$$\zeta_n(t) = \left[\psi\left(\frac{t}{n^{1/2}}\right) \right]^n.$$

In this problem, $\psi'(0) = E(Y_i) = 0$ and $\psi''(0) = E(Y_i^2) = 1$. Therefore, the Taylor series expansion of $\psi(t)$ about the point $t = 0$ has the following form:

$$\psi(t) = \psi(0) + t\psi'(0) + \frac{t^2}{2!}\psi''(0) + \frac{t^3}{3!}\psi'''(0) + \dots$$

$$= 1 + \frac{t^2}{2} + \frac{t^3}{3!}\psi'''(0) + \dots$$

Also,

$$\zeta_n(t) = \left[1 + \frac{t^2}{2n} + \frac{t^3\psi'''(0)}{3!n^{3/2}} + \dots \right]^n. \quad (6.3.15)$$

Apply Theorem 5.3.3 with $1 + a_n/n$ equal to the expression inside brackets in (6.3.15) and $c_n = n$. Since

$$\lim_{n \rightarrow \infty} \left[\frac{t^2}{2} + \frac{t^3\psi'''(0)}{3!n^{1/2}} + \dots \right] = \frac{t^2}{2},$$

it follows that

$$\lim_{n \rightarrow \infty} \zeta_n(t) = \exp\left(\frac{1}{2}t^2\right). \quad (6.3.16)$$

Since the right side of Eq. (6.3.16) is the m.g.f. of the standard normal distribution, it follows from Theorem 6.3.5 that the asymptotic distribution of Z_n must be the standard normal distribution.

An outline of the proof of the central limit theorem of Liapounov can also be given by proceeding along similar lines, but we shall not consider this problem further here.

Summary

Two versions of the central limit theorem were given. They conclude that the distribution of the average of a large number of independent random variables is close to a normal distribution. One theorem requires that the random variables all have the same distribution with finite variance. The other theorem does not require that the random variables be identically distributed, but instead requires that their third moments exist and satisfy condition (6.3.9). The delta method lets us find the approximate distribution of a smooth function of a sample average.

1 **Exercise:** What is the practical value of the CLT?

2 _____

3 _____

4 _____

5 _____

6 _____

7 **Exercise:** Let $X_1, X_2, \dots, X_n$ be i.i.d. random variables, each with the
8 Exponential(1) distribution. What is the distribution of $\bar{X}$?

9 _____

10 _____

11 _____

12 _____

13 _____

14 _____

15 _____

16 _____

17 _____

18 _____

19 _____

1 **Exercise:** Suppose that on any day the price of a particular stock
2 either increases by 2% (with probability 0.5) or decreases by 1% (with
3 probability 0.5). Each day's change is independent of the other days.
4 What is the probability that the stock's price is more than twice its
5 current price 100 trading days from now?

6 _____

7 _____

8 _____

9 _____

10 _____

11 _____

12 _____

13 _____

14 _____

15 _____

16 _____

17 _____

18 _____

19 _____

20 _____

21 _____

22 _____

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20

1
2
3
4
5
6
7
8
9
10
11
12

Exercise: A fair coin is tossed 1000 times. What is the probability that the number of heads is between 475 and 525?

The Delta Method

A formal statement of the Delta Method is as follows:

Assume that $Y_1, Y_2, \dots$ are such that

$$\frac{Y_n - \mu}{\sigma_n} \xrightarrow{D} Z$$

where $Z \sim N(0, 1)$, μ is a constant, and σ_n is such that $\sigma_n > 0$ and $\sigma_n \downarrow 0$ as $n \rightarrow \infty$. Then, assuming $g(\cdot)$ is a function that is differentiable at μ and $g'(\mu) \neq 0$, it holds that

$$\frac{g(Y_n) - g(\mu)}{|g'(\mu)|\sigma_n} \xrightarrow{D} Z.$$

A more heuristic, but useful, version is as follows:

If Y_n is approximately $N(\mu, \sigma_n^2)$ and $\sigma_n^2 \downarrow 0$ as $n \rightarrow \infty$, then $g(Y_n)$ is approximately

$$N(g(\mu), (g'(\mu))^2 \sigma_n^2)$$

assuming that $g(\cdot)$ is a function that is differentiable at μ and $g'(\mu) \neq 0$.

This is often (but not always) applied with $Y_n = \bar{X}_n$ and $\sigma_n^2 = \sigma^2/n$ where $\sigma^2 > 0$.

Exercise: Suppose that $X_1, X_2, \dots, X_{50}$ are i.i.d. random variables with the Exponential($\lambda = 2$) distribution. One may hope that $1/\bar{X}$ is a good estimator of λ (imagining a case where λ is not known). What is the probability that $1/\bar{X}$ is within 0.1 of λ ?

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17


18

19


20

21


With $n = 5$



With $n = 30$



With $n = 15$



With $n = 100$

