

Copper-catalyzed reductive coupling of tosylhydrazones with amines: A convenient route to α -branched amines†

Abdallah Hamze,* Bret Tréguier, Jean-Daniel Brion and Mouâd Alami*

Received 28th April 2011, Accepted 6th July 2011

DOI: 10.1039/c1ob05664f

A general procedure for the reductive coupling of *N*-tosylhydrazones with amines in the presence of $\text{Cu}(\text{acac})_2$ and Cs_2CO_3 has been developed. The protocol is very effective and chemoselective with various primary and secondary aliphatic amines, aminoalcohols as well as azole derivatives to give α -branched amines in good yields.

α -Branched amines, including aryl- and diarylmethylamine derivatives represent an important class of organic compounds regarding their various biological and pharmacological activities (Fig. 1).¹ For instance, the diarylmethylamine moiety is a common pharmacophore for multiple drug classes such as histamine H_1 receptor antagonists (cetirizine),² calcium channel blockers (lomerizine),³ and aromatase inhibitors (letrozole).⁴ In addition, these structural motifs are highly valuable intermediates in organic synthesis.⁵ Thus, efficient syntheses of α -branched amines are of interest in medicinal chemistry and for the synthesis of screening libraries. The most common route for their preparation is undoubtedly the reductive amination of carbonyl compounds with metal hydrides.⁶ Alternative procedures consist of addition

of carbon nucleophiles to imines⁷ or related derivatives,⁸ displacement of polymer-supported benzotriazole using Grignard reagents,⁹ addition of organo-lithium and -magnesium reagents to selenoamides¹⁰ and thioformamides,¹¹ or acid-catalyzed reaction of diarylmethanols with phenyl carbamate followed by an extra cleavage step under alkaline media.¹² Systems related to three-component coupling of aromatic zinc reagents, secondary amines and aromatic aldehydes has also been reported.¹³ Although the aforementioned transformations are suitable procedures, however, most of them require the use of stoichiometric amounts of organometallic compounds or metal hydrides, thus generating waste from reagents. Therefore, the development of alternative and general protocols for the preparation of α -branched amines, which operate under environmentally friendly conditions, is still desirable.¹⁴

In recent years, *N*-tosylhydrazones have attracted extensive attention because of their various useful applications in organic synthesis.¹⁵ In particular, they are valuable and readily available reagents in C–C,¹⁶ C–O,¹⁷ and C–S¹⁸ bond-forming reactions through metal-catalyzed and metal-free processes. Although significant progress has been achieved on the aforementioned transformations, application in $\text{C}(\text{sp}^3)$ –N bond-forming reactions is still a relatively unexplored process. To our knowledge, the formation of amines from tosylhydrazones had been previously reported by Cuevas-Yañez and co-workers¹⁹ by employing $\text{Cu}(\text{acac})_2$ as the catalyst and a large excess of tetra-*n*-butylammonium bromide. In their study, the $\text{C}(\text{sp}^3)$ –N bond coupling reaction was only described with imidazole and benzimidazole derivatives, and therefore, its scope seems to be very limited. A supplementary drawback of this methodology was the need to pre-form a THF-solution of tosylhydrazone sodium salts with NaH, followed by a change of the reaction solvent to toluene, thus limited the practicality of the process.

As part of our continuing studies on the Pd-catalyzed coupling of *N*-tosylhydrazones with aryl halides,²⁰ combined with our interest in C–N bond forming reactions,²¹ we decided to explore the ability of *N*-tosylhydrazones to participate in metal-catalyzed C–N coupling reactions with various nitrogen nucleophiles. In this study, we report our success in achieving the reductive coupling of tosylhydrazones with amines in the presence of Cs_2CO_3 as the base and a catalytic amount of easily available $\text{Cu}(\text{acac})_2$ in dioxane. Under these simple and general reaction conditions, various arylsulfonylhydrazones **1** derived from ketones and aldehydes reacted with primary and secondary aliphatic amines as well as




Fig. 1 α -Branched drugs with aryl- and diarylmethylamine moieties.

Univ Paris-Sud 11, CNRS, BioCIS UMR 8076, Laboratoire de Chimie Thérapeutique, Faculté de Pharmacie, 5 rue J.-B. Clément, Châtenay-Malabry, F-92296, France. E-mail: abdallah.hamze@u-psud.fr, mouad.alami@u-psud.fr; Fax: +33-1-46835828; Tel: +33-1-46835887

† Electronic supplementary information (ESI) available. See DOI: 10.1039/c1ob05664f

Table 1 (Contd.)

$\text{R}^1\text{-C}_6\text{H}_4\text{-C(=NHTs)-R}^2 + \text{R}^3\text{-N(R}^4\text{)-H} \xrightarrow[\text{dioxane, 100 }^\circ\text{C}]{\text{Cu(acac)}_2, \text{Cs}_2\text{CO}_3} \text{R}^1\text{-C}_6\text{H}_4\text{-C(R}^2\text{)(N(R}^3\text{)R}^4\text{))-R}^2$				
Tosylhydrazones used in this study				
		1c: R = H 1d: R = Cl		
		1g: R = H 1h: R = Me		
Entry	1	Amine 2	Compound 3	Yield (%) ^b
9	1f	2a		71
10	1f	2e		60
11	1a	2f		50
12	1a	2g		62
13	1a	2h		53
14	1c	2i		42
15	1a	2j		0
16	1a	2k		60

Table 1 (Contd.)


$\text{R}^1\text{-C}_6\text{H}_4\text{-C(=NHTs)-R}^2 + \text{R}^3\text{-N(R}^4\text{)-H} \xrightarrow[\text{dioxane, 100 }^\circ\text{C}]{\text{Cu(acac)}_2, \text{Cs}_2\text{CO}_3} \text{R}^1\text{-C}_6\text{H}_4\text{-C(R}^2\text{)(N(R}^3\text{)R}^4\text{))-R}^2$				
Tosylhydrazones used in this study				
		1c: R = H 1d: R = Cl		
		1g: R = H 1h: R = Me		
Entry	1	Amine 2	Compound 3	Yield (%) ^b
17	1a	2l		84
18	1a	2m		61
19	1a	2n		65
20	1c	2o		64
21	1c	2p		70
22	1g	2n		63
23	1h	2n		40

^a Reactions conditions: tosylhydrazone **1** (0.25 mmol), amine **2** (2 equiv), Cu(acac)₂ (10 mol%), Cs₂CO₃ (2.2 equiv), dioxane (2.5 mL) in a sealed tube at 100 °C. ^b Isolated yield of **3**. ^c A 72% yield was obtained when the reaction was carried out on a 2 gram scale of tosylhydrazone **1a**. ^d Using open-vessel reaction conditions, a 71% yield was obtained within 3 h heating at 100 °C. ^e Reaction conducted in the presence of 20 mol% *N,N'*-dimethylethylenediamine (DMEDA); less than 10% of **3g** was obtained in the absence of DMEDA.

ether products resulting from C–O bond-forming reaction was observed, although tosylhydrazones have been recently reported to provide reductive etherification with alcohols.¹⁷

Finally, extending this chemistry to include azoles **2n–p** as substrates proved to be quite successful, with tosylhydrazones derived from aromatic ketones (entries 19–21) as well as aliphatic aldehydes (entry 22) and ketones (entry 23). Thus, imidazole, benzimidazole and pyrazole also underwent a C–N bond forming reaction providing coupling products **3s–w** in satisfactory to good yields.

Although there is no clear experimental evidence, we suppose that the reaction proceeds as shown in Scheme 2. Initially, *N*-tosylhydrazone **1** in basic media undergoes thermal decomposition through the Bamford–Stevens reaction to generate diazo compound **II** followed by *in situ* reduction of Cu(II) by **II** into the active catalytic Cu(I) species **III**.²³ This latter would react with **II** to give the copper(I) carbene complex²⁴ **IV**. Further copper carbene N–H insertion might proceed with attack of the nitrogen atom on the electrophilic carbene to generate copper species **V**, followed by hydrogen transfer²⁵ to produce amine **3**.



Scheme 2 Proposed mechanism for the Cu-catalyzed reductive coupling of *N*-tosylhydrazones **1** with amines.

In summary, we have demonstrated that the reductive coupling of tosylhydrazones with amines in presence of Cs_2CO_3 and a catalytic amount of $\text{Cu}(\text{acac})_2$ represents a general, operationally simple, inexpensive and efficient approach for the synthesis of aryl- and diarylmethylamine derivatives. A variety of amines, including primary and secondary aliphatic amines as well as azole derivatives were used successfully. It is particularly noteworthy that this protocol is chemoselective allowing amino alcohols to react with hydrazones to give exclusively C–N bond forming products. Further development of this coupling methodology is under way.

Notes and references

- (a) W. N. Washburn, C. Q. Sun, G. Bisacchi, G. Wu, P. T. Cheng, P. M. Sher, D. Ryono, A. V. Gavai, K. Poss, R. N. Girotra, P. J. McCann, A. B. Mikkilineni, T. C. Dejneka, T. C. Wang, Z. Merchant, M. Morella, C. M. Arbeen, T. W. Harper, D. A. Slusarchyk, S. Skwish, A. D. Russell, G. T. Allen, B. Tesfamariam, B. H. Frohlich, B. E. Abboa-Offei, M. Cap, T. L. Waldron, R. J. George, D. Young, K. E. Dickinson and A. A. Seymour, *Bioorg. Med. Chem. Lett.*, 2004, **14**, 3525–3529; (b) W. N. Washburn, P. M. Sher, K. M. Poss, R. N. Girotra, P. J. McCann, A. V. Gavai, A. B. Mikkilineni, A. Mathur, P. Cheng, T. C. Dejneka, C. Q. Sun, T. C. Wang, T. W. Harper, A. D. Russell, D. A. Slusarchyk, S. Skwish, G. T. Allen, D. E. Hillyer, B. H. Frohlich, B. E. Abboa-Offei, M. Cap, T. L. Waldron, R. J. George, B. Tesfamariam, C. P. Ciosek, D. Ryono, D. A. Young, K. E. Dickinson, A. A. Seymour,

- C. M. Arbeen and R. E. Gregg, *Bioorg. Med. Chem. Lett.*, 2001, **11**, 3035–3039.
- M. P. Curran, L. J. Scott and C. M. Perry, *Drugs*, 2004, **64**, 523–561.
- H. Hara, N. Toriu and M. Shimazawa, *Cardiovasc. Drug Rev.*, 2004, **22**, 199–214.
- H. Mouridsen, M. Gershanovich, Y. Sun, R. n. Perez-Carrion, C. Boni, A. Monnier, J. Apffelstaedt, R. Smith, H. P. Sleeboom, F. Jaenicke, A. Pluzanska, M. Dank, D. Becquart, P. P. Bapsy, E. Salminen, R. Snyder, H. Chaudri-Ross, R. Lang, P. Wyld and A. Bhatnagar, *J. Clin. Oncol.*, 2003, **21**, 2101–2109.
- (a) B. Hatano, J.-I. Kadokawa and H. Tagaya, *Tetrahedron Lett.*, 2002, **43**, 5859–5861; (b) F. Shibahara, R. Sugiura, E. Yamaguchi, A. Kitagawa and T. Murai, *J. Org. Chem.*, 2009, **74**, 3566–3568.
- (a) A. F. Abdel-Magid, K. G. Carson, B. D. Harris, C. A. Maryanoff and R. D. Shah, *J. Org. Chem.*, 1996, **61**, 3849–3862; (b) T. H. Jones and M. S. Blum, *J. Org. Chem.*, 1980, **45**, 4778–4780; (c) O. Provot, J. P. Célrier, H. Petit and G. Lhommet, *J. Org. Chem.*, 1992, **57**, 2163–2166.
- (a) N. Cabello, J.-C. Kizirian and A. Alexakis, *Tetrahedron Lett.*, 2004, **45**, 4639–4642; (b) D. Ferraris, *Tetrahedron*, 2007, **63**, 9581–9597.
- (a) H. Böhme and P. Plappert, *Chem. Ber.*, 1975, **108**, 2827–2833; (b) N. Plobeck and D. Powell, *Tetrahedron: Asymmetry*, 2002, **13**, 303–310.
- K. Schiemann and H. D. H. Showalter, *J. Org. Chem.*, 1999, **64**, 4972–4975.
- M. Sekiguchi, A. Ogawa, S.-I. Fujiwara, I. Ryu, N. Kambe and N. Sonoda, *Chem. Lett.*, 1990, **19**, 2053–2056.
- T. Murai, K. Ui and Narengerile, *J. Org. Chem.*, 2009, **74**, 5703–5706.
- S. Jolidon, D. Alberati, A. Dowle, H. Fischer, D. Hainzl, R. Narquizian, R. Norcross and E. Pinard, *Bioorg. Med. Chem. Lett.*, 2008, **18**, 5533–5536.
- (a) C. Haurena, E. Le Gall, S. p. Sengmany, T. Martens and M. Troupel, *J. Org. Chem.*, 2010, **75**, 2645–2650; (b) E. Le Gall, C. Haurena, S. p. Sengmany, T. Martens and M. Troupel, *J. Org. Chem.*, 2009, **74**, 7970–7973; (c) E. Le Gall, M. Troupel and J.-Y. Nédélec, *Tetrahedron*, 2006, **62**, 9953–9965; (d) E. Le Gall, M. Troupel and J.-Y. Nédélec, *Tetrahedron Lett.*, 2006, **47**, 2497–2500; (e) S. Sengmany, E. Le Gall, C. Le Jean, M. Troupel and J.-Y. Nédélec, *Tetrahedron*, 2007, **63**, 3672–3681.
- For transition metal-catalyzed hydrominoration of styrenes, see: M. Utsunomiya and J. F. Hartwig, *J. Am. Chem. Soc.*, 2003, **125**, 14286–14287.
- J. R. Fulton, V. K. Aggarwal and J. de Vicente, *Eur. J. Org. Chem.*, 2005, 1479–1492.
- (a) J. Barluenga, P. Moriel, C. Valdes and F. Aznar, *Angew. Chem., Int. Ed.*, 2007, **46**, 5587–5590; (b) J. Barluenga, M. Tomas-Gamasa, P. Moriel, F. Aznar and C. Valdes, *Chem.–Eur. J.*, 2008, **14**, 4792–4795; (c) W.-Y. Yu, Y.-T. Tsoi, Z. Zhou and A. S. C. Chan, *Org. Lett.*, 2008, **11**, 469–472; (d) J. Barluenga, M. Escrivano, P. Moriel, F. Aznar and C. Valdes, *Chem.–Eur. J.*, 2009, **15**, 13291–13294; (e) J. Barluenga, M. Tomas-Gamasa, F. Aznar and C. Valdes, *Nat. Chem.*, 2009, **1**, 494–499; (f) Q. Xiao, J. Ma, Y. Yang, Y. Zhang and J. Wang, *Org. Lett.*, 2009, **11**, 4732–4735; (g) J. Barluenga, M. Tomas-Gamasa, F. Aznar and C. Valdes, *Chem.–Eur. J.*, 2010, **16**, 12801–12803; (h) J. Barluenga, M. Tomás-Gamasa, F. Aznar and C. Valdés, *Adv. Synth. Catal.*, 2010, **352**, 3235–3240; (i) X. Zhao, J. Jing, K. Lu, Y. Zhang and J. Wang, *Chem. Commun.*, 2010, **46**, 1724–1726; (j) X. Zhao, G. Wu, C. Yan, K. Lu, H. Li, Y. Zhang and J. Wang, *Org. Lett.*, 2010, **12**, 5580–5583; (k) L. Zhou, F. Ye, Y. Zhang and J. Wang, *J. Am. Chem. Soc.*, 2010, **132**, 13590–13591; (l) J. Barluenga, L. Florentino, F. Aznar and C. Valdes, *Org. Lett.*, 2011, **13**, 510–513; (m) X. Zhao, G. Wu, Y. Zhang and J. Wang, *J. Am. Chem. Soc.*, 2011, **133**, 3296–3299; (n) L. Zhou, F. Ye, J. Ma, Y. Zhang and J. Wang, *Angew. Chem., Int. Ed.*, 2011, **50**, 3510–3514.
- J. Barluenga, M. Tomas-Gamasa, F. Aznar and C. Valdes, *Angew. Chem., Int. Ed.*, 2010, **49**, 4993–4996.
- (a) X.-W. Feng, J. Wang, J. Zhang, J. Yang, N. Wang and X.-Q. Yu, *Org. Lett.*, 2010, **12**, 4408–4411; (b) J. Barluenga, M. Tomás-Gamasa, F. Aznar and C. Valdés, *Eur. J. Org. Chem.*, 2011, 1520–1526; (c) Q. Ding, B. Cao, J. Yuan, X. Liu and Y. Peng, *Org. Biomol. Chem.*, 2011, **9**, 748–751.
- E. Cuevas-Yañez, J. M. Serrano, G. Huerta, J. M. Muchowski and R. Cruz-Almanza, *Tetrahedron*, 2004, **60**, 9391–9396.
- (a) B. Treguer, A. Hamze, O. Provot, J.-D. Brion and M. Alami, *Tetrahedron Lett.*, 2009, **50**, 6549–6552; (b) E. Brachet, A. Hamze, J.-F. Peyrat, J.-D. Brion and M. Alami, *Org. Lett.*, 2010, **12**, 4042–4045; (c) E. Rasolofonjatovo, B. Tréguier, O. Provot, A. Hamze, E. Morvan, J.-D. Brion and M. Alami, *Tetrahedron Lett.*, 2011, **52**,

- 1036–1040; (d) S. Messaoudi, B. Treguier, A. Hamze, O. Provot, J.-F. Peyrat, J. R. De Losada, J.-M. Liu, J. Bignon, J. Wdzieczak-Bakala, S. Thoret, J. Dubois, J.-D. Brion and M. Alami, *J. Med. Chem.*, 2009, **52**, 4538–4542; (e) S. Messaoudi, A. Hamze, O. Provot, B. Tréguier, J. R. De Losada, J. Bignon, J.-M. Liu, J. Wdzieczak-Bakala, S. Thoret, J. Dubois, J.-D. Brion and M. Alami, *ChemMedChem*, 2011, **6**, 488–497.
- 21 (a) D. Audisio, S. Messaoudi, J.-F. Peyrat, J.-D. Brion and M. Alami, *Tetrahedron Lett.*, 2007, **48**, 6928–6932; (b) S. Messaoudi, D. Audisio, J.-D. Brion and M. Alami, *Tetrahedron*, 2007, **63**, 10202–10210; (c) S. Messaoudi, J.-D. Brion and M. Alami, *Adv. Synth. Catal.*, 2010, **352**, 1677–1687; (d) S. Messaoudi, J.-D. Brion and M. Alami, *Tetrahedron Lett.*, 2011, **52**, 2687–2691; (e) D. Audisio, S. Messaoudi, L. Ceigeikowski, J.-F. Peyrat, J.-D. Brion, D. Methy-Gonnot, C. Radanyi, J.-M. Renoir and M. Alami, *ChemMedChem*, 2011, **6**, 804–815.
- 22 W. R. Bamford and T. S. Stevens, *J. Chem. Soc.*, 1952, 4735–4740.
- 23 R. G. Salomon and J. K. Kochi, *J. Am. Chem. Soc.*, 1973, **95**, 3300–3310.
- 24 (a) W. Kirmse, *Angew. Chem., Int. Ed.*, 2003, **42**, 1088–1093; (b) B. F. Straub and P. Hofmann, *Angew. Chem., Int. Ed.*, 2001, **40**, 1288–1290.
- 25 (a) M. P. Doyle and D. C. Forbes, *Chem. Rev.*, 1998, **98**, 911–935; (b) Z. Zhang and J. Wang, *Tetrahedron*, 2008, **64**, 6577–6605; (c) Y.-Z. Zhang, S.-F. Zhu, Y. Cai, H.-X. Mao and Q.-L. Zhou, *Chem. Commun.*, 2009, 5362–5364.