

# Weiner Reconstruction in 2D

Vikrant Kamble

September 10, 2018

Given a data

$$\mathbf{d} = \mathbf{s}_d + \mathbf{e}$$

where  $\mathbf{s}_d$  is the underlying signal and  $\mathbf{e}$  is the measurement uncertainty, our aim is to obtain estimates of the signal  $\mathbf{s}_{map}$  at a set of positions  $\mathbf{x}_{map}$ :

$$\mathbf{s}_{map} = \mathbf{M} \mathbf{d} \quad (1)$$

Using the matrix  $\mathbf{M}$  that minimizes the objective function

$$\mathbf{E}[|\mathbf{s}_{map} - \mathbf{s}_{actual}|^2]$$

the above equation becomes:

$$\mathbf{s}_{map} = \mathbf{C}_{map,d} [\mathbf{C}_{d,d} + \mathbf{N}]^{-1} \mathbf{d} \quad (2)$$


where  $\mathbf{C}_{map,d}$  is the map-data covariance matrix and  $\mathbf{C}_{d,d}$  is the data-data covariance matrix.

The covariance matrix of the mapped data  $\mathbf{s}_{map}$  that has been Wiener reconstructed is given by:

$$\mathbf{C} = \mathbf{C}_{map,map} - \mathbf{C}_{map,d} [\mathbf{C}_{d,d} + \mathbf{N}]^{-1} \mathbf{C}_{map,d}^T$$

Example:

$$A = 2.0, \quad a = 5.0$$



In 2D, for isotropic field, the correlation function  $\xi(r)$  and the power spectrum  $P(k)$  are related by:

$$P(k) = \int_0^\infty \xi(r) J_0(kr) 2\pi r dr, \quad (3)$$

where  $J_0(kr)$  is the Bessel function of the first kind.

Let's create a perturbation field  $\delta(\mathbf{x})$  in 2D using the correlation function:

$$\xi(r) = A \exp \left[ -\frac{r^2}{2a^2} \right] \quad (4)$$


The power spectrum becomes:

$$P(k) = 2\pi A a^2 \exp \left[ -\frac{k^2 a^2}{2} \right] \quad (5)$$

Its much quicker to create fields in fourier space  $\delta_{\mathbf{k}}$  and then use the efficient IFFT algorithm to get  $\delta(\mathbf{x})$ . **Actually, this method gives 2 independent fields with one run!!!**

$\text{num\_points} = 2400 \times 2400$

$dx_{\perp} = dx_{\parallel} = 0.052$



$$A = 1.0, a = 8.0$$

