

Mean of datasets

TOTAL POINTS 6

1. What is the mean of the dataset $\mathcal{D} = \{1, 2, 3\}$?

1 / 1 point

Do the exercises using pen and paper.

- ☐ 3
- ☐ 6
- ☒ 2

✓ **Correct**
That's it. Good job!

2. Compute the mean of the following dataset:

1 / 1 point

$$\mathcal{D} = \left\{ \begin{bmatrix} 1 \\ 4 \\ 7 \end{bmatrix}, \begin{bmatrix} 2 \\ 5 \\ 8 \end{bmatrix}, \begin{bmatrix} 3 \\ 6 \\ 9 \end{bmatrix} \right\}$$

Do the exercises using pen and paper.

- ☒ $\begin{bmatrix} 2 \end{bmatrix}$
- ☐ $\begin{bmatrix} 3 \\ 3 \\ 3 \end{bmatrix}$
- ☐ $\begin{bmatrix} 18 \\ 18 \\ 18 \end{bmatrix}$

✓ **Correct**
Well done!

5. Assuming that we know the mean \bar{x}_{n-1} of a dataset \mathcal{D}_{n-1} with $n - 1$ data points. Now, suppose that we collect another data point, which we denote by x_* . Select the correct formula that computes the correct new mean \bar{x}_n of the full data set $\mathcal{D}_n = \mathcal{D}_{n-1} \cup \{x_*\}$, i.e., we add x_* to the dataset \mathcal{D} .

1 / 1 point

- ☐ $\bar{x}_n = \bar{x}_{n-1} + \frac{1}{n+1}(\bar{x}_{n-1} - x_*)$
- ☐ $\bar{x}_n = \bar{x}_{n-1} + \frac{1}{n-1}(x_* - \bar{x}_{n-1})$
- ☒ $\bar{x}_n = \bar{x}_{n-1} + \frac{1}{n}(x_* - \bar{x}_{n-1})$
- ☐ $\bar{x}_n = \bar{x}_{n-1} + \frac{1}{n+1}(x_* - \bar{x}_{n-1})$

✓ **Correct**
Excellent!

6. Assuming you are given an image as a two dimensional array of shape 28 x 28. Write a small piece of python code to reshape this image to a vector of length 784 (=28 x 28).

1 / 1 point

Hint: This can be a one-liner.

```
1 import numpy as np
2
3 def reshape(x):
4     """return x_reshaped as a flattened vector of the multi-dimensional array x"""
5     x_reshaped = x.reshape(x.shape[0]*x.shape[1])
6     return x_reshaped
7
```

Run

Reset

✓ **Correct**

Good job!