

Matching functions and approximations

TOTAL POINTS 5


- As mentioned in the previous video, the Taylor series approximation can also be viewed as a power series, in which these approximations are used to build functions that are often simpler and easier to evaluate, particularly when using numerical methods. In the following questions, we are looking at developing our understanding of how the increasing order of a power series allows us to develop further information of a function.

1 / 1 point

Below are three graphs highlighting the zeroth, second and fourth order approximations of a common trigonometric function. Observe how increasing the number of approximations in the power series begins to build a better approximation, and determine which function these approximations represent.


Zeroth order approximation:

$$f_0(x) = 1$$



Fourth order approximation:

$$f_4(x) = 1 - \frac{x^2}{2} + \frac{x^4}{24}$$



☒ $f(x) = \cos(x)$



☐ $f(x) = \cos^2(x)$

The function $f(x) = \cos(x)$ is symmetric about the line $x = 0$. Furthermore our approximation of this function is when $x = 0$, known as a Maclaurin series. At the point $x = 0$, $f(0) = 1$ which is shown in the zeroth order approximation.

- Below are three graphs highlighting the first, third and fifth order approximations of a common trigonometric function. Observe how the power series begins to build the function, and determine which function these approximations represent.

1 / 1 point

First Order:

$$f_1(x) = 2x$$



Fifth Order:

$$f_5(x) = 2x - \frac{4x^3}{3} + \frac{4x^5}{15}$$

☐ $f(x) = \sin(2x)$



☐ $f(x) = \sin(\frac{1}{2}x)$



✓ Correct

The function $f(x) = \sin(2x)$ has rotational symmetry about the origin. Furthermore, we can see that the period is much shorter, also evident from the three approximations shown.

- The graph below shows the function $f(x) = \tan^{-1}(x)$, select all the power series approximations that can be used to obtain an approximation for this function.

1 / 1 point



☐ $f(x) = -\frac{x^2}{5} + \frac{x^4}{5} \dots$



✓ Correct

We can see this approximation goes through the origin and also looks as if it fits the function well between $-0.5 < x < 0.5$.

☐ $f(x) = -x^2 \dots$



☒ $f(x) = x \dots$

✓ Correct


We can see this approximation goes through the origin and also looks as if it fits the function well between $-0.5 < x < 0.5$. As this is a linear function, this is a first-order approximation.

- The sinusoidal function $f(x) = \sin(x)$ (green line) centered at $x = 0$ is shown in the graph below. The approximation for this function is shown through the series $f(x) = x - \frac{x^3}{6} \dots$ (orange line). Determine what polynomial order is represented by the orange line.

1 / 1 point



- ☐ Zeroth Order
☐ First Order
☒ Third Order
☐ Fifth Order
☐ None of the above



- ☐ First Order
☐ Second Order
☐ Third Order
☒ Not a correct approximation

✓ Correct

The approximation shown is not tangent to this point and is, therefore, a poor approximation of the function.