

for postbronchodilator testing (8), or the different geographic location (South America), may explain some of the differences.

Overall, this study provides valuable new information, as well as the opportunity for the authors and others to initiate additional studies to examine diverse populations and settings and to determine if the findings in this study are relevant beyond Sweden. Then we will know with greater certainty whether postbronchodilator reference values should be the new norm. ■

**Author disclosures** are available with the text of this article at [www.atsjournals.org](http://www.atsjournals.org).

Lewis J. Smith, M.D.  
Feinberg School of Medicine  
Northwestern University  
Chicago, Illinois

ORCID ID: 0000-0002-4728-1562 (L.J.S.).

## References

1. Johannessen A, Lehmann S, Omenaas ER, Eide GE, Bakke PS, Gulsvik A. Post-bronchodilator spirometry reference values in adults and

- implications for disease management. *Am J Respir Crit Care Med* 2006; 173:1316–1325.
2. Quanjer PH, Brazzale DJ, Boros PW, Pretto JJ. Implications of adopting the Global Lungs Initiative 2012 all-age reference equations for spirometry. *Eur Respir J* 2013;42:1046–1054.
3. Moreira GL, Manzano BM, Gazzotti MR, Nascimento OA, Perez-Padilla R, Menezes AM, *et al.*; PLATINO. PLATINO, a nine-year follow-up study of COPD in the city of São Paulo, Brazil: the problem of underdiagnosis. *J Bras Pneumol* 2014;40:30–37.
4. Pérez-Padilla R, Torre Bouscoulet L, Vázquez-García JC, Muño A, Márquez M, López MV, *et al.*; Grupo PLATINO. Spirometry reference values after inhalation of 200 microg of salbutamol [in Spanish]. *Arch Bronconeumol* 2007;43:530–534.
5. Kato B, Gulsvik A, Vollmer W, Janson C, Studnicka M, Buist S, *et al.* Can spirometric norms be set using pre- or post-bronchodilator test results in older people? *Respir Res* 2012;13:102.
6. Malinowski A, Zhou X, Andersson A, Backman H, Bake B, Blomberg A, *et al.* Consequences of using post- or pre-bronchodilator reference values in interpreting spirometry. *Am J Respir Crit Care Med* 2023;208: 461–471.
7. Bergström G, Berglund G, Blomberg A, Brandberg J, Engström G, Engvall J, *et al.* The Swedish Cardiopulmonary Bioimage Study: objectives and design. *J Intern Med* 2015;278:645–659.
8. Menezes AM, Perez-Padilla R, Jardim JRB, Muño A, Lopez MV, Valdivia G, *et al.*; PLATINO Team. Chronic obstructive pulmonary disease in five Latin American cities (the PLATINO study): a prevalence study. *Lancet* 2005;366:1875–1881.

Copyright © 2023 by the American Thoracic Society



# Unraveling the Distal Lung Destruction in Emphysema

The landmark study by Booth and colleagues (pp. 472–486) in this issue of the *Journal* hones our understanding of the loss of the terminal conducting airways and gas exchange areas in chronic obstructive pulmonary disease (COPD) by applying single-cell analyses of lung tissue from well-phenotyped individuals (1). The investigation builds on this group's previous discovery (2) that the terminal bronchioles, the smallest conducting airways, are the early battlegrounds of tissue destruction in COPD.

The study's ambitious objective was to create a comprehensive blueprint that sheds light on the structural, cellular, and extracellular matrix changes underpinning terminal bronchiole loss in COPD. The cross-sectional evaluation of more than 200 terminal bronchioles from 109 lungs of individuals with mild to moderate COPD and 24 with severe COPD, as gauged by Global Initiative for Chronic Obstructive Lung Disease criteria, compared with 82 lungs from ex-smokers without lung dysfunction used a multifaceted approach that included stereology, micro-computed tomography, nonlinear optical microscopy, imaging mass cytometry, and transcriptomics. Their findings unveiled that in addition to a net loss, the terminal

bronchioles progressively narrow as COPD severity increases, a phenomenon marked by the loss of elastin fibers within alveolar attachments. The pathology of alveolar untethering was noticeable even in the early stages of the disease in the absence of emphysematous alveolar loss, suggesting that this step is one of the first events in the hallmark distal lung destruction of emphysema. Although the concept of matrix elastin and collagen fiber degradation has been the forefront paradigm of protease–antiprotease imbalance of emphysema pathogenesis for decades (3, 4), this report refines our understanding of its spatial and temporal association with centrilobular distal lung destruction.

Furthermore, the single-cell atlas identified inflammatory and immune cells concentrated in this region of interest, with proinflammatory M1-like macrophages and neutrophils being located within disrupted alveolar attachments, whereas adaptive immune cells such as naive T cells, CD4 and CD8 T cells, and B cells were adjacent to terminal bronchiole wall remodeling. The genetic landscape was also consistent with upregulation of genes involved in both innate and adaptive immune responses, IFN response, and neutrophil degranulation. The proximity of macrophages and degranulating neutrophils to areas of matrix disruption is consistent with the mechanistic involvement of matrix metalloproteinases and neutrophil elastase and other proteinases in distal lung destruction (5–7). With our increased appreciation of the lung macrophage heterogeneity, future studies will have to build on expanding on the phenotyping of macrophages associated with disruption of alveolar tethers. Potential candidates are CD206<sup>+</sup>/CD43<sup>−</sup> interstitial

Ⓢ This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0. For commercial usage and reprints, please e-mail Diane Gern ([dgern@thoracic.org](mailto:dgern@thoracic.org)).

Supported by NHLBI grant R01HL077328.

Originally Published in Press as DOI: 10.1164/rccm.202307-1198ED on July 14, 2023

macrophages, which were found to be increased in the alveolar septae of the lungs of smokers compared with nonsmokers (8). The study by Booth and colleagues indicates progressive changes in the cellular heterogeneity of terminal bronchioles, enriched in inflammatory and immune cells in milder disease with a relative paucity of these cells in more severe COPD. The finding of an apparent exuberant immune cell activity in the distal lung early in the disease course substantiates the importance of the immune response in its development (9). However, because not all individuals with mild COPD progress to severe emphysema, this cross-sectional evaluation leaves unanswered the question of whether all the herein identified immune effector cells are causal to the untethering and remodeling of the terminal bronchiole or whether they contribute to the maintenance and repair of the distal lung. Although it is remarkable that the present study of 33 cell markers enabled the mass cytometric identification of 21 cell types composing the distal lung, their functionality as well as that of other cells that have not been systematically evaluated in the present study remains to be explored in more detail by future investigations. Indeed, many of the cellular responses implicated in distal lung cell destruction and repair (10, 11), such as oxidative stress, autophagy, mitochondrial dysfunction, senescence, progenitor cell exhaustion, and various types of programmed cell death, have not been explored. Also notable is the enrichment of the terminal bronchiolar region in unique cell types that may have a role in protecting this vulnerable and critical area at the junction of conducting airways and the alveolar gas exchange surfaces, including secretory cells and various types of progenitor or reparative cells (12–15).

This report marks a significant step forward in our quest to unravel the complexities of COPD pathogenesis and underscores the urgency to better understand the mechanisms and develop strategies that target the pathological changes involving the terminal bronchiole. ■

**Author disclosures** are available with the text of this article at [www.atsjournals.org](http://www.atsjournals.org).

Irina Petrache, M.D.  
National Jewish Health and the University of Colorado  
Denver, Colorado

ORCID ID: 0000-0003-1094-2600 (I.P.).

## References

- Booth S, Hsieh A, Mostaco-Guidolin L, Koo HK, Wu K, Aminazadeh F, et al. A single-cell atlas of small airway disease in chronic obstructive pulmonary disease: a cross-sectional study. *Am J Respir Crit Care Med* 2023;208:472–486.
- Hogg JC, Chu F, Utokaparch S, Woods R, Elliott WM, Buzatu L, et al. The nature of small-airway obstruction in chronic obstructive pulmonary disease. *N Engl J Med* 2004;350:2645–2653.
- Stone PJ. The elastase-antielastase hypothesis of the pathogenesis of emphysema. *Clin Chest Med* 1983;4:405–412.
- Evans MD, Pryor WA. Cigarette smoking, emphysema, and damage to alpha 1-proteinase inhibitor. *Am J Physiol* 1994;266:L593–L611.
- Gharib SA, Manicone AM, Parks WC. Matrix metalloproteinases in emphysema. *Matrix Biol* 2018;73:34–51.
- Atkinson JJ, Lutey BA, Suzuki Y, Toennies HM, Kelley DG, Kobayashi DK, et al. The role of matrix metalloproteinase-9 in cigarette smoke-induced emphysema. *Am J Respir Crit Care Med* 2011;183:876–884.
- Finlay GA, O'Driscoll LR, Russell KJ, D'Arcy EM, Masterson JB, FitzGerald MX, et al. Matrix metalloproteinase expression and production by alveolar macrophages in emphysema. *Am J Respir Crit Care Med* 1997;156:240–247.
- Hume PS, Gibbins SL, Jakubzick CV, Tudor RM, Curran-Everett D, Henson PM, et al. Localization of macrophages in the human lung via design-based stereology. *Am J Respir Crit Care Med* 2020;201:1209–1217.
- Kheradmand F, Zhang Y, Corry DB. Contribution of adaptive immunity to human COPD and experimental models of emphysema. *Physiol Rev* 2023;103:1059–1093.
- Tudor RM, Petrache I. Pathogenesis of chronic obstructive pulmonary disease. *J Clin Invest* 2012;122:2749–2755.
- Ryter SW, Nakahira K, Haspel JA, Choi AM. Autophagy in pulmonary diseases. *Annu Rev Physiol* 2012;74:377–401.
- Rustam S, Hu Y, Mahjour SB, Rendeiro AF, Ravichandran H, Urso A, et al. A unique cellular organization of human distal airways and its disarray in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2023;207:1171–1182.
- Ji CM, Plopper CG, Pinkerton KE. Clara cell heterogeneity in differentiation: correlation with proliferation, ultrastructural composition, and cell position in the rat bronchiole. *Am J Respir Cell Mol Biol* 1995;13:144–151.
- Wang C, de Mochel NSR, Christenson SA, Cassandras M, Moon R, Brumwell AN, et al. Expansion of hedgehog disrupts mesenchymal identity and induces emphysema phenotype. *J Clin Invest* 2018;128:4343–4358.
- Summers ME, Richmond BW, Kropski JA, Majka SA, Bastarache JA, Hatzopoulos AK, et al. Balanced Wnt/Dickkopf-1 signaling by mesenchymal vascular progenitor cells in the microvascular niche maintains distal lung structure and function. *Am J Physiol Cell Physiol* 2021;320:C119–C131.

Copyright © 2023 by the American Thoracic Society