Name: _____

Instructions: All solutions should be prepared carefully and neatly. Any questions that ask you to explain or justify should have answers written in complete sentences. All solution sets shall be completed on this packet and handed in at the beginning of class on the due date below. Careless presentation (e.g. bad handwriting, pen scribbles, doodles, wasted space, etc) will result in a deduction of points at my discretion. If the question asks to submit a graph or plot, be sure to print the plot and include it in your final lab report. Use a PENCIL and if you make a mistake, use an eraser. Each student should prepare their own solutions that demonstrate their own comprehension of the material. Submitted work that does not demonstrate clearly the process by which one arrived at the answer will not receive credit of any kind. Academic dishonesty will not be tolerated.

Desmos Lab III

MATH 181 – Calculus I

Due: Tuesday, November 25

Read: Section 4.2 and 4.6

Problem Number	Available Points	Your Points
1	10	
2	10	
Total	20	

1. The Mean Value Theorem In this part, we'll consider a simple function that satisfy both the Mean Value Theorem

   (a) Open a new graph in Desmos and save it as "The Mean Value Theorem."

   (b) Define the following function on the first line in Desmos: $f(x) = x^3 - x$. On the next two lines, define $a = -2$ and $b = 3$, but allow it to have the slider option. Graph this function only on the set $[a, b]$ by typing $\{a \leq x \leq b\}$ right next to the function on the first line. Change the window to $x \in [-3, 4]$ and $y \in [-10, 25]$. Plot the curve in black with width size 4.

   (c) In the space below, find the secant line slope that connects the points $(-2, f(-2))$ and $(3, f(3))$. In Desmos, on a new line, define the value m_{sec} to be this slope, but instead of using -2 and 3, define it in terms of a and b. Does Desmos output the same answer that you got?

   (d) Create a new folder in Desmos, call it "Secant Line". In this folder, plot the two points $(a, f(a))$ and $(b, f(b)$. Plot them in orange, size 15, and plot a label with them. In the space below, derive the equation for the line that connects the two points. Your answer should only involve m_{sec}, a, and $f(a)$. In the folder, define this line and plot it as a dashed green line. If you change the values of a and b, the graph should maintain this line connecting the two points.

   (e) Outside the folder, on a new line, define a new slider value c that ranges from a to b. On the line below, define m_{tan} as the slope of the tangent line at c as $m_{tan} = f'(c)$.

   (f) In a new folder called "Tangent Line," plot the point $(c, f(c))$ in red with size 20. In the space below, write down the equation for the tangent line at $x = c$. Your answer should only involve m_{tan}, c, and $f(c)$. Define this line in the folder and plot it only on the range $[a, b]$.

(g) Slide the value of c until m_{tan} is the same (or very close) to m_{sec}. What is this value? In the space below, find the exact value of this number using algebra and calculus. Show all work.. Once you have the answer, change c to be that value in Desmos to check your answer.

(h) When you submit this assignment on D2L, share the url to this graph.

2. Optimization: In this part, we'll maximize the area of a fenced in rectangular region given that we only have so much fencing. This will build a code to essentially solve any optimization problem of this form.
   (a) Open a new graph in Desmos and save it as "Optimization."
   (b) Suppose we have 40 ft of fencing and we wish to enclose a rectangular area A. We want to maximize the area. Draw a picture below, write down the objective function, and the constraint.

   (c) Use the constraint to eliminate one of your variables and write down the area function as a function of x only below. Define this function on the first line of Desmos as $A(x)$. Plot it in black and adjust the window so you can see it appropriately. Does this function have a maximum for positive values of x? Where does it seem to occur?

   (d) In the space below, confirm your guess above by finding the critical points of A. Confirm that it is a maximum by performing the second derivative test below.

(e) In the space below, find the dimensions of the rectangle and give the maximum area.

(f) Now, let's generalize this by changing the perimeter value from 40 to P in the function $A(x)$. Create a slider for P in Desmos and allow it to range from 0 to 100 with a stepsize of 0.1. Redo the problem, and find the critical point in terms of P below. What are the new values of x and y? What is the maximum area in terms of P. Plot this point on the curve. On a new line in Desmos, output the value of $A''(x)$ evaluated at the critical point in terms of P. Push play on P. As P is changing is the second derivative always negative? Explain why this is a good thing!

(g) When you submit this assignment on D2L, share the url to this graph.