# Non-Parametric Least Squares

## Based on Chapter 13 of *High Dimensional Statistics* by Martin Wainwright

Behrad Moniri

Department of Electrical Engineering
Sharif University of Technology

Statistics Reading Group
Tehran Institute for Advanced Studies (TeIAS)

**First Session:**


1. Introduction
2. Examples
3. Critical Radius: Bounding the Prediction Error
4. How to Compute the Critical Radius?


**Second Session:**


5. Review and Examples
6. Oracle Inequalities
7. Regularized Estimators

# Part I

## First Session

# Section 1

## Introduction

# Non-Parametric Least Squares

- **MMSE Estimation**:

$$\bar{\mathcal{L}}_f = \mathbb{E}_{X,Y}[(Y - f(X))^2] \implies f^\star(x) = \mathbb{E}[Y|X = x]$$

- In practice we are given a collection of samples $\{(\mathbf{x}_i, y_i)\}_{i=1}^n$, which can be used to compute an empirical analog of the MSE:

$$\widehat{\mathcal{L}}_f = \frac{1}{n}\sum_{i=1}^{n}\big(y_i - f(x_i)\big)^2$$

- **Non-Parametric Least Squares**: Minimizing this criterion over some suitably controlled function class.

# Different Measures of Quality

- **Excess Risk**:

$$||f - f^\star||_{L^2(\mathbb{P})} = \mathbb{E}_{X\sim\mathbb{P}}\Big[\big(f(X) - f^\star(X)\big)^2\Big]$$

where $\mathbb{P}$ denotes the distribution over covariates.

- **This Chapter**: Let $\{\mathbf{x}_i\}_{i=1}^n$ be the set of fixed covariates and $\mathbb{P}_n$ be their Empirical measure. Define:

$$||f - f^\star||_{L^2(\mathbb{P}_n)} = \left[\frac{1}{n}\sum_{i=1}^n \big(f(\mathbf{x}_i) - f^\star(\mathbf{x}_i)\big)^2\right]^{1/2}.$$

In this talks, we will use this measure and denote it as $||f - f^\star||_n$.

# Fixed or Stochastic Covariates?

**Note**

- We will view the samples $\{\mathbf{x}_i\}_{i=1}^n$ as being fixed, a set-up known as *regression with a fixed design*.
- Results from Chapter 14 to follow can be used to translate these bounds into equivalent results in the population $L_2(\mathbb{P})$-norm.

# Estimation via Least Squares

Given a fixed collection $\{\mathbf{x}_i\}_{i=1}^n$, model the responses as

$$y_i = f^\star(\mathbf{x}_i) + \nu_i, \quad \text{for } i = 1, 2, \ldots, n.$$

where $\nu_i = \sigma w_i$ in which $w_i \sim \mathcal{N}(0,1)$. The least squares estimate is given by the function

$$\widehat{f} \in \arg\min_{f \in \mathcal{F}} \left\{ \frac{1}{n} \sum_{i=1}^n \big(y_i - f(x_i)\big)^2 \right\}.$$

- When $\nu_i \sim \mathcal{N}(0, \sigma^2)$, the LS estimate is equivalent to the constrained maximum likelihood.
- When $\mathcal{F}$ is an RKHS, it can also be convenient to use regularized estimators of the form:

$$\widehat{f} \in \arg\min_{f \in \mathcal{F}} \left\{ \frac{1}{n} \sum_{i=1}^n \big(y_i - f(x_i)\big)^2 + \lambda_n ||f||_{\mathcal{F}}^2 \right\}.$$

Section 2

# Examples

# Parametric: Linear Regression

- For a given vector $\boldsymbol{\theta} \in \mathbb{R}^d$, define $f_{\boldsymbol{\theta}}(\mathbf{x}) = \langle \boldsymbol{\theta}, \mathbf{x} \rangle$ and consider the function class $\mathcal{F}_C = \{f_{\boldsymbol{\theta}} : \mathbb{R}^d \to \mathbb{R} | \boldsymbol{\theta} \in C\}$ for a compact $C$.
- The least squares estimate:

$$\widehat{\boldsymbol{\theta}} \in \arg\min_{\boldsymbol{\theta} \in C} \left\{ \frac{1}{n} ||y - \mathbf{X}\boldsymbol{\theta}||_2^2 \right\}$$

  where $\mathbf{X} \in \mathbb{R}^{n \times d}$ is the design matrix.
- The constrained form of Ridge Regression:

$$C = \left\{ \boldsymbol{\theta} \in \mathbb{R}^d \;\middle|\; ||\boldsymbol{\theta}||_2^2 \leq R \right\}.$$

- The constrained form of LASSO:

$$C = \left\{ \boldsymbol{\theta} \in \mathbb{R}^d \;\middle|\; ||\boldsymbol{\theta}||_1 \leq R \right\}.$$

# Non-Parametric: Cubic Smoothing Spline

Consider the class of twice cont. differentiable functions $f\colon [0,1] \to \mathbb{R}$ and for a given radius $R$, define:

$$\mathcal{F}(R) = \left\{ f\colon [0,1] \to \mathbb{R} \;\middle|\; \int_0^1 \left(f''(x)\right)^2 dx \leq R \right\}.$$

The integral can be understood as a Hilbert norm bound in the second-order Sobolev space.

Section 3

# Critical Radius: Bounding the Prediction Error

# Critical Radius: Bounding the Prediction Error

- Intuitively, the difficulty of estimating the function $f^\star$ should depend on the complexity of the function class $\mathcal{F}$ in which it lies.
- **a localized form of Gaussian complexity:** it measures the complexity of the function class $\mathcal{F}$, locally in a neighborhood around the true regression function $f^\star$.

## Local Gaussian Complexity

Define $f^\star = \mathcal{F} - \{f^\star\} = \{f - f^\star | f \in \mathcal{F}\}$. For a given radius $\delta > 0$, *the local Gaussian complexity* around $f^\star$ at scale $\delta$ is given by:

$$\mathcal{G}_n(\delta; f^\star) = \mathbb{E}_w \left[ \sup_{g \in \mathcal{F}^\star, \, ||g||_n \leq \delta} \left| \frac{1}{n} \sum_{i=1}^{n} w_i \, g(\mathbf{x}_i) \right| \right]$$

# Critical Inequality

## Critical Inequality

A central object in our analysis is the set of positive scalars $\delta$ that satisfy the *critical inequality*:

$$\frac{\delta}{2\sigma} \geq \frac{\mathcal{G}_n(\delta; \mathcal{F}^\star)}{\delta}$$

- We refer to any $\delta > 0$ satisfying the critical inequality as being valid.
- We use $\delta_n^\star$ to denote the smallest positive radius for which the critical inequality holds.

# Some Intuition

- Note that $\frac{1}{2n}\sum_{i=1}^{n}\left(y_i - \widehat{f}(\mathbf{x}_i)\right)^2 \leq \frac{1}{2n}\sum_{i=1}^{n}\left(y_i - f^\star(\mathbf{x}_i)\right)^2$.
- After some computations: $\frac{1}{2}||\widehat{f} - f^\star||_n^2 \leq \frac{\sigma}{n}\sum_{i=1}^{n} w_i\left(\widehat{f}(\mathbf{x}_i) - f^\star(\mathbf{x}_i)\right)$.
- Note that $\widehat{f} - f^\star \in f^\star$. Reasoning hueristically, this observation suggests that

$$\frac{\delta^2}{2} \leq \sigma\mathcal{G}_n(\delta;\mathcal{F}^\star) \text{ or equivalently } \frac{\delta}{2\sigma} \leq \frac{\mathcal{G}_n(\delta;\mathcal{F}^\star)}{\delta}.$$

- We will present a rigorous result later in this talk.

# The Main Result

## Star-Shaped Classes

a function class $\mathcal{F}$ is star-shaped if for any $\alpha \in [0, 1]$ we have

$$f \in \mathcal{F} \implies \alpha f \in \mathcal{F}.$$

## Theorem

Suppose that the shifted function class $\mathcal{F}^\star$ is star-shaped, and let $\delta$ be any solution to the critical inequality. Then for any $t \geq \delta$, the nonparametric least-squares estimate $\widehat{f}_n$ satisfies the bound

$$\mathbb{P}\left[||\widehat{f}_n - f^\star||_n^2 \geq 16t\delta\right] \leq \exp\left(\frac{-nt\delta}{2\sigma^2}\right) \tag{1}$$

# Properties of Star-Shaped Classes

- The star shaped condition on $\mathcal{F}^\star$ is needed in various parts of the proof, including ensuring a valid radii $\delta$ exists.
- $\mathcal{F}^\star$ Star shaped $=$ $\mathcal{F}$ is star shaped around $f^\star$.

### Property

If $\mathcal{F}$ is convex, it is star-shaped around any $f^\star$.

### Proof

Let $\widetilde{f} = f - f^\star \in \mathcal{F}^\star$ be an arbitrary point in $\mathcal{F}^\star$ with $f, f^\star \in \mathcal{F}$.
For any $\alpha \in [0, 1]$, by convexity we have $g = \alpha f + (1 - \alpha) f^\star \in \mathcal{F}$.
Hence $\alpha \widetilde{f} = \alpha(f - f^\star) = g - f^\star \in \mathcal{F}^\star$, proving $\mathcal{F}^\star$ is star-shaped.

### Conversely

Similarly if $\mathcal{F}$ is not convex, there must exists a $f^\star$ such that $\mathcal{F}$ is not star shaped around $f^\star$.

# Properties of Star-Shaped Classes

- If the star-shaped condition fails to hold, then the main Theorem can instead by applied with $\delta$ defined in terms of the star hull (we will see next session.)

$$\mathrm{star}(\mathcal{F}^\star) = \{\alpha(f - f^\star) | f \in \mathcal{F}, \alpha \in [0,1]\}.$$

## Existence of Critical Radius

For any star shaped class $\mathcal{F}^\star$, the function $\delta \to \frac{\mathcal{G}_n(\delta, \mathcal{F}^\star)}{\delta}$ is non-increasing on $(0, \infty)$. Consequently, for any $c > 0$, the inequality

$$\frac{\mathcal{G}_n(\delta, \mathcal{F}^\star)}{\delta} \leq c\delta$$

has a smallest positive solution.

# Proof of the Existence of Critical Radius

Given $0 < \delta \leq t$, we should show that $\frac{\delta}{t}\mathcal{G}_n(t) \leq \mathcal{G}_n(\delta)$.
Given any $h \in \mathcal{F}^\star$ with $||h||_n \leq t$, define $\widetilde{h} = \frac{\delta}{t}h \in \mathcal{F}^\star$ (by star-shaped assumption) and write

$$\frac{1}{n}\Big\{\frac{\delta}{t}\sum_{i=1}^{n} w_i h(\mathbf{x}_i)\Big\} = \frac{1}{n}\Big\{\sum_{i=1}^{n} w_i \widetilde{h}(\mathbf{x}_i)\Big\}.$$

By construction, $||\widetilde{h}||_n \leq \delta$. Consequently:

- The RHS is at most $\mathcal{G}_n(\delta)$ in expectation.
- Taking supremum over the set $\mathcal{F} \cap \{||h||_n \leq t\}$ following by expectation, yields $\frac{\delta}{t}\mathcal{G}_n(t)$ on the left hand side.

This concludes the proof. $\square$




**Figure 13.2** Illustration of the critical radius for sample size $n = 100$ and two different function classes. (a) A first-order Sobolev space. (b) A Gaussian kernel class. In both cases, the function $\delta \mapsto \frac{\mathcal{G}_n(\delta;\mathscr{F})}{\delta}$, plotted as a solid line, is non-increasing, as guaranteed by Lemma 13.6. The critical radius $\delta_n^*$, marked by a gray dot, is determined by finding its intersection with the line of slope $1/(2\sigma)$ with $\sigma = 1$, plotted as the dashed line. The set of all valid $\delta_n$ consists of the interval $[\delta_n^*, \infty)$.

# Proof of the main theorem

**Theorem**

Suppose that the shifted function class $\mathcal{F}^\star$ is star-shaped, and let $\delta$ be any solution to the critical inequality. Then for any $t \geq \delta$, the nonparametric least-squares estimate $\widehat{f}_n$ satisfies the bound

$$\mathbb{P}\big[||\widehat{f}_n - f^\star||_n^2 \geq 16t\delta\big] \leq \exp\left(\frac{-nt\delta}{2\sigma^2}\right)$$

- Denote $\widehat{\Delta} = \widehat{f} - f^\star \in \mathcal{F}^\star$, we have $\frac{1}{2}||\widehat{\Delta}||_n^2 \leq \frac{\sigma}{n}\sum_{i=1}^n w_i \widehat{\Delta}(\mathbf{x}_i)$. We need to control the stochastic component of the right-hand side.

## Auxiliary Lemma

Let $\mathcal{F}^\star$ be a star-shaped class and $\delta$ satisfy the critical inequality. Define the bad event $A(u)$ as

$$A(u) = \left\{ \exists g \in \mathcal{F}^\star, ||g||_n \geq u \ : \ \left| \frac{\sigma}{n} \sum_{i=1}^n w_i g(\mathbf{x}_i) \right| \geq 2u||g||_n \right\}.$$

For a given $u \geq \delta$, we have $\mathbb{P}[A(u)] \leq \exp\left(\frac{-nu^2}{2\sigma^2}\right)$.

Now consider two cases:

- **Case 1:** $||\widehat{\Delta}||_n < \sqrt{t\delta}$ which implies $||\widehat{\Delta}||_n^2 \leq t\delta$ trivially.
- **Case 2:** $||\widehat{\Delta}||_n \geq \sqrt{t\delta}$. We condition on $A^c(\sqrt{t\delta})$. Set $u = \sqrt{t\,\delta}$, so that we have $\mathbb{P}[A^c(\sqrt{t\,\delta})] \geq 1 - \exp(\frac{-nt\delta}{2\sigma^2})$. Hence:

$$||\widehat{\Delta}||_n^2 \leq 2\left|\frac{\sigma}{n}\sum_{i=1}^n w_i \widehat{\Delta}(\mathbf{x}_i)\right| \leq 4||\widehat{\Delta}||_n\sqrt{t\delta} \implies ||\Delta||_n^2 \leq 16t\delta. \quad \square$$

# Proof of the Auxiliary Lemma

Recall $A(u) = \left\{ \exists g \in \mathcal{F}^\star, ||g||_n \geq u \ : \ \left| \frac{\sigma}{n} \sum_{i=1}^n w_i g(\mathbf{x}_i) \right| \geq 2u||g||_n \right\}$.

**Step 1**: reduce to controlling a sup over a subset with $||\widetilde{g}||_n \leq u$.

- Suppose that $A(u)$ is true, hence there exists $g \in \mathcal{F}^\star$ with $||g||_n \geq u$ such that

$$\left| \frac{\sigma}{n} \sum_{i=1}^n w_i g(\mathbf{x}_i) \right| \geq 2||g||_n u.$$

- Define $\widetilde{g} = \frac{u}{||g||_n} g \in \mathcal{F}^\star$, we observe that $||\widetilde{g}||_n = u$. We have

$$\left| \frac{\sigma}{n} \sum_{i=1}^n w_i \widetilde{g}(\mathbf{x}_i) \right| = \frac{u}{||g||_n} \left| \frac{\sigma}{n} \sum_{i=1}^n w_i g(\mathbf{x}_i) \right| \geq 2u^2.$$

- We thus conclude that:

$$\mathbb{P}[A(u)] \leq \mathbb{P}[Z_n(u) \geq 2u^2], \ Z_n(u) := \sup_{\widetilde{g} \in \mathcal{F}^\star, \ ||\widetilde{g}||_n \leq u} \left| \frac{\sigma}{n} \sum_{i=1}^n w_i \widetilde{g}(\mathbf{x}_i) \right|.$$

# Proof of the Auxiliary Lemma

$\mathbb{P}[A(u)] \leq \mathbb{P}[Z_n(u) \geq 2u^2]$, $Z_n(u) := \sup_{\widetilde{g} \in \mathcal{F}^\star, \quad ||\widetilde{g}||_n \leq u} \left| \frac{\sigma}{n} \sum_{i=1}^n w_i \widetilde{g}(\mathbf{x}_i) \right|$.

**Step 2**: Concentration of suprema.

- Recall that $w_i \sim \mathcal{N}(0,1)$, hence $\frac{\sigma}{n} \sum_{i=1}^n w_i \widetilde{g}(\mathbf{x}_i) \sim \mathcal{N}\left(0, \frac{\sigma^2}{n} ||\widetilde{g}||_n^2\right)$.
- $Z_n(u) = h(w_1, \ldots, w_n) := \sup_{\widetilde{g} \in \mathcal{F}^\star, \quad ||\widetilde{g}||_n \leq u} \left| \frac{\sigma}{n} \sum_{i=1}^n w_i \widetilde{g}(\mathbf{x}_i) \right|$.
- The Lipschitz constant of $h$ is at most $\frac{\sigma u}{\sqrt{n}}$.
- We know that for $Z_i \sim \mathcal{N}(0, \sigma^2)$ and $h$ a L-Lipschitz function:

$$\mathbb{P}[h(Z) - \mathbb{E}h(Z) \geq t] \leq e^{-\frac{s^2}{2L^2\sigma^2}}.$$

- Let $s = u^2$, we obtain $\mathbb{P}(Z_n(u) \geq \mathbb{E}[Z_n(u)] + u^2) \leq \exp\left(\frac{-nu^2}{2\sigma^2}\right)$.

# Proof of the Auxiliary Lemma

**Step 3**: Bounding the expectation.

- Note that $\mathbb{E}[Z_n(u)] = \sigma\, \mathcal{G}_n(u)$.
- We know that $\nu \to \frac{\mathcal{G}_n(\nu)}{\nu}$ is non-increasing. We have also assumed that $u \geq \delta$.

$$\sigma \frac{\mathcal{G}_n(u)}{u} \leq \sigma \frac{\mathcal{G}_n(\delta)}{\delta} \leq \frac{\delta}{2} \leq \delta \implies \mathbb{E}[Z_n(u)] \leq u\delta \leq u^2.$$

Comibining our results, we obtain

$$\mathbb{P}[A(u)] \leq \mathbb{P}[Z_n(u) \geq 2u^2] \leq \mathbb{P}(Z_n(u) \geq \mathbb{E}[Z_n(u)] + u^2) \leq \exp\left(\frac{-nu^2}{2\sigma^2}\right),$$

Which concludes the proof. $\square$

Section 4

# How to Compute the Critical Radius?

# Bounds via Metric Entropy

For any star-shaped function class $\mathcal{F}^\star$, define:

- $B(\delta\,;\mathcal{F}^\star) = \{h \in \mathcal{F}^\star \mid ||h||_n \leq \delta\}$
- $\mathcal{N}\left(t; B_n(\delta; \mathcal{F}^\star)\right)$ be the $t$-convering number of $B_n(\delta; \mathcal{F})$ in norm $||.||_n$.

## Critical Inequality via Metric Entropy

Any $\delta \in (0, \infty]$ satisfying

$$\frac{16}{\sqrt{n}} \int_{\frac{\delta^2}{4\sigma}}^{\delta} \sqrt{\log \mathcal{N}\left(t\,; B(\delta; \mathcal{F}^\star)\right)}\, dt \leq \frac{\delta^2}{4\sigma},$$

satisfies the critical inequality.

# Bounds via Metric Entropy: Proof

- Construct a $\frac{\delta^2}{4\sigma}$-covering of the set $B(\delta;\mathcal{F})$, say $\{g^1,\ldots,g^M\}$. For any function $g \in B(\delta;\mathcal{F})$, there is a $j \in [M]$ such that $||g^j - g||_n \leq \frac{\delta^2}{4\sigma}$.
- We have

$$\begin{aligned}
\left|\frac{1}{n}\sum_{i=1}^n w_i\, g(\mathbf{x}_i)\right| &\leq \left|\frac{1}{n}\sum_{i=1}^n w_i\, g^j(\mathbf{x}_i)\right| + \left|\frac{1}{n}\sum_{i=1}^n w_i\left(g(\mathbf{x}_i) - g^j(\mathbf{x}_i)\right)\right| \\
&\leq \max_{j=1,\ldots,M}\left|\frac{1}{n}\sum_{i=1}^n w_i\, g^j(\mathbf{x}_i)\right| + \sqrt{\frac{\sum_{i=1}^n {w_i}^2}{n}}\sqrt{\frac{\sum_{i=1}^n (g(\mathbf{x}_i) - g^k(\mathbf{x}_i))^2}{n}} \\
&\leq \max_{j=1,\ldots,M}\left|\frac{1}{n}\sum_{i=1}^n w_i\, g^j(\mathbf{x}_i)\right| + \frac{\delta^2}{4\sigma}\sqrt{\frac{\sum_{i=1}^n {w_i}^2}{n}}.
\end{aligned}$$

Hence

$$\mathcal{G}_n(\delta) \leq \mathbb{E}\left[\max_{j=1,\ldots,M}\left|\frac{1}{n}\sum_{i=1}^n w_i\, g^j(\mathbf{x}_i)\right|\right] + \frac{\delta^2}{4\sigma}.$$

# Bounds via Metric Entropy: Proof

- To upper bound the first term, we use Dudley's Integral. Define $Z(g^j) = \frac{1}{\sqrt{n}} \sum_{i=1}^{n} w_i g^j(\mathbf{x}_i)$ for $j = 1, \ldots, M$.
- The induced metric: $\rho_Z^2(g^j, g^k) := \mathrm{Var}\big(Z(g^j) - Z(g^k)\big) = ||g^j - gk||_n^2$
- Since $||g||_n \leq \delta$ for all $g \in B_n(\delta, \mathcal{F}^*)$, the coarsest resolution of the chaining can be set to $\delta$. We can terminate it at $\frac{\delta^2}{4\sigma}$ since any member of our finite set can be reconstructed at this resolution.

$$\mathbb{E}\left[\max_{j=1,\ldots,M}\left|\frac{1}{n}\sum_{i=1}^{n} w_i g^j(\mathbf{x}_i)\right|\right] = \mathbb{E}\left[\max_{j=1,\ldots,M}\frac{\left|Z(g^j)\right|}{\sqrt{n}}\right]$$
$$\leq \frac{16}{\sqrt{n}}\int_{\frac{\delta^2}{4\sigma}}^{\delta}\sqrt{\log N_n(t; B_n(\delta; \mathcal{F}))}dt$$

Hence $\mathcal{G}_n(\delta) \leq \frac{\delta^2}{4\sigma} + \frac{16}{\sqrt{n}}\int_{\frac{\delta^2}{4\sigma}}^{\delta}\sqrt{\log N_n(t; B_n(\delta; \mathcal{F}))}dt$. □

# Part II

## Second Session

# Section 5

# Review and Examples

# Review: Estimator, Models, and Definitions

## Model

Let $\mathcal{F}$ be a function class and $f^\star \in \mathcal{F}$ and $\{\mathbf{x}_i\}_{i=1}^n$ be a set of $n$ covariates. Assume that $y_i = f^\star(\mathbf{x}_i) + \sigma w_i$ where $w_i \sim \mathcal{N}(0,1)$.

## Estimator

Given the set $\{(\mathbf{x}_1, y_1), \ldots, (\mathbf{x}_n, y_n)\}$:

$$\widehat{f} \in \arg\min_{f \in \mathcal{F}} \frac{1}{n} \sum_{i=1}^n \Big(y_i - f(\mathbf{x}_i)\Big)^2.$$

## Measure of Goodness

Let $\{\mathbf{x}_i\}_{i=1}^n$ be the set of fixed covariates:

$$||f - f^\star||_n := \left[\frac{1}{n} \sum_{i=1}^n \big(f(\mathbf{x}_i) - f^\star(\mathbf{x}_i)\big)^2\right]^{1/2}.$$

## Star-Shaped Classes

A function class $\mathcal{F}$ is star-shaped if for any $\alpha \in [0,1]$ we have

$$f \in \mathcal{F} \implies \alpha f \in \mathcal{F}.$$

## Critical Inequality

A central object in our analysis is the set of positive scalars $\delta$ that satisfy the *critical inequality*:

$$\frac{\delta}{2\sigma} \geq \frac{\mathcal{G}_n(\delta; \mathcal{F}^\star)}{\delta}$$

## Existence of Critical Radius

For any star shaped class $\mathcal{F}^\star$, the function $\delta \to \frac{\mathcal{G}_n(\delta, \mathcal{F}^\star)}{\delta}$ is non-increasing on $(0, \infty)$. Consequently, for any $c > 0$, the inequality $\frac{\mathcal{G}_n(\delta, \mathcal{F}^\star)}{\delta} \leq c\delta$ has a smallest positive solution.

# Review: Main Results

### Theorem

Suppose that the shifted function class $\mathcal{F}^\star$ is star-shaped, and let $\delta$ be any solution to the critical inequality. Then for any $t \geq \delta$, the nonparametric least-squares estimate $\widehat{f}_n$ satisfies the bound

$$\mathbb{P}\left[||\widehat{f}_n - f^\star||_n^2 \geq 16t\delta\right] \leq \exp\left(\frac{-nt\delta}{2\sigma^2}\right)$$

### Auxiliary Lemma

Let $\mathcal{F}^\star$ be a star-shaped class and $\delta$ satisfy the critical inequality. Define the bad event $A(u)$ as

$$A(u) = \left\{\exists g \in \mathcal{F}^\star, ||g||_n \geq u \ : \ \left|\frac{\sigma}{n}\sum_{i=1}^{n} w_i g(\mathbf{x}_i)\right| \geq 2u||g||_n\right\}.$$

For a given $u \geq \delta$, we have $\mathbb{P}[A(u)] \leq \exp\left(\frac{-nu^2}{2\sigma^2}\right)$.

# Review: How to Compute a Critical Radius

For any star-shaped function class $\mathcal{F}^\star$, define:

- $B(\delta\,;\mathcal{F}^\star) = \{h \in \mathcal{F}^\star \mid ||h||_n \leq \delta\}$
- $\mathcal{N}\left(t; B(\delta; \mathcal{F}^\star)\right)$ be the $t$-convering number of $B(\delta; \mathcal{F})$ in norm $||.||_n$.

## Critical Inequality via Metric Entropy

Any $\delta \in (0, \infty]$ satisfying

$$\frac{16}{\sqrt{n}} \int_{\frac{\delta^2}{4\sigma}}^{\delta} \sqrt{\log \mathcal{N}\left(t\,; B(\delta; \mathcal{F}^\star)\right)}\, dt \leq \frac{\delta^2}{4\sigma},$$

satisfies the critical inequality.

Section 6

# Oracle Inequalities

# Oracle Inequalities

- In our analysis so far, we have assumed that $f^\star \in \mathcal{F}$. If we lift this assumption, we have:
  - Prediction Error
  - Approximation Error
- It is natural to measure the approximation error in terms of the best approximating function in $\mathcal{F}$, i.e. $\inf_{f \in \mathcal{F}} ||f - f^\star||_n$.
- Only an oracle that has access to $f^\star(\mathbf{x}_i)$ can calculate $\inf_{f \in \mathcal{F}} ||f - f^\star||_n$. Hence we call inequalities with $\inf_{f \in \mathcal{F}} ||f - f^\star||_n$, the *Oracle Inequalities*.

# Oracle Inequalities

Assume that $y_i = f^\star(\mathbf{x}_i) + \sigma w_i$, where $w_i \sim \mathcal{N}(0,1)$.
Define $\partial\mathcal{F} = \{f_1 - f_2 | f_1, f_2 \in \mathcal{F}\}$. We assume that this set is star-shaped.

## Oracle Inequality

Let $\delta$ be any positive solution to the critical inequality $\frac{\mathcal{G}_n(\delta,\partial\mathcal{F})}{\delta} \leq \frac{\delta}{2\sigma}$.
For any $t \geq \delta$, the non-parametric least-square estimate $\widehat{f}$ satisfies the bound

$$||\widehat{f} - f^\star||_n^2 \leq \inf_{\gamma\in(0,1)} \left\{ \frac{1+\gamma}{1-\gamma} ||f - f^\star||_n^2 + \frac{c_0}{\gamma(1-\gamma)} t\delta \right\} \text{ for all } f \in \mathcal{F},$$

with probability at least $1 - c_1 \exp(-\frac{c_2 n t \delta}{\sigma^2})$.

When $f^\star \in \mathcal{F}$, we can set $f = f^\star$. Hence, $||\widehat{f} - f^\star||_n^2 \lesssim t\delta$, recovering the previous result up to constants.

# Oracle Inequalities

$$||\widehat{f} - f^\star||_n^2 \leq \inf_{\gamma \in (0,1)} \left\{ \frac{1+\gamma}{1-\gamma} ||f - f^\star||_n^2 + \frac{c_0}{\gamma(1-\gamma)} t\delta \right\} \text{ for all } f \in \mathcal{F}.$$

- When $f^\star \notin \mathcal{F}$, setting $t = \delta$ and taking infimum over $f \in \mathcal{F}$, yields

$$||\widehat{f} - f^\star||_n^2 \lesssim \inf_{f \in \mathcal{F}} ||\widehat{f} - f^\star||_n^2 + \delta^2$$

with high probability. The term $\delta^2$ can be viewed as the estimation error and $\inf_{f \in \mathcal{F}} ||\widehat{f} - f^\star||_n^2$ as the approximation error.

# Oracle Inequalities: Proof

- Define $\widehat{\Delta} = \widehat{f} - f^\star$ and $\widetilde{\Delta} = \widehat{f} - \widetilde{f}$.
- Given an arbitrary $\widetilde{f} \in \mathcal{F}$, we have

$$0 \geq \frac{1}{2n}\sum_{i=1}^{n}\left(y_i - \widehat{f}_i\right)^2 - \frac{1}{2n}\sum_{i=1}^{n}\left(y_i - \widetilde{f}_i\right)^2$$

$$= \frac{1}{2}||\widehat{\Delta}||_n^2 + \frac{1}{2n}\sum_{i=1}^{n}\left(y_i - f_i{}^\star\right)^2 + \frac{1}{n}\sum_{i=1}^{n}(y_i - f_i{}^\star)(f_i{}^\star - \widehat{f}_i) - \frac{1}{2n}\sum_{i=1}^{n}\left(y_i - \widetilde{f}_i\right)^2$$

Hence $\frac{1}{2}||\widehat{\Delta}||_n^2 \leq \frac{1}{2n}\sum_{i=1}^{n}\left(\widetilde{\Delta}_i - \widehat{\Delta}_i\right)(2y_i - \widetilde{f}_i - f_i{}^\star) - \frac{1}{n}\sum_{i=1}^{n}\sigma w_i(f_i^\star - \widehat{f}_i)$.

$$\frac{1}{2}||\widehat{\Delta}||_n^2 \leq \frac{1}{2n}\sum_{i=1}^{n}\left[\left(\widetilde{\Delta}_i - \widehat{\Delta}_i\right)(2\sigma w_i - \widetilde{\Delta}_i - \widehat{\Delta}_i) + 2\sigma w_i\widehat{\Delta}_i\right]$$

$$\leq \frac{1}{n}\sum_{i=1}^{n}\sigma w_i\widetilde{\Delta}_i + \frac{1}{2}||f_i^\star - \widetilde{f}_i||_n^2 \leq \frac{1}{2}\,||f_i^\star - \widetilde{f}_i\,||_n^2 + \frac{\sigma}{n}\left|\sum_{i=1}^{n} w_i\widetilde{\Delta}_i\right|.$$

# Oracle Inequalities: Proof

To bound the error, we consider two cases:

- **Case 1**: Suppose that $||\widehat{\Delta}||_n^2 \leq \sqrt{t\delta}$.

$$||\widehat{\Delta}||_n^2 = ||(\widetilde{f} - f^\star) + \widetilde{\Delta}||_n^2 \leq \left(||\widetilde{f} - f^\star||_n + \sqrt{t\delta}\right)^2$$
$$\leq (1 + 2\beta)||\widetilde{f} - f^\star||_n^2 + \left(1 + \frac{2}{\beta}\right) t\delta.$$

Set $\beta = \frac{\gamma}{1-\gamma}$. This yields

$$||\widehat{f} - f^\star||_n^2 \leq \inf_{\gamma \in (0,1)} \left\{ \frac{1+\gamma}{1-\gamma} ||\widetilde{f} - f^\star||_n^2 + \frac{c_0}{\gamma(1-\gamma)} t\delta \right\}.$$

- **Case 2**: Suppose that $||\widehat{\Delta}||_n^2 > \sqrt{t\delta}$.

### Auxiliary Lemma

Let $\mathcal{H}$ be a star-shaped class and $\delta$ satisfy the critical inequality:

$$\forall u \geq \delta : \quad \mathbb{P}\left[\exists\, g \in \mathcal{H}, ||g||_n \geq u \ : \ \left|\frac{\sigma}{n}\sum_{i=1}^{n} w_i g(\mathbf{x}_i)\right| \geq 2u||g||_n\right] \leq \exp\left(\frac{-nu^2}{2\sigma^2}\right).$$

Let $u = \sqrt{t\delta}$ and $\mathcal{H} = \partial\mathcal{F}$. We have

$$\mathbb{P}\left[2\left|\frac{\sigma}{n}\sum_{i=1}^{n} w_i \widetilde{\Delta}(\mathbf{x}_i)\right| \geq 4\sqrt{t\delta}||\widetilde{\Delta}||_n\right] \leq \exp\left(-\frac{nt\delta}{2\sigma^2}\right).$$

Hence

$$||\widehat{\Delta}||_n^2 \leq ||\widetilde{f} - f^\star||_n^2 + 4\sqrt{t\delta}||\widetilde{\Delta}||_n^2 \leq ||\widetilde{f} - f^\star||_n^2 + 4\sqrt{t\delta}\left\{||\widehat{\Delta}||_n^2 + ||\widetilde{f} - f^\star||_n^2\right\}$$

$$\leq ||\widetilde{f} - f^\star||_n^2 + \left[4\beta||\widehat{\Delta}||_n^2 + \frac{4}{\beta}t\delta_n\right] + \left[4\beta||\widehat{\Delta}||_n^2 + \frac{4}{\beta}t\delta\right].$$

Rearrenging terms: $||\widehat{\Delta}||_n^2 \leq \frac{1+4\beta}{1-4\beta}||\widetilde{f} - f^\star||_n^2 + \frac{8}{\beta(1-4\beta)}t\delta.$ $\square$

Section 7

# Regularized Estimators

# Oracle Inequalities for Regularized Estimators

- Given a space $\mathcal{F}$ of real-valued functions, an an associated norm $||.||_{\mathcal{F}}$, consider the family of regularized least-square problems:

$$\widehat{f} \in \arg\min_{\mathcal{F}} \left\{ \frac{1}{2n} \sum_{i=1}^{n} \left(y_i - f(\mathbf{x}_i)\right)^2 + \lambda_n ||f||_{\mathcal{F}}^2 \right\}.$$

- A Local Gaussian Complexity Measure:

$$\mathcal{G}_n(\delta) \triangleq \mathcal{G}_n\big(\delta; B_{\partial\mathcal{F}}(3)\big) = \mathbb{E}_w \left[ \sup_{\substack{g \in \partial\mathcal{F} \\ ||g||_n \leq \delta \\ ||g||_{\mathcal{F}} \leq 3}} \left| \frac{1}{n} \sum_{i=1}^{n} w_i f(\mathbf{x}_i) \right| \right]$$

- The critical inequality for a user defined $R$:

$$\frac{\mathcal{G}_n(\delta)}{\delta} \leq \frac{R}{2\sigma}\delta.$$

# Oracle Inequalities for Regularized Estimators

**Theorem**

Consider a convex function class $\mathcal{F}$ and assume that $\delta$ satisfies the critical inequality

$$\frac{\mathcal{G}_n(\delta)}{\delta} \leq \frac{R}{2\sigma}\delta.$$

If $\lambda_n \geq 2\delta^2$. There exists universal constants $c_0, c_1, c_2, c_3$ such that

$$||\widehat{f} - f^\star||_n^2 \leq c_0 \inf_{||f||_{\mathcal{F}} \leq R} ||f - f^\star||_n^2 + c_1 R^2(\delta^2 + \lambda)$$

with probability grater than $1 - c_2 \exp\big(-c_3 \frac{nR^2\delta^2}{\sigma^2}\big)$.

# Oracle Inequalities for Regularized Estimators: Proof

- $y_i = f^\star(\mathbf{x}_i) + \sigma w_i$. Rescale the model by $R$.
  - Rescaled Noise variance: $\tilde{\sigma}^2 = \left(\frac{\sigma}{R}\right)^2$.
  - Rescaled approx. error: $\inf_{||f||_{\mathcal{F}} \leq 1} ||f - f^\star||_n^2$.
  - The final MSE should be multiplied by $R^2$.
- Let $\widetilde{f}$ be an arbitrary element in $\mathcal{F}$ with $||\widetilde{f}||_{\mathcal{F}} \leq 1$.
- We have

$$\frac{1}{2}\sum_{i=1}^{n}\left(y_i - \widehat{f}(\mathbf{x}_i)\right)^2 + \lambda_n ||\widehat{f}||_{\mathcal{F}}^2 \leq \frac{1}{2}\sum_{i=1}^{n}\left(y_i - \widetilde{f}(\mathbf{x}_i)\right)^2 + \lambda_n ||\widetilde{f}||_{\mathcal{F}}^2.$$

- Denote $\widehat{\Delta} = \widehat{f} - f^\star$ and $\widetilde{\Delta} = \widehat{f} - \widetilde{f}$. With a simple calculation (next slide):

$$\frac{1}{2}||\widehat{\Delta}||_n^2 \leq \frac{1}{2}||\widetilde{f} - f^\star||_n^2 + \frac{\widetilde{\sigma}}{n}\left|\sum_{i=1}^{n} w_i \widetilde{\Delta}(\mathbf{x}_i)\right| + \lambda_n \underbrace{\left[||\widetilde{f}||_{\mathcal{F}}^2 - ||\widehat{f}||_{\mathcal{F}}^2\right]}_{\leq 1}.$$

- Given an arbitrary $\widetilde{f} \in \mathcal{F}$, we have

$$0 \geq \frac{1}{2n}\sum_{i=1}^{n}\left(y_i - \widehat{f}_i\right)^2 + \lambda_n||\widehat{f}||_n^2 - \frac{1}{2n}\sum_{i=1}^{n}\left(y_i - \widetilde{f}_i\right)^2 - \lambda_n||\widetilde{f}||_n^2$$

$$= \frac{1}{2}||\widehat{\Delta}||_n^2 + \frac{1}{2n}\sum_{i=1}^{n}\left(y_i - f_i{}^\star\right)^2 + \frac{1}{n}\sum_{i=1}^{n}(y_i - f_i{}^\star)(f_i{}^\star - \widehat{f}_i)$$

$$- \frac{1}{2n}\sum_{i=1}^{n}\left(y_i - \widetilde{f}_i\right)^2 + \lambda_n||\widehat{f}||_n^2 - \lambda_n||\widetilde{f}||_n^2$$

- Rearranging terms:

$$\frac{1}{2}||\widehat{\Delta}||_n^2 \leq \frac{1}{2n}\sum_{i=1}^{n}(\widetilde{\Delta}_i - \widehat{\Delta}_i)(2y_i - \widetilde{f}_i - f_i{}^\star) - \frac{1}{n}\sum_{i=1}^{n}\sigma w_i(f_i^\star - \widehat{f}_i) + \lambda_n||\widetilde{f}||_n^2 - \lambda_n$$

$$\leq \frac{1}{2n}\sum_{i=1}^{n}\left[(\widetilde{\Delta}_i - \widehat{\Delta}_i)(2\sigma w_i - \widetilde{\Delta}_i - \widehat{\Delta}_i) + 2\sigma w_i\widehat{\Delta}_i\right] + \lambda_n||\widetilde{f}||_n^2 - \lambda_n||\widehat{f}||_n^2$$

$$\leq \frac{1}{n}\sum_{i=1}^{n}\sigma w_i\widetilde{\Delta}_i + \frac{1}{2}||f_i^\star - \widetilde{f}_i||_n^2 + \lambda_n||\widetilde{f}||_n^2 - \lambda_n||\widehat{f}||_n^2$$

$$\leq \frac{1}{2}||f_i^\star - \widetilde{f}_i||_n^2 + \frac{\sigma}{n}\left|\sum_{i=1}^{n} w_i\widetilde{\Delta}_i\right| + \lambda_n||\widetilde{f}||_n^2 - \lambda_n||\widehat{f}||_n^2.$$

# Parts of Proof

- If $||\widetilde{\Delta}||_n \leq \sqrt{t\delta}$ : It is trivial.
- Assume that $||\widetilde{\Delta}||_n > \sqrt{t\delta}$ .
  - **Case 1** Suppose that $||\widehat{f}||_{\mathcal{F}} \leq 2$.
  - **Case 2** Suppose that $||\widehat{f}||_{\mathcal{F}} > 2$.

**Case 1**: $||\widehat{f}||_{\mathcal{F}} \leq 2$.

$$\tfrac{1}{2}||\widehat{\Delta}||_n^2 \leq \tfrac{1}{2}\,||f_i^\star - \widetilde{f}_i\,||_n^2 + \tfrac{\sigma}{n}\Big|\sum_{i=1}^n w_i\widetilde{\Delta}_i\Big| + \lambda_n||\widetilde{f}||_n^2 - \lambda_n||\widehat{f}||_n^2.$$

## Auxiliary Lemma

Let $\mathcal{H}$ be a star-shaped class and $\delta$ satisfy the critical inequality:

$$\forall u \geq \delta: \quad \mathbb{P}\Big[\exists\, g \in \mathcal{H}, ||g||_n \geq u \;:\; \Big|\frac{\sigma}{n}\sum_{i=1}^n w_i g(\mathbf{x}_i)\Big| \geq 2u||g||_n\Big] \leq \exp\big(\frac{-nu^2}{2\sigma^2}\big).$$

Let $||\widetilde{f}||_{\mathcal{F}} \leq 1$, hence $||\widetilde{\Delta}||_{\mathcal{F}} \leq 3$. Apply lemma to $\{g \in \partial\mathcal{F} \;\big|\; ||g||_{\mathcal{F}} \leq 3\}$:

$$\frac{\widetilde{\sigma}}{n}\Big|\sum_{i=1}^n w_i\widetilde{\Delta}(\mathbf{x}_i)\Big| \leq c_0\sqrt{t\delta}||\widetilde{\Delta}||_n, \text{ with prob. at least } 1 - \exp\big(-\frac{t^2}{2\tilde{\sigma}^2}\big).$$

By triangle inequality:

$$\begin{aligned} 2\sqrt{t\delta}\;||\widetilde{\Delta}||_n^2 &\leq 2\sqrt{t\delta}\;||\widehat{\Delta}||_n^2 + 2\sqrt{t\delta}||\widetilde{f} - f^\star||_n \\ &\leq 2\sqrt{t\delta}\;||\widehat{\Delta}||_n^2 + 2t\delta + \frac{||\widetilde{f} - f^\star||_n^2}{2} \end{aligned}$$

Hence $\frac{1}{2}||\widehat{\Delta}||_n^2 \leq \frac{1}{2}(1+c_0)||\widetilde{f} - f^\star||_n^2 + 2c_0 t\delta + 2c_0\sqrt{t\delta}||\widehat{\Delta}||_n + \lambda$

**Case 2**: $||\widehat{f}||_{\mathcal{F}} > 2 > 1 \geq ||\widetilde{f}||_{\mathcal{F}}$. We have

$$||\widetilde{f}||_{\mathcal{F}}^2 - ||\widehat{f}||_{\mathcal{F}}^2 = \underbrace{\Big[||\widetilde{f}||_{\mathcal{F}} + ||\widehat{f}||_{\mathcal{F}}\Big]}_{>1} \cdot \underbrace{\Big[||\widetilde{f}||_{\mathcal{F}} - ||\widehat{f}||_{\mathcal{F}}\Big]}_{<0} \leq \underbrace{\Big[||\widetilde{f}||_{\mathcal{F}} + ||\widehat{f}||_{\mathcal{F}}\Big]}_{<0}.$$

Writing $\widehat{f} = \widetilde{f} + \widetilde{\Delta} \implies ||\widehat{f}||_{\mathcal{F}} \geq ||\widetilde{\Delta}||_{\mathcal{F}} - ||\widetilde{f}||_{\mathcal{F}}$:

$$\begin{aligned}\lambda\Big[||\widetilde{f}||_{\mathcal{F}}^2 - ||\widehat{f}||_{\mathcal{F}}^2\Big] &\leq \lambda\Big[||\widetilde{f}||_{\mathcal{F}} - ||\widehat{f}||_{\mathcal{F}}\Big] \\ &\leq \lambda\Big[2||\widetilde{f}||_{\mathcal{F}} - ||\widetilde{\Delta}||_{\mathcal{F}}\Big] \leq \lambda\Big[2 - ||\widetilde{\Delta}\,||_{\mathcal{F}}\Big].\end{aligned}$$

Substituting in the basic inequality:

$$\begin{aligned}\frac{1}{2}||\widehat{\Delta}||_n^2 &\leq \frac{1}{2}\,||f_i^\star - \widetilde{f}_i\,||_n^2 + \frac{\sigma}{n}\Big|\sum_{i=1}^n w_i\widetilde{\Delta}_i\Big| + \lambda_n||\widetilde{f}||_n^2 - \lambda_n||\widehat{f}||_n^2 \\ &\leq \frac{1}{2}\,||f_i^\star - \widetilde{f}_i\,||_n^2 + \frac{\sigma}{n}\Big|\sum_{i=1}^n w_i\widetilde{\Delta}_i\Big| + 2\lambda_n - \lambda_n||\widetilde{\Delta}||_{\mathcal{F}}.\end{aligned}$$

## Auxiliary Lemma 2

There exists positive constants $c_1, c_2$ such that with prob. at least $1 - c_1 \exp\left(\frac{-n\delta^2}{c_2\tilde{\sigma}^2}\right)$, we have

$$\left|\frac{\tilde{\sigma}}{n}\sum_{i=1}^{n} w_i\,\Delta(\mathbf{x}_i)\right| \leq 2\delta||\Delta||_n + 2\delta^2||\Delta||_{\mathcal{F}} + \frac{1}{16}||\Delta||_n^2,$$

for all $\Delta \in \partial\mathcal{F}$ with $||\Delta||_{\mathcal{F}} \geq 1$.

$||\widetilde{f}||_{\mathcal{F}} \leq 1, ||\widehat{f}||_{\mathcal{F}} > 2$, the trig. ineq. yields: $||\widetilde{\Delta}||_{\mathcal{F}} \geq ||\widehat{f}||_{\mathcal{F}} - ||\widetilde{f}||_{\mathcal{F}} > 1$. This yields:

$$\begin{aligned}\frac{1}{2}||\widehat{\Delta}||_n^2 &\leq \frac{1}{2}\,||f_i^\star - \widetilde{f}_i\,||_n^2 + 2\delta||\widetilde{\Delta}||_n + (2\delta^2 - \lambda_n)||\widetilde{\Delta}||_{\mathcal{F}} + 2\lambda_n + \frac{||\widetilde{\Delta}||_n^2}{16} \\ &\leq \frac{1}{2}\,||f_i^\star - \widetilde{f}_i\,||_n^2 + \underbrace{2\delta||\widetilde{\Delta}||_n}_{\leq 2\delta||\widetilde{f}-f^\star||_n + 2\delta||\widehat{\Delta}||_n} + 2\lambda_n + \underbrace{\frac{||\widetilde{\Delta}||_n^2}{16}}_{\leq \frac{1}{8}\left[||\widetilde{f}-f^\star||_n^2 + ||\widehat{\Delta}||_n^2\right]} . \quad \square\end{aligned}$$

# Proof of Auxiliary Lemma 2

## Auxiliary Lemma 2

There exists positive constants $c_1, c_2$ such that with prob. at least $1 - c_1 \exp\left(\frac{-n\delta^2}{c_2\tilde{\sigma}^2}\right)$, we have

$$\left|\frac{\tilde{\sigma}}{n}\sum_{i=1}^{n} w_i\,\Delta(\mathbf{x}_i)\right| \leq 2\delta||\Delta||_n + 2\delta^2||\Delta||_{\mathcal{F}} + \frac{1}{16}||\Delta||_n^2,$$

for all $\Delta \in \partial\mathcal{F}$ with $||\Delta||_{\mathcal{F}} \geq 1$.

It suffices to prove the theorem for $g \in \partial\mathcal{F}$ such that $||g||_{\mathcal{F}} = 1$. Given $\Delta \in \partial\mathcal{F}$ with $||\Delta|| > 1$. Apply the lemma for $g = \frac{\Delta}{||\Delta||_{\mathcal{F}}} \in \mathcal{F}$. Hence:

$$\left|\frac{\tilde{\sigma}}{n}\sum_{i=1}^{n} w_i\,\Delta(\mathbf{x}_i)\right| \leq c_1\delta||\Delta||_n + c_2\delta^2||\Delta||_n + \frac{1}{16}\overbrace{\frac{||\Delta||_n^2}{||\Delta||_{\mathcal{F}}}}^{\leq||\Delta||_n^2}.$$

# Proof of Auxiliary Lemma 2

- We first consider it over $\{||g||_n \leq t\}$. Define

$$Z_n(t) = \sup_{\substack{||g||_{\mathcal{F}} \leq 1 \\ ||g||_n \leq t}} \left| \frac{\tilde{\sigma}}{n} \sum_{i=1}^{n} w_i \, g(\mathbf{x}_i) \right|.$$

Lipschitz with constant $\frac{\tilde{\sigma}}{\sqrt{n}}$. Consequently,

$$\mathbb{P}\big[Z_n(t) \geq \mathbb{E}[Z_n(t)] + u\big] \leq \exp\left(\frac{-nu^2}{2t^2\tilde{\sigma}^2}\right).$$

- We first derive a bound for $t = \delta$. Note that $\mathbb{E}[Z_n(\delta)] \leq \tilde{\sigma}\mathcal{G}_n(\delta) \leq \delta^2$.

$$\mathbb{P}\big[Z_n(\delta) \geq 2\delta^2\big] \leq \exp\left(\frac{-n\delta^2}{2\tilde{\sigma}^2}\right).$$

- Also note that $\mathbb{E}[Z_n(t)] = \tilde{\sigma} \leq \tilde{\sigma}\mathcal{G}_n(t) = t\frac{\tilde{\sigma}\mathcal{G}_n(t)}{t} \leq t\frac{\tilde{\sigma}\mathcal{G}_n(t)}{\delta} \leq t\delta$:

$$\mathbb{P}\left[Z_n(t) \geq t\delta + \frac{t^2}{32}\right] \leq \exp\left(\frac{-c_2 n t^2}{\tilde{\sigma}^2}\right), \text{ for } t > \delta.$$

# Proof of Auxiliary Lemma 2

## Auxiliary Lemma 2

$$\left|\frac{\tilde{\sigma}}{n}\sum_{i=1}^{n} w_i\,\Delta(\mathbf{x}_i)\right| \leq 2\delta||\Delta||_n + 2\delta^2||\Delta||_{\mathcal{F}} + \frac{1}{16}||\Delta||_n^2, \tag{2}$$

for all $\Delta \in \partial\mathcal{F}$ with $||\Delta||_{\mathcal{F}} \geq 1$ with prob. at least $1 - c_1 \exp\left(\frac{-n\delta^2}{c_2\tilde{\sigma}^2}\right)$.

- We will complete the proof by a *peeling* argument.
- Let $\mathcal{E}$ the event that (2) is violated for some $g \in \partial\mathcal{F}$ with $||g||_{\mathcal{F}} = 1$.
- For $a, b \in \mathbb{R}$, let $\mathcal{E}(a, b)$ be the event that (2) is violated for some function such that $||g||_n \in [a, b]$ and $||g||_{\mathcal{F}} = 1$.
- For $m \in \mathbb{I}$, define $t_m = 2^m\delta$. We have $\mathcal{E} = \mathcal{E}(0, t_0) \cup \left(\bigcup_{m=0}^{\infty} \mathcal{E}(t_m, t_{m+1})\right)$.
- Hence, $\mathbb{P}[\mathcal{E}] \leq \mathbb{P}[\mathcal{E}(0, t_0)] + \sum_{m=0}^{\infty} \mathbb{P}[\mathcal{E}(t_m, t_{m+1})]$.
- Since $t = \delta$, we have $\mathbb{P}[\mathcal{E}(0, t_0)] \leq \mathbb{P}\left[Z_n(\delta) \geq 2\delta^2\right] \leq \exp\left(\frac{-n\delta^2}{2\tilde{\sigma}^2}\right)$.

# Proof of Auxiliary Lemma 2

- Assume that $\mathcal{E}(t_m, t_{m+1})$ holds. Meaning there exists $g$ with $||g||_{\mathcal{F}} = 1$, and $||g||_n \in [t_m, t_{m+1}]$, such that

$$\begin{aligned}\left|\frac{\tilde{\sigma}}{n}\sum_{i=1}^{n} w_i\, g(\mathbf{x}_i)\right| &\geq 2\delta||g||_n + 2\delta^2 + \frac{1}{16}||g||_n^2 \\ &\geq 2\delta t_m + 2\delta^2 + \frac{1}{8}t_m^2 \\ &= \delta t_{m+1} + 2\delta^2 + \frac{1}{32}t_{m+1}^2.\end{aligned}$$

- This lower bound implies that $Z_n(t_{m+1}) \geq \delta t_{m+1} + \frac{t_{m+1}^2}{32}$. Thus

$$\mathbb{P}[\mathcal{E}(t_m, t_{m+1})] \leq \exp\left(\frac{-c_2 n 2^{2m+2}\delta^2}{\tilde{\sigma}^2}\right)$$

- Wrapping up:

$$\mathbb{P}[\mathcal{E}] \leq \exp\left(\frac{-n\delta^2}{2\tilde{\sigma}^2}\right) + \sum_{m=0}^{\infty}\exp\left(\frac{-c_2 n 2^{2m+2}\delta^2}{\tilde{\sigma}^2}\right) \leq c_1 \exp\left(\frac{-c_2 n\delta^2}{\tilde{\sigma}^2}\right).$$