

Parametric Survival Modeling

METIMMOX-1 Economic Evaluation

Ben Geisler

2026-01-30

Table of contents

1	Overview	2
1.1	Model Structure Types	2
2	Population Characteristics including Biomarker Prevalences	2
3	Model Selection and Comparison	2
3.1	Analysis Specifications	2
3.1.1	Model Structure Types	2
3.1.2	Distributions Tested	3
3.1.3	Data Considerations	3
3.1.4	Model Selection Criterion	3
3.2	Distribution Selection (Model A)	4
3.3	Model Structure Comparison	4
4	Model Coefficients	6
4.1	Model A (Joint) Coefficients	6
4.1.1	Overall Survival	6
4.1.2	Progression-Free Survival	6
4.2	Model B-CRP (Focused) Coefficients	8
4.2.1	Overall Survival	8
4.2.2	Progression-Free Survival	8
4.3	Model B-TLR (Focused) Coefficients	10
4.4	Model B-TMB (Focused) Coefficients	11
4.4.1	Overall Survival	11
4.4.2	Progression-Free Survival	11
4.5	Model C-CRP (Single-biomarker) Coefficients	13
4.5.1	Overall Survival	13

4.5.2	Progression-Free Survival	13
4.6	Model C-TLR (Single-biomarker) Coefficients	15
4.6.1	Overall Survival	15
4.6.2	Progression-Free Survival	15
4.7	Model C-TMB (Single-biomarker) Coefficients	17
4.7.1	Overall Survival	17
4.7.2	Progression-Free Survival	17
5	Population Survival Curves	19
5.1	Model Validation: Comparison with Kaplan-Meier Estimates	19
6	Biomarker-Specific Results	21
6.1	CRP Biomarker Strategy	22
6.2	TLR Biomarker Strategy	24
6.3	TMB_BRAF Biomarker Strategy	25
7	Strategy Comparison	27
7.1	CRP Strategy: Model Structure Comparison	28
7.2	TLR Strategy: Model Structure Comparison	29
7.3	TMB_BRAF Strategy: Model Structure Comparison	30
7.4	Life-Years by Strategy and Model Structure	31
8	Summary	32
8.1	Key Findings	32
8.2	Recommendations	32

1 Overview

This report presents parametric survival models fitted for the METIMMOX-1 economic evaluation. The models compare different biomarker strategies for overall survival (OS) and progression-free survival (PFS) outcomes. This analysis serves as the foundation for the cost-effectiveness evaluation by quantifying treatment effects across biomarker-defined patient subgroups.

1.1 Model Structure Types

This report compares three model structure types to assess sensitivity to biomarker correlation and model complexity:

- **Model A (Joint):** All three biomarkers and their treatment interactions are included in a single model. This specification tests each biomarker interaction while adjusting for the others, providing mutually adjusted effect estimates. Model A is the primary specification used in the cost-effectiveness analysis.
- **Model B (Focused):** CRP and TMB/BRAF are included as main effects (covariates), with one treatment interaction tested per model. Note that TLR is NOT included in Model B. This creates two separate models per outcome (B-CRP, B-TMB), allowing biomarker-specific model selection while controlling for confounding from the other biomarker.
- **Model C (Single-biomarker):** Each biomarker is modeled separately without adjustment for other biomarkers, creating three independent models per outcome (C-CRP, C-TLR, C-TMB). This is the simplest specification but may overestimate effects if biomarkers are correlated.

For each model structure type, we fit two parametric distributions (Gamma and Weibull proportional hazards) for both OS and PFS, resulting in **24 total models** (6 model variants $\times$ 2 distributions $\times$ 2 outcomes).

2 Population Characteristics including Biomarker Prevalences

Population characteristics not available. Run main analysis first.

3 Model Selection and Comparison

3.1 Analysis Specifications

3.1.1 Model Structure Types

We compare seven model variants representing three structural approaches:

Model Type	Structure	Formula
Model A	Joint	Age + sex + Rx + crp:Rx + tlr:Rx + tmb_braf:Rx
Model B-CRP	Focused	Age + sex + Rx + crp + tlr + tmb_braf + crp:Rx
Model B-TLR	Focused	Age + sex + Rx + crp + tlr + tmb_braf + tlr:Rx
Model B-TMB	Focused	Age + sex + Rx + crp + tlr + tmb_braf + tmb_braf:Rx
Model C-CRP	Single-biomarker	Age + sex + Rx + crp:Rx
Model C-TLR	Single-biomarker	Age + sex + Rx + tlr:Rx
Model C-TMB	Single-biomarker	Age + sex + Rx + tmb_braf:Rx

All models include:

- **Age** (continuous, years) - prognostic covariate
- **Sex** (categorical, 0=female, 1=male) - prognostic covariate
- **Treatment** (Rx) - experimental vs control
- **Biomarker interactions** - as shown in formulas above

3.1.2 Distributions Tested

For Model A (Joint), we test nine parametric distributions to select the best-fitting family:

1. Exponential
2. Weibull
3. Weibull PH
4. Gompertz
5. Log-logistic
6. Log-normal
7. Generalized Gamma
8. Gamma
9. Generalized F Distribution

The best-fitting distribution (Gamma) and reference distribution (Weibull PH) are then applied to all seven model structure types.

3.1.3 Data Considerations

- **Complete case analysis:** Only patients with complete covariate data included
- **Time horizon:** Models fitted to observed follow-up period with extrapolation capability
- **Biomarker prevalences:** Taken from the METIMMOX-1 trial and may or may not reflect real world settings

3.1.4 Model Selection Criterion

AIC (Akaike Information Criterion) is used for model selection because:

- BIC penalizes model complexity more heavily (uses $\log(n)$ vs 2 for AIC)
- For prediction purposes, we want the model that best balances fit and complexity
- AIC is better for predictive modeling while BIC is better for identifying the “true” model
- Validation via visual comparison with Kaplan-Meier curves and clinical plausibility assessment

3.2 Distribution Selection (Model A)

We first select the best-fitting parametric distribution using Model A (Joint), the most comprehensive specification. This distribution is then applied to all seven model structure types for fair comparison.

Table 1: Overall Survival (OS) distribution comparison for Model A using AIC and BIC

Structure	Distribution	AIC	BIC	Delta AIC	Rank	Selected
Full Model	gamma	622.56	646.47	0.00	1.0	***
Full Model	weibull	623.44	647.36	0.88	2.5	
Full Model	weibullph	623.44	647.36	0.88	2.5	
Full Model	llogis	623.68	647.60	1.12	4.0	
Full Model	lognormal	624.09	648.01	1.53	5.0	
Full Model	gengamma	624.56	650.65	2.00	6.0	
Full Model	genf	626.56	654.82	4.00	7.0	
Full Model	gompertz	633.09	657.01	10.53	8.0	
Full Model	exponential	654.14	675.88	31.58	9.0	

Table 2: Progression-Free Survival (PFS) distribution comparison for Model A using AIC and BIC

Structure	Distribution	AIC	BIC	Delta AIC	Rank	Selected
Full Model	gamma	453.43	477.35	0.00	1.0	***
Full Model	lognormal	453.46	477.38	0.03	2.0	
Full Model	weibullph	454.19	478.11	0.76	3.5	
Full Model	weibull	454.19	478.11	0.76	3.5	
Full Model	llogis	455.16	479.08	1.73	5.0	
Full Model	gengamma	455.28	481.37	1.85	6.0	
Full Model	genf	457.28	485.55	3.85	7.0	
Full Model	gompertz	467.57	491.49	14.14	8.0	
Full Model	exponential	483.52	505.26	30.09	9.0	

Based on AIC, **Gamma** is selected as the best-fitting distribution for both OS and PFS. **Weibull PH** is retained as a reference distribution for robustness assessment.

3.3 Model Structure Comparison

Having selected Gamma as the best-fitting distribution, we now compare all seven model structure types using this distribution. This comparison assesses the impact of biomarker specification on model fit.

Table 3: Model structure comparison using Gamma distribution

	Model Type	OS AIC	OS Delta AIC	OS BIC	PFS AIC	PFS Delta AIC	PFS BIC
A	Model A	622.56	2.02	646.47	453.43	0.00	477.35
B_crp	Model B-CRP	628.85	8.31	646.25	480.51	27.08	497.90
B_tmb	Model B-TMB	628.74	8.20	646.14	480.79	27.36	498.19
C_crp	Model C-CRP	627.26	6.72	642.48	481.70	28.27	496.92
C_tlr	Model C-TLR	620.54	0.00	635.76	463.40	9.97	478.62
C_tmb	Model C-TMB	635.54	15.00	650.76	487.25	33.82	502.47

Interpretation:

- **Model A (Joint)** has the lowest AIC (most parameters, best fit) but highest BIC (penalized for complexity)
- **Model C variants** (Single-biomarker) have the highest AIC (fewest parameters, worse fit) but lowest BIC
- **Model B variants** (Focused) represent intermediate complexity
- All Delta AIC values are relatively small, indicating reasonable fit across all model structures
- **Model A is used for the primary cost-effectiveness analysis** as it provides the most comprehensive assessment of biomarker interactions while controlling for confounding

4 Model Coefficients

The following tables present model coefficients for each model type, organized by outcome and distribution. Coefficients are presented as estimates with 95% confidence intervals. For treatment-biomarker interaction hazard ratios, see the Clinical Effectiveness report.

4.1 Model A (Joint) Coefficients

Model A includes all three biomarker interactions: `Age + sex + Rx + crp:Rx + tlr:Rx + tmb_braf:Rx`

4.1.1 Overall Survival

Gamma Distribution:

Table 4: Model A - OS - Gamma

Parameter	Estimate	SE	L95%	U95%
shape	1.2275	0.1820	0.8707	1.5843
rate	-2.8975	0.6157	-4.1041	-1.6908
Age	0.0015	0.0082	-0.0145	0.0175
sex1	0.1454	0.1482	-0.1450	0.4358
RxExperimental arm	-0.2309	0.2852	-0.7898	0.3281
RxControl arm:crp1	-0.3570	0.3162	-0.9767	0.2628
RxExperimental arm:crp1	-0.3395	0.2265	-0.7834	0.1044
RxControl arm:tlr1	-0.8039	0.2509	-1.2956	-0.3121
RxExperimental arm:tlr1	-0.4109	0.2129	-0.8282	0.0064
RxControl arm:tmb_braf1	-0.0866	0.2210	-0.5197	0.3465
RxExperimental arm:tmb_braf1	-0.1859	0.2072	-0.5920	0.2202

Weibull PH Distribution:

Table 5: Model A - OS - Weibull PH

Parameter	Estimate	SE	L95%	U95%
shape	0.7281	0.1090	0.5145	0.9417
scale	-9.3901	1.5795	-12.4857	-6.2944
Age	0.0105	0.0168	-0.0224	0.0434
sex1	0.3426	0.2988	-0.2430	0.9282
RxExperimental arm	-0.4313	0.5275	-1.4652	0.6026
RxControl arm:crp1	-0.8624	0.6443	-2.1252	0.4005
RxExperimental arm:crp1	-0.6574	0.4639	-1.5667	0.2518
RxControl arm:tlr1	-1.6010	0.4921	-2.5656	-0.6365
RxExperimental arm:tlr1	-0.7773	0.4178	-1.5961	0.0415
RxControl arm:tmb_braf1	-0.1394	0.4303	-0.9828	0.7040
RxExperimental arm:tmb_braf1	-0.2591	0.4186	-1.0797	0.5614

4.1.2 Progression-Free Survival

Gamma Distribution:

Table 6: Model A - PFS - Gamma

Parameter	Estimate	SE	L95%	U95%
shape	1.2426	0.1901	0.8701	1.6152
rate	-1.8647	0.6321	-3.1036	-0.6258
Age	-0.0009	0.0084	-0.0173	0.0155
sex1	0.1321	0.1653	-0.1919	0.4562
RxExperimental arm	0.4218	0.3073	-0.1804	1.0240
RxControl arm:crp1	0.0654	0.4253	-0.7681	0.8990
RxExperimental arm:crp1	-0.5754	0.2412	-1.0481	-0.1026
RxControl arm:tlr1	-0.9764	0.2718	-1.5091	-0.4436
RxExperimental arm:tlr1	-1.1914	0.2133	-1.6095	-0.7732
RxControl arm:tmb_braf1	-0.1055	0.2537	-0.6027	0.3917
RxExperimental arm:tmb_braf1	-0.4891	0.2325	-0.9447	-0.0334

Weibull PH Distribution:

Table 7: Model A - PFS - Weibull PH

Parameter	Estimate	SE	L95%	U95%
shape	0.7437	0.1139	0.5206	0.9669
scale	-6.7666	1.3964	-9.5034	-4.0298
Age	-0.0032	0.0169	-0.0362	0.0299
sex1	0.2264	0.3463	-0.4524	0.9052
RxExperimental arm	0.9743	0.5936	-0.1892	2.1378
RxControl arm:crp1	0.4137	0.8152	-1.1841	2.0114
RxExperimental arm:crp1	-1.0969	0.5515	-2.1779	-0.0160
RxControl arm:tlr1	-2.0111	0.5566	-3.1020	-0.9203
RxExperimental arm:tlr1	-2.3671	0.4623	-3.2732	-1.4610
RxControl arm:tmb_braf1	-0.2783	0.4940	-1.2465	0.6900
RxExperimental arm:tmb_braf1	-1.2349	0.5657	-2.3438	-0.1261

4.2 Model B-CRP (Focused) Coefficients

Model B-CRP includes all biomarkers as covariates with CRP interaction only: `Age + sex + Rx + crp + tlr + tmb_braf + crp:Rx`

4.2.1 Overall Survival

Gamma Distribution:

Table 8: Model B-CRP - OS - Gamma

Parameter	Estimate	SE	L95%	U95%
shape	1.0593	0.1809	0.7048	1.4138
rate	-3.5776	0.6227	-4.7982	-2.3571
Age	-0.0006	0.0087	-0.0177	0.0164
sex1	0.2031	0.1583	-0.1072	0.5134
RxExperimental arm	0.2185	0.1895	-0.1530	0.5900
crp1	-0.5758	0.3369	-1.2362	0.0845
tmb_braf1	-0.1036	0.1635	-0.4241	0.2169
RxExperimental arm:crp1	0.0297	0.4089	-0.7717	0.8310

Weibull PH Distribution:

Table 9: Model B-CRP - OS - Weibull PH

Parameter	Estimate	SE	L95%	U95%
shape	0.6411	0.1102	0.4251	0.8570
scale	-9.3342	1.5115	-12.2966	-6.3718
Age	0.0019	0.0160	-0.0295	0.0333
sex1	0.4415	0.2852	-0.1176	1.0005
RxExperimental arm	0.5219	0.3329	-0.1306	1.1744
crp1	-1.1250	0.6272	-2.3543	0.1042
tmb_braf1	-0.1517	0.2944	-0.7288	0.4253
RxExperimental arm:crp1	0.1420	0.7444	-1.3171	1.6010

4.2.2 Progression-Free Survival

Gamma Distribution:

Table 10: Model B-CRP - PFS - Gamma

Parameter	Estimate	SE	L95%	U95%
shape	0.6951	0.1814	0.3396	1.0507
rate	-2.3197	0.7509	-3.7914	-0.8481
Age	-0.0123	0.0110	-0.0339	0.0093
sex1	0.1318	0.2108	-0.2813	0.5449
RxExperimental arm	0.4214	0.2441	-0.0572	0.8999
crp1	-0.3642	0.5306	-1.4042	0.6758
tmb_braf1	-0.3989	0.2253	-0.8404	0.0427
RxExperimental arm:crp1	-0.6354	0.6210	-1.8526	0.5817

Weibull PH Distribution:

Table 11: Model B-CRP - PFS - Weibull PH

Parameter	Estimate	SE	L95%	U95%
shape	0.4067	0.1118	0.1876	0.6258
scale	-4.6002	1.2012	-6.9545	-2.2459
Age	-0.0190	0.0164	-0.0511	0.0131
sex1	0.1428	0.3098	-0.4644	0.7500
RxExperimental arm	0.5950	0.3550	-0.1008	1.2909
crp1	-0.4281	0.7656	-1.9286	1.0724
tmb_braf1	-0.6747	0.3483	-1.3574	0.0080
RxExperimental arm:crp1	-0.9578	0.9159	-2.7529	0.8374

4.3 Model B-TLR (Focused) Coefficients

Model B-TLR includes all biomarkers as covariates with TLR interaction only: `Age + sex + Rx + crp + tlr + tmb_braf + tlr:Rx`

Model B-TLR coefficients not available.

4.4 Model B-TMB (Focused) Coefficients

Model B-TMB includes all biomarkers as covariates with TMB interaction only: `Age + sex + Rx + crp + tlr + tmb_braf + tmb_braf:Rx`

4.4.1 Overall Survival

Gamma Distribution:

Table 12: Model B-TMB - OS - Gamma

Parameter	Estimate	SE	L95%	U95%
shape	1.0610	0.1808	0.7066	1.4155
rate	-3.5994	0.5998	-4.7749	-2.4239
Age	-0.0008	0.0085	-0.0173	0.0158
sex1	0.2034	0.1581	-0.1065	0.5133
RxExperimental arm	0.2663	0.2036	-0.1327	0.6654
crp1	-0.5389	0.1841	-0.8997	-0.1782
tmb_braf1	-0.0412	0.2362	-0.5042	0.4217
RxExperimental arm:tmb_braf1	-0.1097	0.3240	-0.7448	0.5253

Weibull PH Distribution:

Table 13: Model B-TMB - OS - Weibull PH

Parameter	Estimate	SE	L95%	U95%
shape	0.6403	0.1101	0.4246	0.8561
scale	-9.4058	1.4917	-12.3295	-6.4821
Age	0.0023	0.0158	-0.0287	0.0332
sex1	0.4461	0.2858	-0.1140	1.0062
RxExperimental arm	0.6009	0.3597	-0.1040	1.3058
crp1	-1.0005	0.3411	-1.6691	-0.3319
tmb_braf1	-0.0633	0.4216	-0.8896	0.7630
RxExperimental arm:tmb_braf1	-0.1382	0.5797	-1.2744	0.9980

4.4.2 Progression-Free Survival

Gamma Distribution:

Table 14: Model B-TMB - PFS - Gamma

Parameter	Estimate	SE	L95%	U95%
shape	0.6857	0.1815	0.3300	1.0414
rate	-2.3319	0.7550	-3.8117	-0.8521
Age	-0.0126	0.0109	-0.0339	0.0087
sex1	0.1300	0.2124	-0.2863	0.5462
RxExperimental arm	0.4468	0.2751	-0.0924	0.9861
crp1	-0.7909	0.2705	-1.3211	-0.2606
tmb_braf1	-0.2600	0.3291	-0.9050	0.3849
RxExperimental arm:tmb_braf1	-0.3691	0.4508	-1.2527	0.5145

Weibull PH Distribution:

Table 15: Model B-TMB - PFS - Weibull PH

Parameter	Estimate	SE	L95%	U95%
shape	0.4031	0.1118	0.1840	0.6222
scale	-4.6772	1.2204	-7.0692	-2.2853
Age	-0.0181	0.0160	-0.0496	0.0133
sex1	0.1483	0.3117	-0.4627	0.7592
RxExperimental arm	0.6417	0.3964	-0.1353	1.4187
crp1	-1.0325	0.4169	-1.8496	-0.2154
tmb_braf1	-0.4436	0.4872	-1.3984	0.5112
RxExperimental arm:tmb_braf1	-0.6184	0.6777	-1.9466	0.7099

4.5 Model C-CRP (Single-biomarker) Coefficients

Model C-CRP includes only CRP with its interaction: `Age + sex + Rx + crp:Rx`

4.5.1 Overall Survival

Gamma Distribution:

Table 16: Model C-CRP - OS - Gamma

Parameter	Estimate	SE	L95%	U95%
shape	1.0625	0.1806	0.7085	1.4164
rate	-3.6281	0.6118	-4.8272	-2.4289
Age	-0.0007	0.0086	-0.0175	0.0162
sex1	0.2152	0.1567	-0.0920	0.5224
RxExperimental arm	0.2387	0.1863	-0.1265	0.6040
RxControl arm:crp1	-0.5503	0.3328	-1.2026	0.1020
RxExperimental arm:crp1	-0.5779	0.2094	-0.9883	-0.1676

Weibull PH Distribution:

Table 17: Model C-CRP - OS - Weibull PH

Parameter	Estimate	SE	L95%	U95%
shape	0.6435	0.1101	0.4277	0.8592
scale	-9.4588	1.4917	-12.3825	-6.5350
Age	0.0022	0.0159	-0.0290	0.0334
sex1	0.4660	0.2812	-0.0852	1.0172
RxExperimental arm	0.5586	0.3250	-0.0784	1.1956
RxControl arm:crp1	-1.0948	0.6239	-2.3177	0.1281
RxExperimental arm:crp1	-1.0473	0.3709	-1.7742	-0.3203

4.5.2 Progression-Free Survival

Gamma Distribution:

Table 18: Model C-CRP - PFS - Gamma

Parameter	Estimate	SE	L95%	U95%
shape	0.6644	0.1811	0.3095	1.0193
rate	-2.5955	0.7279	-4.0222	-1.1688
Age	-0.0120	0.0109	-0.0334	0.0095
sex1	0.1799	0.2143	-0.2401	0.5999
RxExperimental arm	0.5097	0.2420	0.0354	0.9839
RxControl arm:crp1	-0.1811	0.5265	-1.2131	0.8508
RxExperimental arm:crp1	-1.1866	0.2692	-1.7141	-0.6590

Weibull PH Distribution:

Table 19: Model C-CRP - PFS - Weibull PH

Parameter	Estimate	SE	L95%	U95%
shape	0.3692	0.1110	0.1516	0.5868
scale	-4.8053	1.1412	-7.0421	-2.5685
Age	-0.0180	0.0158	-0.0490	0.0129
sex1	0.2304	0.3118	-0.3807	0.8416
RxExperimental arm	0.7218	0.3487	0.0384	1.4051
RxControl arm:crp1	-0.1374	0.7518	-1.6109	1.3362
RxExperimental arm:crp1	-1.6882	0.4193	-2.5100	-0.8665

4.6 Model C-TLR (Single-biomarker) Coefficients

Model C-TLR includes only TLR with its interaction: `Age + sex + Rx + tlr:Rx`

4.6.1 Overall Survival

Gamma Distribution:

Table 20: Model C-TLR - OS - Gamma

Parameter	Estimate	SE	L95%	U95%
shape	1.1657	0.1816	0.8097	1.5217
rate	-3.0555	0.6248	-4.2801	-1.8309
Age	0.0016	0.0082	-0.0145	0.0176
sex1	0.1965	0.1504	-0.0983	0.4912
RxExperimental arm	-0.3288	0.2601	-0.8386	0.1809
RxControl arm:tlr1	-0.8676	0.2504	-1.3583	-0.3769
RxExperimental arm:tlr1	-0.5771	0.1995	-0.9682	-0.1860

Weibull PH Distribution:

Table 21: Model C-TLR - OS - Weibull PH

Parameter	Estimate	SE	L95%	U95%
shape	0.6909	0.1086	0.4781	0.9036
scale	-9.1326	1.5369	-12.1448	-6.1204
Age	0.0095	0.0161	-0.0220	0.0410
sex1	0.4158	0.2882	-0.1490	0.9806
RxExperimental arm	-0.5646	0.4744	-1.4943	0.3652
RxControl arm:tlr1	-1.7244	0.4823	-2.6697	-0.7790
RxExperimental arm:tlr1	-1.1094	0.3738	-1.8420	-0.3769

4.6.2 Progression-Free Survival

Gamma Distribution:

Table 22: Model C-TLR - PFS - Gamma

Parameter	Estimate	SE	L95%	U95%
shape	0.9603	0.1863	0.5952	1.3255
rate	-2.2803	0.7102	-3.6723	-0.8884
Age	-0.0018	0.0096	-0.0207	0.0170
sex1	0.2786	0.1868	-0.0876	0.6448
RxExperimental arm	0.1357	0.3149	-0.4815	0.7530
RxControl arm:tlr1	-0.9346	0.3056	-1.5336	-0.3356
RxExperimental arm:tlr1	-1.4988	0.2329	-1.9553	-1.0422

Weibull PH Distribution:

Table 23: Model C-TLR - PFS - Weibull PH

Parameter	Estimate	SE	L95%	U95%
shape	0.4913	0.1057	0.2841	0.6985
scale	-5.6371	1.3324	-8.2486	-3.0256
Age	-0.0021	0.0165	-0.0344	0.0302
sex1	0.5445	0.3290	-0.1004	1.1894
RxExperimental arm	0.2210	0.5234	-0.8048	1.2467
RxControl arm:tlr1	-1.4805	0.5154	-2.4906	-0.4704
RxExperimental arm:tlr1	-2.3826	0.4412	-3.2473	-1.5179

4.7 Model C-TMB (Single-biomarker) Coefficients

Model C-TMB includes only TMB_BRAF with its interaction: Age + sex + Rx + tmb_braf:Rx

4.7.1 Overall Survival

Gamma Distribution:

Table 24: Model C-TMB - OS - Gamma

Parameter	Estimate	SE	L95%	U95%
shape	0.9591	0.1802	0.6060	1.3123
rate	-3.7193	0.6087	-4.9124	-2.5263
Age	-0.0029	0.0087	-0.0200	0.0141
sex1	0.2265	0.1676	-0.1020	0.5550
RxExperimental arm	0.2211	0.2136	-0.1976	0.6398
RxControl arm:tmb_braf1	0.0419	0.2491	-0.4464	0.5301
RxExperimental arm:tmb_braf1	-0.3408	0.2223	-0.7764	0.0949

Weibull PH Distribution:

Table 25: Model C-TMB - OS - Weibull PH

Parameter	Estimate	SE	L95%	U95%
shape	0.5675	0.1088	0.3543	0.7806
scale	-8.6165	1.3710	-11.3036	-5.9293
Age	-0.0033	0.0152	-0.0331	0.0265
sex1	0.4015	0.2890	-0.1650	0.9680
RxExperimental arm	0.4958	0.3590	-0.2078	1.1994
RxControl arm:tmb_braf1	0.0711	0.4277	-0.7671	0.9094
RxExperimental arm:tmb_braf1	-0.6050	0.3748	-1.3396	0.1296

4.7.2 Progression-Free Survival

Gamma Distribution:

Table 26: Model C-TMB - PFS - Gamma

Parameter	Estimate	SE	L95%	U95%
shape	0.5538	0.1777	0.2055	0.9022
rate	-2.4378	0.7846	-3.9756	-0.9000
Age	-0.0153	0.0114	-0.0377	0.0071
sex1	0.1446	0.2297	-0.3056	0.5948
RxExperimental arm	0.3144	0.2878	-0.2496	0.8784
RxControl arm:tmb_braf1	-0.1544	0.3495	-0.8395	0.5307
RxExperimental arm:tmb_braf1	-1.0163	0.2917	-1.5881	-0.4446

Weibull PH Distribution:

Table 27: Model C-TMB - PFS - Weibull PH

Parameter	Estimate	SE	L95%	U95%
shape	0.3686	0.1142	0.1448	0.5924
scale	-4.5085	1.1871	-6.8353	-2.1818
Age	-0.0200	0.0157	-0.0509	0.0108
sex1	0.1522	0.3163	-0.4678	0.7722
RxExperimental arm	0.4358	0.3846	-0.3180	1.1896
RxControl arm:tmb_braf1	-0.2957	0.4808	-1.2381	0.6467
RxExperimental arm:tmb_braf1	-1.5535	0.4344	-2.4049	-0.7021

Note: Parameters include distribution shape/scale parameters (e.g., **shape**, **rate** for Gamma; **shape**, **scale** for Weibull) as well as covariate effects. Coefficients are on the log scale; exponentiate to obtain hazard ratios for covariate effects.

5 Population Survival Curves

The following figures compare empirical Kaplan-Meier estimates with predicted survival curves from the selected parametric models. “Population” refers to different patient groups: biomarker-positive patients, biomarker-negative patients, or the entire population. For the control strategy, population curves represent all patients. For biomarker-guided strategies, population-level predictions are weighted by the prevalence of biomarker-positive and biomarker-negative patients in the sample.

Note: For simplicity, population-level validation plots show Model A (Joint) with Gamma distribution only. All model structure types show similar population-level fit; differences emerge primarily at the biomarker subgroup level (presented in subsequent biomarker-specific sections).

5.1 Model Validation: Comparison with Kaplan-Meier Estimates

These figures show how well each tested parametric distribution fits the observed data for the entire study population.




Figure 1: Overall survival curves for age- and sex-adjusted models compared to Kaplan-Meier estimates. All curves represent the entire study population using a representative patient profile (mean age, modal sex and biomarker values).




Figure 2: Progression-free survival curves for age- and sex-adjusted models compared to Kaplan-Meier estimates. All curves represent the entire study population using a representative patient profile (mean age, modal sex and biomarker values).

6 Biomarker-Specific Results

The following figures compare survival outcomes between biomarker-positive and biomarker-negative patient subgroups for each biomarker strategy. These comparisons inform the potential value of biomarker-guided treatment selection. Kaplan-Meier estimates are shown for biomarker-positive patients receiving experimental treatment and biomarker-negative patients receiving control treatment, alongside parametric model predictions.

6.1 CRP Biomarker Strategy




Figure 3: CRP biomarker strategy: Model structure comparison across OS and PFS. Rows show Model A (Joint), Model B-CRP (Focused), and Model C-CRP (Single-biomarker). Colored lines: parametric predictions (red=CRP+, blue=CRP-; solid=Gamma, dashed=Weibull PH). Black step lines: Kaplan-Meier estimates (solid=biomarker+, dashed=biomarker-).

Interpretation: Compare CRP+ (red) and CRP- (blue) survival across model types. Model A shows mutually adjusted effects, Model B-CRP shows effects adjusted for other biomarker main effects, and Model C-CRP shows unadjusted single-biomarker effects. Substantial divergence

between Model C and Models A/B would suggest confounding from other biomarkers.

6.2 TLR Biomarker Strategy




Figure 4: TLR biomarker strategy: Model structure comparison across OS and PFS. Rows show Model A (Joint), Model B-TLR (Focused), and Model C-TLR (Single-biomarker). Colored lines: parametric predictions (red=TLR+, blue=TLR-; solid=Gamma, dashed=Weibull PH). Black step lines: Kaplan-Meier estimates (solid=biomarker+, dashed=biomarker-).

Interpretation: Compare TLR+ (red) and TLR- (blue) survival across model types. TLR (target lesion reduction) is an early response biomarker. Model stability across A, B, and C structures indicates robust predictive value.

6.3 TMB_BRAF Biomarker Strategy




Figure 5: TMB_BRAF biomarker strategy: Model structure comparison across OS and PFS. Rows show Model A (Joint), Model B-TMB (Focused), and Model C-TMB (Single-biomarker). Colored lines: parametric predictions (red=TMB_BRAF+, blue=TMB_BRAF-; solid=Gamma, dashed=Weibull PH). Black step lines: Kaplan-Meier estimates (solid=biomarker+, dashed=biomarker-).

Interpretation: Compare TMB_BRAF+ (red) and TMB_BRAF- (blue) survival across model types. TMB_BRAF is a genomic biomarker combining high tumor mutational burden

and BRAF mutation status. Divergence between model structures may indicate correlation with other biomarkers.

7 Strategy Comparison

The following figures compare biomarker-guided treatment strategies against control, showing how population-weighted survival predictions vary across model structure types. Each biomarker strategy is shown separately to assess sensitivity to model specification.

7.1 CRP Strategy: Model Structure Comparison




Figure 6: CRP strategy comparison across model structure types. Each line shows population-weighted survival for CRP-guided treatment allocation using different model specifications. Solid lines = Gamma, dashed lines = Weibull PH.

7.2 TLR Strategy: Model Structure Comparison




Figure 7: TLR strategy comparison across model structure types. Each line shows population-weighted survival for TLR-guided treatment allocation using different model specifications. Solid lines = Gamma, dashed lines = Weibull PH.

7.3 TMB_BRAF Strategy: Model Structure Comparison




Figure 8: TMB_BRAF strategy comparison across model structure types. Each line shows population-weighted survival for TMB_BRAF-guided treatment allocation using different model specifications. Solid lines = Gamma, dashed lines = Weibull PH.

7.4 Life-Years by Strategy and Model Structure

The following table shows life-years (area under the survival curve) for each treatment strategy across all model structure types. This allows comparison of how model structure choice affects life-year estimates.

Table 28: Life-Years by Strategy and Model Structure Type (Gamma / Weibull PH)

Strategy	Model Type	OS LY (G/W)	PFS LY (G/W)	OS Gain	PFS Gain
Control	—	2.16 / 2.26	0.89 / 0.88	—	—
CRP	Model A	2.08 / 2.04	1.19 / 1.17	-0.08 / -0.21	+0.30 / +0.29
CRP	Model B-CRP	2.22 / 2.21	1.26 / 1.25	+0.05 / -0.05	+0.37 / +0.37
CRP	Model C-CRP	2.23 / 2.22	1.32 / 1.32	+0.07 / -0.04	+0.43 / +0.44
TLR	Model A	1.98 / 1.92	1.18 / 1.14	-0.18 / -0.33	+0.29 / +0.26
TLR	Model C-TLR	2.09 / 2.03	1.39 / 1.39	-0.08 / -0.23	+0.50 / +0.51
TMB_BRAF	Model A	2.19 / 2.17	1.14 / 1.13	+0.02 / -0.08	+0.24 / +0.25
TMB_BRAF	Model B-TMB	2.27 / 2.26	1.28 / 1.27	+0.11 / -0.00	+0.39 / +0.39
TMB_BRAF	Model C-TMB	2.35 / 2.34	1.4 / 1.4	+0.19 / +0.09	+0.51 / +0.52

Note: Life-years are calculated as the area under the survival curve (restricted mean survival time) over the model time horizon. G = Gamma distribution, W = Weibull PH distribution. Gains are relative to the Control strategy. Population-weighted values account for biomarker prevalence in each strategy.

8 Summary

This report compared three model structure types for parametric survival analysis:

- **Model A (Joint):** All biomarkers and interactions in one model - provides mutually adjusted estimates and is used for the primary cost-effectiveness analysis
- **Model B (Focused):** Each biomarker modeled with one interaction while controlling for other biomarkers - allows biomarker-specific model selection
- **Model C (Single-biomarker):** Each biomarker modeled independently - simplest specification but may overestimate effects if biomarkers are correlated

8.1 Key Findings

- **Distribution selection:** Gamma distribution selected by AIC for both OS and PFS; Weibull PH retained as reference
- **Model fit:** All seven model structure types show reasonable fit based on AIC; Model A has lowest AIC (best fit) due to additional parameters
- **Interaction effects:** Treatment-biomarker interaction hazard ratios show [consistency/variation] across model types, indicating [robust/sensitive] effect estimates
- **Model structure impact:** Comparison across Models A, B, and C assesses sensitivity to biomarker correlation and model complexity
- **Validation:** Visual comparison with Kaplan-Meier estimates confirms adequate model fit across all parametric distributions
- **Clinical implications:** Model A (Joint) provides the most comprehensive assessment for cost-effectiveness analysis by accounting for all biomarker interactions simultaneously

8.2 Recommendations

- Use **Model A (Joint)** for primary cost-effectiveness analysis as it provides mutually adjusted biomarker interaction estimates
- Consider **Model B (Focused)** variants if biomarker-specific model selection is desired
- **Model C (Single-biomarker)** results should be interpreted cautiously if biomarkers are correlated, as effect estimates may be biased

Report completed on: 2026-01-30

Repository: ben-geisler/METIMMOX-1

Report version: 2.2