

# The Impact of Anthropogenic Forcing on ENSO Amplitude

Ben Goldman

March 8, 2022

# Climate Change: Global Warming vs. Climate Variability




Figure 1: Global mean land air temperature in GISSTEMP 4 dataset. (Team et al., 2019) and (Lenssen et al., 2019)

# Climate Forcing

- **Forcing**: any external factor that affects climate.
  - GHG** Greenhouse gasses
  - AER** Aerosols (natural: volcanic ash, artificial: smoke)
  - BMB** Biomass burning
  - LULC** Land use/cover (deforestation, desertification)





Figure 2: Factors that contribute to the greenhouse effect.

# El Niño, La Niña, and ENSO

- ENSO = El Niño/Southern Oscillation
- El Niño is the warm phase, and La Niña is the cool phase.
- Warming and cooling of the Pacific Ocean.
- Affects human societies through temperature and rainfall. (Ropelewski and Halpert, 1987)

**Figure 3:** Comparison of SST anomaly between 1975 La Niña event and 1997 El Niño event in HadISST 1 dataset. (Rayner et al., 2003)



- ENSO's properties observed vary across different decades. (Lübbecke and McPhaden, 2014).
- Weakened ENSO during the Ice Age due to reduced CO<sub>2</sub> levels (Zhu et al., 2017).
- Models show possible increasing ENSO activity in the future (Zheng et al., 2017) and (Maher et al., 2018).

## Research Gap

- Little research using a large ensemble to examine the effect of individual factors on ENSO.
- Considerable disagreement between studies on whether ENSO will strengthen or weaken due to global warming

## Questions


1. Do the CESM1 and CESM2 climate models predict increased or decreased ENSO intensity in the future?
2. Is the predicted increase (or decrease) due to human activities?

# Methods and Results

- Community Earth System Model (CESM) Versions 1 and 2 (Kay et al., 2015) (Danabasoglu et al., 2020).
- Predicts climate over 21st century with global warming.
- Ensemble: collection of multiple simulations.
- Single forcing ensembles that represent influence of single factor.
- Software used: R, ncd4, zoo, dplyr, ggplot2, WaveletComp, reshape2, nco.




# Measuring ENSO Intensity



# ENSO is Becoming Stronger

- Increase in ENSO intensity in both ensembles. (Exceeds 2 standard errors)
- Increase slows down in CESM1 and decreases in CESM2 after around 2050.




**Figure 4:** ENSO intensity ensemble mean and standard error for CESM1 and CESM2

# Influence of Aerosols and Greenhouse Gases

- Influence of each factor on ENSO amplitude.
- Increased variance due to greenhouse gas emissions.
- Somewhat increased variance from aerosol emissions, but not linear.

**Takeaway:** Human activities are triggering predicted strengthening of ENSO.



**Figure 5:** Influence of GHG, AER, and BMB forcing on ENSO amplitude in CESM1

- Separate ENSO record into changes in period over time.
- In CESM1, increase in ENSO intensity is mainly strengthening of longer-period cycle.
- In CESM2, longer-period ENSO weakens after 2025.
- Indicates that longer frequency bands are more susceptible to climate change.




Figure 6: Wavelet power spectrum for the Niño 3.4 index in the fully-forced CESM1 and CESM2 ensembles

## Student:

- Analyze raw data on computer
- Produce graphics for analysis and publication
- Write documentation
- Identify key features of results

## Mentor:

- Review student writing
- Interpret results in the context of climatology
- Conduct parallel analysis
- Provide raw data from facility

1. Do the CESM1 and CESM2 climate models predict increased or decreased ENSO intensity in the future?
2. Is the predicted increase (or decrease) due to human activities?
1. Yes, ENSO variance increases significantly over the 21<sup>st</sup> century
2. Yes, the major factor contributing to this increase is greenhouse gasses, with aerosol emissions playing a secondary role.

### Limitations:

- Niño 3.4 index shown to be inaccurate for some models (Cai et al., 2018).
- CESM may contain biases.
- Models are only an approximation of the Earth's actual climate.

Application: to improve our ability to predict ENSO and help people prepare for increased likelihood of extreme weather.

## Acknowledgments

- This material is based upon work supported by the National Center for Atmospheric Research, which is a major facility sponsored by the National Science Foundation under Cooperative Agreement No. 1852977.
- Thank you to my teacher, my family, and my mentor!



# References

- Cai, W., Wang, G., Dewitte, B., Wu, L., Santoso, A., Takahashi, K., Yang, Y., Carréric, A., and McPhaden, M. J. (2018). Increased variability of eastern pacific el niño under greenhouse warming. *Nature*, 564(7735):201–206.
- Danabasoglu, G., Lamarque, J.-F., Bacmeister, J., Bailey, D., DuVivier, A., Edwards, J., Emmons, L., Fasullo, J., Garcia, R., Gettelman, A., et al. (2020). The community earth system model version 2 (cesm2). *Journal of Advances in Modeling Earth Systems*, 12(2).
- Kay, J. E., Deser, C., Phillips, A., Mai, A., Hannay, C., Strand, G., Arblaster, J. M., Bates, S. C., Danabasoglu, G., Edwards, J., Holland, M., Kushner, P., Lamarque, J.-F., Lawrence, D., Lindsay, K., Middleton, A., Munoz, E., Neale, R., Oleson, K., Polvani, L., and Vertenstein, M. (2015). The community earth system model (CESM) large ensemble project: A community resource for studying climate change in the presence of internal climate variability. *Bulletin of the American Meteorological Society*, 96(8):1333–1349.
- Lenssen, N. J., Schmidt, G. A., Hansen, J. E., Menne, M. J., Persin, A., Ruedy, R., and Zyss, D. (2019). Improvements in the gistemp uncertainty model. *Journal of Geophysical Research: Atmospheres*, 124(12):6307–6326.
- Lübbecke, J. F. and McPhaden, M. J. (2014). Assessing the twenty-first-century shift in ENSO variability in terms of the bjerknes stability index. *Journal of Climate*, 27(7):2577–2587.
- Maher, N., Matei, D., Milinski, S., and Marotzke, J. (2018). ENSO change in climate projections: Forced response or internal variability? *Geophysical Research Letters*, 45(20).
- Rayner, N., Parker, D. E., Horton, E., Folland, C. K., Alexander, L. V., Rowell, D., Kent, E. C., and Kaplan, A. (2003). Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century. *Journal of Geophysical Research: Atmospheres*, 108(D14).
- Ropelewski, C. F. and Halpert, M. S. (1987). Global and regional scale precipitation patterns associated with the el niño/southern oscillation. *Monthly weather review*, 115(8):1606–1626.
- Team, G. et al. (2019). Giss surface temperature analysis (gistemp), version 4. *NASA Goddard Institute for Space Studies*.
- Zheng, X.-T., Hui, C., and Yeh, S.-W. (2017). Response of ENSO amplitude to global warming in CESM large ensemble: uncertainty due to internal variability. *Climate Dynamics*, 50(11-12):4019–4035.
- Zhu, J., Liu, Z., Brady, E., Otto-Bliesner, B., Zhang, J., Noone, D., Tomas, R., Nusbaumer, J., Wong, T., Jahn, A., et al. (2017). Reduced enso variability at the lgm revealed by an isotope-enabled earth system model. *Geophysical Research Letters*, 44(13):6984–6992.
- <https://www.coolaustralia.org/the-greenhouse-effect-secondary>