

The Impact of Industrial Emissions on El Niño Intensity

Ben Goldman

March 4, 2021

What is El Niño/Southern Oscillation (ENSO)?

- ▶ Drives extreme weather around the world
- ▶ Oscillation between warm and cold temperature in the Pacific Ocean
- ▶ Some events are more strong than others
- ▶ Significant effect on people
- ▶ Major issue is prediction




Figure 1: Changes to tropical Pacific climate during El Niño

Short-Term vs. Long-Term Change

- ▶ Weather is not climate!
- ▶ Climate varies on short and long term
- ▶ Short term changes: ENSO and more
- ▶ Long term changes: Global Warming




Figure 2: Global average temperature changes since 1880. Red line: smoothed average, black line: unsmoothed average

Review of Literature

Chen et. al. 2017	Why are the results of climate studies disagreeing about whether ENSO will intensify or weaken?
Maher et. al. 2018	Used a dataset of climate predictions, finding that ENSO intensity may increase in the future.
Cai et. al. 2018	Analyzed climate predictions, finding that changes in ocean stratification are causing increase in ENSO strength.
Gap	Little research comparing the effects of greenhouse gasses vs. aerosol emissions.

Research Goals

Problem	ENSO is serious, so we need to predict it.
What: Verify past research	Does the data show that ENSO will become stronger?
Why: Human impact	Figure out which human activities are causing the increase.
How: Ocean structure	Determine what changes are taking place to make ENSO stronger.

Data: the CESM1 Large Ensemble

- ▶ How do we explore hypothetical scenarios? With a computer model!
- ▶ Estimation of how the earth's climate actually works, but can receive hypothetical input of increasing greenhouse gas and aerosol levels.
- ▶ Can now find how the climate will appear in the future.
- ▶ Models may be inaccurate, so researchers use an large group of them – an ensemble.
- ▶ Control run with input as if there were no industrial emissions.

Methodology Step 1: Niño 3.4 Variance

- ▶ How to calculate ENSO intensity in the model output?
- ▶ Step 1: Calculate sea temperature in Niño 3.4 region of tropical Pacific Ocean.
- ▶ Step 2: Convert temperature dataset to dataset representing change in temperature variation over time.
- ▶ Calculate variance around one point, move point forward slightly, repeat




Figure 3: Niño 3.4 region is the shaded box.

Butterfly Effect: The Need for a Large Ensemble

- ▶ Small differences in initial conditions can blow up to big differences in end result: Butterfly effect!
- ▶ Each simulation by its self is inaccurate.
- ▶ Repeat simulation with slightly different initial conditions.
- ▶ Eventually, real trends may appear.




Figure 4: Niño 3.4 20-year variance for individual members in full forcing ensemble.

Model Predictions: ENSO in the Future

- ▶ Calculate mean and standard error for ENSO intensity in ensemble and control.
- ▶ ENSO is predicted to intensify in the 21st century!
- ▶ This means more extreme weather!




Figure 5: 20-year variance of Niño 3.4 index for fully-forced ensemble. Grey bar shows control mean and standard errors

Human Impact: Analysis of Individual Factors

- ▶ Why is ENSO predicted to intensify? What human impacts play the largest role?
 - ▶ Factors include: Greenhouse gasses, aerosols, natural factors.
- ▶ Separate out individual influences in model output.
- ▶ Single forcing ensembles: forced by all factors except for 1.
- ▶ Subtract “all-but-one” ensembles from original “full-forcing” ensemble.
- ▶ Resulting data represents influence of only one factor.

Role of Greenhouse and Aerosol Emissions

- ▶ Greenhouse gasses and aerosols contribute to increase in variance.
- ▶ Sum of individual factors is different from full forcing mean. This means things are complicated.
- ▶ Greenhouse gasses and aerosols are both human-produced!




Figure 6: Influence of individual human factors. Yellow is greenhouse gasses, Green is aerosols.


Why: Correlation With Changes in Ocean Temperature

- ▶ Why do greenhouse gasses and aerosols strengthen ENSO?
- ▶ Calculate correlation coefficient between ENSO intensity and ocean temperature
- ▶ Detrend and smooth
- ▶ Pearson's correlation coefficient at each grid-point

Physical Mediator: Heating Difference

- ▶ Strong negative correlation in fully forced ensemble below surface.
- ▶ Positive correlation in greenhouse ensemble
- ▶ Weak/zero correlation in aerosols ensemble
- ▶ Rising temperatures heat different layers of ocean at different rates, modifying transfer of heat.

Figure 7: Correlation between ENSO intensity and ocean temperature in a) fully forced b) greenhouse and c) aerosol ensembles



Summary

Estimate future changes	There is likely to be an increase in ENSO strength over the next 100 years.
Role of single factors	Increase is likely caused by the combined influence of greenhouse gasses and aerosols.
Correlation with ocean temperature	Global warming increases ENSO intensity by warming upper layers of the Pacific faster than central layers.

Discussion

- ▶ ENSO intensification is bad for human societies.
- ▶ Improve prediction ability to help people prepare.
- ▶ Reduce danger by switching to renewable energy.
- ▶ Next steps:
 - ▶ Work with other datasets, such as the new CESM2
 - ▶ Examine other variables to further analyze mediator process




Figure 8: Regions affected by ENSO

Acknowledgements

- ▶ This material is based upon work supported by the National Center for Atmospheric Research, which is a major facility sponsored by the National Science Foundation under Cooperative Agreement No. 1852977.
- ▶ Thank you to my teacher, my family, and my mentor!
- ▶ Role of mentor:
 - ▶ Provide raw data from his facility
 - ▶ Suggest methods and interpretations
 - ▶ Provide feedback on results
 - ▶ Make similar calculations to check student's results

References

- ▶ Chen, L., Li, T., Yu, Y., and Behera, S. K. (2017). A possible explanation for the divergent projection of enso amplitude change under global warming. *Climate Dynamics*, 49(11-12):3799–3811.
- ▶ Deser, C., Phillips, A. S., Simpson, I. R., Rosenbloom, N., Coleman, D., Lehner, F., Pendergrass, A. G., DiNezio, P., and Stevenson, S. (2020). Isolating the evolving contributions of anthropogenic aerosols and greenhouse gases: a new cesm1 large ensemble community resource. *Journal of Climate*, 33(18):7835–7858.
- ▶ Kay, J. E., Deser, C., Phillips, A., Mai, A., Hannay, C., Strand, G., Arblaster, J. M., Bates, S., Danabasoglu, G., Edwards, J., et al. (2015). The community earth system model (cesm) large ensemble project: A community resource for studying climate change in the presence of internal climate variability. *Bulletin of the American Meteorological Society*, 96(8):1333–1349.
- ▶ Maher, N., Matei, D., Milinski, S., and Marotzke, J. (2018). Enso change in climate projections: Forced response or internal variability? *Geophysical Research Letters*, 45(20):11–390.
- ▶ Stevenson, S., Fox-Kemper, B., Jochum, M., Neale, R., Deser, C., and Meehl, G. (2012). Will there be a significant change to el nino in the twenty-first century? *Journal of Climate*, 25(6):2129–2145.

Image Sources

- ▶ https://www.esa.int/ESA_Multimedia/Images/2018/08/El_Nino
- ▶ https://data.giss.nasa.gov/gistemp/graphs_v4
- ▶ <https://www.cpc.ncep.noaa.gov>
- ▶ https://www.cesm.ucar.edu/working_groups