

# A Monte Carlo method to estimate cell population heterogeneity

Ben Lambert<sup>1,2\*</sup>, David J. Gavaghan<sup>3</sup>, Simon Tavener<sup>4</sup>.

**1** Department of Zoology, University of Oxford, Oxford, Oxfordshire, U.K.

**2** MRC Centre for Outbreak Analysis and Modelling, Infectious Disease Epidemiology, Imperial College London, London W2 1PG, UK.

**3** Department of Computer Science, University of Oxford, Oxford, U.K.

**4** Department of Statistics, Colorado State University, Fort Collins, Colorado, U.S.A.

\*ben.c.lambert@gmail.com.

Revision date & time: 2019-05-21 16:19

# 1 Abstract

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26

Variation is characteristic of all living systems. Laboratory techniques such as flow cytometry can probe individual cells and, after decades of experimentation, it is clear that even members of seemingly homogeneous cell populations can exhibit differences. To understand whether this variation is biologically meaningful, it is essential to discern its source. Mathematical models of biological systems are tools that can be used to investigate causes of cell-to-cell variation. From mathematical analysis and simulation of these models, biological hypotheses can be posed and investigated, then parameter inference can determine which of these is most compatible with experimental data. Data from laboratory experiments often takes the form of “snapshots” representing distributions of cellular properties at different points in times, rather than individual cell trajectories. This data is not straightforward to fit using hierarchical Bayesian methods since it requires inferring the identities of the groups to which individual cells belong. Here, we introduce a computational sampling method we call “Contour Monte Carlo” for estimating mathematical model parameters from snapshot distributions which is straightforward to implement and does not require explicitly assigning cells to categories. Our method is most applicable to systems where the dominant source of uncertainty is heterogeneity in cellular processes rather than experimental measurement error which, due to the increasingly finescale resolution of laboratory techniques, may be the case for a wide class of systems. In this paper, we illustrate the use of our method by quantifying cellular variation for two biological systems of interest and provide code in the form of a Julia notebook which allows others to apply this approach to their problem.

# 2 Introduction

27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

Variation rather than homogeneity is the rule rather than exception in biology. Indeed, without variation, biology as a discipline would not exist, since as evolutionary biologist JBS Haldane wrote, variation is the “raw material” of evolution. The Red Queen Hypothesis asserts that organisms must continually evolve in order to survive when pitted against other - also evolving - organisms [1]. A corollary of this hypothesis is that multicellular organisms may evolve cellular phenotypic heterogeneity to allow for faster adaptation to changing environments, which may explain the observed variation in a range of biological systems [2]. Whilst cell population variation can confer evolutionary advantages, it can also be costly in other circumstances. In biotechnological processes, heterogeneity in cellular function can lead to reduced yields of biochemical products [3]. In human biology, variation across cells can enable pathologies to develop and also prevents effective medical treatment, since medical interventions typically aim to steer modal cellular properties and hence fail to influence key subpopulations. For example, cellular heterogeneity likely contributes to the persistence of some cancerous tumours [4] and may also allow them to evolve resistance to chemotherapies over time [5]. Identifying and quantifying sources of variation in populations of cells is important for a wide range of applications because it allows us to determine whether this variability is benign or alternatively requires remedy.

Mathematical models are essential tools for making sense of cellular systems, whose emergent properties are the result of complex interactions

between various actors. Perhaps the simplest flavour of mathematical model used in biological systems are ordinary differential equations (ODEs) that lump individual actors into partitions according to structure or function, and seek to model the mean behaviour of each partition. Data from population-averaged experimental assays can be a powerful resource to understand whether such models faithfully reproduce system behaviours and can allow quantification of the interactions of various cellular components of complex metabolic, signalling and transcriptional networks. The worth of such models however is determined by whether averages mask differences in behaviour of individual cells that result in functional consequences [6]. In some cases, differences in cellular protein abundances due to biochemical “noise” may not be meaningful biologically [7] and so mean cell behaviour suffices as a description of the system, whereas in others there are functional consequences. For example, a recent study demonstrated that subpopulations of clonally derived hematopoietic progenitor cells with low or high expression of a particular stem cell marker produced different blood lineages [8].

To accommodate cell population heterogeneity in mathematical models, a variety of modelling choices are available, each posing different challenges for parameter inference, and are described in a recent review [9]. These include modelling biochemical processes stochastically, with properties of ensembles of cells represented by probability distributions evolving according to chemical master equations (see [10] for a tutorial on stochastic reaction-diffusion processes; RDEs). Alternatively, population balance equations (PBEs) can be used to dictate the evolution of the “number density” of differing cell types, whose properties are represented as points in  $\mathbb{R}^n$  which, in turn, affect their function, including their rate of death and cell division (see [11] for an introduction to PBEs). In a PBE approach, variation in measured quantities results primarily due to differing functional properties of heterogeneous cell types and variable initial densities of each type.

The approach we follow here is similar to that of [12], wherein dynamic cellular variation is generated by describing the evolution of each cell’s state using an ODE, but with individual cell differences in the rate parameters of the process. To our knowledge, this flavour of model is unnamed and so, for sake of reference, we term them “heterogenous ODE” models (HODEs). In HODEs, the aim of inference is to estimate the distributions of parameter values across cells consistent with observed distributions of measurements at various timepoints. A benefit of using HODEs to model cell heterogeneity is that these models are computationally straightforward to simulate and, arguably, simpler to parameterise than PBEs. In these models the predominant source of variation is due to differences in biological processes across cells not inherent stochasticity in biochemical reactions within cells, as in stochastic RDEs.

The difficulty of parameter inference for HODEs is partly due to experimental hurdles in generating data of sufficient quality to allow identification. Unlike models which represent a population by a single scalar ODE, since HODEs are individual-based they ideally require individual cell data for estimation. A widely-used method for generating data for individual cells is flow cytometry, where a large number of cells are streamed individually through a laser beam and, for example, abundance measurements are made of proteins labelled with fluorescent markers [13]. Alternatively, experimental techniques such as Western blotting and cytometric fluorescence microscopy can generate single cell measurements [14, 15]. A property of

these experimental methods is that they are destructive, meaning that individual cells are sacrificed as part of the measurement process. This means that the measurements of cell properties conducted at a certain point in time represent what are termed “snapshots” of the underlying population [15]. These snapshots are often described by histograms [12] or density functions [9] fit to the underlying data at different points in time. Since HODEs represent the underlying state of individual cells as evolving continuously through time, corresponding data showing individual cell trajectories constitutes a richer data resource. The demands of obtaining this data are higher however and typically involve either tracking individual cells through imaging methods [citation] or trapping cells in a spatial position where their individual dynamics can be readily monitored [citation]. These techniques impose restrictions on experimental practices meaning that they cannot be realised in all circumstances, including for online monitoring of biotechnological processes or analysis of *in vivo* studies. For this reason, snapshot data continues to play an important role for determining cell level variability in a wide variety of cases.

A variety of approaches have been proposed to estimate cell-to-cell variability by fitting HODE models to snapshot data. In HODEs, parameter values vary across cells according to a to-be-determined probability distribution meaning that in order to solve the exact inverse problem, the underlying ODE system needs to be simulated for each individual. Since the numbers of cells in these experiments are typically  $>\sim 10^4$  [15], this usually precludes exact inference due to its computational burden and instead the raw snapshot data is approximated by probability densities [12, 15–17]. Hasenauer et al. (2011) presents a Bayesian approach to inference for HODEs, which models the input parameter space using mixtures of ansatz densities, and use their method to reproduce population substructure on synthetic data generated from a model of tumour necrosis factor stimulus. Hasenauer et al. (2014) uses mixture-models to model the subpopulation structure in the snapshot data and uses multiple-start local optimisation to maximise the non-convex likelihood, which they then apply to a range of synthetic and real reaction data and signalling pathway examples. Loos et al. (2018) uses also uses mixture models to represent subpopulation structure and a maximum likelihood approach that allows for estimation of within- and between-subpopulation variability which also allows fitting to multivariate dependent output distributions. Dixit et al. (2018) discretises cell abundances into bins, then uses a maximum entropy approach as part of a Bayesian framework to fit the distribution representing cell-to-cell variability.

The framework we present here is Bayesian although is distinct from the traditional Bayesian inferential paradigm used to fit dynamic models since the source of stochasticity arises solely due to cell-to-cell parameter variation not measurement noise. Our approach is hence most suitable when measurement error is a minor contributor to observed experimental variability. Our computational method is a two-step Monte Carlo approach which, for reasons described in §3, we term “Contour Monte Carlo” (CMC). Unlike many of the existing methods however CMC is relatively computationally straightforward to implement and does not require extensive computation time. CMC uses MCMC in its second step to sample from the posterior distribution over parameter values and hence does not require specification of ansatz densities. It also does not require *a priori* representation of subpopulation structure using mixture components rather subpopulations appear


naturally as modes in the posterior parameter distributions. Like [17] CMC can fit multivariate snapshot data and unlike [12], does not require this data to be discretised into bins. As more experimental techniques are developed which elucidate single cell behaviour, there is likely to be more interest in methods which can be used to recapitulate the observed snapshots. We argue that due to its simplicity and generality, CMC is a useful addition to the modeller’s toolkit, which has a role to play in the analysis of the proliferation of rich single cell data.

Outline of the paper: In §3, we present the details of our methodological framework and detail the CMC algorithm we use to sample from the posterior parameter distribution. In §4, we then use CMC to estimate cell population heterogeneity in three systems of biological interest.

### 3 Method

In this section, we describe the first describe the probabilistic framework that underlies the CMC algorithm, before introducing CMC in pseudocode (Algorithm 1). We also detail the workflow we have found useful in applying this approach to analyse cell snapshot data and suggest practical remedies to issues we have encountered in using CMC (Figure 4). A glossary of all the variables used in this paper is included as Table 1.

Experimental methods such as flow cytometry can measure single cell characteristics at a given point in time. Cells are typically destroyed by the measurement process and so rather than providing time series for each individual cell, the data consists of cross-sections or “snapshots” of sampled individuals from the population (Figure 1).



**Figure 1: Time series data (A) versus snapshot data (B) typical of single cell experiments.** In A that the cell identities are retained at each measurement point (indicated by given plot markers) whereas in the snapshot data in B, either this information is lost or, more often, cells are destroyed by the measurement process and so each observation corresponds to a distinct cell.

We model the processes of an individual cell using a system of ordinary differential equations (ODEs), where each element of the system describes the governing dynamics of a particular quantity of interest (for example, protein levels, RNA concentrations and so on),

$$\dot{\mathbf{x}}(t) = f(\mathbf{x}(t); \boldsymbol{\theta}). \quad (1)$$


Here  $\mathbf{x}(t) = (x_1(t), x_2(t), \dots, x_k(t))$  is a vector of states for each compartment in the system at time  $t$  and  $f(\cdot)$  is a function of these states and parameters

$\theta \in \mathbb{R}^p$ . Note that in most circumstances, the initial state of the system,  $\mathbf{x}(0)$ , is unknown and it is convenient to include these as elements of  $\theta$  to be estimated. The solution of eq. (1) is given by  $\mathbf{x}(t) = g(t; \theta)$ , where  $\mathbf{x}(t) \in \mathbb{R}^k$  is a vector of outputs at time  $t$  and  $g(\cdot)$  is a function that typically won't be analytically-determined; instead approximated via a numerical integration scheme.

In this paper, we assume variation characterised by snapshot data arises due to between-cell heterogeneity in the underlying parameters  $\theta$ . Therefore, the evolution of the underlying state of cell  $i$  is described by an idiosyncratic ODE,

$$\dot{\mathbf{x}}^i(t) = f(\mathbf{x}^i(t); \theta^i), \quad (2)$$

with solution  $\mathbf{x}^i(t) = g(t; \theta^i)$ . The traditional (non-hierarchical) state-space approach to modelling dynamic systems supposes that measurement randomness generates output variation (Figure 2A). Our approach, by contrast, relies on the assumption that stochasticity in outputs is solely the result of variability in parameter values ( $\theta$ ) between cells (Figure 2B). Whether the assumption of “perfect” measurements is reasonable depends on the experimental details of the system under investigation but we argue that our method nevertheless provides a useful approximation in many cases where the signal to noise ratio is high.



**Figure 2: Two ways to generate randomness in measured outputs: the state-space model (A) versus the parameter heterogeneity model (B).** For non-hierarchical state-space models (A), there is assumed to be a single “true” latent state where observations result from a noisy measurement process (grey histograms). For models with parameter heterogeneity (B), the uncertainty is generated by differences in cellular processes (black lines) between cells. Note that in both cases, individual cells are measured only once in their lifetime.

Our generative model used to produce output observations for a single cell consists of two stages: (i) sample  $\theta^i \sim p(\theta)$ , where  $p(\theta)$  is a probability distribution characterising heterogeneity in cellular processes and, (ii) calculate output values using  $\mathbf{x}_j^i(t') = g(t'; \theta)$  where, for each cell, a measurement of a subset  $j$  of output states  $\mathbf{x}_j^i(t') \in \mathbf{x}^i$  is made at a single point in time  $t' \in (t_1, t_2, \dots, t_o)$ . The experimental observations for a single time point then consists of the collection of individual cell measurements  $\mathbf{X}_j(t') = (\mathbf{x}_j^1(t'), \mathbf{x}_j^2(t'), \dots, \mathbf{x}_j^n(t'))$ , where  $n$  is the number of cells measured at time  $t'$ . The entire observation dataset is the union of such sets across all measured time points  $\mathbf{X}(t) = (\mathbf{X}_{j_1}(t_1), \mathbf{X}_{j_2}(t_2), \dots, \mathbf{X}_{j_o}(t_o))$ , where  $t = (t_1, t_2, \dots, t_o)$  is the vector of observation times and the subscript  $a$  on each  $j_a$  allows for measurement of different elements of the system at distinct timepoints.

Variable	Definition	Dimension
$\boldsymbol{x}(t)$	state of modelled system at time $t$	$\mathbb{R}^k$
$x_j(t) \in \boldsymbol{x}(t)$	individual state $j$ of modelled system at time $t$	$\mathbb{R}$
$\boldsymbol{\theta}$	parameters of ODE system	$\mathbb{R}^p$
$f(\boldsymbol{x}(t); \boldsymbol{\theta})$	function specifying RHS of ODE system	$\mathbb{R}^k$
$g(t; \boldsymbol{\theta})$	solution of ODE at time $t$	$\mathbb{R}^k$
$g_j(t; \boldsymbol{\theta})$	solution of ODE for state $j$ at time $t$	$\mathbb{R}$
$\boldsymbol{x}^i(t)$	state of modelled system of cell $i$ at time $t$	$\mathbb{R}^k$
$\boldsymbol{x}_j^i(t) \in \boldsymbol{x}^i(t)$	state of subset $j$ of modelled system of cell $i$ at time $t$	$\mathbb{R}^{[j] \times 1}$
$\mathbf{X}_j(t) = (\boldsymbol{x}_j^1(t), \dots, \boldsymbol{x}_j^n(t))$	collection of $n$ individual cell measurements at time $t$	$\mathbb{R}^{[j] \times n}$
$\mathbf{t} = (t_1, t_2, \dots, t_o)$	unique observation times	$\mathbb{R}^o$
$\mathbf{X}(\mathbf{t}) = (\mathbf{X}_{j_1}(t_1), \dots, \mathbf{X}_{j_o}(t_o))$	all observations collected at times $\mathbf{t}$	$\dim(\mathbf{X}_{j_1}) \times \dots \times \dim(\mathbf{X}_{j_o})$
$\Phi$	parameters characterising output target distribution $p(\boldsymbol{x} \Phi)$	-
$\hat{a}$	estimates of any quantity $a$	-
$\tilde{\mathbf{t}} = (t_1, \dots, t_m)$	times when each observable is recorded	$\mathbb{R}^m$
$\tilde{\boldsymbol{x}}(\tilde{\mathbf{t}}) = (x_{j_1}(t_1), \dots, x_{j_m}(t_m))$	system observables	$\mathbb{R}^m$
$\mathcal{V}(\tilde{\boldsymbol{x}})$	the “volume” of parameter space mapping to an output of value $\tilde{\boldsymbol{x}}$	$\mathbb{R}^+$
$\mathbf{g}(\boldsymbol{\theta}) = \mathbf{g}(\tilde{\mathbf{t}}; \boldsymbol{\theta}) = (g_{j_1}(t_1; \boldsymbol{\theta}), \dots, g_{j_m}(t_m; \boldsymbol{\theta}))$	solution of ODE for each observable at respective times $\tilde{\mathbf{t}}$	$\mathbb{R}^m$
$V$	total volume of parameter space with uniform priors used for all parameters	$\mathbb{R}^+$
$\Omega(\tilde{\boldsymbol{x}}) = \{\boldsymbol{\theta} : \mathbf{g}(\boldsymbol{\theta}) = \tilde{\boldsymbol{x}}\}$	region of parameter space mapping to output $\tilde{\boldsymbol{x}}$	$\mathbb{R}^p$
$\Psi$	parameters characterising output prior distribution $p(\tilde{\boldsymbol{x}} \Psi)$	-
$\Xi$	parameters characterising parameter prior distribution $p(\boldsymbol{\theta} \Xi)$	-

Table 1: **Glossary of variable names used in this paper.** The dimensions of  $\Phi$ ,  $\Psi$  and  $\Xi$  are listed as “-” since they depend on the form of the density used to represent the process and can be anywhere from  $\mathbb{R}^1$  to  $\mathbb{R}^\infty$ . The variables are listed in the approximate order in which they appear in the text.

Raw snapshot data consists of measurements of individual cells with exact inference requiring simulating the underlying ODE system for each individual. This is cumbersome and impractical for the numbers of cells sampled in typical experimental setups and so, instead, we follow previous work and instead represent snapshot data using probability distributions [12, 15–17]. The snapshots themselves can either be distributions of a single species or multiple species, which can be approximated by univariate and multivariate probability distributions respectively. These probability distributions are characterised by parameter estimates  $\hat{\Phi}$  determined by the output observations  $\mathbf{X}(\mathbf{t})$ . The dimensionality of these probability distributions depends on the set of  $m$  distinct observables  $\tilde{\boldsymbol{x}}(\tilde{\mathbf{t}}) = (x_{j_1}(t_1), x_{j_2}(t_2), \dots, x_{j_m}(t_m))$  recorded by experimental measurements. Note that,  $\tilde{\boldsymbol{x}}(\tilde{\mathbf{t}})$  corresponds to a particular set of measurements from a hypothetical cell and is distinct from  $\mathbf{X}(\mathbf{t})$ , which represents the full set of experimental outputs. The vector  $\tilde{\boldsymbol{x}}(\tilde{\mathbf{t}})$  is hypothetical because in reality each cell is measured at a single timepoint (although we suppose measurements of different cellular attributes are possible contemporaneously).

The goal of our inference process is to characterise the probability distribution  $p(\boldsymbol{\theta}|\mathbf{X}(\mathbf{t}))$  representing heterogeneity in cellular processes. The first step in our inference workflow is to fit the output distributions using probability distributions (Figure 4(i)). We assume that the volume of observational data means the estimated probability distributions are approximate sufficient statistics of the outputs, meaning  $p(\boldsymbol{\theta}|\hat{\Phi}) \approx p(\boldsymbol{\theta}|\mathbf{X}(\mathbf{t}))$ .




Figure 3: The non-linear mapping from parameter values (left panel) to outputs (black lines; right panel) means different sized regions of parameter space (orange and blue surfaces; left panel) correspond to distinct output values (orange triangle and blue square; right panel). In the right panel, each black line represents a distinct model simulation  $g(t; \theta_1, \theta_2, \theta_3)$  and the triangle and square indicate outputs at a given point in time.

The models we seek to fit to snapshot data mostly cannot be identified from the observations. This is often because the number of model parameters exceeds the dimensionality of the output distribution (that is,  $m > k$ ) meaning there typically exist non-singular sets of parameter values mapping to a single set of output values. That is, each vector of observed outputs  $\tilde{\mathbf{x}} \in \mathbb{R}^m$ , can often be caused by many combinations of parameters although, due to the non-linearity of the map from parameters to outputs, the “volume” of these regions of parameter space,  $\mathcal{V}(\tilde{\mathbf{x}})$ , is a function of output (Figure 3). In what follows, we make clear the distinction between observables  $\tilde{\mathbf{x}}(\tilde{\mathbf{t}})$  and the vector-valued function representing modelled outputs  $\mathbf{g}(\tilde{\mathbf{t}}; \boldsymbol{\theta}) = (g_{j_1}(t_1; \boldsymbol{\theta}), g_{j_2}(t_2; \boldsymbol{\theta}), \dots, g_{j_m}(t_m; \boldsymbol{\theta})) \in \mathbb{R}^m$  since the latter is a function whereas that latter is a numeric value; we also drop the  $\tilde{\mathbf{t}}$  notation from following expressions to minimise clutter.

A consequence of this non-linear parameter to output geometry is that any target output distribution  $p(\tilde{\mathbf{x}}|\hat{\Phi})$  does not correspond to a unique parameter distribution. For example, suppose  $g(\theta_1, \theta_2) = \theta_1 + \theta_2$ : the target distribution  $\tilde{\mathbf{x}} \sim \mathcal{N}(0, 1)$  can be generated by any member of the set of parameter distributions  $\sqrt{\eta}\theta_1 + \sqrt{1 - \eta}\theta_2$ , where  $\eta \in [0, 1]$  and  $\theta_1, \theta_2 \sim \mathcal{N}(0, 1)$ . This means that in order to ensure uniqueness of the “posterior” parameter distributions, we are required to specify “prior” distributions for the parameters, as in more traditional Bayesian inference. An additional consequence of the degeneracy of the mapping from parameters to outputs is that any sampling algorithm aimed at exploring posterior parameter space must account for the differential volumes of iso-output contours. Whilst we refer the interested reader to our companion paper on this subject [citation for tutorial paper published in Open Science], we provide a quick derivation of the posterior parameter distribution which accounts for the non-linear mapping.

To derive the posterior distribution of parameter values  $p(\boldsymbol{\theta}|\hat{\Phi})$ , we consider the joint density of parameters and outputs  $p(\boldsymbol{\theta}, \tilde{\mathbf{x}}|\hat{\Phi})$ . This can be decomposed in two ways,

$$p(\boldsymbol{\theta}|\tilde{\mathbf{x}}, \hat{\Phi}) \times p(\tilde{\mathbf{x}}|\hat{\Phi}) = p(\boldsymbol{\theta}, \tilde{\mathbf{x}}|\hat{\Phi}) = p(\tilde{\mathbf{x}}|\boldsymbol{\theta}, \hat{\Phi}) \times p(\boldsymbol{\theta}|\hat{\Phi}). \quad (3)$$

The left and right hand sides of eq. (3) can be equated and rearranged to

obtain the posterior parameter distribution,

274

$$p(\boldsymbol{\theta}|\hat{\Phi}) = \frac{p(\boldsymbol{\theta}|\tilde{\mathbf{x}}, \hat{\Phi}) \times p(\tilde{\mathbf{x}}|\hat{\Phi})}{p(\tilde{\mathbf{x}}|\boldsymbol{\theta}, \hat{\Phi})}. \quad (4)$$

Given parameters  $\boldsymbol{\theta}$ , the mapping from parameters to outputs is deterministic meaning  $p(\tilde{\mathbf{x}}|\boldsymbol{\theta}, \hat{\Phi}) = \delta(\mathbf{g}(\boldsymbol{\theta}))$  is the Dirac delta function centred at  $\tilde{\mathbf{x}} = \mathbf{g}(\boldsymbol{\theta})$ . In what follows, we assume that the conditional distribution  $p(\boldsymbol{\theta}|\tilde{\mathbf{x}}, \hat{\Phi})$  is independent of the data, meaning it represents a conditional “prior”, which can be manipulated by Bayes’ rule,

275

276

277

278

279

$$p(\boldsymbol{\theta}|\mathbf{g}(\boldsymbol{\theta})) = \frac{p(\boldsymbol{\theta})}{p(\mathbf{g}(\boldsymbol{\theta}))}, \quad (5)$$

where we have used the Dirac delta function for  $p(\tilde{\mathbf{x}}|\boldsymbol{\theta})$ . This results in the form of the posterior parameter distribution targeted by our sampling algorithm,

280

281

282

$$p(\boldsymbol{\theta}|\hat{\Phi}) = \frac{p(\boldsymbol{\theta})}{p(\mathbf{g}(\boldsymbol{\theta}))} p(\mathbf{g}(\boldsymbol{\theta})|\hat{\Phi}). \quad (6)$$

Again, we refer to our companion piece [citation] for detailed explanation of eqs. (5) & (6) and instead here provide brief interpretation when considering a uniform prior on parameter space. In this case,  $p(\boldsymbol{\theta}) = \frac{1}{V}$ , where  $V$  is the total volume of parameter space. The denominator term of eq. (5) is the prior induced on output space by the prior over parameter space. For a uniform prior on parameter values, this is just proportion of parameter space where  $\mathbf{g}(\boldsymbol{\theta}) = \tilde{\mathbf{x}}$ , meaning,

283

284

285

286

287

288

289

$$p(\boldsymbol{\theta}|\mathbf{g}(\boldsymbol{\theta})) = \frac{1}{\mathcal{V}(\mathbf{g}(\boldsymbol{\theta}))}, \quad (7)$$

where  $\mathcal{V}(\mathbf{g}(\boldsymbol{\theta}))$  is the volume of parameter space occupied by the iso-output region  $\Omega(\tilde{\mathbf{x}}) = \{\boldsymbol{\theta} : \mathbf{g}(\boldsymbol{\theta}) = \tilde{\mathbf{x}}\}$ . Therefore a uniform prior over parameter space implies a prior structure where all parameter values resulting in the same output  $\tilde{\mathbf{x}}$  are given equal weighting.

290

291

292

293

The denominator term of eq. (5) cannot be calculated apart from for some toy examples, meaning that exact sampling from the posterior parameter distribution of eq. (6) is not, in general, possible. We propose instead a computationally efficient sampling method to estimate  $p(\mathbf{g}(\boldsymbol{\theta}))$ , which forms the first step of our so-called “Contour Monte Carlo” (CMC) algorithm (Algorithm 1; Figure 4(ii)), where we estimate the volume of iso-output contours with output value  $\mathbf{g}(\boldsymbol{\theta})$ . This step involves repeated independent sampling from the prior distribution of parameters  $\boldsymbol{\theta}^i \sim p(\boldsymbol{\theta}|\Xi)$ , where, for completeness, we have conditioned on  $\Xi$  parameterising our probability density. Each parameter sample is then converted into an output value  $\tilde{\mathbf{x}}^i = \mathbf{g}(\boldsymbol{\theta}^i)$ . The collection of output samples is then fitted using a vine copula kernel density estimator (KDE) [18],  $(\tilde{\mathbf{x}}^1, \tilde{\mathbf{x}}^2, \dots, \tilde{\mathbf{x}}^{N_1}) \sim p(\tilde{\mathbf{x}}|\hat{\Psi})$ . Throughout the course of development of CMC, we have tested many forms of KDE and have found vine copula KDE is best suited to approximating the higher dimensional probability distributions required in practice.

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

The second step in our algorithm then uses Markov chain Monte Carlo (MCMC) to sample from an approximate version of eq. (6) with the estimated density  $p(\mathbf{g}(\boldsymbol{\theta})|\hat{\Psi})$  replacing its corresponding estimand (Algorithm 1; Figure 4(iii)),

$$p(\boldsymbol{\theta}|\hat{\Phi}, \Xi, \hat{\Psi}) = \frac{p(\boldsymbol{\theta}|\Xi)}{p(\mathbf{g}(\boldsymbol{\theta})|\hat{\Psi})} p(\mathbf{g}(\boldsymbol{\theta})|\hat{\Phi}). \quad (8)$$

---

**Algorithm 1** Pseudocode for the Contour Monte Carlo algorithm for sampling from the posterior parameter distribution of eq. (8). Here we provide code for the Random Walk Metropolis algorithm for the MCMC sampling but for the examples in §4, we use an adaptive MCMC algorithm [19]. A definition of all variables is provided in Table 1.

---

```

procedure CMC( $\mathbf{X}(t), \Xi, N_1, N_2$ )  $\triangleright$  Sample from posterior parameter distribution
     $\hat{\Phi} = \text{SNAPSHOTESTIMATOR}(\mathbf{X}(t))$ 
     $\hat{\Psi} = \text{CONTOURVOLUMEESTIMATOR}(\Xi)$ 
     $(\boldsymbol{\theta}_1, \dots, \boldsymbol{\theta}_{N_2}) = \text{MCMC}(\hat{\Phi}, \Xi, \hat{\Psi})$ 
    converged = COMPAREOUTPUTTOTARGET(( $\boldsymbol{\theta}_1, \dots, \boldsymbol{\theta}_{N_2}$ ),  $\hat{\Psi}$ )
    if converged  $\neq 1$  then
         $(\boldsymbol{\theta}_1, \dots, \boldsymbol{\theta}_{N_2}) = \text{CONTOURVOLUMEESTIMATOR}(\hat{\Phi}, \Xi, N'_1, N'_2)$ 
        where,  $N'_1 > N_1$  and/or  $N'_2 > N_2$   $\triangleright$  Rerun contour volume estimation and/or
        MCMC with larger sample sizes
    end if
    return ( $\boldsymbol{\theta}_1, \dots, \boldsymbol{\theta}_{N_2}$ )
end procedure

procedure SNAPSHOTESTIMATOR( $\mathbf{X}(t)$ )  $\triangleright$  Fit density to snapshot observations
     $\mathbf{X}(t) \sim p(\tilde{\mathbf{x}}|\hat{\Phi})$ 
    return  $\hat{\Phi}$ 
end procedure

procedure CONTOURVOLUMEESTIMATOR( $\Xi$ )  $\triangleright$  Estimate volume of contours
    for  $i$  in  $1 : N_1$  do
         $\boldsymbol{\theta}_i \sim p(\boldsymbol{\theta}|\Xi)$   $\triangleright$  Sample from prior density
         $\tilde{\mathbf{x}}_i = \mathbf{g}(\boldsymbol{\theta}_i)$   $\triangleright$  Calculate corresponding output value
    end for
     $(\tilde{\mathbf{x}}_1, \tilde{\mathbf{x}}_2, \dots, \tilde{\mathbf{x}}_{N_1}) \sim p(\tilde{\mathbf{x}}|\hat{\Psi})$   $\triangleright$  Fit vine copula kernel density estimator to output
    values.
    return  $\hat{\Psi}$ 
end procedure

procedure MCMC( $\hat{\Phi}, \Xi, \hat{\Psi}$ )  $\triangleright$  Random Walk Metropolis algorithm targeting posterior
    parameter distribution.
     $\boldsymbol{\theta}_0 \sim \pi(\cdot)$   $\triangleright$  Sample from arbitrary initialisation distribution
    for  $i$  in  $1 : N_2$  do
         $\boldsymbol{\theta}'_i \sim \mathcal{N}(\boldsymbol{\theta}_{i-1}, \Sigma)$   $\triangleright$  Propose new parameter values for parameters
         $r = \left[ p(\boldsymbol{\theta}'|\Xi) p(\mathbf{g}(\boldsymbol{\theta})|\hat{\Psi}) p(\mathbf{g}(\boldsymbol{\theta}')|\hat{\Phi}) \right] / \left[ p(\boldsymbol{\theta}|\Xi) p(\mathbf{g}(\boldsymbol{\theta})|\hat{\Psi}) p(\mathbf{g}(\boldsymbol{\theta})|\hat{\Phi}) \right]$   $\triangleright$  Metropolis
        acceptance ratio.
         $u \sim U(0, 1)$   $\triangleright$  Sample from uniform distribution
        if  $r > u$  then
             $\boldsymbol{\theta}_i = \boldsymbol{\theta}'_i$   $\triangleright$  Accept proposal
        else
             $\boldsymbol{\theta}_i = \boldsymbol{\theta}_{i-1}$   $\triangleright$  Reject proposal
        end if
    end for
    return ( $\boldsymbol{\theta}_1, \dots, \boldsymbol{\theta}_{N_2}$ )
end procedure

procedure COMPAREOUTPUTTOTARGET(( $\boldsymbol{\theta}_1, \dots, \boldsymbol{\theta}_{N_2}$ ),  $\hat{\Psi}$ )  $\triangleright$  Check output distribution
    close to target
    for  $i$  in  $1 : N_2$  do
         $\tilde{\mathbf{x}}_i = \mathbf{g}(\boldsymbol{\theta}_i)$   $\triangleright$  Compute output for each parameter sample
    end for
    if  $(\tilde{\mathbf{x}}_1, \tilde{\mathbf{x}}_2, \dots, \tilde{\mathbf{x}}_{N_2}) \sim p(\tilde{\mathbf{x}}|\hat{\Psi})$ ? then
        return 1  $\triangleright$  Compare outputs with target
         $\triangleright$  If outputs sufficiently close then converged
    else
        return 0
    end if
end procedure

```

---




Figure 4: **The workflow for using Contour Monte Carlo to estimate cell population heterogeneity.** In (iii), the distribution targeted is given by eq. (8) but with  $p(\tilde{x}|\hat{\Psi}) \rightarrow p(g(\theta))$ . The variables used in this figure are defined in the text and Table 1.

The final step in CMC is to transform parameter samples from the MCMC into outputs, then compare the sampled outputs with the target distribution (Figure 4(iv)). Asymptotically (in terms of the sample size of both sampling steps), CMC produces a sample of parameter values  $(\theta^1, \theta^2, \dots)$  which, when transformed to outputs, corresponds to the target distribution  $p(\tilde{x}|\hat{\Psi})$ . In developing CMC, we have found that a finite sample of modest size for both steps of CMC results in parameter samples that, when transformed, often represent reasonable approximations of the target. There are however occasions when this is not the case and we have found this final confirmatory step indispensable since it frequently highlights inadequacies in the contour volume estimation or the MCMC, meaning more samples from either or both of these steps are required. It may also be necessary to tweak hyperparameters of the KDE to ensure reasonable approximation in the contour volume estimation step. If the target distribution is sensitive to the contour volume estimates, this may also indicate that the target snapshot distribution is incompatible with the model: here, we make no claims on existence of a solution to the inverse problem, only that, if one should exist, Contour Monte Carlo is a pragmatic approach to approximate it by sampling. A useful way to diagnose whether the target distribution can be produced from the model and specified priors is to examine the output values from the contour volume estimation step of CMC. If the majority of probability mass of the target lies outside the bounds of the bulk simulated output values obtained by independent sampling from the prior, then the model and/or chosen prior is unlikely to be invertible to this particular target.

For the contour volume estimation step, we assumed sample sizes were sufficient if the output samples from the MCMC provided a reasonable approximation to the target, although we recognise that future work should refine this process further. For the MCMC step, we use adaptive covariance MCMC (see SOM of [19]) to sample from the target distribution, as we have found that it provides a considerable speed-up over Random Walk Metropolis [20, 21]. We also use the Gelman-Rubin convergence statistic  $\hat{R}$  which provides a heuristic measurement of convergence [21, 22], and use a

313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345

## 4 Results

In this section, we use CMC to estimate the posterior parameter distribution for three biological systems of interest targeting synthetic parametric densities. That is, we assume that the first step of CMC (“SnapshotEstimator” within Algorithm 1) has already been undertaken and we are faced with inferring a parameter distribution which, when transformed to outputs, recapitulates the target density. Alongside the text, we also provide a Julia notebook used to generate the results, which we hope will be of use to others wanting to apply CMC to estimate cell population heterogeneity.

Model	Target density	Parameter	Prior density	Prior parameter 1	Prior parameter 2
Growth factor	two-dimensional normal	$R_T$	uniform	$2.5 \times 10^5$	$8 \times 10^5$
		$k_1$	uniform	0.25	3.0
		$k_{-1}$	uniform	2.0	20.0
		$k_{deg}$	uniform	0.005	0.03
		$k_{deg}^*$	uniform	0.1	0.5
Growth factor	two-dimensional normal	$R_T$	normal	$5 \times 10^5$	$1 \times 10^5$
		$k_1$	normal	0.5	0.1
		$k_{-1}$	normal	3.0	1.0
		$k_{deg}$	normal	0.02	0.005
		$k_{deg}^*$	normal	0.3	0.1
Michaelis-Menten kinetics	bimodal normal	$k_f$	uniform	0.2	15
		$k_r$	uniform	0.2	2.0
		$k_{cat}$	uniform	0.5	3.0
Michaelis-Menten kinetics	four-dimensional normal	$k_f$	uniform	0.2	15
		$k_r$	uniform	0.2	2.0
		$k_{cat}$	uniform	0.2	3.0
		$E_0$	uniform	3.0	5.0
		$S_0$	uniform	5.0	10.0
		$ES_0$	uniform	0.0	0.2
		$P_0$	uniform	0.0	0.2
TNF signalling	bivariate normal	$a_1$	uniform	0.4	0.8
		$a_2$	uniform	0.1	0.7
		$a_3$	uniform	0.3	0.7
		$a_4$	uniform	0.1	0.3
		$b_1$	uniform	0.5	0.7
		$b_2$	uniform	0.4	0.6
		$b_3$	uniform	0.4	0.6
		$b_4$	uniform	0.2	0.4
		$b_5$	uniform	0.2	0.4
TNF signalling	bimodal normal	$a_1$	uniform	0.5	0.7
		$a_2$	uniform	0.1	0.3
		$a_3$	uniform	0.1	0.3
		$a_4$	uniform	0.4	0.6
		$b_1$	uniform	0.3	0.5
		$b_2$	uniform	0.6	0.8
		$b_3$	uniform	0.2	0.4
		$b_4$	uniform	0.4	0.6
		$b_5$	uniform	0.3	0.5

Table 2: The priors used for each problem in §4.

### 4.1 Growth factor model

Here we consider the “growth factor model” introduced by [12], which concerns the dynamics of inactive ligand-free cell surface receptors  $R$  and active ligand-bound cell surface receptors  $P$ , modulated by an exogenous

ligand  $L$ . The governing dynamics are determined by the following system,

$$\dot{R}(t) = R_T k_{deg} + k_1 L R(t) + k_{-1} P(t) - k_{deg} R(t) \quad (9)$$


$$\dot{P}(t) = k_1 L R(t) - k_{-1} P(t) - k_{deg}^* P(t), \quad (10)$$

where  $\theta = (R_T, k_1, k_{-1}, k_{deg}, k_{deg}^*)$  are parameters to be determined. In this example, we use measurements of the active ligand-bound receptors  $P$  to estimate cellular heterogeneity in processes. We denote the solution of eq. (9) as  $P(t; \theta, L)$  and seek to determine the parameter distribution consistent with an output distribution,

$$\begin{pmatrix} P(10; \theta, 2) \\ P(10; \theta, 10) \end{pmatrix} \sim \mathcal{N} \left[ \begin{pmatrix} 2 \times 10^4 \\ 3 \times 10^4 \end{pmatrix}, \begin{pmatrix} 1 \times 10^5 & 0 \\ 0 & 1 \times 10^5 \end{pmatrix} \right]. \quad (11)$$

To start, we specify a uniform prior for each of the five parameters, with bounds given in Table 2. To estimate the posterior parameter distribution, we use CMC, with adaptive covariance MCMC [19] for the second step.

In Figure 5A, we show the sampled outputs (blue points) versus the contours of the target distribution (black solid closed curves), illustrating a good correspondence between the sampled and target densities. Above and to the right of the main panel, we also display the marginal target densities (solid black lines) versus kernel density estimator reconstructions of the output marginals from the CMC samples (dashed blue lines), which again highlights the fidelity of the CMC sampled density to the target.



**Figure 5: The target joint output distribution (solid contour lines) and target marginal distributions (solid) versus outputs sampled by CMC (blue points and dashed lines) for (A) uniform and (B) normal parameter priors.** In the CMC, 100,000 independent samples were used in the “ContourVolumeEstimator” step and 10,000 MCMC samples across each of 4 Markov chains were used in the second step, with the first half of the chains discarded as “warm-up” [21]. For the reconstructed marginal densities in the plots, we use Mathematica’s “SmoothKernelDistribution” function specifying bandwidths of 100 with Gaussian kernels [23].

In Figure 6A, we plot the joint posterior parameter distribution for  $k_1$ , the rate of ligand binding to inactive receptors, and  $k_{-1}$ , which dictates the rate of the reverse reaction, where the ligands unbind. The output measurements we used to fit the model correspond to levels of the bound

ligands, which can be generated whenever the ratio of  $k_1$  to  $k_{-1}$  approximately equals their steady state ratio. Because of this, the distribution representing cell process heterogeneity contains linear positive correlations between these parameters. In Figure 6B, we show the posterior parameter distribution for  $k_{deg}$ , the rate of degradation of ligand-free cell surface receptors and  $R_T$ , the steady state level of this species, which shows a greater concentration of posterior probability mass across prior space. Why is it that we are better able to resolve  $(k_{deg}, R_T)$  compared to  $(k_1, k_{-1})$  from our measurements? To answer this, it is useful to calculate the sensitivity of  $P(t; \theta, L)$  to changes in each of the parameters. To account for the differing magnitudes of each parameter, we calculate elasticities, the proportional changes in measured output for a proportional change in parameter values, using the forward sensitivities method described in [24], which are shown in Figure 7. When the exogenous ligand is set  $L = 2$ , these indicate the active ligand-bound receptor concentration is most elastic to changes in  $R_T$  and  $k_{deg}$ , meaning that their range is more restricted by the output measurement than for  $k_1$  and  $k_{-1}$ , which have elasticities at  $t = 10$  closer to 0. In Table 3, we show the posterior quantiles for the estimated parameters and, in the last column, indicate the ratio of the 25%-75% posterior interval widths to the uniform prior range for each parameter. These were strongly negatively correlated with the magnitude of the elasticities for each parameter ( $\rho = 0.95$ ,  $t = -5.22$ ,  $df = 3$ ,  $p = 0.01$  Pearson's product-moment correlation), indicating the utility of sensitivity analyses for optimal experimental design. We would suggest however that CMC can also be used for this purpose, using synthetic data in place of real measurements.




Figure 6: The joint distributions of  $(k_1, k_{-1})$  (left-column) and  $(k_{\text{deg}}, R_T)$  for the growth factor model using uniform priors (top row) and normal priors (bottom row). See Figure 5 caption for CMC details and Table 2 for the priors used.




Figure 7: The elasticities of the measured concentration of active ligand-bound receptors  $P$  versus time when  $L = 2$ . When calculating the elasticities of each parameter, the other parameters were set to their posterior medians given in Table 3.

For an unidentified model, there are typically a multitude of possible probability distributions over parameters which map to the same target output distribution. To reduce the space of posterior parameter distributions

402

403

404

Prior	Parameter	2.5%	25%	Quantiles 50%	75%	97.5%	Posterior 25%-75% concentration
Uniform	$R_T$	441,006	548,275	606,439	677,055	772,484	23%
	$k_1$	0.90	1.69	2.17	2.56	2.95	32%
	$k_{-1}$	4.35	8.35	11.23	14.23	18.71	33%
	$k_{deg}$	0.013	0.019	0.021	0.024	0.029	20%
	$k_{deg}^*$	0.20	0.34	0.40	0.44	0.49	27%
Normal	$R_T$	408,396	487,372	529,558	577,970	678,632	16%
	$k_1$	0.39	0.49	0.54	0.60	0.70	4%
	$k_{-1}$	1.39	1.92	2.26	2.63	3.35	4%
	$k_{deg}$	0.016	0.020	0.022	0.024	0.027	16%
	$k_{deg}^*$	0.22	0.29	0.33	0.38	0.46	21%

Table 3: **Estimated quantiles from CMC samples for the growth factor model with uniform and normal priors.** The last column indicates the proportion of the uniform prior bounds occupied by the 25%-75% posterior interval in each case. The particular priors used in each case are given in Table 2.

to one, it is therefore necessary to specify a prior parameter distribution. CMC accommodates the effect of changes to a prior distribution directly, through changes to both the ‘‘ContourVolumeEstimation’’ step and the acceptance ratio in the ‘‘MCMC’’ step (Algorithm 1), such that the posterior parameter distribution maps to the same output target. We now use CMC to estimate the posterior parameter distribution when changing from uniform to normal priors (prior hyperparameters shown in Table 2). As desired, the target output distribution appears invariant (Figure 5B) although with substantial changes in the posterior parameter distributions (Figure 6C&D). In particular, the posterior distributions obtained from shifting to the normal prior are more concentrated in parameter space compared to the uniform case (rightmost column of Table 3).

## 4.2 Michaelis-Menten kinetics

In this section, we use CMC to invert output measurements from the Michaelis-Menten model of enzyme kinetics (see, for example, [25]); illustrating the capability of CMC to resolve population substructure from multimodality of the output distribution. The Michaelis-Menten model of enzyme kinetics describes the dynamics of concentrations of an enzyme ( $E$ ), a substrate ( $S$ ), an enzyme-substrate complex ( $ES$ ), and a product ( $P$ ). Specifically,

$$\begin{aligned}\dot{E}(t) &= -k_f E(t)S(t) + k_r ES(t) + k_{cat}ES(t), \\ \dot{S}(t) &= -k_f E(t)S(t) + k_r ES(t), \\ \dot{ES}(t) &= k_f E(t)S(t) - k_r ES(t) - k_{cat}ES(t), \\ \dot{P}(t) &= k_{cat}ES(t),\end{aligned}\tag{12}$$

with initial conditions

$$E(0) = E_0, S(0) = S_0, ES(0) = ES_0, P(0) = P_0,\tag{13}$$

where  $k_f$  is the rate constant for the forward reaction  $E + S \rightarrow ES$ ,  $k_r$  is the rate of the reverse reaction  $ES \rightarrow E + S$ , and  $k_{cat}$  is the catalytic rate at which the product is formed by the reaction  $ES \rightarrow E + P$ .

When subpopulations of cells, each with distinct dynamics, are thought to exist, determining their characteristics - proportions of overall cell number,

likely parameter values, and so on - is often of key interest [15,17]. Before formal inference occurs, multi-modality of the output distribution may signal the existence of fragmented subpopulations of cells. Here we target the following bimodal bivariate normal distribution,

$$f(\mathbf{x}; \boldsymbol{\mu}_1, \Sigma_1, \boldsymbol{\mu}_2, \Sigma_2) = \frac{1}{2} (\mathcal{N}(\mathbf{x}; \boldsymbol{\mu}_1, \Sigma_1) + \mathcal{N}(\mathbf{x}; \boldsymbol{\mu}_2, \Sigma_2)), \quad (14)$$

where  $E(2; \boldsymbol{\theta})$  and  $S(1; \boldsymbol{\theta})$  corresponds to the solutions of eq. (12) for the enzyme and substrate at times  $t = 2$  and  $t = 1$ , respectively, and  $\boldsymbol{\theta} = (k_f, k_r, k_{cat})$ . The parameters of the mixture normal output distribution we target are  $\boldsymbol{\mu}_1 = [2.2, 1.6]', \Sigma_1 = \begin{pmatrix} 0.018 & -0.013 \\ -0.013 & 0.010 \end{pmatrix}$ ,  $\boldsymbol{\mu}_2 = [2.8, 1.0]'$  and  $\Sigma_2 = \begin{pmatrix} 0.020 & -0.010 \\ -0.010 & 0.020 \end{pmatrix}$ . In what follows, we specify uniform priors on each of the elements of  $\boldsymbol{\theta}$  (see Table 2).

Using a modest number of samples in each step, CMC was able to well-replicate the output target distribution (Figure 8A). Without specifying *a priori* information on the subpopulations of cells, two distinct clusters of cells emerged from application of CMC (orange and blue points in Figure 8B), each corresponding to distinct modes of the output distribution (corresponding coloured points in Figure 8A). It is worth noting however that the issues inherent with MCMC sampling of multimodal distributions similarly apply here and so, whilst here adaptive MCMC [19] sufficed to explore the posterior surface, it may be necessary to use MCMC methods known to be robust to such geometries (for example, population MCMC [26]).




Figure 8: **(A) bimodal target distribution (solid contour lines) versus output samples (points) for the Michaelis-Menten model and (B) posterior parameter samples (points).** The orange (blue) points in A were generated by the orange (blue) parameter samples in B. See Figure 5 caption for CMC details. For the reconstructed marginal densities in the plots, we use Mathematica’s “SmoothKernelDistribution” function with Gaussian kernels with (A) default bandwidths and (B) bandwidths of 0.3 (horizontal axis) and 0.03 (vertical axis) [23]. The clusters were identified by applying Mathematica’s “ClusteringComponents” function to the pairs of parameter samples displayed in B [23].

Loos et al. (2018) consider a multidimensional output distribution, with correlations between system characteristics that evolve over time. Our

approach allows arbitrary covariance structure between measurements, and to exemplify this, we now target a four-dimensional output distribution, with paired measurements of enzyme and substrate at  $t = 1$  and  $t = 2$ , 448  
449  
450

$$\begin{pmatrix} E(1.0; \theta) \\ S(1.0; \theta) \\ E(2.0; \theta) \\ S(2.0; \theta) \end{pmatrix} \sim \mathcal{N} \left[ \begin{pmatrix} 0.5 \\ 2.8 \\ 0.9 \\ 1.4 \end{pmatrix}, \begin{pmatrix} 0.02 & -0.05 & 0.04 & -0.05 \\ -0.05 & 0.30 & -0.15 & 0.20 \\ 0.04 & -0.15 & 0.12 & -0.17 \\ -0.05 & 0.20 & -0.17 & 0.30 \end{pmatrix} \right]. \quad (15)$$

Since this system has four output measurements, and the Michaelis-Menten model has three rate parameters ( $k_f, k_r, k_{cat}$ ), the system is over-identified and so CMC cannot be straightforwardly applied. Instead, we allow the four initial states ( $E_0, S_0, ES_0, P_0$ ) to be uncertain quantities, bringing the total number of parameters to 7, and ensuring that the system is in the unidentified regime where CMC applies. We set uniform priors on all parameters (see Table 2) and to check that the model and priors were consistent with the output distribution given by eq. (15), we plotted the output measurements used to estimate contour volumes (in the first step of the “ContourVolumeEstimator” method in Algorithm 1) against the target (Figure 9). Since the main support of the densities (black contours) lies within a region of output space reached by independent sampling of the priors (blue points), this indicated that the distribution given by eq. (15) could feasibly be generated from this model and priors, and we proceeded to estimation by CMC. 451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465




Figure 9: The output measurements (blue points) of enzyme and substrate at times  $t = 1$  (left panel) and  $t = 2$  (right panel) obtained by independently sampling the priors of the 7 parameter Michaelis-Menten model versus the target distribution (black solid contours). In this plot, we show 20,000 output samples, where each set of four measurements was obtained from a single sample of the 7 parameters. The output target distribution shown by the contours corresponds to the marginal densities of each pair of enzyme-substrate measurements given by eq. (15).

Figure 10 plots the output samples of enzyme and substrate from the last step of CMC for  $t = 1$  (blue points) and  $t = 2$  (orange points) versus the contours (black lines) of the joint marginal distributions of 466  
467  
468

eq. (15). The distribution of paired enzyme-substrate samples illustrates that the CMC output samples approximated the target density, itself representing dynamic evolution of the covariance between enzyme and substrate measurements. The target marginal distributions (solid lines) along with their approximations from kernel density estimation (dashed lines) are also shown above and beside the main panel of Figure 10, and largely indicate correspondence.

469  
470  
471  
472  
473  
474  
475




Figure 10: **Posterior output samples from CMC (coloured points) versus the contour plots of the joint marginal distributions of eq. (15) (black solid lines).** The blue and orange points indicate output observations of enzyme (horizontal axis) and substrate (vertical axis) at times  $t = 1$  and  $t = 2$  respectively. The solid and dotted coloured lines outside of the panels indicate the true target marginals of eq. (15) and those estimated from CMC, respectively. The line colours correspond with the point colours and indicate the marginals for  $t = 1$  (blue) and  $t = 2$  (orange). In the CMC, 200,000 independent samples were used in the “ContourVolumeEstimator” step and 10,000 MCMC samples across each of 4 Markov chains were used in the second step, with the first half of the chains discarded as “warm-up” [21]. For the reconstructed marginal densities in the plots, we use Mathematica’s “SmoothKernelDistribution” function with Gaussian kernels [23] of bandwidths varying from 0.1 to 0.4.

476  
477  
478  
479  
480  
481  
482

### 4.3 TNF signalling pathway

We now illustrate how CMC can be applied to an ODE system of larger size, the tumour necrosis factor (TNF) signalling pathway model introduced in [27] and used by [15] to illustrate a Bayesian approach to cell population variability estimation. The model incorporates known activating and inhibitory interactions between four key species within the TNF pathway: active caspase 8 ( $X_1$ ) and active caspase 3 ( $X_2$ ), a nuclear transcription

factor ( $X_3$ ) and its inhibitor ( $X_4$ ),

483

$$\begin{aligned}\dot{X}_1(t) &= -X_1(t) + \frac{1}{2} (\text{inh}_4(X_3(t))\text{act}_1(u(t)) + \text{act}_3(X_2(t))) \\ \dot{X}_2(t) &= -X_2(t) + \text{act}_2(X_1(t))\text{inh}_3(X_3(t)) \\ \dot{X}_3(t) &= -X_3(t) + \text{inh}_2(X_2(t))\text{inh}_5(X_4(t)) \\ \dot{X}_4(t) &= -X_4(t) + \frac{1}{2} (\text{inh}_1(u(t)) + \text{act}_4(X_3(t))),\end{aligned}\tag{16}$$

where the functions  $\text{act}_i$  and  $\text{inh}_j$  represent activating and inhibitory interactions respectively,

484  
485

$$\begin{aligned}\text{act}_i(X) &= \frac{X^2}{a_i^2 + X^2} \\ \text{inh}_j(X) &= \frac{b_j^2}{b_j^2 + X^2},\end{aligned}\tag{17}$$

and the parameters  $a_i$  for  $i \in (1, 2, 3, 4)$  and  $b_j$  for  $j \in (1, 2, 3, 4, 5)$  represent activation and inhibition thresholds. We assume that the initial states of the system are  $(X_1(0), X_2(0), X_3(0), X_4(0)) = (0.0, 0.0, 0.29, 0.625)$ , which correspond to the steady state of the system when  $X_2 = 0$ . The function  $u(t)$  represents a TNF stimulus which is given by a top hat function,

486  
487  
488  
489  
490

$$u(t) = \begin{cases} 1, & \text{if } t \in [0, 2], \\ 0, & \text{otherwise.} \end{cases}\tag{18}$$

When there are fewer output measurements than parameters, models tend to be underdetermined meaning that many combinations of parameters can lead to the same combination of output values. A consequence of this undeterminacy is that we cannot perform “full circle” inference: that is, using a known parameter distribution to generate an output distribution does not result in that parameter distribution being recapitulated through inference. We illustrate this idea by generating an output distribution by varying a single parameter value between runs of the forward model corresponding to the solution of eq. (16) and performing inference on all nine system parameters, whilst collecting only two output measurements. Specifically, we vary  $a_1 \sim \mathcal{N}(0.6, 0.05)$ , whilst holding the other parameters constant  $(a_2, a_3, a_4, b_1, b_2, b_3, b_4, b_5) = (0.2, 0.2, 0.5, 0.4, 0.4, 0.7, 0.3, 0.5, 0.4)$  and measure  $X_1(2.0)$  and  $X_2(1.0)$ . By running the forward model, we obtain an output distribution that is well approximated by the bivariate normal distribution,

491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505

$$\begin{pmatrix} X_1(2.0) \\ X_2(1.0) \end{pmatrix} \sim \mathcal{N} \left[ \begin{pmatrix} 0.26 \\ 0.07 \end{pmatrix}, \begin{pmatrix} 2.1 \times 10^{-4} & 5.9 \times 10^{-5} \\ 5.9 \times 10^{-5} & 1.8 \times 10^{-5} \end{pmatrix} \right].\tag{19}$$

We now apply CMC to the target output distribution given by eq. (19) to estimate a posterior distribution over all nine parameters of eq. (16). Apart than for  $a_1$ , the priors for each parameter were chosen to exclude the values that were used to generate the output distribution (see Table 2), to illustrate the non-equivalence between the recovered posterior distribution and the data generating process. In Figure 11A, we plot the actual parameter values (horizontal axis) used in the true data generating process versus the inferred values (vertical axis). This illustrates that apart from  $a_1$ , where the estimated parameter values correspond well with the range of values used to generate the data, due to the choice of priors there is a disjunction

506  
507  
508  
509  
510  
511  
512  
513  
514  
515

between the actual and estimated values. Despite these differences, due to the model being underdetermined, it is nonetheless possible to use CMC to sample from an output distribution that well approximates the target (Figure 11B). 516  
517  
518  
519




Figure 11: In (A) we plot the actual parameter values versus estimated quantiles for the output distribution for the TNF signalling pathway model with output distribution given and (B) the marginal output target (solid lines) given by eq. (19) and sampled output distribution (histograms). In (A), the vertical direction, the red points indicate the 50% posterior quantiles and the upper and lower whiskers indicate the 97.5% and 2.5% quantiles, respectively; in the horizontal direction, with the exception of  $a_1$  where 95% quantiles are given, the parameter values used to generate the data had no uncertainty. In the CMC, 10,000 independent samples were used in the “ContourVolumeEstimator” step and 5,000 MCMC samples across each of 4 Markov chains were used in the second step, with the first half of the chains discarded as “warm-up” [21]. 520  
521  
522  
523  
524  
525

Cell populations may be well described by subpopulations which each evolve along characteristic trajectories over time. We now apply CMC to investigate a bimodal output distribution for the TNF signalling pathway model similar to that investigated by [15]. In particular, we aim to find a distribution over parameter values which, when fed through the solution to the system of ODEs, results in the following output distribution, 526  
527  
528  
529  
530  
531  
532  
533

$$\begin{aligned} \mathbf{X}_2(1.0) &\sim \mathcal{N}(0.06, 0.01) \\ \mathbf{X}_2(2.0) &\sim \frac{1}{2} (\mathcal{N}(0.1, 0.01) + \mathcal{N}(0.14, 0.01)) \\ \mathbf{X}_2(4.0) &\sim \frac{1}{2} (\mathcal{N}(0.1, 0.01) + \mathcal{N}(0.20, 0.01)), \end{aligned} \quad (20)$$

where via a slight abuse of notation, the target distributions for  $\mathbf{X}_2(2.0)$  and  $\mathbf{X}_2(4.0)$  indicate mixtures of univariate normals, and the priors used are shown in Table 2. This target distribution, along with the unique trajectories obtained by applying the CMC algorithm for 5,000 MCMC steps, are shown in Figure 12. This Figure illustrates that given by bimodality of the output distribution, CMC estimates a corresponding subpopulation structure in the parameter distribution without *a priori* specification of the number of clusters. 526  
527  
528  
529  
530  
531  
532  
533




Figure 12: **The target output distribution (dashed plots with grey filling) and unique trajectories (black solid lines) obtained from the posterior parameter distribution.** In the CMC, 10,000 independent samples were used in the ‘ContourVolumeEstimator’ step and 5,000 MCMC samples across each of 4 Markov chains were used in the second step, with the first half of the chains discarded as ‘warm-up’ [21].

## 5 Discussion

Determining the cause of variability in cellular processes is crucial in many applications, ranging from bioengineering to drug development. In this paper, we introduce a Bayesian method for estimating cellular heterogeneity from ‘snapshot’ measurements of cellular properties, taken at discrete intervals throughout the experimental course. Our approach assumes what we term a ‘heterogeneous ordinary differential equation’ (HODE) framework, in which biochemical processes in individual cells are assumed to follow dynamics governed by a common ODE, although with idiosyncratic differences in parameter values. In this framework, estimating heterogeneity in cellular processes amounts to determining the probability distributions over parameter values of the governing ODE. Our method of estimation is a two-step Monte Carlo sampling process we term ‘Contour Monte Carlo’ (CMC) which does not require *a priori* specification of cell population substructure unlike other approaches and can handle high volumes of individual cellular measurements by fitting a kernel density estimator to these data. CMC also allows for arbitrary multivariate structure in the measurement space, meaning it can capture correlations that occur between the same cellular species at different time points or, for example, contemporaneous correlations between different cellular compartments. Being a Bayesian approach, CMC uses prior distributions over parameter values to ensure uniqueness of the posterior distribution, allowing pre-experimental knowledge to be used to improve estimation robustness. The flexible and robust framework that CMC provides means it can be used to do automatic inference for wide-ranging systems of practical interest.

As well as providing a framework for estimating cellular variation,

our approach also provides a natural way to test that it is working as  
560  
desired. By feeding the posterior parameter samples obtained by CMC into  
561  
the forward model, this results in a distribution over output values that  
562  
can be compared to the target. Indeed, we have found this comparison  
563  
indispensable in applying CMC in practice since discrepancies can occur  
564  
either as a result of poor estimates of the ‘contour volume distribution’  
565  
in the first stage of the algorithm or due to insufficient MCMC samples  
566  
in the second. Either of these issues can often be easily addressed and  
567  
although kernel density estimation in high dimensional spaces remains an  
568  
open research problem, we have found vine copula kernel density estimation  
569  
works well in the size of problems we investigate here [18].  
570

Failure to reproduce a given output distribution can also indicate that  
571  
the generating model (the priors and the forward model) are incongruent  
572  
with experimental results. This may either be due to misspecification of the  
573  
ODE system or the inadequacy of the assumed deterministic framework.  
574  
Our approach currently assumes that output stochasticity is dominated by  
575  
cellular variation in the parameter values of the underlying ODE, with mea-  
576  
surement noise making a minor contribution. Whether this is a reasonable  
577  
assumption depends on the system under investigation and, more impor-  
578  
tantly, on experimental details. We recognise that neglecting measurement  
579  
noise when it is an important determinant of the observed data is likely  
580  
to mean we overstate the degree of cellular variation. It may also mean  
581  
that some output distributions cannot be obtained through our assumed  
582  
model system. Future work allowing inclusion of a stochastic noise process  
583  
or, more generally, including stochastic cellular mechanisms is thus likely  
584  
to be worthwhile.  
585

Whilst we have labelled the approach we follow here as Bayesian, since it  
586  
involves explicit estimation of probability distributions and involves priors  
587  
over parameter values, we recognise that it is not in the form typically  
588  
utilised by exponents of this framework. This is because rather than aiming  
589  
to formulate a model that describes output observations, instead, our  
590  
approach aims to recapitulate output distributions. Others [28], (including  
591  
us [29]), have considered this problem before; perhaps most notably by  
592  
Albert Tarantola in his landmark work on inverse problem theory (see,  
593  
for example, [30]). In Tarantola’s framework, we consider a joint input  
594  
parameter & output space, where prior knowledge and experimental theory  
595  
combine elegantly to produce a posterior distribution whose marginal output  
596  
distribution matches the experimentally obtained one. This work has seen  
597  
considerable interest in areas such as geosciences [31,32], and we propose that  
598  
these methods may prove useful for the biosciences. In particular, we posit  
599  
that this framework may allow for a more generalised Bayesian inference  
600  
that can more easily encompass output data and output distributions as  
601  
their outcome measures.  
602

The natural world is rife with variation. Mathematical models represent  
603  
frameworks for understanding the causes of such variation. Typically, the  
604  
state of biological knowledge is such that one effect, a given pattern of  
605  
variation, has many possible causes, and observational or experimental  
606  
data are necessary to apportion weight to each of them, in a process  
607  
which amounts to solving inverse an problem. The approach we describe  
608  
here follows the Bayesian paradigm of inverse problem solving whereby  
609  
uncertainty in potential causes is reflected by probability distributions. Here,  
610  
we illustrate the utility of our method by applying it to estimate cellular  
611  
heterogeneity in biochemical processes however, it could equally well be used  
612

to understand the inversion of deterministic systems more generally. Whilst  
describing the inversion process of deterministic models using probability  
distributions may sound contradictory, it is worth acknowledging that many  
ODE systems are structurally unidentified meaning there is incompressible  
uncertainty over some regions of parameter space. Contour Monte Carlo  
provides an automatic framework for doing inference on these deterministic  
systems and the use of priors allows for robust and precise parameter  
estimation unattainable through the data alone. 613  
614  
615  
616  
617  
618  
619  
620

## 6 Author contributions

BL, DJG and SJT conceived the study. BL carried out the analysis. All  
authors helped to write and edit the manuscript. 621  
622  
623

## References

- [1] Matt Ridley. *The red queen: Sex and the evolution of human nature*. Penguin UK, 1994.
- [2] Dawn Fraser and Mads Kaern. A chance at survival: gene expression noise and phenotypic diversification strategies. *Molecular microbiology*, 71(6):1333–1340, 2009.
- [3] Frank Delvigne, Quentin Zune, Alvaro R Lara, Waleed Al-Soud, and Søren J Sørensen. Metabolic variability in bioprocessing: implications of microbial phenotypic heterogeneity. *Trends in Biotechnology*, 32(12):608–616, 2014.
- [4] RA Gatenby, K Smallbone, PK Maini, F Rose, J Averill, Raymond B Nagle, L Worrall, and RJ Gillies. Cellular adaptations to hypoxia and acidosis during somatic evolution of breast cancer. *British journal of cancer*, 97(5):646, 2007.
- [5] Philipp M Altrock, Lin L Liu, and Franziska Michor. The mathematics of cancer: integrating quantitative models. *Nature Reviews Cancer*, 15(12):730, 2015.
- [6] Steven J Altschuler and Lani F Wu. Cellular heterogeneity: do differences make a difference? *Cell*, 141(4):559–563, 2010.
- [7] Michael B Elowitz, Arnold J Levine, Eric D Siggia, and Peter S Swain. Stochastic gene expression in a single cell. *Science*, 297(5584):1183–1186, 2002.
- [8] Hannah H Chang, Martin Hemberg, Mauricio Barahona, Donald E Ingber, and Sui Huang. Transcriptome-wide noise controls lineage choice in mammalian progenitor cells. *Nature*, 453(7194):544, 2008.
- [9] Steffen Waldherr. Estimation methods for heterogeneous cell population models in systems biology. *Journal of The Royal Society Interface*, 15(147):20180530, 2018.
- [10] Radek Erban, Jonathan Chapman, and Philip Maini. A practical guide to stochastic simulations of reaction-diffusion processes. *arXiv preprint arXiv:0704.1908*, 2007.

- [11] Doraiswami Ramkrishna and Meenesh R Singh. Population balance modeling: current status and future prospects. *Annual review of chemical and biomolecular engineering*, 5:123–146, 2014.
- [12] Purushottam Dixit, Eugenia Lyashenko, Mario Niepel, and Dennis Vitkup. Maximum entropy framework for inference of cell population heterogeneity in signaling network dynamics. *bioRxiv*, page 137513, 2018.
- [13] William G Telford, Teresa Hawley, Fedor Subach, Vladislav Verkhusha, and Robert G Hawley. Flow cytometry of fluorescent proteins. *Methods*, 57(3):318–330, 2012.
- [14] Alex J Hughes, Dawn P Spelke, Zhuchen Xu, Chi-Chih Kang, David V Schaffer, and Amy E Herr. Single-cell western blotting. *Nature methods*, 11(7):749, 2014.
- [15] Jan Hasenauer, Steffen Waldherr, Małgorzata Doszczak, Nicole Radde, Peter Scheurich, and Frank Allgöwer. Identification of models of heterogeneous cell populations from population snapshot data. *BMC bioinformatics*, 12(1):125, 2011.
- [16] Jan Hasenauer, Christine Hasenauer, Tim Hucho, and Fabian J Theis. Ode constrained mixture modelling: a method for unraveling sub-population structures and dynamics. *PLoS computational biology*, 10(7):e1003686, 2014.
- [17] Carolin Loos, Katharina Moeller, Fabian Fröhlich, Tim Hucho, and Jan Hasenauer. A hierarchical, data-driven approach to modeling single-cell populations predicts latent causes of cell-to-cell variability. *Cell systems*, 6(5):593–603, 2018.
- [18] Thomas Nagler and Claudia Czado. Evading the curse of dimensionality in nonparametric density estimation with simplified vine copulas. *Journal of Multivariate Analysis*, 151:69–89, 2016.
- [19] Ross H Johnstone, Eugene TY Chang, Rémi Bardenet, Teun P De Boer, David J Gavaghan, Pras Pathmanathan, Richard H Clayton, and Gary R Mirams. Uncertainty and variability in models of the cardiac action potential: Can we build trustworthy models? *Journal of molecular and cellular cardiology*, 96:49–62, 2016.
- [20] Nicholas Metropolis, Arianna W Rosenbluth, Marshall N Rosenbluth, Augusta H Teller, and Edward Teller. Equation of state calculations by fast computing machines. *The Journal of Chemical Physics*, 21(6):1087–1092, 1953.
- [21] Ben Lambert. *A Student’s Guide to Bayesian Statistics*. Sage Publications Ltd., 2018.
- [22] Andrew Gelman and Donald B Rubin. Inference from iterative simulation using multiple sequences. *Statistical Science*, pages 457–472, 1992.
- [23] Inc. Wolfram Research. Mathematica 8.0. <https://www.wolfram.com>.

- [24] A.C. Daly, D. Gavaghan, J. Cooper, and S.J. Tavener. Inference-based assessment of parameter identifiability in nonlinear biological models. *Journal of the Royal Society Interface*, 15, 2018.
- [25] James D. Murray. *Mathematical biology: I. An Introduction (interdisciplinary applied mathematics)(Pt. 1)*. New York, Springer, 2007.
- [26] Ajay Jasra, David A Stephens, and Christopher C Holmes. On population-based simulation for static inference. *Statistics and Computing*, 17(3):263–279, 2007.
- [27] Madalena Chaves, Thomas Eissing, and Frank Allgower. Bistable biological systems: A characterization through local compact input-to-state stability. *IEEE Transactions on Automatic Control*, 53(Special Issue):87–100, 2008.
- [28] T. Butler, D. Estep, S.J. Tavener, C. Dawson, and J.J. Westerink. A measure-theoretic computational method for inverse sensitivity problems iii: Multiple quantities of interest. *SIAM/ASA Journal on Uncertainty Quantification*, 2(1):174–202, 2014.
- [29] Ben Lambert, David Gavaghan, and Simon Tavener. Inverse sensitivity analysis of mathematical models avoiding the curse of dimensionality. *BioRxiv*, page 432393, 2018.
- [30] Albert Tarantola. *Inverse problem theory and methods for model parameter estimation*, volume 89. siam, 2005.
- [31] Klaus Mosegaard and Albert Tarantola. Monte carlo sampling of solutions to inverse problems. *Journal of Geophysical Research: Solid Earth*, 100(B7):12431–12447, 1995.
- [32] Tomislava Vukicevic and Derek Posselt. Analysis of the impact of model nonlinearities in inverse problem solving. *Journal of the Atmospheric Sciences*, 65(9):2803–2823, 2008.