

Contents

Homotopy theory 11

Knot Theory 13

4-manifolds 15

Bibliography 17

Introduction

Homotopy theory

CW-complexes

Definition 0.0.1. A map $f: X \rightarrow Y$ is called a *weak homotopy equivalence* if it induces isomorphisms

$$\pi_n(X, x_0) \rightarrow \pi_n(Y, f(x_0))$$

for all $n \geq 0$ and all choices of basepoints x_0 in X .

Theorem 0.0.2 (Whitehead's Theorem). *A weak homotopy equivalence between CW-complexes is a homotopy equivalence.*

Proposition 0.0.3 (Geometric interpretation of n -connectedness). *If (X, A) is an n -connected CW-pair, then there exists a CW-pair $(Z, A) \sim_{\text{rel } A} (X, A)$ such that all cells of $Z \setminus A$ have dimension greater than n .*

Homology

Definition 0.0.4 (Acyclic). A space X is called *acyclic* if $\tilde{H}_i(X) = 0$ for all i , i.e. if its reduced homology vanishes.

Example 0.0.5. *Removing a point from a homology sphere yields an acyclic space. This example for the Poincare homology sphere is described in TODO*
Insert proof.

Knot Theory

Definitions

Definition 0.0.6. If K is an oriented knot, then

- the *reverse* \bar{K} is K with the opposite orientation
- the *obverse* rK is the reflection of K in a plane
- the *inverse* $r\bar{K}$ is the concordance inverse of K .

Proposition 0.0.7. For $K \subset S^3$ we have that $K \# r\bar{K}$ is slice, even ribbon.

Alexander polynomial

Definition 0.0.8. L oriented link with Seifert matrix A , then the first homology of the infinite cyclic covering of the link complement, $H_1(X_\infty; \mathbb{Z})$, has square presentation matrix $tA - A^T$.

The *Alexander polynomial* of L is given by

$$\Delta_L(t) \doteq \det(tA - A^T)$$

where \doteq means “up to a multiplication with a unit $\{\pm t^{\pm n}\}$ of the Laurent ring $\mathbb{Z}[t, t^{-1}]$ ”.

Remark 0.0.9. $\mathbb{Z}[t^{\pm 1}]$ is **not** a PID.

Invariants

Definition 0.0.10. The tunnel number $t(K)$ of a knot $K \subset S^3$ is the minimal number of arcs that must be added to the knot (forming a graph with three edges at a vertex) so that its complement in S^3 is a handlebody. The same definition is valid for links.

The boundary will be a minimal Heegaard splitting of the knot complement (The knot complement is a manifold with boundary, so what is the definition of a Heegaard splitting in that case?).

Remark 0.0.11. Every link has a tunnel number, this can be seen by adding a “vertical” tunnel at every crossing in a link diagram. This

shows that the tunnel number of a knot is always less than or equal to the crossing number, $t(K) \leq c(K)$.

Example 0.0.12. • *The unknot is the only knot with tunnel number 0.*
(Why?)

- *The trefoil knot has tunnel number 1.*
- *The figure eight knot has tunnel number 1.*

Open questions

Open question 1. *Is the crossing number of a satellite knot bigger than that of its companion?*

4-manifolds

TODO

Bibliography

Index

acyclic, 5

Alexander

 polynomial, 7

homotopy equivalence

 weak, 5

knot

 inverse, 7

 obverse, 7

 reverse, 7