



A nonlocal boundary condition for domain truncation in frequency-domain Helmholtz problems

Robert Kirby¹

Andreas Klöckner²

Ben Sepanski³

¹Baylor University

²University of Illinois at Urbana-Champaign

³University of Texas at Austin

25 March 2021



Helmholtz scattering

A new boundary condition

Quo vadis, Firedrake? (Future work)




Thanks to...

- ▶ NSF 1525697, 1909176
- ▶ The U.S. Department of Energy, Office of Science, Office of Advanced Scientific Computing Research, Department of Energy Computational Science Graduate Fellowship under Award Number DE-SC0021110
- ▶ Luke Olson (UIUC)



Model problem



Real problem:


$$-\Delta u + \kappa^2 u = 0, \quad \mathbb{R}^d \setminus \Omega^c$$

$$-\frac{\partial u}{\partial n} = f, \quad \Gamma$$

$$\lim_{r \rightarrow \infty} r^{\frac{d-1}{2}} \left(\frac{\partial u}{\partial r} - i\kappa u \right) = 0,$$



Model problem



Computational problem:


$$-\Delta u + \kappa^2 u = 0, \quad \Omega'$$

$$-\frac{\partial u}{\partial n} = f, \quad \Gamma$$

$$????? \quad \Sigma$$



Perfectly Matched Layers (PML)



Computational problem:


$$-\nabla \cdot \beta(x) \nabla u + \kappa^2 u = 0, \quad \Omega'$$

$$-\frac{\partial u}{\partial n} = f, \quad \Gamma$$

$$u = 0, \quad \Sigma$$



Perfectly Matched Layers (PML)



Computational problem:

$$-\nabla \cdot \beta(x) \nabla u + \kappa^2 u = 0, \quad \Omega'$$


$$-\frac{\partial u}{\partial n} = f, \quad \Gamma$$

$$u = 0, \quad \Sigma$$

► $\beta = I$ in Ω'



Perfectly Matched Layers (PML)



Computational problem:

$$-\nabla \cdot \beta(x) \nabla u + \kappa^2 u = 0, \quad \Omega'$$


$$-\frac{\partial u}{\partial n} = f, \quad \Gamma$$

$$u = 0, \quad \Sigma$$

- ▶ $\beta = I$ in Ω'
- ▶ β is complex-valued, eats waves in Ω_S



Perfectly Matched Layers (PML)



Computational problem:

$$-\nabla \cdot \beta(x) \nabla u + \kappa^2 u = 0, \quad \Omega'$$


$$-\frac{\partial u}{\partial n} = f, \quad \Gamma$$

$$u = 0, \quad \Sigma$$

- ▶ $\beta = I$ in Ω'
- ▶ β is complex-valued, eats waves in Ω_S
- ▶ Solution is right in Ω'



Perfectly Matched Layers (PML)



Computational problem:

$$-\nabla \cdot \beta(x) \nabla u + \kappa^2 u = 0, \quad \Omega'$$

$$-\frac{\partial u}{\partial n} = f, \quad \Gamma$$

$$u = 0, \quad \Sigma$$

- ▶ $\beta = I$ in Ω'
- ▶ β is complex-valued, eats waves in Ω_S
- ▶ Solution is right in Ω'
- ▶ **Solvers are a pain!**



Integral form of the solution

With \mathcal{K} the Green's function, the *true* solution satisfies:

$$u(x) = \int_{\Gamma} \left(\frac{\partial}{\partial n} \mathcal{K}(x - y) \right) u(y) - \left(\frac{\partial}{\partial n} u(y) \right) \mathcal{K}(x - y) dy,$$

where



Integral form of the solution

With \mathcal{K} the Green's function, the *true* solution satisfies:

$$u(x) = \int_{\Gamma} \left(\frac{\partial}{\partial n} \mathcal{K}(x - y) \right) u(y) - \left(\frac{\partial}{\partial n} u(y) \right) \mathcal{K}(x - y) dy,$$

where

$$D(u)(x) = \text{PV} \int_{\Gamma} \left(\frac{\partial}{\partial n} \mathcal{K}(x - y) \right) u(y) dy,$$



Integral form of the solution

With \mathcal{K} the Green's function, the *true* solution satisfies:

$$u(x) = \int_{\Gamma} \left(\frac{\partial}{\partial n} \mathcal{K}(x - y) \right) u(y) - \left(\frac{\partial}{\partial n} u(y) \right) \mathcal{K}(x - y) dy,$$


where

$$D(u)(x) = \text{PV} \int_{\Gamma} \left(\frac{\partial}{\partial n} \mathcal{K}(x - y) \right) u(y) dy,$$

$$S(u)(x) = \int \mathcal{K}(x - y) u(y) dy$$



Model problem



Real problem

$$-\Delta u + \kappa^2 u = 0, \quad \Omega'$$

$$-\frac{\partial u}{\partial n} = f, \quad \Gamma$$

$$(i\kappa - \frac{\partial}{\partial n})(u - D(u) + S(f)) = 0, \quad \Sigma$$



Variational form

$$a(u, v) = F(v)$$

for all $v \in H^1(\Omega')$,



Variational form

$$a(u, v) = F(v)$$

for all $v \in H^1(\Omega')$, where

$$a(u, v) = (\nabla u, \nabla v) - \kappa^2 (u, v) - i\kappa \langle u, v \rangle_{\Sigma} + \langle (i\kappa - \frac{\partial}{\partial n}) D(u), v \rangle_{\Sigma}$$



Variational form

$$a(u, v) = F(v)$$

for all $v \in H^1(\Omega')$, where

$$a(u, v) = (\nabla u, \nabla v) - \kappa^2 (u, v) - i\kappa \langle u, v \rangle_\Sigma + \langle (i\kappa - \frac{\partial}{\partial n}) D(u), v \rangle_\Sigma$$

and

$$F(v) = \langle f, v \rangle_\Gamma + \langle (i\kappa - \frac{\partial}{\partial n}) S(f), v \rangle_\Sigma$$



Note



Note

- ▶ Not DtN (Keller/Givoli): No Fourier series



Note

- ▶ Not DtN (Keller/Givoli): No Fourier series
- ▶ Not Poincare-Steklov operator: No singular integrals.



Note

- ▶ Not DtN (Keller/Givoli): No Fourier series
- ▶ Not Poincare-Steklov operator: No singular integrals.

BUT still requires layer potentials on boundary.



Theory



Theory

- ▶ a is a bounded bilinear form on $H^1 \times H^1$



Theory

- ▶ a is a bounded bilinear form on $H^1 \times H^1$
- ▶ F is a bounded linear functional on H^1



Theory

- ▶ a is a bounded bilinear form on $H^1 \times H^1$
- ▶ F is a bounded linear functional on H^1
- ▶ Gårding inequality. There exist M and an $\alpha > 0$ such that

$$\operatorname{Re}(a(u, u)) + M \|u\|^2 \geq \alpha \|u\|_{H^1(\Omega)}^2.$$



Theory

- ▶ a is a bounded bilinear form on $H^1 \times H^1$
- ▶ F is a bounded linear functional on H^1
- ▶ Gårding inequality. There exist M and an $\alpha > 0$ such that

$$\operatorname{Re}(a(u, u)) + M \|u\|^2 \geq \alpha \|u\|_{H^1(\Omega)}^2.$$

- ▶ For $h \leq h_0$, we have optimal-order H^1 and L^2 error estimates.



Linear system $Ax = b$

Separate out FEM/transmission from layer potential terms:

$$A = A^L + A^{NL}$$



Linear system $Ax = b$

Separate out FEM/transmission from layer potential terms:

$$A = A^L + A^{NL}$$

- ▶ A^L is built with Firedrake



Linear system $Ax = b$

Separate out FEM/transmission from layer potential terms:

$$A = A^L + A^{NL}$$

- ▶ A^L is built with Firedrake
- ▶ A^{NL} is evaluated matrix-free



Linear system $Ax = b$

Separate out FEM/transmission from layer potential terms:

$$A = A^L + A^{NL}$$

- ▶ A^L is built with Firedrake
- ▶ A^{NL} is evaluated matrix-free
 - Firedrake – pyntential marshalling



Linear system $Ax = b$

Separate out FEM/transmission from layer potential terms:

$$A = A^L + A^{NL}$$

- ▶ A^L is built with Firedrake
- ▶ A^{NL} is evaluated matrix-free
 - Firedrake – pyntial marshalling
 - Pyntial constructs layer potential via FMM, evaluates at edge quad points



Linear system $Ax = b$


Separate out FEM/transmission from layer potential terms:

$$A = A^L + A^{NL}$$

- ▶ A^L is built with Firedrake
- ▶ A^{NL} is evaluated matrix-free
 - Firedrake – pytential marshalling
 - Pytential constructs layer potential via FMM, evaluates at edge quad points
 - Firedrake integrates against test function







Marshalling issue





Numerical results





Preconditioning

Flexible GMRES requires right-preconditioning:

$$A(P^{-1}y) = b; \quad y = Px$$



Preconditioning

Flexible GMRES requires right-preconditioning:

$$A(P^{-1}y) = b; \quad y = Px$$

Our choice of preconditioner...



Preconditioning

Flexible GMRES requires right-preconditioning:


$$A(P^{-1}y) = b; \quad y = Px$$

Our choice of preconditioner...

$$P = A^L$$




Using LU on A^L






Scalable – gamg?






Scalable – gamg?




And explodes for higher κ .



W-Cycles: 1



W-Cycles: 3





Had NUFL of vaporware?

Can we do this?



More generally



More generally

- ▶ Extend UFL with nonlocal operators: Layer/volume potentials, fractional derivatives, etc



More generally

- ▶ Extend UFL with nonlocal operators: Layer/volume potentials, fractional derivatives, etc
- ▶ Compiler should segregate local and nonlocal terms:



More generally

- ▶ Extend UFL with nonlocal operators: Layer/volume potentials, fractional derivatives, etc
- ▶ Compiler should segregate local and nonlocal terms:
 - Matrix free, hybrid, HSS, etc



More generally

- ▶ Extend UFL with nonlocal operators: Layer/volume potentials, fractional derivatives, etc
- ▶ Compiler should segregate local and nonlocal terms:
 - Matrix free, hybrid, HSS, etc
 - TSFC + pytentinal/sumpy/loopy



More generally

- ▶ Extend UFL with nonlocal operators: Layer/volume potentials, fractional derivatives, etc
- ▶ Compiler should segregate local and nonlocal terms:
 - Matrix free, hybrid, HSS, etc
 - TSFC + pytentinal/sumpy/loopy
- ▶ Extend solver infrastructure