



Veech Fibrations

Sam Freedman (Brown)



Joint with Trent Lucas (Brown)
Gatsby Spring 2024





The Nielsen realization problem for K3 surfaces


Benson Farb and Eduard Looijenga *





The Nielsen realization problem for K3 surfaces

Benson Farb and Eduard Looijenga *



PERIODIC POINTS ON VEECH SURFACES AND THE MORDELL-WEIL GROUP OVER A TEICHMÜLLER CURVE

MARTIN MÖLLER






The Nielsen realization problem for K3 surfaces

Benson Farb and Eduard Looijenga *



PERIODIC POINTS ON VEECH SURFACES AND THE MORDELL-WEIL GROUP OVER A TEICHMÜLLER CURVE

MARTIN MÖLLER



Me

VEECH FIBRATIONS

SAM FREEDMAN AND TRENT LUCAS





General Veech Fibrations

General Veech Fibrations

(Generically) minimal complex surfaces of general type + smooth 4-manifolds

General Veech Fibrations


(Generically) minimal complex surfaces of general type + smooth 4-manifolds




C.f. Möller *Variations of Hodge structures of a Teichmüller curve*, 2005

General Veech Fibrations

(Generically) minimal complex surfaces of general type + smooth 4-manifolds



Caution: base is cover of Teichmüller curve



C.f. Möller *Variations of Hodge structures of a Teichmüller curve*, 2005

From general fibrations to congruence fibrations

From general fibrations to congruence fibrations

Problem:

Formulas for Euler characteristic,
signature, etc. require **explicit**
computation of cover $B \rightarrow C$.

$$e(\tilde{X}) = \boxed{\chi(\bar{B})} \chi(X) + \boxed{T}$$
$$\sigma(\tilde{X}) = -2\kappa_\mu \boxed{\chi(B)} - \frac{2}{3} \boxed{T}$$
$$T := \sum_{c \in \Delta} \sum_{i=1}^{\ell_c} k_i^c$$

From general fibrations to congruence fibrations

Problem:

Formulas for Euler characteristic, signature, etc. require **explicit** computation of cover $B \rightarrow C$.

$$e(\tilde{X}) = \boxed{\chi(\bar{B})} \chi(X) + \boxed{T}$$
$$\sigma(\tilde{X}) = -2\kappa_\mu \boxed{\chi(B)} - \frac{2}{3} \boxed{T}$$

$$T := \sum_{c \in \Delta} \sum_{i=1}^{\ell_c} k_i^c$$

Solution:

Restrict to *congruence Veech fibrations* built from congruence subgroups of the MCG.

$$\begin{array}{ccc} B = \mathbb{H}^2 / \Gamma(n) & \longrightarrow & \mathcal{M}_1^{[n]} \\ \downarrow & & \downarrow \\ C = \mathbb{H}^2 / \mathrm{SL}(2, \mathbb{Z}) & \longrightarrow & \mathcal{M}_1 \end{array}$$

(Generalizing elliptic modular surfaces)

From general fibrations to congruence fibrations

Problem:

Formulas for Euler characteristic, signature, etc. require **explicit** computation of cover $B \rightarrow C$.

$$e(\tilde{X}) = \boxed{\chi(\bar{B})} \chi(X) + \boxed{T}$$
$$\sigma(\tilde{X}) = -2\kappa_\mu \boxed{\chi(B)} - \frac{2}{3} \boxed{T}$$

$$T := \sum_{c \in \Delta} \sum_{i=1}^{\ell_c} k_i^c$$

Solution:

Restrict to *congruence Veech fibrations* built from congruence subgroups of the MCG.

$$\begin{array}{ccc} B = \mathbb{H}^2/\Gamma(n) & \longrightarrow & \mathcal{M}_1^{[n]} \\ \downarrow & & \downarrow \\ C = \mathbb{H}^2/\text{SL}(2, \mathbb{Z}) & \longrightarrow & \mathcal{M}_1 \end{array}$$


(Generalizing elliptic modular surfaces)

Problem:


Requires describing the representation $\rho_m : \text{Aff}^+(X, \omega) \rightarrow \text{Sp}(H_1(X; \mathbb{Z}/m\mathbb{Z}))$. Difficult for any **specific choices** of surface and level m!

Case of algebraically primitive examples


Case of algebraically primitive examples



Genus 2 “L-tables”
(non-square discriminant)




Regular n-gons
($n = p, 2p, 2^k$)




Sporadic E_7 and E_8 examples


Case of algebraically primitive examples



Genus 2 “L-tables”
(non-square discriminant)



Regular n-gons
($n = p, 2p, 2^k$)




Sporadic E_7 and E_8 examples

Theorem (F. – Lucas):

For surfaces in the above three families, and for primes p in a computable infinite set depending on the surface, we have:

1. $\pi_1(\widetilde{\mathbb{X}}_p) \rightarrow \pi_1(\overline{B}_p)$ is an isomorphism,
2. $\widetilde{\mathbb{X}}_p$ is a minimal complex surface of general type, and
3. explicit formulas for $e(\widetilde{\mathbb{X}}_p)$ and $\sigma(\widetilde{\mathbb{X}}_p)$.


Case of algebraically primitive examples



$$e(\tilde{\mathbb{X}}_{q,p}) = dg + d(q-3) \left(\frac{1}{2} - \frac{1}{q} - \frac{1}{p} \right)$$

$$\sigma(\tilde{\mathbb{X}}_{q,p}) = -\frac{d(q^2-1)}{4q}$$

Genus 2 “L-tables”
(non-square discriminant)



Theorem (F. – Lucas):

For surfaces in the above three families, and for primes p in a computable infinite set depending on the surface, we have:

1. $\pi_1(\tilde{\mathbb{X}}_p) \rightarrow \pi_1(\overline{B}_p)$ is an isomorphism,
2. $\tilde{\mathbb{X}}_p$ is a minimal complex surface of general type, and
3. explicit formulas for $e(\tilde{\mathbb{X}}_p)$ and $\sigma(\tilde{\mathbb{X}}_p)$.

Conclusions

Conclusions

- New class of smooth 4-manifolds and complex surfaces that we can study explicitly using flat geometry of Veech surfaces.

Conclusions

- New class of smooth 4-manifolds and complex surfaces that we can study explicitly using flat geometry of Veech surfaces.
- The structure of a Veech fibration lets us “see” interesting mapping classes (e.g. translation automorphisms on the fibers)

Conclusions

- New class of smooth 4-manifolds and complex surfaces that we can study explicitly using flat geometry of Veech surfaces.
- The structure of a Veech fibration lets us “see” interesting mapping classes (e.g. translation automorphisms on the fibers)
- Future directions:
 - Can complex surfaces fiber as a Veech fibration in more than one way?
 - Which complex surfaces have the structure of a Veech fibration?
 - Can the Veech fibration structure be used to distinguish between complex surface?

Conclusions

- New class of smooth 4-manifolds and complex surfaces that we can study explicitly using flat geometry of Veech surfaces.
- The structure of a Veech fibration lets us “see” interesting mapping classes (e.g. translation automorphisms on the fibers)
- Future directions:
 - Can complex surfaces fiber as a Veech fibration in more than one way?
 - Which complex surfaces have the structure of a Veech fibration?
 - Can the Veech fibration structure be used to distinguish between complex surface?

arXiv:2310.02325



Nielsen-Thurston types and Teichmüller translation lengths of mapping classes along random walks

Dongryul M. Kim

Dept. of Mathematics, Yale University

GATSBY 2024 Spring

Random walk on MCG

S : closed connected orientable surface of genus ≥ 2 .

μ : prob on $\text{MCG}(S)$

Random walk on MCG

S : closed connected orientable surface of genus ≥ 2 .

μ : prob on $\text{MCG}(S)$

For $n \geq 1$,

$$\omega_n = g_1 \cdots g_n, \quad i.i.d. \quad g_i \sim \mu$$

Random walk on MCG

S : closed connected orientable surface of genus ≥ 2 .

μ : prob on $\text{MCG}(S)$

For $n \geq 1$,

$$\omega_n = g_1 \cdots g_n, \quad i.i.d. \quad g_i \sim \mu$$

This gives a prob \mathbb{P} on $\text{MCG}(S)^{\mathbb{N}}$,

“Random walk on MCG”

Always assume that $\langle \text{supp } \mu \rangle < \text{MCG}(S)$ is non-elementary.

Type and translation length along RW

Theorem (Maher-Tiozzo 2018)

Suppose that $\# \text{ supp } \mu < \infty$.

Type and translation length along RW

Theorem (Maher-Tiozzo 2018)

Suppose that $\# \text{ supp } \mu < \infty$. Then \mathbb{P} -a.e. $(\omega_n) \in \text{MCG}(S)^{\mathbb{N}}$, there exists $N > 0$ such that

ω_n is pseudo-Anosov for all $n \geq N$.

Type and translation length along RW

Theorem (Maher-Tiozzo 2018)

Suppose that $\# \text{ supp } \mu < \infty$. Then \mathbb{P} -a.e. $(\omega_n) \in \text{MCG}(S)^{\mathbb{N}}$, there exists $N > 0$ such that

ω_n is pseudo-Anosov for all $n \geq N$.

Theorem (Dahmani-Horbez 2018)

Suppose that $\# \text{ supp } \mu < \infty$.

Type and translation length along RW

Theorem (Maher-Tiozzo 2018)

Suppose that $\# \text{ supp } \mu < \infty$. Then \mathbb{P} -a.e. $(\omega_n) \in \text{MCG}(S)^{\mathbb{N}}$, there exists $N > 0$ such that

ω_n is pseudo-Anosov for all $n \geq N$.

Theorem (Dahmani-Horbez 2018)

Suppose that $\# \text{ supp } \mu < \infty$. Then \mathbb{P} -a.e. $(\omega_n) \in \text{MCG}(S)^{\mathbb{N}}$,

$$\lim_{n \rightarrow \infty} \frac{\log \lambda(\omega_n)}{n} = L_\mu$$

where λ is the stretch factor of p-A and L_μ is a const determined by μ (called 'drift').

Note: for p-A $g \in \text{MCG}(S)$, $\log \lambda(g) = \text{Teich. tr. length} = \text{top. entropy}$.

Type and translation length along RW

Theorem (Baik-Choi-K. 2021 (published in 2024))

Without any moment condition on μ ,

Type and translation length along RW

Theorem (Baik-Choi-K. 2021 (published in 2024))

Without any moment condition on μ , for \mathbb{P} -a.e. $(\omega_n) \in \text{MCG}(S)^{\mathbb{N}}$, there exists $N > 0$ such that

ω_n is pseudo-Anosov for all $n \geq N$.

Type and translation length along RW

Theorem (Baik-Choi-K. 2021 (published in 2024))

Without any moment condition on μ , for \mathbb{P} -a.e. $(\omega_n) \in \text{MCG}(S)^{\mathbb{N}}$, there exists $N > 0$ such that

ω_n is pseudo-Anosov for all $n \geq N$.

Moreover, under finite first moment, i.e. $\sum_g d_{\text{Teich}}(o, go)\mu(g) < \infty$,

Type and translation length along RW

Theorem (Baik-Choi-K. 2021 (published in 2024))

Without any moment condition on μ , for \mathbb{P} -a.e. $(\omega_n) \in \text{MCG}(S)^{\mathbb{N}}$, there exists $N > 0$ such that

ω_n is pseudo-Anosov for all $n \geq N$.

Moreover, under finite first moment, i.e. $\sum_g d_{\text{Teich}}(o, go)\mu(g) < \infty$, we have for \mathbb{P} -a.e. $(\omega_n) \in \text{MCG}(S)^{\mathbb{N}}$,

$$\lim_{n \rightarrow \infty} \frac{\log \lambda(\omega_n)}{n} = L_\mu$$

where λ is the stretch factor of p -A and L_μ is a const determined by μ (called ‘drift’).

This establishes the SLLN for translation lengths.



Finite quotients of hyperbolic 3-manifold groups

Tam Cheetham-West

Yale University

GATSBY Spring 2024

Introduction...

- An infinite group G is residually finite if every non-trivial element of G survives to some finite quotient of G .

Introduction...

- An infinite group G is residually finite if every non-trivial element of G survives to some finite quotient of G .
- For a group G , let

$$\mathcal{C}(G) = \{Q : G \twoheadrightarrow Q, |Q| < \infty\}$$

be the set of finite quotients of G . One can ask to what extent $\mathcal{C}(G)$ determines G . This is a well-studied question for several classes of groups.

Profinite rigidity

- A *profinitely rigid* group G is one for which $\mathcal{C}(G) = \mathcal{C}(H)$
 $\implies G \cong H$ (G, H f.g., r.f.)

Profinite rigidity

- A *profinitely rigid* group G is one for which $\mathcal{C}(G) = \mathcal{C}(H)$
 $\implies G \cong H$ (G, H f.g., r.f.)
- A *profinitely rigid* group is a residually finite group that is completely determined (among residually finite groups) up to isomorphism by its collection of finite quotients

Profinite rigidity

- A *profinitley rigid* group G is one for which $\mathcal{C}(G) = \mathcal{C}(H)$
 $\implies G \cong H$ (G, H f.g., r.f.)
- A *profinitley rigid* group is a residually finite group that is completely determined (among residually finite groups) up to isomorphism by its collection of finite quotients
- We do not know whether free groups or surface groups are profinitely rigid

Profinite rigidity for Kleinian groups

Theorem (Bridson-McReynolds-Reid-Spitler)

The following Kleinian groups are profinitely rigid

- $\pi_1(\text{The Weeks manifold})$
- $PSL(2, \mathbb{Z}[\omega])$ where $\omega^2 + \omega + 1 = 0$

Theorem (C)

The following fibered 3-manifolds have profinitely rigid π_1

- 0-surgery on 6_2
- 0-surgery on 6_3

The knots 6_2 and 6_3




Figure 1: 6_2






Figure 2: 6_3

And...

Thanks for listening

Are these curves different?

Sayantika Mondal
Graduate Center, CUNY



Filling Curves

Cut along curve —→ Disks and annuli

Alternate Characterization:

A closed curve on a surface is said to be filling if it intersects every essential simple, non-peripheral closed curve on the surface.


Filling Curves

Cut along curve —> Disks and annuli

Alternate Characterization:

A closed curve on a surface is said to be filling if it intersects every essential simple, non-peripheral closed curve on the surface.

Example:




Topological Types:

Two curves are said to be of the same topological type if there is a mapping class group element (think Dehn twists!) taking one to the other.





Topological Types:

Two curves are said to be of the same topological type if there is a mapping class group element (think Dehn twists!) taking one to the other.







Topological Types:

Two curves are said to be of the same topological type if there is a mapping class group element (think Dehn twists!) taking one to the other.



Topological Types:


Two curves are said to be of the same topological type if there is a mapping class group element (think Dehn twists!) taking one to the other.




Different

Topological Types:


Two curves are said to be of the same topological type if there is a mapping class group element (think Dehn twists!) taking one to the other.



Different



Same



A first example...

Minimal filling curve

Filling curve with the minimum self intersection number on a given surface.

A first example...


Minimal filling curve

Filling curve with the minimum self intersection number on a given surface.

- Exists!
- This is sort of the simplest example on a surface.

Building filling curves (an infinite family)


Start with a (minimal) filling curve γ_0



Building filling curves (an infinite family)

Start with a (minimal) filling curve γ_0

Cut at an intersection point with a separating curve.




Building filling curves (an infinite family)

Start with a (minimal) filling curve γ_0

Cut at an intersection point with a separating curve.

Given a pair of positive integers (m, n) , let γ be the curve $\eta^m * \gamma_0^n$ (based at the intersection point)




Building filling curves (an infinite family)

Start with a (minimal) filling curve γ_0

Cut at an intersection point with a separating curve.

Given a pair of positive integers (m, n) , let γ be the curve $\eta^m * \gamma_0^n$ (based at the intersection point)

A (2,2) curve.



Are these different?

For every n, m we get a curve.
Are the different types?
How can we tell?

Are these different?

For every n, m we get a curve.
Are the different types?
How can we tell?



A first guess...

Self -Intersection number

Are these different?

For every n, m we get a curve.
Are the different types?
How can we tell?



A first guess...

Self -Intersection number

Yes! But there are curves with
same self-intersection
number within the family

Are these different?

For every n, m we get a curve.
Are the different types?
How can we tell?




A first guess...

Self -Intersection number

Yes! But there are curves with
same self-intersection
number within the family

$$i(\gamma, \gamma) = i(\gamma_0, \gamma_0)n^2 + (i(\gamma_0, \eta)n - 1)m$$



Solve for k

A better invariant...

For (ϕ, X) in $\text{Teich}(\Sigma)$. Let $\ell_\gamma(X)$ denote the ‘ X -length’ of the geodesic in the free homotopy class of $\phi(\gamma)$.

We define the length infimum of γ as follows:

$$m_\gamma = \inf \{ \ell_\gamma(X) : (\phi, X) \text{ in } \text{Teich}(\Sigma) \}$$

A better invariant...

For (ϕ, X) in $\text{Teich}(\Sigma)$. Let $\ell_\gamma(X)$ denote the ‘ X -length’ of the geodesic in the free homotopy class of $\phi(\gamma)$.

We define the length infimum of γ as follows:

$$m_\gamma = \inf \{ \ell_\gamma(X) : (\phi, X) \text{ in } \text{Teich}(\Sigma) \}$$

OR,

For every hyperbolic metric measure length of curve and make a list and take the smallest of the list!

A better invariant...

For (ϕ, X) in $\text{Teich}(\Sigma)$. Let $\ell_\gamma(X)$ denote the ‘ X -length’ of the geodesic in the free homotopy class of $\phi(\gamma)$.

We define the length infimum of γ as follows:




$$m_\gamma = \inf \{ \ell_\gamma(X) : (\phi, X) \text{ in } \text{Teich}(\Sigma) \}$$

OR,

For every hyperbolic metric measure length of curve and make a list and take the smallest of the list!




! Realized uniquely and
is a MCG invariant

Back to slide 1





Back to slide 1

Both have same intersection number (k) ;(




Back to slide 1

Both have same intersection number (k) ;(





$$k \lesssim m_{\beta_k}$$




$$m_{\alpha_k} \lesssim \log k$$

Back to slide 1

Both have same intersection number (k) ;(



$$k \lesssim m_{\beta_k}$$



$$m_{\alpha_k} \lesssim \log k$$

So these are of different topological types!

Theorem (ish):

For any finite type surface and any choice of natural number k (some minor restrictions),

Can build curves with same intersection numbers (k) but of different infimum length and as a result topological types.

....and some more info on the inf metrics.

Joint work with A. Basmajian



Thanks!

smondal@gradcenter.cuny.edu

CREDITS: This presentation template was created by [Slidesgo](#), including icons by [Flaticon](#) and infographics & images by [Freepik](#)

Please keep this slide for attribution

Octonions and exceptional Lie groups G_2 and F_4

Christa Ishimwe

Wesleyan University

GATSBY, April 20, 2024

Introduction

Classical Lie groups can be realized as isometry groups of projective spaces over skew fields \mathbb{R} , \mathbb{C} , and \mathbb{H} . When they were first discovered, the exceptional Lie groups could not be realized as isometry groups of any known mathematical structure. Now we know that $\text{Aut}(\mathbb{O}) \cong G_2$ and $\text{Aut}(\mathbb{O}P^2) \cong F_4$.

Constructing \mathbb{O} and $\mathbb{O}P^2$

- We construct \mathbb{O} as an extension of \mathbb{C} through the Cayley-Dickson method.
- Given a division algebra A over the reals, we usually define the n -dimensional projective space as the quotient A^{n+1}/\sim where $x \sim y$ if $\exists \lambda \in \mathbb{R} \setminus \{0\}$ such that $x = \lambda y$. This gives us **homogeneous coordinates**.
- The above method does not work for the octonions because they are not associative. Instead of using **homogeneous coordinates**, we use **Veronese Coordinates**. We can also obtain a space isomorphic to $\mathbb{O}P^2$ from the exceptional Jordan algebra $\mathfrak{h}_3(\mathbb{O})$.

Constructing \mathbb{O} and $\mathbb{O}P^2$

- We construct \mathbb{O} as an extension of \mathbb{C} through the Cayley-Dickson method.
- Given a division algebra A over the reals , we usually define the n -dimensional projective space as the quotient A^{n+1}/ \sim where $x \sim y$ if $\exists \lambda \in \mathbb{R} \setminus \{0\}$ such that $x = \lambda y$. This gives us **homogeneous coordinates**.
- The above method does not work for the octonions because they are not associative. Instead of using **homogeneous coordinates**, we use **Veronese Coordinates**. We can also obtain a space isomorphic to $\mathbb{O}P^2$ from the exceptional Jordan algebra $\mathfrak{h}_3(\mathbb{O})$.

Constructing \mathbb{O} and $\mathbb{O}P^2$

- We construct \mathbb{O} as an extension of \mathbb{C} through the Cayley-Dickson method.
- Given a division algebra A over the reals , we usually define the n -dimensional projective space as the quotient A^{n+1}/ \sim where $x \sim y$ if $\exists \lambda \in \mathbb{R} \setminus \{0\}$ such that $x = \lambda y$. This gives us **homogeneous coordinates**.
- The above method does not work for the octonions because they are not associative. Instead of using **homogeneous coordinates**, we use **Veronese Coordinates**. We can also obtain a space isomorphic to $\mathbb{O}P^2$ from the exceptional Jordan algebra $\mathfrak{h}_3(\mathbb{O})$.

Key steps for proving $\text{Aut}(\mathbb{O}) \cong G_2$

- Prove that if $f \in \text{Aut}(\mathbb{O})$ then f is \mathbb{R} -linear and preserves length.
- Prove that $\text{Aut}(\mathbb{O})$ is a compact lie group by showing that it is a closed subgroup of the compact Lie group $O(7)$.
- Prove that $\text{Aut}(\mathbb{O})$ is simple.
- Prove that $\text{Aut}(\mathbb{O})$ is connected and simply connected.
- Prove that the dimension of $\text{Aut}(\mathbb{O})$ is 14.
- By the classification of simple compact connected and simply connected Lie groups, conclude that $\text{Aut}(\mathbb{O}) \cong G_2$.

Key steps for proving $\text{Aut}(\mathbb{O}) \cong G_2$

- Prove that if $f \in \text{Aut}(\mathbb{O})$ then f is \mathbb{R} -linear and preserves length.
- Prove that $\text{Aut}(\mathbb{O})$ is a compact lie group by showing that it is a closed subgroup of the compact Lie group $O(7)$.
- Prove that $\text{Aut}(\mathbb{O})$ is simple.
- Prove that $\text{Aut}(\mathbb{O})$ is connected and simply connected.
- Prove that the dimension of $\text{Aut}(\mathbb{O})$ is 14.
- By the classification of simple compact connected and simply connected Lie groups, conclude that $\text{Aut}(\mathbb{O}) \cong G_2$.

Key steps for proving $\text{Aut}(\mathbb{O}) \cong G_2$

- Prove that if $f \in \text{Aut}(\mathbb{O})$ then f is \mathbb{R} -linear and preserves length.
- Prove that $\text{Aut}(\mathbb{O})$ is a compact lie group by showing that it is a closed subgroup of the compact Lie group $O(7)$.
- Prove that $\text{Aut}(\mathbb{O})$ is simple.
 - Prove that $\text{Aut}(\mathbb{O})$ is connected and simply connected.
 - Prove that the dimension of $\text{Aut}(\mathbb{O})$ is 14.
 - By the classification of simple compact connected and simply connected Lie groups, conclude that $\text{Aut}(\mathbb{O}) \cong G_2$.

Key steps for proving $\text{Aut}(\mathbb{O}) \cong G_2$

- Prove that if $f \in \text{Aut}(\mathbb{O})$ then f is \mathbb{R} -linear and preserves length.
- Prove that $\text{Aut}(\mathbb{O})$ is a compact lie group by showing that it is a closed subgroup of the compact Lie group $O(7)$.
- Prove that $\text{Aut}(\mathbb{O})$ is simple.
- Prove that $\text{Aut}(\mathbb{O})$ is connected and simply connected.
- Prove that the dimension of $\text{Aut}(\mathbb{O})$ is 14.
- By the classification of simple compact connected and simply connected Lie groups, conclude that $\text{Aut}(\mathbb{O}) \cong G_2$.

Key steps for proving $\text{Aut}(\mathbb{O}) \cong G_2$

- Prove that if $f \in \text{Aut}(\mathbb{O})$ then f is \mathbb{R} -linear and preserves length.
- Prove that $\text{Aut}(\mathbb{O})$ is a compact lie group by showing that it is a closed subgroup of the compact Lie group $O(7)$.
- Prove that $\text{Aut}(\mathbb{O})$ is simple.
- Prove that $\text{Aut}(\mathbb{O})$ is connected and simply connected.
- Prove that the dimension of $\text{Aut}(\mathbb{O})$ is 14.
- By the classification of simple compact connected and simply connected Lie groups, conclude that $\text{Aut}(\mathbb{O}) \cong G_2$.

Key steps for proving $\text{Aut}(\mathbb{O}) \cong G_2$

- Prove that if $f \in \text{Aut}(\mathbb{O})$ then f is \mathbb{R} -linear and preserves length.
- Prove that $\text{Aut}(\mathbb{O})$ is a compact lie group by showing that it is a closed subgroup of the compact Lie group $O(7)$.
- Prove that $\text{Aut}(\mathbb{O})$ is simple.
- Prove that $\text{Aut}(\mathbb{O})$ is connected and simply connected.
- Prove that the dimension of $\text{Aut}(\mathbb{O})$ is 14.
- By the classification of simple compact connected and simply connected Lie groups, conclude that $\text{Aut}(\mathbb{O}) \cong G_2$.

The Lie group F_4

- The exceptional Jordan algebra $\mathfrak{h}_3(\mathbb{O})$ consists of 3×3 hermitian matrices with octonion entries.
- $\mathbb{O}P^2$ correspond to trace 1 projections in $\mathfrak{h}_3(\mathbb{O})$.
- The Lie group F_4 is $\text{Aut}(\mathfrak{h}_3(\mathbb{O}))$.

The Lie group F_4

- The exceptional Jordan algebra $\mathfrak{h}_3(\mathbb{O})$ consists of 3×3 hermitian matrices with octonion entries.
- $\mathbb{O}P^2$ correspond to trace 1 projections in $\mathfrak{h}_3(\mathbb{O})$.
- The Lie group F_4 is $\text{Aut}(\mathfrak{h}_3(\mathbb{O}))$.

The Lie group F_4

- The exceptional Jordan algebra $\mathfrak{h}_3(\mathbb{O})$ consists of 3×3 hermitian matrices with octonion entries.
- $\mathbb{O}P^2$ correspond to trace 1 projections in $\mathfrak{h}_3(\mathbb{O})$.
- The Lie group F_4 is $\text{Aut}(\mathfrak{h}_3(\mathbb{O}))$.



BROWN

Yale

Surfaces Subgroups of Cocompact Lattices of Isometries of Hyperbolic Spaces

Zhenghao Rao

Geometry and Topology Seminar at Brown and Yale

April 20, 2024

Department of Mathematics

Brown University

Definitions

- Kleinian group: discrete subgroup of $\mathrm{PSL}(2, \mathbb{C})$
- Fuchsian group: finitely generated discrete subgroup of $\mathrm{PSL}(2, \mathbb{R})$
(the limit set is contained in a round circle)
- quasi-Fuchsian group: finitely generated discrete subgroup of $\mathrm{PSL}(2, \mathbb{C})$ with the limit set contained in a topological circle.
(\Leftrightarrow quasiconformally conjugate to a Fuchsian group)

Surface Subgroup Theorem

Theorem (Kahn-Marković, 2012)

Let $M^3 = \mathbb{H}^3/\mathcal{G}$ denote a closed hyperbolic three manifold where \mathcal{G} is a Kleinian group, and let $\epsilon > 0$. Then there exists a Riemann surface $S_\epsilon = \mathbb{H}^2/F_\epsilon$ where F_ϵ is a Fuchsian group, and F_ϵ is $(1 + \epsilon)$ -quasiconformally conjugate to a subgroup of \mathcal{G} .

Main Results

Theorem (Rao, 2023)

Let Γ be a genus-2 quasi-Fuchsian group and G be a cocompact Kleinian group. For any $K > 1$, there is a surface subgroup $H < G$ that is K -quasiconformally conjugate to a finite index subgroup $F < \Gamma$.

Main Results

Theorem (Rao, 2023)

Let Γ be a genus-2 quasi-Fuchsian group and G be a cocompact Kleinian group. For any $K > 1$, there is a surface subgroup $H < G$ that is K -quasiconformally conjugate to a finite index subgroup $F < \Gamma$.

Theorem (Rao, 2023)

For any cocompact Kleinian group G , the set of the Hausdorff dimension of limit sets of all surface subgroups of G is dense in $[1, 2]$.

More Generalization

Theorem (Hamenstädt, 2015)

Let \mathcal{G} be a simple rank one Lie group of non-compact type distinct from $\mathrm{SO}(2m, 1)$ for some $m \geq 1$ and $\Gamma < \mathcal{G}$ be a cocompact lattice. Then Γ contains surface subgroups.

Thank You!

$$S^2 \times S^2 \xrightarrow{1_6} \mathbb{C}P^2 \xrightarrow{1c} S^4$$

$$\textcolor{orange}{\zeta}(x,y) = (y,x)$$

$$\textcolor{violet}{C}[z_1:z_2:z_3] = [\bar{z}_1:\bar{z}_2:\bar{z}_3]$$

$$S^2 \times S^2 \xrightarrow{1_6} \mathbb{C}P^2 \xrightarrow{1c} S^4$$

$$\iota(x, y) = (y, x)$$

$$C[z_1 : z_2 : z_3] = [\bar{z}_1 : \bar{z}_2 : \bar{z}_3]$$


alg. geo

Massey '72
Kuiper '73

$$S^2 \times S^2 \xrightarrow{1_6} \mathbb{C}P^2 \xrightarrow{1_C} S^4$$

\zeta(x,y) = (y,x)


C[z_1, z_2, z_3] = [\bar{z}_1, \bar{z}_2, \bar{z}_3]



$$S^2 \times S^2 \xrightarrow{1_6} \mathbb{C}P^2 \xrightarrow{1c} S^4$$

\iota(x,y) = (y,x)


C[z_1, z_2, z_3] = [\bar{z}_1, \bar{z}_2, \bar{z}_3]



$$S^2 \times S^2 \xrightarrow{1_0} \mathbb{C}P^2 \xrightarrow{1_C} S^4$$

$$\iota(x,y) = (y,x)$$


$$C[z_1:z_2:z_3] = [\bar{z}_1:\bar{z}_2:\bar{z}_3]$$



$$S^2 \times S^2 \xrightarrow{1_0} \mathbb{C}P^2 \xrightarrow{1_C} S^4$$

$$\zeta(x,y) = (y,x)$$


$$C[z_1:z_2:z_3] = [\bar{z}_1:\bar{z}_2:\bar{z}_3]$$



$$S^2 \times S^2 \xrightarrow{1_6} \mathbb{C}P^2 \xrightarrow{1_C} S^4$$

$\zeta(x,y) = (y,x)$

$$C[z_1:z_2:z_3] = [\bar{z}_1:\bar{z}_2:\bar{z}_3]$$



Flows vs foliations: universal circles and ideal boundaries

Junzhi Huang

Yale University

GATSBY, Spring 2024

Depth-1 foliation from spinning

Let M be a fibered closed hyperbolic 3-manifold with fiber F and pseudo-Anosov monodromy $f : F \rightarrow F$:

$$M = F \times [0, 1] / (x, 1) \sim (f(x), 0).$$

Depth-1 foliation from spinning

Let M be a fibered closed hyperbolic 3-manifold with fiber F and pseudo-Anosov monodromy $f : F \rightarrow F$:

$$M = F \times [0, 1] / (x, 1) \sim (f(x), 0).$$

We denote the suspension flow by ϕ and let Σ be a closed embedded taut non-fiber surface in M that is transverse to ϕ .


Depth-1 foliation from spinning

Let M be a fibered closed hyperbolic 3-manifold with fiber F and pseudo-Anosov monodromy $f : F \rightarrow F$:

$$M = F \times [0, 1] / (x, 1) \sim (f(x), 0).$$

We denote the suspension flow by ϕ and let Σ be a closed embedded taut non-fiber surface in M that is transverse to ϕ .

We can “spin” the fibration of M around Σ to obtain a taut depth-1 foliation \mathcal{F} in M transverse to ϕ . The foliation \mathcal{F} has a unique compact leaf Σ , and all the other leaves are surfaces of infinite type spiraling towards Σ .



A circle from the foliation

Let \tilde{M} be the universal cover of M and let $\tilde{\mathcal{F}}$ be the lift of \mathcal{F} to \tilde{M} .

A circle from the foliation

Let \tilde{M} be the universal cover of M and let $\tilde{\mathcal{F}}$ be the lift of \mathcal{F} to \tilde{M} .

A construction of Thurston and Calegari-Dunfield associates a circle \mathfrak{S} , called a universal circle, to \mathcal{F} .

A circle from the foliation

Let \tilde{M} be the universal cover of M and let $\tilde{\mathcal{F}}$ be the lift of \mathcal{F} to \tilde{M} .

A construction of Thurston and Calegari-Dunfield associates a circle \mathfrak{S} , called a universal circle, to \mathcal{F} .

The circle \mathfrak{S} admits a faithful $\pi_1(M)$ -action. For any leaf λ of $\tilde{\mathcal{F}}$, there is a monotone map $I_\lambda : \mathfrak{S} \rightarrow \partial_\infty \lambda$ (by identifying λ with \mathbb{H}^2) so that for any $\gamma \in \pi_1(M)$, the following diagram commutes:

$$\begin{array}{ccc} \mathfrak{S} & \xrightarrow{I_\lambda} & \partial_\infty \lambda \\ \gamma \downarrow & & \downarrow \gamma \\ \mathfrak{S} & \xrightarrow{I_{\gamma\lambda}} & \partial_\infty(\gamma\lambda) \end{array}$$

A circle from the foliation

Let \tilde{M} be the universal cover of M and let $\tilde{\mathcal{F}}$ be the lift of \mathcal{F} to \tilde{M} .

A construction of Thurston and Calegari-Dunfield associates a circle \mathfrak{S} , called a universal circle, to \mathcal{F} .

The circle \mathfrak{S} admits a faithful $\pi_1(M)$ -action. For any leaf λ of $\tilde{\mathcal{F}}$, there is a monotone map $I_\lambda : \mathfrak{S} \rightarrow \partial_\infty \lambda$ (by identifying λ with \mathbb{H}^2) so that for any $\gamma \in \pi_1(M)$, the following diagram commutes:

$$\begin{array}{ccc} \mathfrak{S} & \xrightarrow{I_\lambda} & \partial_\infty \lambda \\ \downarrow \gamma & & \downarrow \gamma \\ \mathfrak{S} & \xrightarrow{I_{\gamma\lambda}} & \partial_\infty(\gamma\lambda) \end{array}$$

The universal circle exists and the π_1 -action is faithful for any taut foliation with hyperbolic leaves in an atoroidal 3-manifold. In general, the construction is not canonical and not unique.

The pseudo-Anosov suspension flow

Let ϕ be the pseudo-Anosov suspension flow in M , and let $\mathcal{F}_M^{s/u}$ denotes the 2-dimensional singular stable/unstable foliation of ϕ .

The pseudo-Anosov suspension flow

Let ϕ be the pseudo-Anosov suspension flow in M , and let $\mathcal{F}_M^{s/u}$ denotes the 2-dimensional singular stable/unstable foliation of ϕ .


We lift ϕ and $\mathcal{F}_M^{s/u}$ to $\tilde{\phi}$ and $\tilde{\mathcal{F}}_M^{s/u}$ in \tilde{M} . Quotienting \tilde{M} by $\tilde{\phi}$ we obtain the flow space \mathcal{O} . The foliation $\tilde{\mathcal{F}}_M^{s/u}$ descends to a foliation $\mathcal{F}_{\mathcal{O}}^{s/u}$ in \mathcal{O} .

The pseudo-Anosov suspension flow

Let ϕ be the pseudo-Anosov suspension flow in M , and let $\mathcal{F}_M^{s/u}$ denotes the 2-dimensional singular stable/unstable foliation of ϕ .

We lift ϕ and $\mathcal{F}_M^{s/u}$ to $\tilde{\phi}$ and $\tilde{\mathcal{F}}_M^{s/u}$ in \tilde{M} . Quotienting \tilde{M} by $\tilde{\phi}$ we obtain the flow space \mathcal{O} . The foliation $\tilde{\mathcal{F}}_M^{s/u}$ descends to a foliation $\mathcal{F}_{\mathcal{O}}^{s/u}$ in \mathcal{O} .

By fibration, we can identify \mathcal{O} with the universal cover of the fiber F , and $\mathcal{F}_{\mathcal{O}}^{s/u}$ with the lift of the stable/unstable foliations of f .



The circle from the flow

The flow space \mathcal{O} has a natural ideal boundary $\partial\mathcal{O}$, and every ray of $\mathcal{F}_{\mathcal{O}}^{s/u}$ has a well-defined ideal endpoint. There is a natural $\pi_1(M)$ -action on $\partial\mathcal{O}$.

The circle from the flow

The flow space \mathcal{O} has a natural ideal boundary $\partial\mathcal{O}$, and every ray of $\mathcal{F}_{\mathcal{O}}^{s/u}$ has a well-defined ideal endpoint. There is a natural $\pi_1(M)$ -action on $\partial\mathcal{O}$.


Let p be the projection map from \tilde{M} to \mathcal{O} . For every leaf λ of $\tilde{\mathcal{F}}$, $p(\lambda)$ is a region bounded by leaves of $\mathcal{F}_{\mathcal{O}}^{s/u}$.

The circle from the flow

The flow space \mathcal{O} has a natural ideal boundary $\partial\mathcal{O}$, and every ray of $\mathcal{F}_{\mathcal{O}}^{s/u}$ has a well-defined ideal endpoint. There is a natural $\pi_1(M)$ -action on $\partial\mathcal{O}$.

Let p be the projection map from \tilde{M} to \mathcal{O} . For every leaf λ of $\tilde{\mathcal{F}}$, $p(\lambda)$ is a region bounded by leaves of $\mathcal{F}_{\mathcal{O}}^{s/u}$.

Each boundary leaf correspond to a unique point at infinity of λ .



The identification of the two circles


It turns out that the circle \mathfrak{S} from the foliation \mathcal{F} and the circle $\partial\mathcal{O}$ from the transverse pseudo-Anosov flow ϕ can be identified.

The identification of the two circles

It turns out that the circle \mathfrak{S} from the foliation \mathcal{F} and the circle $\partial\mathcal{O}$ from the transverse pseudo-Anosov flow ϕ can be identified.

Theorem

There is a $\pi_1(M)$ -equivariant homeomorphism $T : \mathfrak{S} \rightarrow \partial\mathcal{O}$. The monotone map $I_\lambda \circ T^{-1}$ is given by collapsing each component of $\partial\mathcal{O} - \overline{p(\lambda)}$ to the ideal point corresponding to the boundary leaf of $p(\lambda)$ it is facing.



Thank you!