

**Graph homology to motivate the generalisation of the Tutte
polynomial to simplicial complexes**

PROJECT IV: COMPUTATIONAL TOPOLOGY,
MICHAELMAS TERM, WEEK 5
BEN NAPIER

Theorem 1. *Let $G = (V, E)$ be a directed graph. Then*

$$H_0(G) \cong \mathbb{Z}^{|C|}, \quad H_1(G) \cong \mathbb{Z}^{|E|-|V|+|C|}$$

where $C \subset G$ denotes the set of components of G .

Proof. Recall that $H_0(G) = \ker \partial_0 / \operatorname{im} \partial_1$. Now we consider a component of G . We see that any vertex can be obtained by another vertex by adding elements of $\operatorname{im} \partial_1$. Thus all vertices belong to the same equivalence class. It is clear that we cannot do the same for two vertices within different components, thus $H_0(G) \cong \mathbb{Z}^{|C|}$. Now we let $T \subset E'$ be a spanning tree of a component $G' = (V', E')$. We observe that if we add another edge (not in T) to T , we get a cycle. Thus each edge in $E' \setminus T$ corresponds to a cycle in G' . $|E' \setminus T| = |E'| - (|V'| - 1)$. Considering every component, we get $H_1(G) \cong \mathbb{Z}^{|E|-|V|+|C|}$. \square