

Quantentheorie II Übung 7

Besprechung: 2021WE23 (KW23)

SS 2021

Prof. Dominik Stöckinger (IKTP)

1. Questions

- (a) In the lecture some remarks were made on the normalization of symmetrized states (see e.g. section 2.2.2 on states obtained via the symmetrization operators). Discuss the different options for normalization for the example states $|nn\rangle^{(+)}$ and $|nm\rangle^{(+)}$ for $n \neq m$.
- (b) Discuss simple examples how you can experimentally determine whether two particles are distinguishable bosons/fermions. (see e.g. lecture 2.3.2)
- (c) What is the difference between the occupation number formalism and the “multi-particle-state” formalism?

2. Creation and annihilation operators: What are the resulting states of the following expressions for bosons and fermions respectively?

- (a) $a_5^\dagger a_3^\dagger a_4^\dagger a_2^\dagger a_7^\dagger |0\rangle$
- (b) $a_4^\dagger a_8^\dagger a_5 a_1 a_3^\dagger a_5^\dagger a_6^\dagger |0\rangle$
- (c) $a_4^\dagger a_7^\dagger a_5 a_1 a_1^\dagger a_3^\dagger a_5^\dagger a_8^\dagger |0\rangle$
- (d) $a_5^\dagger a_3^\dagger a_8^\dagger a_1 a_3^\dagger a_2^\dagger a_1^\dagger a_7^\dagger |0\rangle$

3. Number operator:

- (a) The number operator is defined as

$$\hat{N} \equiv \sum_i a_i^\dagger a_i .$$

Prove that the following commutation relations are valid for both bosons and fermions:

$$[\hat{N}, a_j^\dagger] = a_j^\dagger, \quad [\hat{N}, a_j] = -a_j .$$

Use the identity

$$[\hat{A}\hat{B}, \hat{C}] = \hat{A}\{\hat{B}, \hat{C}\} - \{\hat{A}, \hat{C}\}\hat{B} .$$

- (b) Show that the general Hamiltonian

$$\hat{H} = \sum_{ij} T_{ij} a_i^\dagger a_j + \frac{1}{2} \sum_{ijkl} V_{ijkl} a_i^\dagger a_j^\dagger a_l a_k$$

commutes with the number operator \hat{N} , i.e. $[\hat{N}, \hat{H}] = 0$. What is the physical meaning of this?

4. **Second quantization I:** consider a system of N identical bosons and let $\hat{\mathcal{O}}_i$ be an operator which operates on the variables of the i th particle only.

- (a) Find the expression for an operator of the form $\hat{\mathcal{O}} = \sum_{i=1}^N \hat{\mathcal{O}}_i$ in second quantization. Assume the discrete basis for the one-particle states.
- (b) Repeat this to find the second quantization expression for the total momentum operator of the system:

$$\hat{P} = \sum_{i=1}^N p_i .$$

Use as basis the continuous 1-dimensional momentum eigenstates normalized as $\langle p' | p \rangle = \delta(p' - p)$.

5. **Second quantization II:** similarly to the previous task consider an operator of the form

$$\hat{\mathcal{O}} = \frac{1}{2} \sum_{i,j=1(i \neq j)}^N \hat{\mathcal{O}}_{ij} , \quad (1)$$

where $\hat{\mathcal{O}}_{ij}$ acts simultaneously on the i th and on the j th particles.

- (a) Rewrite $\hat{\mathcal{O}}$ in Eq. (1) in the second quantization formalism.
- (b) Consider a particular Hamiltonian \hat{H} written in terms of the variables ξ_i (the position and spin of the i th particle) in the form

$$\hat{H} = \sum_{i=1}^N \left[-\frac{1}{2m} \nabla_i^2 + V(\xi_i) \right] + \frac{1}{2} \sum_{i,j=1(i \neq j)}^N W(\xi_i, \xi_j) = \sum_{i=1}^N \hat{H}_i + \frac{1}{2} \sum_{i,j=1(i \neq j)}^N W(\xi_i, \xi_j) .$$

As one-particle basis states use the eigenstates $|\psi_k\rangle$, where $\hat{H}_i |\psi_k\rangle = E_k |\psi_k\rangle$.

6. **N bosons in a cube of edge length L :** consider a system of bosons whose Hamiltonian is of the form

$$\hat{H} = \sum_{i=1}^N \left[-\frac{1}{2m} \nabla_i^2 + \frac{1}{2} \sum_{j=1(i \neq j)}^N V(|\vec{r}_i - \vec{r}_j|) \right] , \quad (2)$$

where $V(|\vec{r}_i - \vec{r}_j|)$ is the two-particle interaction energy, and depends only on the distance between the particles.

Choose as one-particle basis states the eigenstates of momentum with eigenvalue \vec{p} with wave function

$$\psi_{\vec{p}}(\vec{r}) = L^{-\frac{3}{2}} e^{i\vec{p} \cdot \vec{r}} ,$$

which are periodic and normalized in a cube of edge length L , and show that the Hamiltonian \hat{H} in Eq. (2) can be written in second quantization in the form

$$\hat{H} = \sum_k \frac{p_k^2}{2m} a_{\vec{p}_k}^\dagger a_{\vec{p}_k} + \frac{1}{2} \sum \frac{1}{L^3} \omega(\vec{p}_l - \vec{p}_i) a_{\vec{p}_l}^\dagger a_{\vec{p}_m}^\dagger a_{\vec{p}_i} a_{\vec{p}_k} , \quad (3)$$

where $\omega(\vec{p}) \equiv \int d\vec{q} V(|\vec{q}|) e^{-i\vec{p} \cdot \vec{q}}$, and the second summation in Eq. (3) is carried out subject to the condition $\vec{p}_l + \vec{p}_m = \vec{p}_i + \vec{p}_k$.

Note: this exercise is useful in describing superfluidity.