




Robust Post-Matching Inference

Alberto Abadie & Jann Spiess


To cite this article: Alberto Abadie & Jann Spiess (2021): Robust Post-Matching Inference, Journal of the American Statistical Association, DOI: [10.1080/01621459.2020.1840383](https://doi.org/10.1080/01621459.2020.1840383)


To link to this article: <https://doi.org/10.1080/01621459.2020.1840383>

 View supplementary material [↗](#)


 Published online: 14 Jan 2021.

 Submit your article to this journal [↗](#)

 Article views: 710

 View related articles [↗](#)

 View Crossmark data [↗](#)

 Citing articles: 5 View citing articles [↗](#)



Robust Post-Matching Inference

Alberto Abadie^a and Jann Spiess^b

^aDepartment of Economics, MIT, Cambridge, MA; ^bGraduate School of Business, Stanford University, Stanford, CA

ABSTRACT

Nearest-neighbor matching is a popular nonparametric tool to create balance between treatment and control groups in observational studies. As a preprocessing step before regression, matching reduces the dependence on parametric modeling assumptions. In current empirical practice, however, the matching step is often ignored in the calculation of standard errors and confidence intervals. In this article, we show that ignoring the matching step results in asymptotically valid standard errors if matching is done without replacement and the regression model is correctly specified relative to the population regression function of the outcome variable on the treatment variable and *all* the covariates used for matching. However, standard errors that ignore the matching step are not valid if matching is conducted with replacement or, more crucially, if the second step regression model is misspecified in the sense indicated above. Moreover, correct specification of the regression model is not required for consistent estimation of treatment effects with matched data. We show that two easily implementable alternatives produce approximations to the distribution of the post-matching estimator that are robust to misspecification. A simulation study and an empirical example demonstrate the empirical relevance of our results. Supplementary materials for this article are available online.

ARTICLE HISTORY

Received January 2019
Accepted August 2020

KEYWORDS

Matching; Robust estimation;
Treatment effects

1. Introduction

Matching methods are widely used to create balance between treatment and control groups in observational studies. Oftentimes, matching is followed by a simple comparison of means between treated and nontreated (Cochran 1953; Rubin 1973; Dehejia and Wahba 1999). In other instances, however, matching is used in combination with regression or with other estimation methods more complex than a simple comparison of means. The combination of matching in a first step with a second-step regression estimator brings together parametric and nonparametric estimation strategies and, as demonstrated in Ho et al. (2007), reduces the dependence of regression estimates on modeling decisions. Moreover, matching followed by regression allows the estimation of elaborate models, such as those that include interaction effects and other parameters that go beyond average treatment effects.

In this article, we develop valid standard error estimates for regression after matching. The large sample properties of average treatment effect estimators that employ a simple comparison of mean outcomes between treated and nontreated after matching on covariates are well understood (see, e.g., Abadie and Imbens 2006). However, studies that employ regression models after matching usually ignore the matching step when performing inference on post-matching regression coefficients. We show that this practice is not generally valid if the second step regression is misspecified in a sense we make precise below. We propose two easily implementable and robust-to-misspecification approaches to the estimation of the standard errors of regression coefficient estimators in matched samples

(with matching done without replacement). First, we show that standard errors that are clustered at the level of the matched sets are valid under misspecification. Second, we show that a nonparametric block bootstrap that resamples matched pairs or matched sets, as opposed to resampling individual observations, also yields valid inference under misspecification. Furthermore, we show that standard errors that ignore the matching step can both underestimate or overestimate the variation of post-matching estimates. The procedures that we propose in this article are straightforward to implement with standard statistical software.

We will consider the following setup. Let W be a binary random variable representing exposure to the treatment or condition of interest (e.g., smoking), so $W = 1$ for the treated, and $W = 0$ for the nontreated. Y is a random variable representing the outcome of interest (e.g., forced expiratory volume) and X is a vector of covariates (e.g., gender or age). We will study the problem of estimating how the treatment affects the outcomes of the individuals in the treated population (i.e., those with $W = 1$). In particular, we will analyze the properties of a two-step (first matching, then regression) estimator often used in empirical practice. This estimation strategy starts with an unmatched sample, \mathcal{S} , from which treated units and their matches are extracted to create a matched sample, \mathcal{S}^* . Matching is done without replacement and on the basis of the values of X . Then, using data for the matched sample only, the researcher runs a regression of Y on Z , where Z is a vector of functions of W and X (e.g., individual variables plus interactions). We aim to obtain valid inferential methods for the coefficients of this regression, possibly under misspecification. To be precise,

by “misspecification” we mean that there is no version of the conditional expectation of Y given W and X that follows the functional form employed in the second-step estimator. For example, as explained below, a difference in means between treated and nontreated in the second step would be “misspecified” if the conditional expectation of Y given X and W depends on X . To simplify the exposition, here we have described a setting where Z depends only on the treatment, W , and on the covariates used in the matching stage, X . Our general framework in Section 2 allows Z to depend on other covariates not in X .

The intuition behind the results in this article is that, if Y depends on X , then matching on X creates dependence between the outcomes of treated units and their matches. This dependence is absorbed by the second-step regression function as long as the regression function is correctly specified relative to the population regression of Y on W and X . However, if the second-step regression is misspecified relative to the population regression of Y on W and X , dependence between treated units and matches remains in the regression residuals. Ignoring this dependence produces biased inference. Clustered standard errors and analogous block bootstrap procedures take into account the dependence between the outcomes of treated units and their matches, restoring valid inference.

A special case of our setup is that of the standard matching estimator for the average treatment effect on the treated, which is given by the regression coefficient on treatment W in a regression of Y on $Z = (1, W)'$. However, the framework allows for richer analysis, such as the analysis of linear interaction effects of the treatment with covariates, $Z = (1, W, WX', X')'$.

To illustrate the implications of our results, consider the simple case when $Z = (1, W)'$. As we mentioned previously, for $Z = (1, W)'$ the sample regression coefficient on W corresponds to the simple matching estimator often employed in applied studies, which is based on a post-matching comparison of means between treated and nontreated. Under well-known conditions this estimator is consistent for the average effect of the treatment on the treated (see, e.g., Abadie and Imbens 2012), irrespective of the true form of the expectation of Y given W and X . Notice, however, that even in this simple scenario, our results imply that regression standard errors that ignore the matching step are not valid in general. Although the expectation of Y given W is linear because W is binary, a linear regression of Y on $Z = (1, W)'$ will be misspecified *relative to the regression of Y on W and X* , unless Y is mean-independent of X given W over a set of probability one.

The rest of the article is organized as follows. Section 2 starts with a detailed description of the setup of our investigation. We then characterize the parameters estimated by the two-step procedure described above. We show that these parameters are equal to the regression coefficients in a regression of Y on Z in a population for which the distribution of matching covariates X in the control group has been modified to coincide with that of the treated. Under selection on observables—that is, if treatment is as good as random conditional on X —post-matching regression estimands are equal to the population regression coefficients in an experiment where the treatment is randomly assigned in a population that has the same distribution of X as the treated. We next establish consistency with respect to this vector of parameters, show asymptotic normality, and describe

the asymptotic variance of the post-matching estimator. In Section 3, we discuss different ways of constructing standard errors. Based on the results of Section 2, we show that standard errors that ignore the matching step are not generally valid if the regression model is misspecified in the sense indicated above, while clustered standard errors or an analogous block bootstrap procedure yield valid inference. Section 4 presents simulation evidence, which confirms our theoretical results. Section 5 applies our results to the analysis of the effect of smoking on pulmonary function. In this application, matching before regression and the use of the robust standard errors proposed in this article substantially affect empirical findings. Section 6 concludes.

The appendix contains the proofs of our main results. A supplementary appendix contains proofs of intermediate results and two extensions. In particular, the standard errors derived in this article are valid for unconditional inference. Alternatively, one could perform inference conditional on the values of the regressors, X and W , in the sample. Notice that, in this case, the first step matches are fixed. We discuss this alternative setting in the supplementary appendix, where we show that, for the conditional case, the usual regression standard errors are not generally valid, but valid standard errors can be calculated using the formulas in Abadie, Imbens, and Zheng (2014). Also, for concreteness and following the vast majority of applied practice, in the main text of this article we restrict our analysis to linear regression after matching. In the supplementary appendix, we provide an extension of our result to general M -estimation after matching.

2. Post-Matching Inference

In this section, we discuss the asymptotic distribution of the least squares estimator obtained from a linear regression of Y on Z after matching on observables, X .

2.1. Post-Matching Least Squares

Consider a standard binary treatment setting along the lines of Rubin (1974) with potential treatment outcomes $Y(1)$ and $Y(0)$, of which we only observe $Y = Y(W)$ for treatment $W \in \{0, 1\}$. Let S be a set of observed covariates.

We will assume that the data consist of random samples of treated and nontreated. This assumption could be easily relaxed, and we adopt it only to simplify the discussion.

Assumption 1 (Random sampling). $\mathcal{S} = \{(Y_i, W_i, S_i)\}_{i=1}^N$ is a pooled sample obtained from N_1 and N_0 independent draws from the population distribution of (Y, S) for the treated ($W = 1$) and nontreated ($W = 0$), respectively, so $N = N_0 + N_1$.

Consider an $(m \times 1)$ vector of covariates $X = f(S) \in \mathcal{X} \subseteq \mathbb{R}^m$, and let $\mathcal{S}^* \subseteq \mathcal{S}$ be the matched sample generated by matching without replacement each treated unit to M nontreated units on the basis of their X -values. We will denote $\mathcal{J}(i)$ the set of nontreated units matched to treated unit i . For simplicity, in our notation we omit the dependence of $\mathcal{J}(i)$ on N and M . Often, for matching without replacement, the sets $\mathcal{J}(i)$ form the collection

of nonoverlapping subsets of $\{j : W_j = 0\}$, each of cardinality M , that minimizes the sum of the matching discrepancies.

$$\sum_{i=1}^N W_i \sum_{j \in \mathcal{J}(i)} d(X_i, X_j), \tag{1}$$

where $d : \mathcal{X} \times \mathcal{X} \mapsto [0, \infty)$ is a metric. More generally, our conditions do not require a matching scheme that directly minimizes (1), as long as Assumption 3 and the Lipschitz conditions in Assumption 4 and Proposition 3 hold for some metric, $d(\cdot, \cdot)$, under the adopted matching scheme.

The matched sample, $\mathcal{S}^* = \bigcup_{W_i=1} (\{i\} \cup \mathcal{J}(i))$, has size $n = (M + 1)N_1$. We use a double subscript notation to refer to the observations in the matched sample. For instance, Y_{n1}, \dots, Y_{nm} refers to the values of the outcome variable for the units in \mathcal{S}^* , with analogous notation for other variables. Within the matched sample, observations will be rearranged so that the first N_1 observations are the treated units.

Let $Z = g(W, S)$ be a $(k \times 1)$ vector of functions of (W, S) , and let $\hat{\beta}$ be the vector of sample regression coefficients obtained from regressing Y on Z in the matched sample,

$$\begin{aligned} \hat{\beta} &= \operatorname{argmin}_{b \in \mathbb{R}^k} \frac{1}{n} \sum_{i=1}^n (Y_{ni} - Z'_{ni}b)^2 \\ &= \left(\frac{1}{n} \sum_{i=1}^n Z_{ni}Z'_{ni} \right)^{-1} \frac{1}{n} \sum_{i=1}^n Z_{ni}Y_{ni}. \end{aligned} \tag{2}$$

In Section 2.3, we will introduce a set of assumptions under which $\hat{\beta}$ exists with probability approaching one.

As we mentioned above, when $Z = (1, W)'$ the regression coefficient on W in the matched sample is given by

$$\begin{aligned} \hat{\tau} &= \frac{1}{N_1} \sum_{i=1}^n W_{ni}Y_{ni} - \frac{1}{MN_1} \sum_{i=1}^n (1 - W_{ni})Y_{ni} \\ &= \frac{1}{N_1} \sum_{i=1}^N W_i \left(Y_i - \frac{1}{M} \sum_{j \in \mathcal{J}(i)} Y_j \right), \end{aligned}$$

which is the usual matching estimator for the average effect of the treatment on the treated.

2.2. Characterization of the Estimand

Before we study the sampling distribution of $\hat{\beta}$, we first characterize its population counterpart, which we will denote by β . That is, our first task is to obtain a precise description of the nature of the parameters estimated by $\hat{\beta}$. Although post-matching regressions are often used in empirical practice, to the best of our knowledge, the precise nature of post-matching estimands has not been previously derived.

The goal of matching is to change the distribution of the covariates in the sample of nontreated units, so that it reproduces the distribution of the covariates among the treated. To do so, it is necessary that the support of the matching variables, X , for the treated is inside the support for the nontreated.

Assumption 2 (Support condition). Let $\mathcal{X}_1 = \operatorname{supp}(X|W = 1)$ and $\mathcal{X}_0 = \operatorname{supp}(X|W = 0)$, then

$$\mathcal{X}_1 \subseteq \mathcal{X}_0.$$

We now describe the population distribution targeted by the matched sample, \mathcal{S}^* . Let $P(\cdot|W = 1)$ and $P(\cdot|W = 0)$ be the *matching source* distributions of (Y, S) from where the treated and nontreated samples in \mathcal{S} are, respectively, drawn, and let $E[\cdot|W = 1]$ and $E[\cdot|W = 0]$ be the corresponding expectation operators. For given $P(\cdot|W = 1)$ and $P(\cdot|W = 0)$ and a given number of matches, M , we define a *matching target* distribution, P^* , over the triple (Y, S, W) , as follows:

$$P^*(W = 1) = \frac{1}{1 + M},$$

and for each measurable set, A ,

$$P^*((Y, S) \in A|W = 1) = P((Y, S) \in A|W = 1),$$

and

$$P^*((Y, S) \in A|W = 0) = E[P((Y, S) \in A|W = 0, X)|W = 1].$$

That is, in the matching target distribution: (i) treatment is assigned in the same proportion as in the matched sample; (ii) the distribution of (Y, S) among the treated is the same as in the matching source; (iii) the distribution of (Y, S) among the nontreated is generated by integrating the conditional distribution of (Y, S) given X and $W = 0$ over the distribution of X given $W = 1$, in the matching source. As a result, under the matching target distribution, the distribution of X given $W = 0$ coincides with the distribution of X given $W = 1$.

Under regularity conditions stated below, estimation on the matched sample, \mathcal{S}^* , asymptotically recovers parameters of the matching target distribution, P^* , in which the treated and nontreated have the same distribution of X , but possibly different outcome and covariate distributions conditional on X . As a result, comparisons of outcomes between treated and nontreated in the matched sample, \mathcal{S}^* , produce the controlled contrasts of the Oaxaca–Blinder decomposition (Blinder 1973; Oaxaca 1973; DiNardo, Fortin, and Lemieux 1996). More generally, under regularity conditions, regression coefficients of Y on Z in the matched sample, \mathcal{S}^* , asymptotically recover the analogous regression coefficients in the target population:

$$\begin{aligned} \beta &= \operatorname{argmin}_{b \in \mathbb{R}^k} E^*[(Y - Z'b)^2] \\ &= (E^*[ZZ'])^{-1} E^*[ZY]. \end{aligned} \tag{3}$$

Matching methods are often motivated by a selection-on-observables assumption, that is, by the assumption that treatment assignment is as good as random conditional on observed covariates. To formalize the assumption of selection on observables and its implications in our framework, consider source populations expressed this time in terms of potential outcomes and covariates, $Q(\cdot|W = 1)$ and $Q(\cdot|W = 0)$, which represent the distributions of $(Y(1), Y(0), S)$ given $W = 1$ and $W = 0$, respectively. These distributions are defined in such a way that $P(\cdot|W = 1)$ and $P(\cdot|W = 0)$ can be obtained by integrating out $Y(0)$ from $Q(\cdot|W = 1)$ and $Y(1)$ from $Q(\cdot|W = 0)$, respectively. For given $Q(\cdot|W = 1)$ and $Q(\cdot|W = 0)$, selection on observables means

$$(Y(1), Y(0), S)|X, W = 1 \sim (Y(1), Y(0), S)|X, W = 0$$

almost surely with respect to the distribution of $X|W = 1$. That is, the joint distribution of covariates and potential outcomes is independent of treatment assignment conditional on the matching variables. Because in this article, we focus on causal parameters defined for a population with distribution of the matching variables equal to $X|W = 1$, for our purposes it is enough that the selection-on-observables assumption holds for the distribution of $(Y(0), S)$ only,

$$(Y(0), S)|X, W = 1 \sim (Y(0), S)|X, W = 0. \quad (4)$$

Proposition 1 (Estimand under selection on observables). Suppose that [Assumption 2](#) holds and that β , as defined in Equation (3), exists. Then if selection on observables, as defined in Equation (4), holds, the coefficients β are the same as the population coefficients that would be obtained from a regression of Y on Z in a setting where:

1. $(Y(1), Y(0), S)$ has distribution $Q(\cdot|W = 1)$,
2. treatment is randomly assigned with probability $1/(M + 1)$.

This result formalizes the notion that matching under selection on observables allows researchers to reproduce an experimental setting under which average treatment effects can be easily evaluated through a least squares regression of Y on Z . The results in this article, however, apply to the general estimand β in Equation (3), regardless of the validity of the selection-on-observables assumption.

2.3. Consistency and Asymptotic Normality

In this section, we will establish large sample properties of $\hat{\beta}$, as $N_1, N_0 \rightarrow \infty$ with $N_0 \geq MN_1$. Throughout this article, we will assume that the sum of matching discrepancies vanishes quickly enough to allow asymptotic unbiasedness and root- n consistency:

Assumption 3 (Matching discrepancies).

$$\frac{1}{\sqrt{N_1}} \sum_{i=1}^N W_i \sum_{j \in \mathcal{J}(i)} d(X_i, X_j) \xrightarrow{p} 0.$$

Abadie and Imbens (2012) derived primitive conditions for [Assumption 3](#), which require $N_1 = O(N_0^{1/r})$ for some r greater than the number of covariates in X (excluding those that take on a finite number of values). This condition highlights the importance of obtaining matches from a large reservoir of untreated units, especially when the dimensionality of X is large. Of course, in concrete empirical settings, the adequacy of matching should not rely on asymptotic results. Instead, the quality of the matches needs to be evaluated for each particular sample. Abadie and Imbens (2011) and Imbens and Rubin (2015) discussed measures of the discrepancy between the distributions of the covariates of treated and nontreated. For example, the normalized difference in Abadie and Imbens (2011) is $(m_1 - m_0) / \sqrt{(s_1^2 + s_0^2)/2}$, where m_w and s_w^2 are the means and standard deviations of a covariate (typically, products of/and powers of the components of X) for the units with $W = w$ in the matched sample.

For any real matrix A , let $\|A\| = \sqrt{\text{tr}(A'A)}$ be the Euclidean norm of A . The next assumption collects regularity conditions on the conditional moments of (Y, Z) given (X, W) .

Assumption 4 (Well-behavedness of conditional expectations). For $w = 0, 1$, and some $\delta > 0$,

$$E[\|Z\|^4 | W = w, X = x] \quad \text{and} \\ E[\|Z(Y - Z'\beta)\|^{2+\delta} | W = w, X = x]$$

are uniformly bounded on \mathcal{X}_w . Furthermore,

$$E[ZZ' | X = x, W = 0], \quad E[ZY | X = x, W = 0] \\ \text{and} \quad \text{var}(Z(Y - Z'\beta) | X = x, W = 0)$$

are componentwise Lipschitz in x with respect to $d(\cdot, \cdot)$.

To ensure the existence of $\hat{\beta}$ with probability approaching one as $n \rightarrow \infty$, we assume invertibility of the Hessian, $H = E^*(ZZ')$. Notice that

$$H = \frac{E[E[ZZ' | X, W=1] + ME[ZZ' | X, W=0] | W=1]}{1 + M}. \quad (5)$$

Assumption 5 (Linear independence of regressors). H is invertible.

The next proposition establishes the asymptotic distribution of $\hat{\beta}$.

Proposition 2 (Asymptotic distribution of the post-matching estimator). Under [Assumptions 1–5](#),

$$\sqrt{n}(\hat{\beta} - \beta) \xrightarrow{d} \mathcal{N}(0, H^{-1}JH^{-1}),$$

where

$$J = \frac{\text{var}\left(E[Z(Y - Z'\beta) | X, W = 1] + ME[Z(Y - Z'\beta) | X, W = 0] | W = 1\right)}{1 + M} \\ + \frac{E\left[\text{var}(Z(Y - Z'\beta) | X, W = 1) + M\text{var}(Z(Y - Z'\beta) | X, W = 0) | W = 1\right]}{1 + M}$$

and H is as defined in Equation (5).

All proofs are in the appendix.

3. Post-Matching Standard Errors

In the previous section, we established that

$$\sqrt{n}(\hat{\beta} - \beta) \xrightarrow{d} \mathcal{N}(0, H^{-1}JH^{-1})$$

for the post-matching estimator obtained from a regression of Y on Z within the matched sample S^* . In this section, our goal is to estimate the asymptotic variance, $H^{-1}JH^{-1}$.

3.1. Standard Errors Ignoring the Matching Step

Ho et al. (2007) argued that matching can be seen as a preprocessing step, prior to estimation, so the matching step can be ignored in the calculation of standard errors. Here, we consider commonly applied “sandwich” standard error estimates for iid data (Eicker 1967; Huber 1967; White 1980a, 1980b, 1982). In an iid setting, sandwich standard errors are valid in large samples even if the regression is misspecified relative to the conditional expectation of Y given Z , in which case the population regression parameters are the coefficients of an L_2 approximation to the conditional expectation. As we will show, however, the assumption of iid data does not apply in matched samples.

Sandwich standard errors can be computed as the square root of the main diagonal of the matrix $\widehat{H}^{-1}\widehat{J}_s\widehat{H}^{-1}/n$, where

$$\widehat{H} = \frac{1}{n} \sum_{i=1}^n Z_{ni}Z'_{ni} \quad (6)$$

and

$$\widehat{J}_s = \frac{1}{n} \sum_{i=1}^n Z_{ni}(Y_{ni} - Z'_{ni}\widehat{\beta})^2 Z'_{ni}. \quad (7)$$

The following proposition derives the probability limit of \widehat{J}_s with data from a matched sample.

Proposition 3 (Convergence of J_s). Suppose that Assumptions 1–5 hold. Assume also that $E[Z(Y - Z'\beta)^2 Z' | X = x, W = 0]$ is Lipschitz on \mathcal{X}_0 and $E[Y^4 | X = x, W = w]$ is uniformly bounded on \mathcal{X}_w for all $w = 0, 1$. Then, $\widehat{J}_s \xrightarrow{P} J_s$, where

$$J_s = \frac{E\left[E[Z(Y - Z'\beta)^2 Z' | X, W = 1] + ME[Z(Y - Z'\beta)^2 Z' | X, W = 0] | W = 1\right]}{1 + M}.$$

Notice that $J_s = E^*[Z(Y - Z'\beta)^2 Z]$. That is, J_s is equal to the inner matrix of the sandwich asymptotic variance when data are iid with distribution P^* . However, since the matched sample \mathcal{S}^* is not an iid sample from P^* , \widehat{J}_s is not generally consistent for J . The difference between the limit of the sandwich standard errors $\widehat{H}^{-1}\widehat{J}_s\widehat{H}^{-1}$ and the actual asymptotic variance $H^{-1}JH^{-1}$ is given by $H^{-1}\Delta H^{-1}$, where

$$\Delta = \frac{-ME[\Gamma_0(X)\Gamma_1(X)' + \Gamma_1(X)\Gamma_0(X)' | W = 1] - (M - 1)ME[\Gamma_0(X)\Gamma_0(X)' | W = 1]}{M + 1}, \quad (8)$$

and

$$\Gamma_w(x) = E[Z(Y - Z'\beta) | X = x, W = w],$$

for $w = 0, 1$.

Therefore, bias in the estimation of the variance may arise when $\Gamma_0(X) \neq 0$. The following example provides a simple instance of this bias.

Example 1 (Inconsistency of sandwich standard errors). Assume the sample outcomes are drawn from

$$Y = \tau W + X + \varepsilon, \quad (9)$$

where X is a scalar random variable with $\text{var}(X | W = 1) = \sigma_X^2$, and ε has mean zero, variance σ_ε^2 , and is independent of W and X . Consider the case where we match the values of X for N_1 treated units to N_1 untreated units ($M = 1$) without replacement. Let $j(i)$ be the index of the untreated observation that serves as a match for treated observation i . For simplicity, suppose that X is discrete and all matches are perfect, $X_i = X_{j(i)}$ for every treated unit i , so we can ignore potential biases generated by matching discrepancies. Within the matched sample, \mathcal{S}^* , we run a linear regression of Y on $Z = (1, W)'$ to obtain the regression coefficient on W ,

$$\widehat{\tau} = \frac{1}{N_1} \sum_{i=1}^N W_i(Y_i - Y_{j(i)}). \quad (10)$$

$\widehat{\tau}$ is the usual matching estimator for the average effect of the treatment on the treated. Notice that, in the previous expression, $Y_i - Y_{j(i)} = \tau + \varepsilon_i - \varepsilon_{j(i)}$, with variance $2\sigma_\varepsilon^2$. Variation in X is taken care of through matching. Therefore, all variation in $\widehat{\tau}$ comes through the error term, ε . Because $n = 2N_1$, it follows that

$$n \text{var}(\widehat{\tau}) = 4\sigma_\varepsilon^2.$$

Consider now the residuals of the ordinary least squares (OLS) regression of Y_{ni} on a constant and W_{ni} in the matched sample:

$$\widehat{\varepsilon}_{ni} = Y_{ni} - \widehat{\mu} - \widehat{\tau}W_{ni} \approx X_{ni} + \varepsilon_{ni},$$

where $\widehat{\mu}$ is the intercept of the sample regression line. For this simple case, the sandwich variance estimator for $\widehat{\tau}$ is

$$n \widetilde{\text{var}}(\widehat{\tau}) = \frac{4}{n} \sum_{i=1}^n \widehat{\varepsilon}_{ni}^2 \approx 4\sigma_X^2 + 4\sigma_\varepsilon^2.$$

That is, in this example, the sandwich variance estimator overestimates the variance of $\widehat{\tau}$ because it does not take into account the dependence generated by matching between the regression residuals of the treated units and their matches.

Sections 3.2 and 3.3 discuss variance estimators that adjust for the matching step by taking into account the dependence of regression errors between treated units and their matches. For matching with $M = 1$ and a second-step regression of Y on a constant and W , the clustered variance estimator of Section 3.2 becomes

$$n \widehat{\text{var}}(\widehat{\tau}) = \frac{2}{n} \sum_{i=1}^n (\widehat{\varepsilon}_i - \widehat{\varepsilon}_{j(i)})^2 \approx 4\sigma_\varepsilon^2,$$

restoring valid inference.

The next example shows that ignoring the matching step may result in *underestimation* of the variance.

Example 2 (Underestimation of the variance). In the same setting as Example 1, assume that data are generated by

$$Y = \tau W + X - 2WX + \varepsilon. \quad (11)$$

The post-matching estimator of τ from a regression of Y on $(1, W)'$ is $\widehat{\tau}$ as in Equation (10). In this case, if all matches are

perfect, so $X_i = X_{j(i)}$, we obtain $Y_i - Y_{j(i)} = \tau - 2X_i + \varepsilon_i - \varepsilon_{j(i)}$. Therefore,

$$n \operatorname{var}(\widehat{\tau}) = 8\sigma_X^2 + 4\sigma_\varepsilon^2.$$

Least squares regression residuals are

$$\begin{aligned} \widehat{\varepsilon}_{ni} &= Y_{ni} - \widehat{\mu} - \widehat{\tau}W_{ni} \approx X_i - 2W_{ni}X_{ni} + \varepsilon_{ni} \\ &= \begin{cases} -X_{ni} + \varepsilon_{ni} & \text{if } W_{ni} = 1, \\ X_{ni} + \varepsilon_{ni} & \text{if } W_{ni} = 0, \end{cases} \end{aligned}$$

implying

$$n\widetilde{\operatorname{var}}(\widehat{\tau}) = \frac{4}{n} \sum_{i=1}^n \widehat{\varepsilon}_{ni}^2 \approx 4\sigma_X^2 + 4\sigma_\varepsilon^2,$$

for the conventional sandwich variance estimator. Again, the sandwich variance estimator does not take into account dependencies between sample units induced by matching. In this example, matching on X induces a negative correlation between the regression residuals of the treated units and their matches. As a result, the sandwich variance estimator underestimates the variance of $\widehat{\tau}$. Once again, the clustered variance estimator of Section 3.2 takes into account the correlation between regression error induced by matching, and produces valid inference,

$$n \widehat{\operatorname{var}}(\widehat{\tau}) = \frac{2}{n} \sum_{i=1}^n (\widehat{\varepsilon}_i - \widehat{\varepsilon}_{j(i)})^2 \approx 8\sigma_X^2 + 4\sigma_\varepsilon^2.$$

Sandwich standard errors would be valid in Examples 1 and 2 if the specifications for the post-matching regressions included the terms containing X in Equations (9) and (11), respectively. Indeed, sandwich standard errors are generally valid if the regression is correctly specified in a specific sense defined in the following result.

Proposition 4 (Validity of sandwich standard errors under correct specification). Assume that the post-matching regression,

$$Y = Z'\beta + \varepsilon,$$

is correctly specified with respect to the conditional distribution of Y given (Z, X, W) , that is, $E[\varepsilon|Z, X, W] = 0$. Then, under the assumptions of Proposition 3, $J_s = J$ and the sandwich variance estimator, $\widehat{H}^{-1}\widehat{J}_s\widehat{H}^{-1}$, is consistent for the asymptotic variance of $\sqrt{n}(\widehat{\beta} - \beta)$.

Notice, however, that correct specification is precisely the condition under which matching would not be required to obtain a consistent estimator of β , since direct estimation without matching would be valid. Moreover, a correct specification (in the sense defined above) of the post-matching regression is not required for consistent estimation of causal parameters. For example, under regularity conditions, a simple difference in means between the treated and a matched sample of untreated units is consistent for the average effect of the treatment on the treated. Consistent estimators of the variance exist for the simple difference in means in a matched samples. These variance estimators are different from the sandwich variance estimator, and do not rely on correct specification of the post-matching regression (see Abadie and Imbens 2012).

Finally, Equation (8) implies that the conditions of Proposition 4 can be slightly weakened to require only that the regression function is correctly specified among the nontreated, in the sense that $E[\varepsilon|Z, X, W = 0] = 0$. This is because for the estimators studied in this article, matching affects only the distribution of the covariates for the nontreated. In addition, for the special case $M = 1$, it is sufficient that the regression function is correctly specified among the treated, in the sense that $E[\varepsilon|Z, X, W = 1] = 0$.

3.2. Match-Level Clustered Standard Errors

We have shown that sandwich standard errors are not generally valid for the post-matching least squares estimator. In this section, we will demonstrate that, when matching is done without replacement, clustered standard errors (Liang and Zeger 1986; Arellano 1987) can be employed to obtain valid estimates of the standard deviation of post-matching regression coefficients. In particular, we will consider standard errors clustered at the level of the matched sets.

Consider an estimator of the asymptotic variance of $\widehat{\beta}$ given by $\widehat{H}^{-1}\widehat{J}\widehat{H}^{-1}$, where \widehat{H} is as in Equation (6) and \widehat{J} is given by the clustered variance formula applied to the matched sets,

$$\begin{aligned} \widehat{J} &= \frac{1}{n} \sum_{i=1}^n W_i \left(Z_i(Y_i - Z_i'\widehat{\beta}) + \sum_{j \in \mathcal{J}(i)} Z_j(Y_j - Z_j'\widehat{\beta}) \right) \\ &\quad \times \left(Z_i(Y_i - Z_i'\widehat{\beta}) + \sum_{j \in \mathcal{J}(i)} Z_j(Y_j - Z_j'\widehat{\beta}) \right)'. \end{aligned}$$

Clustered standard errors can be readily implemented using standard statistical software. The next result shows that match-level clustered standard errors are valid in large samples for the post-matching estimator (provided matching is done without replacement).

Proposition 5 (Validity of clustered standard errors). Under the assumptions of Proposition 3, we obtain that

$$\widehat{J} \xrightarrow{p} J.$$

In particular, the clustered estimator of the variance is consistent, that is,

$$\widehat{H}^{-1}\widehat{J}\widehat{H}^{-1} - n\operatorname{var}(\widehat{\beta}) \xrightarrow{p} 0.$$

The intuition behind this result is that matching on covariates makes regression errors statistically dependent among units in the same matched sets, $\{i\} \cup \mathcal{J}(i)$, $i = 1, \dots, N_1$. Standard errors clustered at the level of the matched set take this dependency into account.

3.3. Matched Bootstrap

Proposition 5 shows that clustered standard errors are valid for the asymptotic variance of the post-matching estimator. In this section, we show that a clustered version of the nonparametric bootstrap (Efron 1979) is also valid. This version of the bootstrap relies on resampling of matched sets instead on individual observations.

Recall that we reordered the observations in our sample, so that the first N_1 observations are the treated. Consider the nonparametric bootstrap that samples treated units together with their M matches partners from \mathcal{S}^* to obtain

$$\widehat{\beta}^* = \left(\frac{1}{n} \sum_{i=1}^n V_{ni} Z_{ni} Z'_{ni} \right)^{-1} \frac{1}{n} \sum_{i=1}^n V_{ni} Z_{ni} Y_{ni},$$

where $(V_{n1}, \dots, V_{nN_1})$ has a multinomial distribution with parameters $(N_1, (1/N_1, \dots, 1/N_1))$, and $V_{nj} = V_{ni}$ if $j > N_1$ and $j \in \mathcal{J}(i)$. In this bootstrap procedure, N_1 units are drawn at random with replacement from the N_1 treated sample units. Untreated units are drawn along with their treated match. Effectively, the matched bootstrap samples matched sets of one treated unit and M untreated units. The next proposition shows validity of the matched bootstrap.

Proposition 6 (Validity of the matched bootstrap). Under the assumptions of Proposition 5, we have that

$$\sup_{r \in \mathbb{R}^k} |P(\sqrt{n}(\widehat{\beta}^* - \widehat{\beta}) \leq r | \mathcal{S}) - P(\mathcal{N}(0, H^{-1} J H^{-1}) \leq r)| \xrightarrow{p} 0.$$

Proposition 6 shows that the bootstrap distribution provides an asymptotically valid approximation of the limiting distribution of the post-matching estimator, but that does not necessarily imply that the associated bootstrap variance is an asymptotically valid estimate of the variance of the estimator.

The formal analysis of the bootstrap variance is complicated by the fact that, in forming the bootstrap estimate $\widehat{\beta}^*$, the empirical analog

$$\widehat{H}^* = \frac{1}{n} \sum_{i=1}^n V_{ni} Z_{ni} Z'_{ni}$$

of the Hessian H for a given bootstrap draw may be ill-conditioned or noninvertible. In fact, because the bootstrap may sample the same matched set N_1 times, noninvertibility of the Hessian may happen with positive probability for any sample size. To circumvent this issue, we fix constants $c > 0$ and $\alpha \in (0, 1/2)$ and consider the alternative bootstrap estimator

$$\widetilde{\beta}^* = \begin{cases} \widehat{\beta}^* & \text{if } \|\widehat{H}^* - \widehat{H}\| \leq c/n^\alpha, \\ \widehat{\beta} & \text{otherwise.} \end{cases}$$

That is, $\widetilde{\beta}^*$ is equal to $\widehat{\beta}^*$ whenever the bootstrap Hessian, \widehat{H}^* , is close to the matched sample Hessian, \widehat{H} . Otherwise, $\widetilde{\beta}^*$ is equal to the post-matching estimator, $\widehat{\beta}$. As the sample size grows, $\widetilde{\beta}^*$ is equal to $\widehat{\beta}^*$ with probability approaching one.

Proposition 7 (Validity of bootstrap standard errors). Under the assumptions of Proposition 5 and $E[\|Z\|^8 | W = w, X = x]$ uniformly bounded on \mathcal{X}_w , the bootstrap distribution given by $\widetilde{\beta}^*$ is valid in the sense of Proposition 6, and yields a valid estimate of the asymptotic variance of $\widehat{\beta}$, that is,

$$\text{nvar}(\widetilde{\beta}^* | \mathcal{S}) \xrightarrow{p} H^{-1} J H^{-1}$$

as $n \rightarrow \infty$.

The use of $\widetilde{\beta}^*$ in Proposition 7 is a formal device to make the outcome of each bootstrap iteration well-defined. For practical purposes, however, bootstrap standard errors based on $\widehat{\beta}^*$ will perform well unless the bootstrap Hessians are ill-conditioned. Bootstrap standard errors based on $\widehat{\beta}^*$ perform very well in our simulations of Section 4.

It is useful to relate the results in this section, which pertain to matching without replacement, to previous results for matching with replacement. In particular, for matching with replacement Abadie and Imbens (2008) showed that the nonparametric bootstrap fails to consistently estimate the standard error of a simple matching estimator. The consistency results that we obtain in this section is for matching without replacement, and do not directly extend to matching with replacement. The reason is that matching with replacement creates dependencies in the data that are not preserved by resampling matched sets.

4. Simulations

In this section, we study the performance of the post-matching standard error estimators from Section 3 in a simulation exercise using two data generating processes (DGPs).

4.1. DGP1: Robustness to Misspecification

Let $\mathcal{U}(a, b)$ be the uniform distribution on $[a, b]$. We generate data according to

$$Y = WX + 5X^2 + \varepsilon,$$

where $X|W = 1 \sim \mathcal{U}(-1, 1)$, $X|W = 0 \sim \mathcal{U}(-1, 2)$, and $\varepsilon \sim \mathcal{N}(0, 1)$. We sample $N_1 = 50$ treated and $N_0 = 200$ non-treated units. We first match treated and untreated units on the covariates, X , without replacement and with $M = 1$ match per treated unit. We consider the following post-matching regression specifications.

Specification 1:

$$Y = \alpha + \tau_0 W + \tau_1 WX + \beta_1 X + \varepsilon.$$

Specification 2:

$$Y = \alpha + \tau_0 W + \tau_1 WX + \beta_1 X + \beta_2 X^2 + \varepsilon.$$

Specification 2 is correct relative to the conditional expectation $E[Y|X, W]$, while specification 1 is not. Regression estimands can always be seen as L_2 approximations to $E[Y|W, X]$, regardless of the specification adopted for estimation (see, e.g., White 1980b). For our simulation results, we will focus on estimators of τ_0 and τ_1 , the regression coefficients on terms involving W . For the DGP and the two specifications adopted for this simulation, it can be shown that $\tau_0 = 0$ and $\tau_1 = 1$ under the matching target distribution.

Table 1 reports the results of the simulation exercise. In a regression that uses the full sample without matching, the estimates of τ_0 and τ_1 are biased under misspecification (specification 1), while they are valid under correct specification (specification 2). After matching, both specifications yield valid estimates for τ_0 and τ_1 . However, sandwich standard error estimates are inflated under misspecification, while average clustered and

Table 1. Monte Carlo results for DGP1 (10,000 iterations).

(a) Target parameter: coefficient $\tau_0 = 0$ on W								
Specification	Full sample		Post-matching		Average standard error			
	Mean of $\hat{\tau}_0$	Std. of $\hat{\tau}_0$	Mean of $\hat{\tau}_0$	Std. of $\hat{\tau}_0$	Sandwich	Cluster	Bootstrap	
1	-0.85	0.404	0.00	0.204	0.359	0.197	0.199	
2	0.00	0.165	0.00	0.204	0.196	0.196	0.199	

(b) Target parameter: coefficient $\tau_1 = 1$ on the interaction WX								
Specification	Full sample		Post-matching		Average standard error			
	Mean of $\hat{\tau}_1$	Std. of $\hat{\tau}_1$	Mean of $\hat{\tau}_1$	Std. of $\hat{\tau}_1$	Sandwich	Cluster	Bootstrap	
1	-4.00	0.646	0.99	0.358	0.728	0.340	0.348	
2	1.00	0.286	1.00	0.356	0.337	0.338	0.346	

matched bootstrap standard errors (with 1000 bootstrap draws) closely approximate the standard deviation of $\hat{\tau}_0$ and $\hat{\tau}_1$. Under correct specification (specification 2), all standard error estimates perform well.

4.2. DGP2: High Treatment-Effect Heterogeneity

In the simulation in the previous section, sandwich standard errors overestimate the variation of the post-matching estimator under misspecification. In this section, we present an example in which sandwich standard errors are too small. We generate data according to

$$Y = WX + 20WX^2 - 10X^2 + \varepsilon$$

with $\varepsilon \sim \mathcal{N}(0, 1)$ as above. For this DGP2, the conditional treatment effect is nonlinear with

$$E[Y|W = 1, X] - E[Y|W = 0, X] = X + 20X^2.$$

Sample sizes, matching settings, and regression specifications are as in DGP1. Notice that both regression specifications are incorrect relative to $E[Y|X, W]$, as they do not capture nonlinear conditional treatment effects. Like in Section 4.1, regression coefficients represent the parameters of an L_2 approximation to $E[Y|W, X]$ over the distribution of (W, X) in Proposition 1. Direct calculations yield $\tau_0 = 6.67$ and $\tau_1 = 1$ for both specifications in the matching target distribution.

Table 2 presents the results of the simulation exercise for DGP2. The large heterogeneity in conditional treatment effects is not captured by either regression specification, and sandwich standard errors that ignore the matching step underestimate the variation of the post-matching estimator. In contrast, the average clustered and matched bootstrap (with 1000 bootstrap draws) standard errors proposed in this article closely reflect the variability of the post-matching estimators.

5. Application

This section reports the results of an empirical application where we look at the effect of smoking on the pulmonary function of youths. The application is based on data originally collected in

Table 2. Monte Carlo results for DGP2 (10,000 iterations).

(a) Target parameter: coefficient $\tau_0 = 6.67$ on W								
Specification	Full sample		Post-matching		Average standard error			
	Mean of $\hat{\tau}_0$	std. of $\hat{\tau}_0$	mean of $\hat{\tau}_0$	std. of $\hat{\tau}_0$	Sandwich	Cluster	Bootstrap	
1	8.25	0.754	6.55	0.883	0.630	0.869	0.897	
2	6.70	0.857	6.55	0.883	0.630	0.869	0.897	

(b) Target parameter: coefficient $\tau_1 = 1$ on the interaction WX								
Specification	Full sample		Post-matching		Average standard error			
	Mean of $\hat{\tau}_1$	Std. of $\hat{\tau}_1$	Mean of $\hat{\tau}_1$	Std. of $\hat{\tau}_1$	Sandwich	Cluster	Bootstrap	
1	11.00	1.209	1.01	1.950	1.330	1.848	1.932	
2	1.90	1.877	1.01	1.950	1.330	1.848	1.933	

Boston, Massachusetts, by Tager et al. (1979, 1983), and subsequently described and analyzed in Rosner (1995) and Kahn (2005). The sample contains 654 youth, $N_1 = 65$ who have ever smoked regularly ($W = 1$) and $N_0 = 589$ who never smoked regularly ($W = 0$). The outcome of interest is the subjects' forced expiratory volume (Y), ranging from 0.791 to 5.793 liters per second (ℓ/sec). In addition, we use data on age (X_1 , ranging from 3 to 19 with the youngest ever-smoker aged 9) and gender (X_2 , with $X_2 = 1$ for males and $X_2 = 0$ for females).

The use of matching to study the causal effect of smoking is motivated by the likely confounding effects of age and gender. For instance, while the causal effect of smoking on respiratory volume is expected to be negative, older children are more likely to smoke and have a larger respiratory volume, which induces a positive association between smoking and respiratory volume.

We first match every smoker in the sample to a nonsmoker ($M = 1$), without replacement, based on age (X_1) and gender (X_2). Within the resulting matched sample of 65 smokers and 65 nonsmokers, we run linear regressions with the following specifications:

Specification 1:

$$Y = \alpha + \tau_0 W + \varepsilon.$$

Specification 2:

$$Y = \alpha + \tau_0 W + \beta_1 X_1 + \beta_2 X_2 + \varepsilon.$$

Specification 3:

$$Y = \alpha + \tau_0 W + \tau_1 W(X_1 - E[X_1]) + \tau_2 W(X_2 - E[X_2]) + \beta_1(X_1 - E[X_1]) + \beta_2(X_2 - E[X_2]) + \varepsilon.$$

The first specification yields the matching estimator for the average treatment effect τ_0 as the regression coefficient on W , while the second adds linear controls in X_1 and X_2 . The third specification also includes interaction terms of smoking with age and gender.

Table 3 reports regression estimates of τ_0 , τ_1 , and τ_2 along with standard errors (regression coefficients on terms not involving W are omitted from Table 3 for brevity). Estimates for the first specification demonstrate the problem of confounding in this application. Without controlling for age and gender, there is a positive correlation between smoking and forced expiratory

Table 3. OLS and post-matching estimates for the smoking dataset.

	Dependent variable: forced expiratory volume								
	Smoker			Smoker \times age			Smoker \times male		
	Coeff.	Std. error		Coeff.	Std. error		Coeff.	Std. error	
		Sandwich	Cluster		Sandwich	Cluster		Sandwich	Cluster
Specification 1:									
OLS	0.711	0.099							
Post-matching	-0.066	0.132	0.095						
Specification 2:									
OLS	-0.154	0.104							
Post-matching	-0.077	0.104	0.096						
Specification 3:									
OLS	0.495	0.187		-0.182	0.036		0.461	0.193	
Post-matching	-0.077	0.102	0.093	-0.092	0.054	0.038	-0.021	0.249	0.212

function. After matching on age and gender, the sign of the regression coefficient on smoking becomes negative. In this specification, the clustered standard error for the post-matching estimate is considerably smaller than the corresponding sandwich standard error.

Specification 2 includes linear controls for age and gender. The sign and magnitude of the least squares estimate of the coefficient on the smoker variable changes substantially between specifications 1 and 2, while the magnitude of the post-matching estimate stays roughly constant. This result illustrates the higher robustness across specifications of the post-matching estimator relative to least squares on the unmatched sample (Ho et al. 2007). When specification 2 is adopted for regression, the sign of the coefficient on the smoker variable is not affected by matching. Also, for this specification, clustered and sandwich standard errors are similar. Both findings are consistent with the adopted regression specification moving closer toward the correct specification of $E[Y|W, X_1, X_2]$.

In specification 3, which includes interactions between the smoker variable and age and gender, the use of matching and the use of robust standard errors matters for the substantive results of the analysis. First, notice that the coefficient on the interaction of gender with treatment is large, significant and positive without matching, suggesting that the effect of smoking is more severe for girls than for boys. After matching, the sign changes, and the estimated coefficient is small and insignificant. This suggests that the large interaction finding with OLS for this coefficient is caused by misspecification. Second, in the post-matching regression we find a negative estimate for the interaction of treatment with age. With sandwich standard errors, this effect is not significant (at the 5% level). The robust standard errors proposed in this article are smaller and result in a rejection of the null hypothesis of a zero interaction coefficient between smoker and age (at the 5% level).

6. Conclusion

This article establishes valid inference for regression on a sample matched without replacement. Standard errors that ignore the matching step are not generally valid if the regression specification is incorrect relative to the expectation of the outcome conditional on the treatment and the matching covariates. However, using a correct specification relative to $E[Y|W, X]$

is not necessary to consistently estimate treatment parameters after matching. For example, under selection on observables, simple differences in means in a matched sample can be used to estimate average treatment effects.

We propose two alternatives—standard errors clustered at the matched set level and an analogous block bootstrap—that are robust to misspecification and easily implementable with standard statistical software. A simulation study and an empirical example demonstrate the usefulness of our results.

To conclude, we outline potential extensions of our results. First, in this article, we discuss only matching without replacement, and the results do not directly carry over to matching with replacement as in Abadie and Imbens (2006). Matching with replacement (i.e., allowing nontreated units to be used as a match more than once) creates additional dependencies between matched sets that are not reflected in sandwich standard errors or in the robust standard errors proposed in this article. While the negative result about post-matching standard errors extend to matching with replacement (standard errors that ignore the matching step are not generally valid for matching is done with replacement, see Abadie and Imbens 2006), the positive results we describe do not directly apply: Even when the linear regression is correctly specified, sandwich standard errors do not correctly capture the variance of the post-matching estimates, since the overlap between matched sets is not accounted for. Clustered standard errors, as well as the analogous block bootstrap that samples treated units with all their matching partners, do not provide an immediate solution since one untreated unit may now be part of multiple such clusters or bootstrap groups.

In addition, our analysis applies to the case when matching is done directly on the covariates, avoiding substantial complications created by the presence of nuisance parameters in the matching step when matching is done on the estimated propensity score (see Rosenbaum and Rubin 1983; Abadie and Imbens 2016). Finally, our analysis assumes that the quality of matches is good enough for matching discrepancies not to bias the asymptotic distribution of the post-matching regression estimator. Post-matching regression adjustments may, in practice, help eliminate the bias as in the bias-corrected matching estimator in Abadie and Imbens (2011). These are angles that we do not explore in this article and interesting avenues for future research.

Appendix: Proofs

Preliminary Lemmas A.1 and A.2 and Propositions A.1–A.3 are in a supplementary appendix.

Proof of Proposition 1. Let $E_{Q(\cdot|W=1)}$ and $E_{Q(\cdot|W=0)}$ be expectation operators for $Q(\cdot|W = 1)$ and $Q(\cdot|W = 0)$. Notice first that for any measurable function q ,

$$E_{Q(\cdot|W=1)}[q(Y(1), S)] = E[q(Y, S)|W = 1]. \quad (\text{A.1})$$

The result holds also replacing $W = 1$ with $W = 0$, and after conditioning on X . In particular,

$$E_{Q(\cdot|W=0)}[q(Y(0), S)|X] = E[q(Y, S)|X, W = 0]. \quad (\text{A.2})$$

The regression coefficient in the population defined by (a) and (b) is the minimizer of

$$\begin{aligned} & \frac{1}{M+1} E_{Q(\cdot|W=1)}[(Y(1) - g(1, S)'b)^2] \\ & + \frac{M}{M+1} E_{Q(\cdot|W=1)}[(Y(0) - g(0, S)'b)^2]. \end{aligned}$$

Notice that

$$\begin{aligned} E_{Q(\cdot|W=1)}[(Y(1) - g(1, S)'b)^2] &= E[(Y - g(1, S)'b)^2|W = 1] \\ &= E^*[(Y - Z'b)^2|W = 1], \end{aligned}$$

where the first equality follows from Equation (A.1) and the second equality follows from the definitions of $P^*(\cdot|W = 1)$ and Z . Similarly,

$$\begin{aligned} & E_{Q(\cdot|W=1)}[(Y(0) - g(0, S)'b)^2] \\ &= E_{Q(\cdot|W=1)}[E_{Q(\cdot|W=1)}[(Y(0) - g(0, S)'b)^2|X]] \\ &= E_{Q(\cdot|W=1)}[E_{Q(\cdot|W=0)}[(Y(0) - g(0, S)'b)^2|X]] \\ &= E[E[(Y - g(W, S)'b)^2|X, W = 0]|W = 1] \\ &= E^*[(Y - Z'b)^2|W = 0]. \end{aligned}$$

In the last equation, the first equality follows from the law of iterated expectations, the second equality follows from selection on observables, the third equality follows from (A.2) and (A.1), and the last equation follows from the definition of $P^*(\cdot|W = 0)$. Therefore,

$$\begin{aligned} & \frac{1}{M+1} E_{Q(\cdot|W=1)}[(Y(1) - g(1, S)'b)^2] \\ & + \frac{M}{M+1} E_{Q(\cdot|W=1)}[(Y(0) - g(0, S)'b)^2] \\ &= \frac{1}{M+1} E^*[(Y - Z'b)^2|W = 1] \\ & + \frac{M}{M+1} E^*[(Y - Z'b)^2|W = 0] = E^*[(Y - Z'b)^2], \end{aligned}$$

which implies the result of the proposition. \square

Proof of Proposition 2. This proof is based on two lemmas in the supplementary appendix about the asymptotic distribution of averages in matched samples based on a martingale representation of matching estimators similar to Abadie and Imbens (2012). Lemma A.1 establishes convergence in probability, while Lemma A.2 deals with root- n consistency and asymptotic normality. By Lemma A.1,

$$\frac{1}{n} \sum_{i \in S^*} Z_i Z_i' \xrightarrow{P} H.$$

By Lemma A.2,

$$\widehat{H} \sqrt{n} (\widehat{\beta} - \beta) = \sqrt{n} \left(\frac{1}{n} \sum_{i \in S^*} (Z_i Y_i - Z_i Z_i' \beta) \right) \xrightarrow{d} \mathcal{N}(\mathbf{0}, J),$$

where we note that $E[Z(Y - ZZ'\beta)|W = 0, X = x]$ is Lipschitz. Hence,

$$\begin{aligned} & \sqrt{n} (\widehat{\beta} - \beta) \\ &= \underbrace{\widehat{H}^{-1}}_{\xrightarrow{d} \mathcal{N}(\mathbf{0}, J)} \sqrt{n} \left(\frac{1}{n} \sum_{i \in S^*} (Z_i Y_i - Z_i Z_i' \beta) \right) \xrightarrow{d} \mathcal{N}(\mathbf{0}, H^{-1} J H^{-1}). \end{aligned}$$

\square

Proof of Proposition 3. We have that

$$\begin{aligned} \widehat{J}_s &= \frac{1}{n} \sum_{i \in S^*} Z_i (Y_i - Z_i' \widehat{\beta})^2 Z_i' \\ &= \frac{1}{n} \sum_{i \in S^*} Z_i (Y_i - Z_i' \beta)^2 Z_i' \\ &+ \frac{1}{n} \sum_{i \in S^*} Z_i \left((Y_i - Z_i' \widehat{\beta})^2 - (Y_i - Z_i' \beta)^2 \right) Z_i'. \end{aligned}$$

Notice that

$$\begin{aligned} & \frac{1}{n} \sum_{i \in S^*} Z_i \left((Y_i - Z_i' \widehat{\beta})^2 - (Y_i - Z_i' \beta)^2 \right) Z_i' \\ &= (\widehat{\beta} - \beta)' \left(\frac{1}{n} \sum_{i \in S^*} Z_i (Z_i' Z_i) Z_i' (\widehat{\beta} + \beta) - 2 \frac{1}{n} \sum_{i \in S^*} Z_i (Z_i' Z_i) Y_i \right). \end{aligned}$$

By assumption, the functions

$$E[\|Z\|^4|X = x, W = w] \quad \text{and} \quad E[|Y|^4|X = x, W = w]$$

are uniformly bounded on \mathcal{X}_w , for $w = 0, 1$. By Hölder's inequality,

$$E \left[\left\| \frac{1}{n} \sum_{i \in S^*} Z_i Z_i' Z_i Z_i' \right\| \right] \quad \text{and} \quad E \left[\left\| \frac{1}{n} \sum_{i \in S^*} Z_i Z_i' Z_i Y_i \right\| \right]$$

are thus finite. Then, for $\epsilon \in (0, 1/2)$, by Markov's inequality, we obtain

$$\begin{aligned} & \frac{1}{n} \sum_{i \in S^*} Z_i \left((Y_i - Z_i' \widehat{\beta})^2 - (Y_i - Z_i' \beta)^2 \right) Z_i' \\ &= n^{1/2-\epsilon} (\widehat{\beta} - \beta)' \left(\frac{\sum_{i \in S^*} Z_i (Z_i' Z_i) Z_i' / n}{n^{1/2-\epsilon}} (\widehat{\beta} + \beta) \right. \\ & \quad \left. - \frac{2 \sum_{i \in S^*} Z_i (Z_i' Z_i) Y_i / n}{n^{1/2-\epsilon}} \right) \xrightarrow{P} 0. \end{aligned}$$

As a result,

$$\widehat{J}_s = \frac{1}{n} \sum_{i \in S^*} Z_i (Y_i - Z_i' \beta)^2 Z_i' + o_p(1),$$

and the claim follows from Lemma A.1 in the supplementary appendix, which deals with consistency of averages in matched samples. \square

Proof of Proposition 4. Under correct specification, we find that

$$\begin{aligned} \Gamma_W(X) &= E[Z(Y - Z'\beta)|W, X] = E[Z\varepsilon|W, X] \\ &= E[E[Z\varepsilon|Z, W, X]|W, X] = E[Z \underbrace{E[\varepsilon|Z, W, X]}_{=0} |W, X] = 0. \end{aligned}$$

\square

Proof of Proposition 5. First, note that

$$\begin{aligned} \hat{J} &= \frac{1}{n} \sum_{W_i=1} \left(Z_i(Y_i - Z_i'\beta) + \sum_{j \in \mathcal{J}(i)} Z_j(Y_j - Z_j'\beta) \right) \\ &\quad \left(Z_i(Y_i - Z_i'\beta) + \sum_{j \in \mathcal{J}(i)} Z_j(Y_j - Z_j'\beta) \right)' + o_p(1), \end{aligned}$$

where we replace $\hat{\beta}$ by β analogous to the proof of Proposition 3. Write

$$G = Z(Y - Z'\beta) \quad \Gamma_w(x) = E[Z(Y - Z'\beta) | W = w, X = x].$$

Note that $\Gamma_0(x)$ is Lipschitz on \mathcal{X} , and that G_i has uniformly bounded fourth moments. We decompose

$$\begin{aligned} \hat{J} &= \frac{1}{n} \sum_{W_i=1} \left(G_i + \sum_{j \in \mathcal{J}(i)} G_j \right) \left(G_i + \sum_{j \in \mathcal{J}(i)} G_j \right)' + o_p(1) \\ &= \frac{1}{n} \sum_{W_i=1} (\Gamma_1(X_i) + M\Gamma_0(X_i)) (\Gamma_1(X_i) + M\Gamma_0(X_i))' \\ &\quad + \frac{1}{n} \sum_{i \in \mathcal{S}^*} (G_i - \Gamma_{W_i}(X_i)) (G_i - \Gamma_{W_i}(X_i))' \\ &\quad + \frac{1}{n} \sum_{W_i=1} \sum_{\ell \neq i \in \mathcal{J}(i) \cup \{i\}} (G_\ell - \Gamma_{W_\ell}(X_\ell)) (G_\ell - \Gamma_{W_\ell}(X_\ell))' \\ &\quad + \frac{1}{n} \sum_{W_i=1} ((\Gamma_1(X_i) + M\Gamma_0(X_i)) \\ &\quad \left(G_i - \Gamma_1(X_i) + \sum_{j \in \mathcal{J}(i)} (G_j - \Gamma_0(X_j)) \right)' \\ &\quad + (G_i - \Gamma_1(X_i) + \sum_{j \in \mathcal{J}(i)} (G_j - \Gamma_0(X_j))) \\ &\quad (\Gamma_1(X_i) + M\Gamma_0(X_i))') + o_p(1). \end{aligned}$$

Here, the o_p terms absorb the deviation due to using $\hat{\beta}$ instead of β , as well as the matching discrepancies in the conditional expectations. The first sum is iid with

$$\begin{aligned} &\frac{1}{n} \sum_{W_i=1} (\Gamma_1(X_i) + M\Gamma_0(X_i)) (\Gamma_1(X_i) + M\Gamma_0(X_i))' \\ &\xrightarrow{p} \frac{E[(\Gamma_1(X) + M\Gamma_0(X))(\Gamma_1(X) + M\Gamma_0(X))' | W = 1]}{1 + M} \\ &= \frac{E[|W=1]=0 \text{ var}(\Gamma_1(X) + M\Gamma_0(X) | W = 1)}{1 + M}, \end{aligned}$$

while the second is a martingale with

$$\begin{aligned} &\frac{1}{n} \sum_{i \in \mathcal{S}^*} (G_i - \Gamma_{W_i}(X_i)) (G_i - \Gamma_{W_i}(X_i))' \\ &\xrightarrow{p} \frac{E[\text{var}(Z(Y - Z'\beta) | W = 1, X) + M\text{var}(Z(Y - Z'\beta) | W = 0, X) | W = 1]}{1 + M} \end{aligned}$$

by Lemma A.1 in the supplementary appendix, which establishes consistency of averages in matched samples. Under appropriate reordering of the individual increments, all other sums can be represented as averages of mean-zero martingale increments. Since the second moments of the increments are uniformly bounded, they vanish asymptotically. \square

Proof of Proposition 6. In this proof, we invoke Proposition A.2 in the supplementary appendix, which establishes a general result on the validity of the matched bootstrap for averages within matched samples. Write

$$\hat{H}^* = \frac{1}{n} \sum_{i \in \mathcal{S}^*} V_{ni} Z_{ni} Z_{ni}'.$$

Note first that

$$\begin{aligned} &H^{-1} \sqrt{n} (\hat{H}^* (\hat{\beta}^* - \beta) - \hat{H} (\hat{\beta} - \beta)) \\ &= H^{-1} \sqrt{n} \left(\frac{1}{n} \sum_{i=1}^n (V_{ni} - 1) Z_{ni} (Y_{ni} - Z_{ni}'\beta) \right) \\ &\xrightarrow{d} \mathcal{N}(\mathbf{0}, H^{-1} J H^{-1}), \end{aligned}$$

conditional on \mathcal{S} , by Proposition A.2. Now,

$$\begin{aligned} \sqrt{n} (\hat{\beta}^* - \hat{\beta}) &= (\hat{H}^*)^{-1} H (H^{-1} \sqrt{n} (\hat{H}^* (\hat{\beta}^* - \beta) - \hat{H} (\hat{\beta} - \beta))) \\ &= \underbrace{(\hat{H}^*)^{-1} H (H^{-1} \sqrt{n} (\hat{H}^* (\hat{\beta}^* - \beta) - \hat{H} (\hat{\beta} - \beta)))}_{\xrightarrow{p} \mathbb{I}} \\ &\quad + \underbrace{((\hat{H}^*)^{-1} \hat{H} - \mathbb{I}) \sqrt{n} (\hat{\beta} - \beta)}_{\xrightarrow{p} \mathbb{O}} \\ &\xrightarrow{d} \mathcal{N}(\mathbf{0}, H^{-1} J H^{-1}), \end{aligned}$$

conditional on \mathcal{S} , where we have used that $\hat{H}^* - \hat{H} \xrightarrow{p} \mathbb{O}$ conditional on \mathcal{S} . \square

Proof of Proposition 7. First, $P(\tilde{\beta}^* = \hat{\beta}^* | \mathcal{S}) \geq P(\|\hat{H}^* - \hat{H}\| \leq \frac{c}{n^\alpha} | \mathcal{S}) \xrightarrow{p} 1$ as $n \rightarrow \infty$. Indeed, since Z has bounded conditional eighth moments, we also have that $E[\|ZZ'\|^4 | W = w, X = s]$ is uniformly bounded in X_w . It follows with Proposition A.2 in the supplementary appendix, which establishes the validity of the matched bootstrap, that

$$\sup_{r \in \mathbb{R}^{(\dim Z)^2}} |P(\sqrt{n} \text{vec}(\hat{H}^* - \hat{H}) \leq r | \mathcal{S}) - P(\mathcal{N}(\mathbf{0}, \Sigma_H) \leq r)| \xrightarrow{p} 0$$

as $n \rightarrow \infty$ and thus in particular $P(n^\alpha \|\hat{H}^* - \hat{H}\| \leq c | \mathcal{S}) \xrightarrow{p} 1$ for all $\alpha \in (0, 1/2)$, $c > 0$.

Second, since for $\tilde{A} \cap B = A \cap B$ generally

$$\begin{aligned} |P(A) - P(\tilde{A})| &\leq \underbrace{|P(A \cap B) - P(\tilde{A} \cap B)|}_{=0} \\ &\quad + \underbrace{|P(A \cap B^c) - P(\tilde{A} \cap B^c)|}_{\leq P(B^c)} \leq 1 - P(B), \end{aligned}$$

for $\Phi(r) = P(\mathcal{N}(\mathbf{0}, H^{-1} J H^{-1}) \leq r)$ we have specifically that

$$\begin{aligned} &\sup_{r \in \mathbb{R}^k} |P(\sqrt{n}(\tilde{\beta}^* - \hat{\beta}) \leq r | \mathcal{S}) - \Phi(r)| \\ &\leq \sup_{r \in \mathbb{R}^k} \left(|P(\sqrt{n}(\hat{\beta}^* - \hat{\beta}) \leq r | \mathcal{S}) - \Phi(r)| \right. \\ &\quad \left. + \underbrace{|P(\sqrt{n}(\hat{\beta}^* - \hat{\beta}) \leq r | \mathcal{S}) - P(\sqrt{n}(\tilde{\beta}^* - \hat{\beta}) \leq r | \mathcal{S})|}_{\leq 1 - P(\hat{\beta}^* = \hat{\beta}^* | \mathcal{S})} \right) \\ &\leq \underbrace{\sup_{r \in \mathbb{R}^k} |P(\sqrt{n}(\hat{\beta}^* - \hat{\beta}) \leq r | \mathcal{S}) - \Phi(r)|}_{\xrightarrow{p} 0} + \underbrace{1 - P(\hat{\beta}^* = \hat{\beta}^* | \mathcal{S})}_{\xrightarrow{p} 0} \xrightarrow{p} 0. \end{aligned}$$

This shows that this alternative bootstrap is valid in the sense of Proposition 6.

Third, for the bootstrap variance, we find

$$\begin{aligned}\widehat{\beta}^* - \widehat{\beta} &= (\widehat{H}^*)^{-1} \left(\frac{1}{n} \sum_{i \in \mathcal{S}^*} V_{ni} Z_{ni} Y_{ni} - \widehat{H}^* \widehat{\beta} \right) \\ &= (\widehat{H}^*)^{-1} \frac{1}{n} \sum_{i \in \mathcal{S}^*} V_{ni} Z_{ni} (Y_{ni} - Z'_{ni} \widehat{\beta}) \\ &= \widehat{H}^{-1} \frac{1}{n} \sum_{i \in \mathcal{S}^*} V_{ni} Z_{ni} (Y_{ni} - Z'_{ni} \widehat{\beta}) \\ &\quad \underbrace{\hspace{10em}}_{=\widehat{\Delta}^*} \\ &\quad + \underbrace{\left((\widehat{H}^*)^{-1} - \widehat{H}^{-1} \right) \frac{1}{n} \sum_{i \in \mathcal{S}^*} V_{ni} Z_{ni} (Y_{ni} - Z'_{ni} \widehat{\beta})}_{=\widehat{R}^*}.\end{aligned}$$

Since $\frac{1}{n} \sum_{i \in \mathcal{S}^*} Z_{ni} (Y_{ni} - Z'_{ni} \widehat{\beta}) = 0$ and thus $n \text{var} \left(\frac{1}{n} \sum_{i \in \mathcal{S}^*} V_{ni} Z_{ni} (Y_{ni} - Z'_{ni} \widehat{\beta}) \mid \mathcal{S} \right) = \widehat{J}$,

$$\begin{aligned}n \text{var} (\widehat{\Delta}^* \mid \mathcal{S}) &= \widehat{H}^{-1} n \text{var} \left(\frac{1}{n} \sum_{i \in \mathcal{S}^*} V_{ni} Z_{ni} (Y_{ni} - Z'_{ni} \widehat{\beta}) \mid \mathcal{S} \right) \widehat{H}^{-1} \\ &= \widehat{H}^{-1} \widehat{J} \widehat{H}^{-1} \xrightarrow{P} H^{-1} J H^{-1},\end{aligned}$$

which is a valid estimate of the asymptotic variance of $\widehat{\beta}$. However, the remainder term \widehat{R}^* generally does not have a bounded second moment since \widehat{H}^* is badly conditioned for some bootstrap draws.

To show that $\widehat{\beta}^*$ yields valid standard errors, we collect a number of preliminary results. Consider the random variables $\widehat{\Delta}^*$ and $\widetilde{\Delta}^* = \widehat{\Delta}^* \mathbf{1}_{n^\alpha \|\widehat{H}^* - \widehat{H}\| \leq c}$. $\sqrt{n} \widetilde{\Delta}^*$ converges in distribution to $\mathcal{N}(\mathbf{0}, \Sigma)$ with $\Sigma = H^{-1} J H^{-1}$, conditional on \mathcal{S} , by Proposition A.2. Since $P(\widetilde{\Delta}^* = \widehat{\Delta}^* \mid \mathcal{S}) \xrightarrow{P} 1$, the same holds true for $\sqrt{n} \widehat{\Delta}^*$ by the above argument. Also, we have established that

$$E(\sqrt{n} \widehat{\Delta}^* \mid \mathcal{S}) = 0, \quad \text{var}(\sqrt{n} \widehat{\Delta}^* \mid \mathcal{S}) \xrightarrow{P} \Sigma$$

and thus $E[n \|\widehat{\Delta}^*\|^2 \mid \mathcal{S}] \xrightarrow{P} \text{tr}(\Sigma)$. Since $E[n \|\widetilde{\Delta}^*\|^2 \mid \mathcal{S}] \leq E[n \|\widehat{\Delta}^*\|^2 \mid \mathcal{S}]$, and $n \|\widetilde{\Delta}^*\|^2$ and $n \|\widehat{\Delta}^*\|^2$ have the same weak limit (with expectation $\text{tr}(\Sigma)$) by the continuous mapping theorem, $E[n \|\widetilde{\Delta}^*\|^2 \mid \mathcal{S}] \xrightarrow{P} \text{tr}(\Sigma)$ by Proposition A.3 in the supplementary appendix. Consequently,

$$\begin{aligned}E[n \|\widehat{\Delta}^*\|^2 \mid \mathcal{S}] - E[n \|\widetilde{\Delta}^*\|^2 \mid \mathcal{S}] &= P(n^\alpha \|\widehat{H}^* - \widehat{H}\| \\ &> c \mid \mathcal{S}) E[n \|\widehat{\Delta}^*\|^2 \mid n^\alpha \|\widehat{H}^* - \widehat{H}\| > c, \mathcal{S}] \xrightarrow{P} 0.\end{aligned}\quad (\text{A.3})$$

Next, note that for conformable random variables A, B if $\text{var}(A \mid \mathcal{S}) \xrightarrow{P} \Sigma$, $E[\|B\|^2 \mid \mathcal{S}] \xrightarrow{P} 0$ then $\text{var}(A + B \mid \mathcal{S}) \xrightarrow{P} \Sigma$. Indeed,

$$\begin{aligned}|\text{var}(A + B \mid \mathcal{S}) - \text{var}(A \mid \mathcal{S})|_{jj} &= |\text{cov}(A_i, B_j \mid \mathcal{S}) \\ &\quad + \text{cov}(A_j, B_i \mid \mathcal{S}) + \text{cov}(B_i, B_j \mid \mathcal{S})| \\ &\leq \sqrt{\text{var}(A_i \mid \mathcal{S})} \sqrt{\text{var}(B_j \mid \mathcal{S})} + \sqrt{\text{var}(A_j \mid \mathcal{S})} \sqrt{\text{var}(B_i \mid \mathcal{S})} \\ &\quad + \sqrt{\text{var}(B_i \mid \mathcal{S})} \sqrt{\text{var}(B_j \mid \mathcal{S})} \xrightarrow{P} 0.\end{aligned}$$

Hence, setting $A = \sqrt{n} \widehat{\Delta}^*$ and $B = \sqrt{n} (\widehat{\beta}^* - \widehat{\beta} - \widehat{\Delta}^*)$, to establish the desired result $\text{var}(\sqrt{n} (\widehat{\beta}^* - \widehat{\beta}) \mid \mathcal{S}) \xrightarrow{P} H^{-1} J H^{-1}$ it suffices to show that

$$E \left[n \|\widehat{\beta}^* - \widehat{\beta} - \widehat{\Delta}^*\|^2 \mid \mathcal{S} \right] \xrightarrow{P} 0 \quad (\text{A.4})$$

as $n \rightarrow \infty$.

Toward establishing (A.4), note first that whenever $n^\alpha \|\widehat{H}^* - \widehat{H}\| \leq c$ then also

$$\begin{aligned}\|(\widehat{H}^*)^{-1} - \widehat{H}^{-1}\| &= \|(\widehat{H}^*)^{-1} (\widehat{H} - \widehat{H}^*) \widehat{H}^{-1}\| \\ &\leq \|(\widehat{H}^*)^{-1}\| \|\widehat{H} - \widehat{H}^*\| \|\widehat{H}^{-1}\| \\ &\leq \lambda_{\min}^{-1}(\widehat{H}^*) \lambda_{\min}^{-1}(\widehat{H}) \|\widehat{H} - \widehat{H}^*\| \dim(Z),\end{aligned}$$

where

$$\begin{aligned}\lambda_{\min}(\widehat{H}^*) &= \lambda_{\min}(\widehat{H} + \widehat{H}^* - \widehat{H}) = \min_{\|x\|=1} x' (\widehat{H} + \widehat{H}^* - \widehat{H}) x \\ &\geq \min_{\|x\|=1} x' \widehat{H} x + \min_{\|x\|=1} x' (\widehat{H}^* - \widehat{H}) x \\ &\geq \lambda_{\min}(\widehat{H}) - \|\widehat{H}^* - \widehat{H}\|\end{aligned}$$

and thus

$$\begin{aligned}\|(\widehat{H}^*)^{-1} - \widehat{H}^{-1}\| &\leq (\lambda_{\min}(\widehat{H}) - \|\widehat{H}^* - \widehat{H}\|)^{-1} \lambda_{\min}^{-1}(\widehat{H}) \|\widehat{H}^* - \widehat{H}\| \dim(Z) \\ &\leq (\lambda_{\min}(\widehat{H}) - cn^{-\alpha})^{-1} \lambda_{\min}^{-1}(\widehat{H}) cn^{-\alpha} \dim(Z).\end{aligned}\quad (\text{A.5})$$

It follows that

$$\begin{aligned}E \left[n \|\widehat{\beta}^* - \widehat{\beta} - \widehat{\Delta}^*\|^2 \mid \mathcal{S} \right] &= P(n^\alpha \|\widehat{H}^* - \widehat{H}\| \leq c \mid \mathcal{S}) E[n \|\underbrace{\widehat{\beta}^*}_{=\widehat{\beta}} - \widehat{\beta} - \widehat{\Delta}^*\|^2 \mid n^\alpha \|\widehat{H}^* - \widehat{H}\| \leq c, \mathcal{S}] \\ &\quad + P(n^\alpha \|\widehat{H}^* - \widehat{H}\| > c \mid \mathcal{S}) E[n \|\underbrace{\widehat{\beta}^*}_{=\widehat{\beta}} - \widehat{\beta} - \widehat{\Delta}^*\|^2 \mid n^\alpha \|\widehat{H}^* - \widehat{H}\| > c, \mathcal{S}] \\ &= P(n^\alpha \|\widehat{H}^* - \widehat{H}\| \leq c \mid \mathcal{S}) \\ &\quad \leq \|(\widehat{H}^*)^{-1} - \widehat{H}^{-1}\|^2 \frac{1}{n} \sum_{i \in \mathcal{S}^*} V_{ni} Z_{ni} (Y_{ni} - Z'_{ni} \widehat{\beta})^2 \\ &\quad E[n \|\underbrace{\widehat{R}^*}_{\|\widehat{R}^*\|^2} \|^2 \mid \mathcal{S}] \\ &\quad + P(n^\alpha \|\widehat{H}^* - \widehat{H}\| > c \mid \mathcal{S}) E[n \|\widehat{\Delta}^*\|^2 \mid n^\alpha \|\widehat{H}^* - \widehat{H}\| > c, \mathcal{S}] \\ (\text{A.5}) \quad &\leq \left(\underbrace{\lambda_{\min}(\widehat{H})}_{\xrightarrow{P} \lambda_{\min}(H) > 0}} - cn^{-\alpha} \right)^{-1} \lambda_{\min}^{-1}(\widehat{H}) cn^{-\alpha} \dim(Z) \\ &\quad + P(n^\alpha \|\widehat{H}^* - \widehat{H}\| \leq c \mid \mathcal{S}) E \left[\underbrace{\|n^{-1/2} \sum_{i \in \mathcal{S}^*} V_{ni} Z_{ni} (Y_{ni} - Z'_{ni} \widehat{\beta})\|^2}_{\|\widehat{R}^*\|^2} \mid n^\alpha \|\widehat{H}^* - \widehat{H}\| \leq c, \mathcal{S} \right] \\ &\quad \leq E \left[\frac{1}{n} \sum_{i \in \mathcal{S}^*} V_{ni} Z_{ni} (Y_{ni} - Z'_{ni} \widehat{\beta})^2 \mid \mathcal{S} \right] = \text{tr}(\widehat{J}) \xrightarrow{P} \text{tr}(J) \\ &\quad + \underbrace{P(n^\alpha \|\widehat{H}^* - \widehat{H}\| > c \mid \mathcal{S}) E[n \|\widehat{\Delta}^*\|^2 \mid n^\alpha \|\widehat{H}^* - \widehat{H}\| > c, \mathcal{S}]}_{\substack{(\text{A.3}) \\ \xrightarrow{P} 0}} \xrightarrow{P} 0.\end{aligned}$$

Hence, $\text{var}(\sqrt{n} (\widehat{\beta}^* - \widehat{\beta}) \mid \mathcal{S})$ and $\text{var}(\sqrt{n} \widehat{\Delta}^* \mid \mathcal{S})$ have the same probability limit $H^{-1} J H^{-1}$, which is also the asymptotic variance of $\widehat{\beta}$. \square

Supplementary Materials

The supplementary appendix contains proofs of intermediate results and extensions.

Acknowledgments

We thank Gary King, seminar participants at Harvard, and the editor (Hongyu Zhao) and referees for helpful comments, and Jaume Vives for expert research assistance.

Funding

Financial support by the NSF through grant SES 0961707 is gratefully acknowledged.

References

- Abadie, A., and Imbens, G. (2006), "Large Sample Properties of Matching Estimators for Average Treatment Effects," *Econometrica*, 74, 235–267. [1,9]
- (2008), "On the Failure of the Bootstrap for Matching Estimators," *Econometrica*, 76, 1537–1557. [7]
- (2011), "Bias-Corrected Matching Estimators for Average Treatment Effects," *Journal of Business & Economic Statistics*, 29, 1–11. [4,9]
- (2012), "A Martingale Representation for Matching Estimators," *Journal of the American Statistical Association*, 107, 833–843. [2,4,6,10]
- (2016), "Matching on the Estimated Propensity Score," *Econometrica*, 84, 781–807. [9]
- Abadie, A., Imbens, G. W., and Zheng, F. (2014), "Inference for Misspecified Models With Fixed Regressors," *Journal of the American Statistical Association*, 109, 1601–1614. [2]
- Arellano, M. (1987), "Computing Robust Standard Errors for Within-Groups Estimators," *Oxford Bulletin of Economics and Statistics*, 49, 431–434. [6]
- Blinder, A. S. (1973), "Wage Discrimination: Reduced Form and Structural Estimates," *Journal of Human Resources*, 8, 436–455. [3]
- Cochran, W. G. (1953), "Matching in Analytical Studies," *American Journal of Public Health and the Nation's Health*, 43, 684–691. [1]
- Dehejia, R. H., and Wahba, S. (1999), "Causal Effects in Nonexperimental Studies: Reevaluating the Evaluation of Training Programs," *Journal of the American Statistical Association*, 94, 1053–1062. [1]
- DiNardo, J., Fortin, N., and Lemieux, T. (1996), "Labor Market Institutions and the Distribution of Wages, 1973–1992: A Semiparametric Approach," *Econometrica*, 64, 1001–1044. [3]
- Efron, B. (1979), "Bootstrap Methods: Another Look at the Jackknife," *The Annals of Statistics*, 7, 1–26. [6]
- Eicker, F. (1967), "Limit Theorems for Regressions With Unequal and Dependent Errors," in *Proceedings of the Fifth Berkeley Symposium on Mathematical Statistics and Probability* (Vol. 1), pp. 59–82. [5]
- Ho, D. E., Imai, K., King, G., and Stuart, E. A. (2007), "Matching as Nonparametric Preprocessing for Reducing Model Dependence in Parametric Causal Inference," *Political Analysis*, 15, 199–236. [1,5,9]
- Huber, P. J. (1967), "The Behavior of Maximum Likelihood Estimates Under Nonstandard Conditions," in *Proceedings of the Fifth Berkeley Symposium on Mathematical Statistics and Probability* (Vol. 1), pp. 221–233. [5]
- Imbens, G. W., and Rubin, D. B. (2015), *Causal Inference for Statistics, Social, and Biomedical Sciences: An Introduction*, Cambridge: Cambridge University Press. [4]
- Kahn, M. (2005), "An Exhale Problem for Teaching Statistics," *The Journal of Statistical Education*, 13. [8]
- Liang, K.-Y., and Zeger, S. L. (1986), "Longitudinal Data Analysis Using Generalized Linear Models," *Biometrika*, 73, 13–22. [6]
- Oaxaca, R. (1973), "Male-Female Wage Differentials in Urban Labor Markets," *International Economic Review*, 14, 693–709. [3]
- Rosenbaum, P. R., and Rubin, D. B. (1983), "The Central Role of the Propensity Score in Observational Studies for Causal Effects," *Biometrika*, 70, 41–55. [9]
- Rosner, B. (1995), *Fundamentals of Biostatistics*, Belmont, CA: Duxbury Press. [8]
- Rubin, D. B. (1973), "Matching to Remove Bias in Observational Studies," *Biometrics*, 29, 159–183. [1]
- (1974), "Estimating Causal Effects of Treatments in Randomized and Nonrandomized Studies," *Journal of Educational Psychology*, 66, 688. [2]
- Tager, I. B., Weiss, S. T., Muñoz, A., Rosner, B., and Speizer, F. E. (1983), "Longitudinal Study of the Effects of Maternal Smoking on Pulmonary Function in Children," *New England Journal of Medicine*, 309, 699–703. [8]
- Tager, I. B., Weiss, S. T., Rosner, B., and Speizer, F. E. (1979), "Effect of Parental Cigarette Smoking on the Pulmonary Function of Children," *American Journal of Epidemiology*, 110, 15–26. [8]
- White, H. (1980a), "A Heteroskedasticity-Consistent Covariance Matrix Estimator and a Direct Test for Heteroskedasticity," *Econometrica*, 48, 817–838. [5]
- (1980b), "Using Least Squares to Approximate Unknown Regression Functions," *International Economic Review*, 21, 149–170. [5,7]
- (1982), "Maximum Likelihood Estimation of Misspecified Models," *Econometrica*, 50, 1–25. [5]