

13. Completeness of QND

1. Completeness of QND

1.1 Semantic vs. Syntactic Consistency

1.2 Proof Sketch

1.3 Stage 0: \exists -Completeness and QL'

1.4 Stage 1: Constructing Γ^*

1.5 Stage 2: Γ^* is M-QND-C & \exists -complete

1.6 Stage 3: Model Construction

The Membership Lemma

Model Construction

Induction on QL' (we be clubbin')

Stage 4? Salvation

Completeness of QND

- **QND is Complete:** For any set Γ of QL-sentences and any QL-sentence \mathcal{P} , if Γ semantically entails \mathcal{P} , then there exists a derivation of \mathcal{P} from Γ in our natural deduction system QND

Completeness of QND

- ▶ **QND is Complete:** For any set Γ of QL-sentences and any QL-sentence \mathcal{P} , if Γ semantically entails \mathcal{P} , then there exists a derivation of \mathcal{P} from Γ in our natural deduction system QND
 - In symbols: If $\Gamma \models \Theta$, then $\Gamma \vdash_{QND} \Theta$

Completeness of QND

- ▶ **QND is Complete:** For any set Γ of QL-sentences and any QL-sentence \mathcal{P} , if Γ semantically entails \mathcal{P} , then there exists a derivation of \mathcal{P} from Γ in our natural deduction system QND
 - In symbols: If $\Gamma \models \Theta$, then $\Gamma \vdash_{QND} \Theta$
 - Note that Γ can be countably infinite

Completeness of QND

- ▶ **QND is Complete:** For any set Γ of QL-sentences and any QL-sentence \mathcal{P} , if Γ semantically entails \mathcal{P} , then there exists a derivation of \mathcal{P} from Γ in our natural deduction system QND
 - In symbols: If $\Gamma \models \Theta$, then $\Gamma \vdash_{QND} \Theta$
 - Note that Γ can be countably infinite
- ▶ Completeness guarantees that for any valid QL-argument, there is at least one corresponding deduction in QND.

Completeness of QND

- ▶ **QND is Complete:** For any set Γ of QL-sentences and any QL-sentence \mathcal{P} , if Γ semantically entails \mathcal{P} , then there exists a derivation of \mathcal{P} from Γ in our natural deduction system QND
 - In symbols: If $\Gamma \models \Theta$, then $\Gamma \vdash_{QND} \Theta$
 - Note that Γ can be countably infinite
- ▶ Completeness guarantees that for any valid QL-argument, there is at least one corresponding deduction in QND.
- ▶ So we need not reason about arbitrary models to determine if a QL-argument is valid; reasoning in QND suffices! WOW COOL

“ \models ”: our Semantic Double Turnstile

- ▶ “ $\Gamma \models \mathcal{P}$ ” means that Γ logically entails \mathcal{P}
In any QL-model \mathfrak{M} where the premises in Γ are true, the conclusion \mathcal{P} is true

“ \models ”: our Semantic Double Turnstile

- ▶ “ $\Gamma \models \mathcal{P}$ ” means that Γ logically entails \mathcal{P}
In any QL-model \mathfrak{M} where the premises in Γ are true, the conclusion \mathcal{P} is true
- ▶ Equivalently: there is no QL-model \mathfrak{M} such that Γ is satisfied while \mathcal{P} is false

“ \models ”: our Semantic Double Turnstile

- ▶ “ $\Gamma \models \mathcal{P}$ ” means that Γ logically entails \mathcal{P}
In any QL-model \mathfrak{M} where the premises in Γ are true, the conclusion \mathcal{P} is true
- ▶ Equivalently: there is no QL-model \mathfrak{M} such that Γ is satisfied while \mathcal{P} is false
- ▶ Equivalently, this means that $\Gamma \cup \{\sim \mathcal{P}\}$ **is unsatisfiable**:
no QL-model makes-true the premises and negated conclusion

“ \models ”: our Semantic Double Turnstile

- ▶ “ $\Gamma \models \mathcal{P}$ ” means that Γ logically entails \mathcal{P}
In any QL-model \mathfrak{M} where the premises in Γ are true, the conclusion \mathcal{P} is true
- ▶ Equivalently: there is no QL-model \mathfrak{M} such that Γ is satisfied while \mathcal{P} is false
- ▶ Equivalently, this means that $\Gamma \cup \{\sim \mathcal{P}\}$ **is unsatisfiable**:
no QL-model makes-true the premises and negated conclusion
- ▶ We'll use this last fact A LOT in our proof that QND is complete!

13. Completeness of QND

a. Semantic vs. Syntactic Consistency

Semantic vs. Syntactic Consistency

- ▶ As with SND, we appeal to two distinct notions of consistency

Semantic vs. Syntactic Consistency

- ▶ As with SND, we appeal to two distinct notions of consistency
- ▶ One is **semantic**:

Semantic vs. Syntactic Consistency

- ▶ As with SND, we appeal to two distinct notions of consistency
- ▶ One is **semantic**:
there is a QL-model that **satisfies** every sentence in the set

Semantic vs. Syntactic Consistency

- ▶ As with SND, we appeal to two distinct notions of consistency
- ▶ One is **semantic**:
there is a QL-model that **satisfies** every sentence in the set
- ▶ We introduce a new **syntactic** notion of consistency relative to QND:

Semantic vs. Syntactic Consistency

- ▶ As with SND, we appeal to two distinct notions of consistency
- ▶ One is **semantic**:
there is a QL-model that **satisfies** every sentence in the set
- ▶ We introduce a new **syntactic** notion of consistency relative to QND:
 - a set of QL sentences is **QND-consistent** provided that you can't derive contradictory sentences from it in QND

Semantic vs. Syntactic Consistency

- ▶ As with SND, we appeal to two distinct notions of consistency
- ▶ One is **semantic**:
there is a QL-model that **satisfies** every sentence in the set
- ▶ We introduce a new **syntactic** notion of consistency relative to QND:
 - a set of QL sentences is **QND-consistent** provided that you can't derive contradictory sentences from it in QND
- ▶ Core proof idea: we'll show that if a set of sentences is **QND-consistent**, then it is also semantically consistent (i.e. **satisfiable**). So by the contrapositive: if a set is **unsatisfiable**, then it is **inconsistent-in-QND**.

Semantic: **Satisfiable** (quantificationally consistent)

- Recall: a set of QL sentences is **satisfiable** provided there is at least one QL-model \mathfrak{M} that makes all of them true

Semantic: **Satisfiable** (quantificationally consistent)

- ▶ Recall: a set of QL sentences is **satisfiable** provided there is at least one QL-model \mathfrak{M} that makes all of them true
- ▶ This is a *semantic* notion of consistency (aka “quantificational consistency”)

Semantic: **Satisfiable** (quantificationally consistent)

- ▶ Recall: a set of QL sentences is **satisfiable** provided there is at least one QL-model \mathfrak{M} that makes all of them true
- ▶ This is a *semantic* notion of consistency (aka “quantificational consistency”)
- ▶ Contrast this with the syntactic notion of **consistency in QND**:

Syntactic: (In)consistent-in-QND (derivationally consistent)

- Let Γ be a (possibly infinite) set of QL sentences

Syntactic: (In)consistent-in-QND (derivationally consistent)

- ▶ Let Γ be a (possibly infinite) set of QL sentences
- ▶ **Inconsistent-in-QND**: from premises in Γ , we can derive contradictory formulas R and $\sim R$ in the scope of the main scope line (i.e. in the scope of these premises)

Syntactic: (In)consistent-in-QND (derivationally consistent)

- ▶ Let Γ be a (possibly infinite) set of QL sentences
- ▶ **Inconsistent-in-QND**: from premises in Γ , we can derive contradictory formulas R and $\sim R$ in the scope of the main scope line (i.e. in the scope of these premises)
- ▶ **Consistent-in-QND**: Γ is not QND-inconsistent, i.e. there is no derivation from premises in Γ resulting in contradictory formulas within the main scope

Syntactic: (In)consistent-in-QND (derivationally consistent)

- ▶ Let Γ be a (possibly infinite) set of QL sentences
- ▶ **Inconsistent-in-QND**: from premises in Γ , we can derive contradictory formulas R and $\sim R$ in the scope of the main scope line (i.e. in the scope of these premises)
- ▶ **Consistent-in-QND**: Γ is not QND-inconsistent, i.e. there is no derivation from premises in Γ resulting in contradictory formulas within the main scope
- ▶ Other words we might use for these concepts: QND-inconsistent, derivationally-inconsistent, QND-consistent, etc.

Syntactic: (In)consistent-in-QND (derivationally consistent)

- ▶ Let Γ be a (possibly infinite) set of QL sentences
- ▶ **Inconsistent-in-QND**: from premises in Γ , we can derive contradictory formulas R and $\sim R$ in the scope of the main scope line (i.e. in the scope of these premises)
- ▶ **Consistent-in-QND**: Γ is not QND-inconsistent, i.e. there is no derivation from premises in Γ resulting in contradictory formulas within the main scope
- ▶ Other words we might use for these concepts: QND-inconsistent, derivationally-inconsistent, QND-consistent, etc.
- ▶ Just remember: this syntactic notion has nothing to do with models or interpretations!

13. Completeness of QND

b. Proof Sketch

Proof Sketch: Just like what we did for SL!

- Goal: prove the completeness of QL: for every QL sentence \mathcal{P} and every set Γ of QL sentences, if $\Gamma \models \mathcal{P}$ then $\Gamma \vdash_{QND} \mathcal{P}$

Proof Sketch: Just like what we did for SL!

- ▶ Goal: prove the completeness of QL: for every QL sentence \mathcal{P} and every set Γ of QL sentences, if $\Gamma \models \mathcal{P}$ then $\Gamma \vdash_{QND} \mathcal{P}$
- ▶ So assume that $\Gamma \models \mathcal{P}$.

Proof Sketch: Just like what we did for SL!

- ▶ Goal: prove the completeness of QL: for every QL sentence \mathcal{P} and every set Γ of QL sentences, if $\Gamma \models \mathcal{P}$ then $\Gamma \vdash_{QND} \mathcal{P}$
- ▶ So assume that $\Gamma \models \mathcal{P}$.
- ▶ This means that $\Gamma \cup \{\sim \mathcal{P}\}$ is **unsatisfiable**:
no QL-model satisfies the premises and negated conclusion
(i.e. $\Gamma \cup \{\sim \mathcal{P}\}$ is *semantically inconsistent*)

Proof Sketch: Just like what we did for SL!

- ▶ Goal: prove the completeness of QL: for every QL sentence \mathcal{P} and every set Γ of QL sentences, if $\Gamma \models \mathcal{P}$ then $\Gamma \vdash_{QND} \mathcal{P}$
- ▶ So assume that $\Gamma \models \mathcal{P}$.
- ▶ This means that $\Gamma \cup \{\sim \mathcal{P}\}$ is **unsatisfiable**:
no QL-model satisfies the premises and negated conclusion
(i.e. $\Gamma \cup \{\sim \mathcal{P}\}$ is *semantically inconsistent*)
- ▶ We now appeal to a **Consistency lemma** that remains the heart of the enterprise: any QND-consistent set of QL sentences is satisfiable (i.e. *semantically consistent*)

Proof Sketch: Using the consistency lemma

- ▶ **Consistency lemma** (CL): any QND-consistent set of QL sentences is satisfiable, i.e. true in some QL-model \mathfrak{M}

Proof Sketch: Using the consistency lemma

- ▶ **Consistency lemma** (CL): any QND-consistent set of QL sentences is satisfiable, i.e. true in some QL-model \mathfrak{M}
- ▶ **Contrapositive** of CL: any set of QL sentences that is Unsatisfiable is QND-Inconsistent

Proof Sketch: Using the consistency lemma

- ▶ **Consistency lemma** (CL): any QND-consistent set of QL sentences is satisfiable, i.e. true in some QL-model \mathfrak{M}
- ▶ **Contrapositive** of CL: any set of QL sentences that is **Unsatisfiable** is QND-**In**consistent
- ▶ From $\Gamma \models \mathcal{P}$ we know that $\Gamma \cup \{\sim \mathcal{P}\}$ is unsatisfiable

Proof Sketch: Using the consistency lemma

- ▶ **Consistency lemma** (CL): any QND-consistent set of QL sentences is satisfiable, i.e. true in some QL-model \mathfrak{M}
- ▶ **Contrapositive** of CL: any set of QL sentences that is **Unsatisfiable** is QND-**In**consistent
- ▶ From $\Gamma \models \mathcal{P}$ we know that $\Gamma \cup \{\sim\mathcal{P}\}$ is unsatisfiable
- ▶ So by the contrapositive of CL, we see that $\Gamma \cup \{\sim\mathcal{P}\}$ is QND-inconsistent

Proof Sketch: Using the consistency lemma

- ▶ **Consistency lemma** (CL): any QND-consistent set of QL sentences is satisfiable, i.e. true in some QL-model \mathfrak{M}
- ▶ **Contrapositive** of CL: any set of QL sentences that is **Unsatisfiable** is QND-**In**consistent
- ▶ From $\Gamma \models \mathcal{P}$ we know that $\Gamma \cup \{\sim \mathcal{P}\}$ is unsatisfiable
- ▶ So by the contrapositive of CL, we see that $\Gamma \cup \{\sim \mathcal{P}\}$ is QND-inconsistent
- ▶ This means that we can derive a pair of contradictory sentences R and $\sim R$ from $\Gamma \cup \{\sim \mathcal{P}\}$! So using the power of negation elimination, we can derive \mathcal{P} from Γ , i.e. $\Gamma \vdash_{QND} \mathcal{P}$. So we are 'done'!

Negation Elimination Refresher (book's Exercise 11.4.2)

- ▶ Claim: if $\Gamma \cup \{\sim\mathcal{P}\}$ is **QND-inconsistent**, then $\Gamma \vdash_{QND} \mathcal{P}$
- ▶ Proof: starting with (finitely-many) premises Δ from Γ , introduce $\sim\mathcal{P}$ as a subproof assumption for negation elimination
- ▶ Since $\Gamma \cup \{\sim\mathcal{P}\}$ is QND-inconsistent, we can derive a contradictory pair R and $\sim R$ within the scope of sentences in $\Delta \cup \{\sim\mathcal{P}\}$
- ▶ Then discharge this assumption $\sim\mathcal{P}$ by negation elimination, writing \mathcal{P} , now in the scope of Δ . So $\Delta \vdash_{QND} \mathcal{P}$
- ▶ Since $\Delta \subseteq \Gamma$, we have $\Gamma \vdash_{QND} \mathcal{P}$

Core subgoal: Prove consistency lemma (book's 11.4.2)

- So all we have to do is prove the **consistency lemma**: any QND-consistent set of QL sentences is satisfiable

Core subgoal: Prove consistency lemma (book's 11.4.2)

- ▶ So all we have to do is prove the **consistency lemma**: any QND-consistent set of QL sentences is satisfiable
- ▶ As with SL, we'll prove this lemma in several 'stages':

Core subgoal: Prove consistency lemma (book's 11.4.2)

- ▶ So all we have to do is prove the **consistency lemma**: any QND-consistent set of QL sentences is satisfiable
- ▶ As with SL, we'll prove this lemma in several 'stages':
- ▶ The first two are straightforward: given a QND-consistent set Γ , we construct a **superset** Γ^* that is *maximally QND-consistent* and *existentially complete* (\exists -complete)

Core subgoal: Prove consistency lemma (book's 11.4.2)

- ▶ So all we have to do is prove the **consistency lemma**: any QND-consistent set of QL sentences is satisfiable
- ▶ As with SL, we'll prove this lemma in several 'stages':
- ▶ The first two are straightforward: given a QND-consistent set Γ , we construct a **superset** Γ^* that is *maximally QND-consistent* and *existentially complete* (\exists -complete)
- ▶ In the third stage, we show that any \exists -complete, maximally QND-consistent set is **satisfiable**: we use maximal consistency and \exists -completeness to construct a model that satisfies every sentence in Γ^* . *Wrinkle*: we work in an extended language QL'!

Core subgoal: Prove consistency lemma (book's 11.4.2)

- ▶ So all we have to do is prove the **consistency lemma**: any QND-consistent set of QL sentences is satisfiable
- ▶ As with SL, we'll prove this lemma in several 'stages':
- ▶ The first two are straightforward: given a QND-consistent set Γ , we construct a **superset** Γ^* that is *maximally QND-consistent* and *existentially complete* (\exists -complete)
- ▶ In the third stage, we show that any \exists -complete, maximally QND-consistent set is **satisfiable**: we use maximal consistency and \exists -completeness to construct a model that satisfies every sentence in Γ^* . *Wrinkle*: we work in an extended language QL' !
- ▶ Since by construction $\Gamma \subseteq \Gamma^*$, this QL' -model satisfies Γ (in QL')

Core subgoal: Prove consistency lemma (book's 11.4.2)

- ▶ So all we have to do is prove the **consistency lemma**: any QND-consistent set of QL sentences is satisfiable
- ▶ As with SL, we'll prove this lemma in several 'stages':
- ▶ The first two are straightforward: given a QND-consistent set Γ , we construct a **superset** Γ^* that is *maximally QND-consistent* and *existentially complete* (\exists -complete)
- ▶ In the third stage, we show that any \exists -complete, maximally QND-consistent set is **satisfiable**: we use maximal consistency and \exists -completeness to construct a model that satisfies every sentence in Γ^* . *Wrinkle*: we work in an extended language QL' !
- ▶ Since by construction $\Gamma \subseteq \Gamma^*$, this QL' -model satisfies Γ (in QL')
- ▶ From our QL' -model, we generate a QL-model that satisfies Γ

13. Completeness of QND

c. Stage 0: \exists -Completeness and QL'

Maximally QND-consistent (no longer enough!)

- ▶ A set Γ^* of QL or QL' sentences is **maximally QND-consistent** provided that:

Maximally QND-consistent (no longer enough!)

- ▶ A set Γ^* of QL or QL' sentences is **maximally QND-consistent** provided that:
 - 1.) Γ^* is QND-consistent (i.e. can't derive contradictory sentences)

Maximally QND-consistent (no longer enough!)

- ▶ A set Γ^* of QL or QL' sentences is **maximally QND-consistent** provided that:
 - 1.) Γ^* is QND-consistent (i.e. can't derive contradictory sentences)
 - 2.) adding **any** additional sentence to Γ^* would result in an QND-**inconsistent** set

Maximally QND-consistent (no longer enough!)

- ▶ A set Γ^* of QL or QL' sentences is **maximally QND-consistent** provided that:
 - 1.) Γ^* is QND-consistent (i.e. can't derive contradictory sentences)
 - 2.) adding **any** additional sentence to Γ^* would result in an QND-**inconsistent** set
- ▶ i.e. for any $P \notin \Gamma^*$, $\{P\} \cup \Gamma^*$ is QND-**inconsistent**

Maximally QND-consistent (no longer enough!)

- ▶ A set Γ^* of QL or QL' sentences is **maximally QND-consistent** provided that:
 - 1.) Γ^* is QND-consistent (i.e. can't derive contradictory sentences)
 - 2.) adding **any** additional sentence to Γ^* would result in an QND-**inconsistent** set
- ▶ i.e. for any $P \notin \Gamma^*$, $\{P\} \cup \Gamma^*$ is QND-**inconsistent**
- ▶ Unlike with SL, maximal derivational consistency is no longer enough to ensure satisfiability

Maximally QND-consistent (no longer enough!)

- ▶ A set Γ^* of QL or QL' sentences is **maximally QND-consistent** provided that:
 - 1.) Γ^* is QND-consistent (i.e. can't derive contradictory sentences)
 - 2.) adding **any** additional sentence to Γ^* would result in an QND-**inconsistent** set
- ▶ i.e. for any $P \notin \Gamma^*$, $\{P\} \cup \Gamma^*$ is QND-**inconsistent**
- ▶ Unlike with SL, maximal derivational consistency is no longer enough to ensure satisfiability
- ▶ Recall that our purely syntactic membership lemma is motivated by the truth-conditions for QL sentences: sentences belong to Γ^* iff the relevant “truth-condition pieces” belong to Γ^* as well

Maximally QND-consistent (no longer enough!)

- ▶ A set Γ^* of QL or QL' sentences is **maximally QND-consistent** provided that:
 - 1.) Γ^* is QND-consistent (i.e. can't derive contradictory sentences)
 - 2.) adding **any** additional sentence to Γ^* would result in an QND-**inconsistent** set
- ▶ i.e. for any $P \notin \Gamma^*$, $\{P\} \cup \Gamma^*$ is QND-**inconsistent**
- ▶ Unlike with SL, maximal derivational consistency is no longer enough to ensure satisfiability
- ▶ Recall that our purely syntactic membership lemma is motivated by the truth-conditions for QL sentences: sentences belong to Γ^* iff the relevant “truth-condition pieces” belong to Γ^* as well
- ▶ To extend our membership lemma to quantified sentences, we require that every existential sentence in Γ^* has a substitution instance also in Γ^* . So we introduce a new property:

Existential-completeness: definition and motivation

- ▶ **\exists -completeness**: a set Γ of QL or QL' sentences is *existentially-complete* just in case for every sentence in Γ of the form $(\exists\chi)\mathcal{P}$, at least one substitution instance $\mathcal{P}[c/\chi]$ is in Γ

Existential-completeness: definition and motivation

- ▶ **\exists -completeness**: a set Γ of QL or QL' sentences is *existentially-complete* just in case for every sentence in Γ of the form $(\exists\chi)\mathcal{P}$, at least one substitution instance $\mathcal{P}[c/\chi]$ is in Γ
- ▶ Motivation: $(\exists\chi)\mathcal{P}$ is true in a model iff some object $r \in D$ is a \mathcal{P}

Existential-completeness: definition and motivation

- ▶ **\exists -completeness**: a set Γ of QL or QL' sentences is *existentially-complete* just in case for every sentence in Γ of the form $(\exists\chi)\mathcal{P}$, at least one substitution instance $\mathcal{P}[c/\chi]$ is in Γ
- ▶ Motivation: $(\exists\chi)\mathcal{P}$ is true in a model iff some object $r \in D$ is a \mathcal{P}
- ▶ To construct an \exists -complete set Γ^* , we need recourse to a countable infinity of unused constants.

Existential-completeness: definition and motivation

- ▶ **\exists -completeness**: a set Γ of QL or QL' sentences is *existentially-complete* just in case for every sentence in Γ of the form $(\exists\chi)\mathcal{P}$, at least one substitution instance $\mathcal{P}[c/\chi]$ is in Γ
- ▶ Motivation: $(\exists\chi)\mathcal{P}$ is true in a model iff some object $r \in D$ is a \mathcal{P}
- ▶ To construct an \exists -complete set Γ^* , we need recourse to a countable infinity of unused constants.
- ▶ Otherwise, new substitution instances that we add could “contradict” sentences already in Γ , spoiling QND-consistency

Existential-completeness: definition and motivation

- ▶ **\exists -completeness**: a set Γ of QL or QL' sentences is *existentially-complete* just in case for every sentence in Γ of the form $(\exists\chi)\mathcal{P}$, at least one substitution instance $\mathcal{P}[c/\chi]$ is in Γ
- ▶ Motivation: $(\exists\chi)\mathcal{P}$ is true in a model iff some object $r \in D$ is a \mathcal{P}
- ▶ To construct an \exists -complete set Γ^* , we need recourse to a countable infinity of unused constants.
- ▶ Otherwise, new substitution instances that we add could “contradict” sentences already in Γ , spoiling QND-consistency
- ▶ *Problem*: our starting Γ might be infinite and so already use infinitely-many constants from QL. What are we to do?

It's a bird! It's a plane! It's ... Language QL'???

- ▶ QL' is exactly like QL except that we allow subscripted *constants* to have **primed-indices**

It's a bird! It's a plane! It's ... Language QL'???

- ▶ QL' is exactly like QL except that we allow subscripted *constants* to have **primed-indices**
- ▶ e.g. $c_{11'}$, $b_{234'}$, $g_{2'}$ (''-symbol always at the end)

It's a bird! It's a plane! It's ... Language QL'???

- ▶ QL' is exactly like QL except that we allow subscripted *constants* to have **primed-indices**
- ▶ e.g. $c_{11'}$, $b_{234'}$, $g_{2'}$ (''-symbol always at the end)
- ▶ Unsubscripted constants remain the same: a thru v

It's a bird! It's a plane! It's ... Language QL'???

- ▶ QL' is exactly like QL except that we allow subscripted *constants* to have **primed-indices**
- ▶ e.g. $c_{11'}$, $b_{234'}$, $g_{2'}$ (''-symbol always at the end)
- ▶ Unsubscripted constants remain the same: a thru v
- ▶ So QL' just adds one new symbol ' ', allowed to occur only at the end of indices for constants

It's a bird! It's a plane! It's ... Language QL'???

- ▶ QL' is exactly like QL except that we allow subscripted *constants* to have **primed-indices**
- ▶ e.g. $c_{11'}$, $b_{234'}$, $g_{2'}$ ('-'symbol always at the end)
- ▶ Unsubscripted constants remain the same: a thru v
- ▶ So QL' just adds one new symbol ' $'$ ', allowed to occur only at the end of indices for constants
- ▶ The recursive structure of truth-in-QL' is defined exactly the same as for QL (using our good friend, satisfaction semantics!)

It's a bird! It's a plane! It's ... Language QL'???

- ▶ QL' is exactly like QL except that we allow subscripted *constants* to have **primed-indices**
- ▶ e.g. $c_{11'}$, $b_{234'}$, $g_{2'}$ ('-'symbol always at the end)
- ▶ Unsubscripted constants remain the same: a thru v
- ▶ So QL' just adds one new symbol ' $'$ ', allowed to occur only at the end of indices for constants
- ▶ The recursive structure of truth-in-QL' is defined exactly the same as for QL (using our good friend, satisfaction semantics!)
- ▶ Note that we do NOT allow primed indices on Predicates

It's a bird! It's a plane! It's ... Language QL'???

- ▶ QL' is exactly like QL except that we allow subscripted *constants* to have **primed-indices**
- ▶ e.g. $c_{11'}$, $b_{234'}$, $g_{2'}$ ('-'symbol always at the end)
- ▶ Unsubscripted constants remain the same: a thru v
- ▶ So QL' just adds one new symbol ' $'$ ', allowed to occur only at the end of indices for constants
- ▶ The recursive structure of truth-in-QL' is defined exactly the same as for QL (using our good friend, satisfaction semantics!)
- ▶ Note that we do NOT allow primed indices on Predicates
- ▶ Moral: reach *for the stars*, **not** drugs

13. Completeness of QND

d. Stage 1: Constructing Γ^*

Stage 1(i): first enumerate the sentences of QL' !

- Let Γ be a QND-consistent set of QL sentences (possibly infinite)

Stage 1(i): first enumerate the sentences of QL' !

- ▶ Let Γ be a QND-consistent set of QL sentences (possibly infinite)
- ▶ To construct Γ^* , we first **enumerate** the QL' sentences, so that every QL' sentence is associated with a unique positive integer $\{1, 2, 3, \dots\}$

Stage 1(i): first enumerate the sentences of QL'

- ▶ Let Γ be a QND-consistent set of QL sentences (possibly infinite)
- ▶ To construct Γ^* , we first **enumerate** the QL' sentences, so that every QL' sentence is associated with a unique positive integer $\{1, 2, 3, \dots\}$
- ▶ As with SL, stipulate an 'alphabetical order' for QL' symbols

Stage 1(i): first enumerate the sentences of QL'

- ▶ Let Γ be a QND-consistent set of QL sentences (possibly infinite)
- ▶ To construct Γ^* , we first **enumerate** the QL' sentences, so that every QL' sentence is associated with a unique positive integer $\{1, 2, 3, \dots\}$
- ▶ As with SL, stipulate an 'alphabetical order' for QL' symbols
- ▶ $\sim, \vee, \&, \supset, \equiv, (,), 0, 1, \dots, 9, A, B, \dots, Z, a, \dots, v, w, x, y, z, \forall, \exists, '$

Stage 1(i): first enumerate the sentences of QL'!

- ▶ Let Γ be a QND-consistent set of QL sentences (possibly infinite)
- ▶ To construct Γ^* , we first **enumerate** the QL' sentences, so that every QL' sentence is associated with a unique positive integer $\{1, 2, 3, \dots\}$
- ▶ As with SL, stipulate an 'alphabetical order' for QL' symbols
- ▶ $\sim, \vee, \&, \supset, \equiv, (,), 0, 1, \dots, 9, A, B, \dots, Z, a, \dots, v, w, x, y, z, \forall, \exists, '$
- ▶ Assign each symbol an **index** between '10' and '84' (skip 17-19)

Stage 1(i): first enumerate the sentences of QL'

- ▶ Let Γ be a QND-consistent set of QL sentences (possibly infinite)
- ▶ To construct Γ^* , we first **enumerate** the QL' sentences, so that every QL' sentence is associated with a unique positive integer $\{1, 2, 3, \dots\}$
- ▶ As with SL, stipulate an 'alphabetical order' for QL' symbols
- ▶ $\sim, \vee, \&, \supset, \equiv, (,), 0, 1, \dots, 9, A, B, \dots, Z, a, \dots, v, w, x, y, z, \forall, \exists, '$
- ▶ Assign each symbol an **index** between '10' and '84' (skip 17-19)
- ▶ Then each QL' sentence corresponds to a unique positive integer, constructed by replacing each symbol in the sentence with its index, from left to right.

Stage 1(i): first enumerate the sentences of QL'

- ▶ Let Γ be a QND-consistent set of QL sentences (possibly infinite)
- ▶ To construct Γ^* , we first **enumerate** the QL' sentences, so that every QL' sentence is associated with a unique positive integer $\{1, 2, 3, \dots\}$
- ▶ As with SL, stipulate an 'alphabetical order' for QL' symbols
- ▶ $\sim, \vee, \&, \supset, \equiv, (,), 0, 1, \dots, 9, A, B, \dots, Z, a, \dots, v, w, x, y, z, \forall, \exists, '$
- ▶ Assign each symbol an **index** between '10' and '84' (skip 17-19)
- ▶ Then each QL' sentence corresponds to a unique positive integer, constructed by replacing each symbol in the sentence with its index, from left to right.
- ▶ So with our ordering, 'A' is the first sentence; 'B' the second ... up to Z, and then we hit $\sim A$ ($\mapsto 1030$), then $\sim B$ ($\mapsto 1031$), etc.

Recall what we did in SL to form Γ^* Max.-SND-Consist.

- We considered the first sentence 'A' in our enumeration.
If A could be added to Γ without the resulting set being SND-inconsistent, then we let $\Gamma_1 := \Gamma \cup \{A\}$.

Recall what we did in SL to form Γ^* Max.-SND-Consist.

- ▶ We considered the first sentence 'A' in our enumeration.
If A could be added to Γ without the resulting set being SND-inconsistent, then we let $\Gamma_1 := \Gamma \cup \{A\}$.
- ▶ Otherwise, let $\Gamma_1 := \Gamma$ (so that Γ_1 stays SND-consistent)

Recall what we did in SL to form Γ^* Max.-SND-Consist.

- ▶ We considered the first sentence 'A' in our enumeration.
If A could be added to Γ without the resulting set being SND-inconsistent, then we let $\Gamma_1 := \Gamma \cup \{A\}$.
- ▶ Otherwise, let $\Gamma_1 := \Gamma$ (so that Γ_1 stays SND-consistent)
- ▶ We proceeded to the 2nd sentence in our enumeration.
If it could be added to Γ_1 without the new set being SND-inconsistent, let Γ_2 be the result. Otherwise, let $\Gamma_2 := \Gamma_1$

Recall what we did in SL to form Γ^* Max.-SND-Consist.

- ▶ We considered the first sentence 'A' in our enumeration.
If A could be added to Γ without the resulting set being SND-inconsistent, then we let $\Gamma_1 := \Gamma \cup \{A\}$.
- ▶ Otherwise, let $\Gamma_1 := \Gamma$ (so that Γ_1 stays SND-consistent)
- ▶ We proceeded to the 2nd sentence in our enumeration.
If it could be added to Γ_1 without the new set being SND-inconsistent, let Γ_2 be the result. Otherwise, let $\Gamma_2 := \Gamma_1$
- ▶ Γ^* was the result of 'doing' this procedure for every SL sentence

Recall what we did in SL to form Γ^* Max.-SND-Consist.

- ▶ We considered the first sentence 'A' in our enumeration.
If A could be added to Γ without the resulting set being SND-inconsistent, then we let $\Gamma_1 := \Gamma \cup \{A\}$.
- ▶ Otherwise, let $\Gamma_1 := \Gamma$ (so that Γ_1 stays SND-consistent)
- ▶ We proceeded to the 2nd sentence in our enumeration.
If it could be added to Γ_1 without the new set being SND-inconsistent, let Γ_2 be the result. Otherwise, let $\Gamma_2 := \Gamma_1$
- ▶ Γ^* was the result of 'doing' this procedure for every SL sentence
- ▶ Now we need to complicate matters a bit, to handle sentences of the form $(\exists \chi)\mathcal{P}$ and ensure we add a suitable substitution instance whenever we can add $(\exists \chi)\mathcal{P}$ while preserving QND-consistency

Building up Γ^*

- Given a QND-consistent set of QL sentences Γ , let $\Gamma_0 := \Gamma$

Building up Γ^*

- ▶ Given a QND-consistent set of QL sentences Γ , let $\Gamma_0 := \Gamma$
- ▶ Consider the k -th sentence P_k in our enumeration of QL'

Building up Γ^*

- ▶ Given a QND-consistent set of QL sentences Γ , let $\Gamma_0 := \Gamma$
- ▶ Consider the k -th sentence P_k in our enumeration of QL'
- ▶ Define Γ_{k+1} as follows:

Building up Γ^*

- ▶ Given a QND-consistent set of QL sentences Γ , let $\Gamma_0 := \Gamma$
- ▶ Consider the k -th sentence P_k in our enumeration of QL'
- ▶ Define Γ_{k+1} as follows:
 - i.) Γ_k if the set $\Gamma_k \cup \{P_k\}$ is QND-**INconsistent**

Building up Γ^*

- ▶ Given a QND-consistent set of QL sentences Γ , let $\Gamma_0 := \Gamma$
- ▶ Consider the k -th sentence P_k in our enumeration of QL'
- ▶ Define Γ_{k+1} as follows:
 - i.) Γ_k if the set $\Gamma_k \cup \{P_k\}$ is QND-**INconsistent**
 - ii.) $\Gamma_k \cup \{P_k\}$ if P_k does NOT have the form $(\exists x)Q$, and $\Gamma_k \cup \{P_k\}$ is QND-consistent

Building up Γ^*

- ▶ Given a QND-consistent set of QL sentences Γ , let $\Gamma_0 := \Gamma$
- ▶ Consider the k -th sentence P_k in our enumeration of QL'
- ▶ Define Γ_{k+1} as follows:
 - i.) Γ_k if the set $\Gamma_k \cup \{P_k\}$ is QND-**INconsistent**
 - ii.) $\Gamma_k \cup \{P_k\}$ if P_k does NOT have the form $(\exists \chi)Q$, and $\Gamma_k \cup \{P_k\}$ is QND-consistent
 - iii.) $\Gamma_k \cup \{P_k, P_k^\dagger\}$ if $\Gamma_k \cup \{P_k\}$ is QND-consistent AND P_k DOES have the form $(\exists \chi)Q$, where P_k^\dagger is a substitution instance $Q[c/\chi]$, and c is the **alphabetically earliest constant** not in P_k or any sentence in Γ_k

Building up Γ^*

- ▶ Given a QND-consistent set of QL sentences Γ , let $\Gamma_0 := \Gamma$
- ▶ Consider the k -th sentence P_k in our enumeration of QL'
- ▶ Define Γ_{k+1} as follows:
 - i.) Γ_k if the set $\Gamma_k \cup \{P_k\}$ is QND-**INconsistent**
 - ii.) $\Gamma_k \cup \{P_k\}$ if P_k does NOT have the form $(\exists \chi)Q$, and $\Gamma_k \cup \{P_k\}$ is QND-consistent
 - iii.) $\Gamma_k \cup \{P_k, P_k^\dagger\}$ if $\Gamma_k \cup \{P_k\}$ is QND-consistent AND P_k DOES have the form $(\exists \chi)Q$, where P_k^\dagger is a substitution instance $Q[c/\chi]$, and c is the **alphabetically earliest constant** not in P_k or any sentence in Γ_k
 - Such a c is guaranteed to exist because Γ_0 belongs to QL .

Building up Γ^*

- ▶ Given a QND-consistent set of QL sentences Γ , let $\Gamma_0 := \Gamma$
- ▶ Consider the k -th sentence P_k in our enumeration of QL'
- ▶ Define Γ_{k+1} as follows:
 - i.) Γ_k if the set $\Gamma_k \cup \{P_k\}$ is QND-**INconsistent**
 - ii.) $\Gamma_k \cup \{P_k\}$ if P_k does NOT have the form $(\exists \chi)Q$, and $\Gamma_k \cup \{P_k\}$ is QND-consistent
 - iii.) $\Gamma_k \cup \{P_k, P_k^\dagger\}$ if $\Gamma_k \cup \{P_k\}$ is QND-consistent AND P_k DOES have the form $(\exists \chi)Q$, where P_k^\dagger is a substitution instance $Q[c/\chi]$, and c is the **alphabetically earliest constant** not in P_k or any sentence in Γ_k
 - Such a c is guaranteed to exist because Γ_0 belongs to QL .
 - So the countable-infinity of primed subscripted constants from QL' are available at each stage if needed.

Building up Γ^*

- ▶ Given a QND-consistent set of QL sentences Γ , let $\Gamma_0 := \Gamma$
- ▶ Consider the k -th sentence P_k in our enumeration of QL'
- ▶ Define Γ_{k+1} as follows:
 - i.) Γ_k if the set $\Gamma_k \cup \{P_k\}$ is QND-**Inconsistent**
 - ii.) $\Gamma_k \cup \{P_k\}$ if P_k does NOT have the form $(\exists \chi)Q$, and $\Gamma_k \cup \{P_k\}$ is QND-consistent
 - iii.) $\Gamma_k \cup \{P_k, P_k^\dagger\}$ if $\Gamma_k \cup \{P_k\}$ is QND-consistent AND P_k DOES have the form $(\exists \chi)Q$, where P_k^\dagger is a substitution instance $Q[c/\chi]$, and c is the **alphabetically earliest constant** not in P_k or any sentence in Γ_k
 - Such a c is guaranteed to exist because Γ_0 belongs to QL .
 - So the countable-infinity of primed subscripted constants from QL' are available at each stage if needed.
- ▶ Then $\Gamma^* := \bigcup_{k=0}^{\infty} \Gamma_k$

13. Completeness of QND

e. Stage 2: Γ^* is M-QND-C &
 \exists -complete

Stage 2: Γ^* is maximally QND-consistent & \exists -complete

- This requires proving three claims (from the definitions):

Stage 2: Γ^* is maximally QND-consistent & \exists -complete

► This requires proving three claims (from the definitions):

1.) Γ^* is consistent in QND

Stage 2: Γ^* is maximally QND-consistent & \exists -complete

- This requires proving three claims (from the definitions):
 - 1.) Γ^* is consistent in QND
 - 2.) Adding any additional sentence to Γ^* would result in a **QND-inconsistent** set

Stage 2: Γ^* is maximally QND-consistent & \exists -complete

- This requires proving three claims (from the definitions):
 - 1.) Γ^* is consistent in QND
 - 2.) Adding any additional sentence to Γ^* would result in a **QND-inconsistent** set
 - 3.) For every QL' sentence of the form $(\exists \chi)Q$ in Γ^* , at least one substitution instance $Q[c/\chi]$ belongs to Γ^*

Stage 2: Γ^* is maximally QND-consistent & \exists -complete

- ▶ This requires proving three claims (from the definitions):
 - 1.) Γ^* is consistent in QND
 - 2.) Adding any additional sentence to Γ^* would result in a **QND-inconsistent** set
 - 3.) For every QL' sentence of the form $(\exists \chi)Q$ in Γ^* , at least one substitution instance $Q[c/\chi]$ belongs to Γ^*
- ▶ We prove these in turn

Stage 2 (i): Γ^* is QND-consistent

- ▶ Assume for *reductio* that Γ^* is inconsistent in QND

Stage 2 (i): Γ^* is QND-consistent

- ▶ Assume for *reductio* that Γ^* is inconsistent in QND
- ▶ Then there would be a QND derivation with finite premise set $\Delta \subset \Gamma^*$ that derives a contradictory pair R and $\sim R$

Stage 2 (i): Γ^* is QND-consistent

- ▶ Assume for *reductio* that Γ^* is inconsistent in QND
- ▶ Then there would be a QND derivation with finite premise set $\Delta \subset \Gamma^*$ that derives a contradictory pair R and $\sim R$
- ▶ Since Δ is finite, there exists some $k + 1 \in \mathbb{N}$ s.t. $\Delta \subset \Gamma_{k+1}$. So then this Γ_{k+1} would be **QND-inconsistent**.

Stage 2 (i): Γ^* is QND-consistent

- ▶ Assume for *reductio* that Γ^* is inconsistent in QND
- ▶ Then there would be a QND derivation with finite premise set $\Delta \subset \Gamma^*$ that derives a contradictory pair R and $\sim R$
- ▶ Since Δ is finite, there exists some $k + 1 \in \mathbb{N}$ s.t. $\Delta \subset \Gamma_{k+1}$. So then this Γ_{k+1} would be **QND-inconsistent**.
- ▶ Yet, each Γ_{k+1} is necessarily **QND-consistent**:

Stage 2 (i): Γ^* is QND-consistent

- ▶ Assume for *reductio* that Γ^* is inconsistent in QND
- ▶ Then there would be a QND derivation with finite premise set $\Delta \subset \Gamma^*$ that derives a contradictory pair R and $\sim R$
- ▶ Since Δ is finite, there exists some $k + 1 \in \mathbb{N}$ s.t. $\Delta \subset \Gamma_{k+1}$. So then this Γ_{k+1} would be **QND-inconsistent**.
- ▶ Yet, each Γ_{k+1} is necessarily **QND-consistent**:
 - If P_k is not existential, it joins Γ_{k+1} only if $\Gamma_k \cup \{P_k\}$ is QND-consistent—by condition (ii)

Stage 2 (i): Γ^* is QND-consistent

- ▶ Assume for *reductio* that Γ^* is inconsistent in QND
- ▶ Then there would be a QND derivation with finite premise set $\Delta \subset \Gamma^*$ that derives a contradictory pair R and $\sim R$
- ▶ Since Δ is finite, there exists some $k + 1 \in \mathbb{N}$ s.t. $\Delta \subset \Gamma_{k+1}$. So then this Γ_{k+1} would be **QND-inconsistent**.
- ▶ Yet, each Γ_{k+1} is necessarily **QND-consistent**:
 - If P_k is not existential, it joins Γ_{k+1} only if $\Gamma_k \cup \{P_k\}$ is QND-consistent—by condition (ii)
 - If P_k is of the form $(\exists x)Q$, it joins only if $\Gamma_k \cup \{P_k\}$ is QND-consistent.

Stage 2 (i): Γ^* is QND-consistent

- ▶ Assume for *reductio* that Γ^* is inconsistent in QND
- ▶ Then there would be a QND derivation with finite premise set $\Delta \subset \Gamma^*$ that derives a contradictory pair R and $\sim R$
- ▶ Since Δ is finite, there exists some $k + 1 \in \mathbb{N}$ s.t. $\Delta \subset \Gamma_{k+1}$. So then this Γ_{k+1} would be **QND-inconsistent**.
- ▶ Yet, each Γ_{k+1} is necessarily **QND-consistent**:
 - If P_k is not existential, it joins Γ_{k+1} only if $\Gamma_k \cup \{P_k\}$ is QND-consistent—by condition (ii)
 - If P_k is of the form $(\exists x)Q$, it joins only if $\Gamma_k \cup \{P_k\}$ is QND-consistent.
 - It remains to show that $\Gamma_k \cup \{(\exists x)Q, Q[c/x]\}$ is QND-consistent

Stage 2 (i): Γ^* is QND-consistent

- ▶ Assume for *reductio* that Γ^* is inconsistent in QND
- ▶ Then there would be a QND derivation with finite premise set $\Delta \subset \Gamma^*$ that derives a contradictory pair R and $\sim R$
- ▶ Since Δ is finite, there exists some $k + 1 \in \mathbb{N}$ s.t. $\Delta \subset \Gamma_{k+1}$. So then this Γ_{k+1} would be **QND-inconsistent**.
- ▶ Yet, each Γ_{k+1} is necessarily **QND-consistent**:
 - If P_k is not existential, it joins Γ_{k+1} only if $\Gamma_k \cup \{P_k\}$ is QND-consistent—by condition (ii)
 - If P_k is of the form $(\exists x)Q$, it joins only if $\Gamma_k \cup \{P_k\}$ is QND-consistent.
 - It remains to show that $\Gamma_k \cup \{(\exists x)Q, Q[c/x]\}$ is QND-consistent
 - **Lemma**: if c does not occur in a QND-C set $\Gamma_k \cup \{(\exists x)Q\}$, then $\Gamma_k \cup \{(\exists x)Q, Q[c/x]\}$ is QND-consistent

Stage 2 (i): Γ^* is QND-consistent

- ▶ Assume for *reductio* that Γ^* is inconsistent in QND
- ▶ Then there would be a QND derivation with finite premise set $\Delta \subset \Gamma^*$ that derives a contradictory pair R and $\sim R$
- ▶ Since Δ is finite, there exists some $k + 1 \in \mathbb{N}$ s.t. $\Delta \subset \Gamma_{k+1}$. So then this Γ_{k+1} would be **QND-inconsistent**.
- ▶ Yet, each Γ_{k+1} is necessarily **QND-consistent**:
 - If P_k is not existential, it joins Γ_{k+1} only if $\Gamma_k \cup \{P_k\}$ is QND-consistent—by condition (ii)
 - If P_k is of the form $(\exists x)Q$, it joins only if $\Gamma_k \cup \{P_k\}$ is QND-consistent.
 - It remains to show that $\Gamma_k \cup \{(\exists x)Q, Q[c/x]\}$ is QND-consistent
 - **Lemma**: if c does not occur in a QND-C set $\Gamma_k \cup \{(\exists x)Q\}$, then $\Gamma_k \cup \{(\exists x)Q, Q[c/x]\}$ is QND-consistent
- ▶ Hence, Γ^* must be QND-consistent, on pain of *reductio*

Stage 2 (ii): Γ^* is **maximally** QND-consistent

- ▶ Assume for *reductio* that Γ^* weren't maximally QND-consistent, despite being QND-consistent
- ▶ i.e. assume *it is not the case that* for all other sentences, adding it to Γ^* would result in a QND-inconsistent set
 - \Rightarrow there exists a sentence \mathcal{Q} that we could add to Γ^* while preserving QND-consistency (i.e. there is some sentence we neglected that could make Γ^* a 'bigger' QND-consistent set)
- ▶ Yet, \mathcal{Q} would appear in our enumeration as some sentence P_k , 'considered' at the k -th stage of our construction of Γ^* .
- ▶ So if \mathcal{Q} isn't in Γ^* , then this is because adding it 'would have' made $\Gamma_k \subset \Gamma^*$ QND-inconsistent.
 - So $\{\mathcal{Q}\} \cup \Gamma^*$ must be QND-inconsistent (*reductio*!)
- ▶ So we can't add any \mathcal{Q} to Γ^* while preserving QND-consistency

Stage 2 (iii): Γ^* is \exists -complete

- We simply need to show that for each sentence of the form $(\exists \chi)Q \in \Gamma^*$, a substitution instance $Q[c/\chi]$ also belongs to Γ^*

Stage 2 (iii): Γ^* is \exists -complete

- ▶ We simply need to show that for each sentence of the form $(\exists \chi)Q \in \Gamma^*$, a substitution instance $Q[c/\chi]$ also belongs to Γ^*
- ▶ Note that this is true by construction: each sentence of the form $(\exists \chi)Q$ occurs in our QL' -enumeration:

Stage 2 (iii): Γ^* is \exists -complete

- ▶ We simply need to show that for each sentence of the form $(\exists \chi)Q \in \Gamma^*$, a substitution instance $Q[c/\chi]$ also belongs to Γ^*
- ▶ Note that this is true by construction: each sentence of the form $(\exists \chi)Q$ occurs in our QL' -enumeration:
- ▶ If we could have “added” $(\exists \chi)Q$ at the k -th stage while preserving QND-consistency, then we also added a substitution instance.

Stage 2 (iii): Γ^* is \exists -complete

- ▶ We simply need to show that for each sentence of the form $(\exists \chi)Q \in \Gamma^*$, a substitution instance $Q[c/\chi]$ also belongs to Γ^*
- ▶ Note that this is true by construction: each sentence of the form $(\exists \chi)Q$ occurs in our QL'-enumeration:
- ▶ If we could have “added” $(\exists \chi)Q$ at the k -th stage while preserving QND-consistency, then we also added a substitution instance.
- ▶ This is so even if $(\exists \chi)Q$ is already in $\Gamma_\emptyset := \Gamma$, since by condition (iii) $\Gamma_{k+1} := \Gamma_k \cup \{(\exists \chi)Q, Q[c/\chi]\}$ which in this case would equal $\Gamma_k \cup \{Q[c/\chi]\}$ (since in this case, $(\exists \chi)Q \in \Gamma_k$)

13. Completeness of QND

f. Stage 3: Model Construction

Stage 3: The Maximal Consistency Lemma (\approx book's 11.4.7)

- \exists -C Maximal Consistency Lemma: every QL' set that is maximally-QND-consistent and \exists -complete is satisfiable

Stage 3: The Maximal Consistency Lemma (\approx book's 11.4.7)

- ▶ \exists -C Maximal Consistency Lemma: every QL' set that is maximally-QND-consistent and \exists -complete is satisfiable
- ▶ (there exists a QL' -model that makes-true every sentence in Γ^*)
We construct this model, calling it " \mathfrak{M}^* " (\approx book's " \mathbf{I}^* ")

Stage 3: The Maximal Consistency Lemma (\approx book's 11.4.7)

- ▶ \exists -C Maximal Consistency Lemma: every QL' set that is maximally-QND-consistent and \exists -complete is satisfiable
- ▶ (there exists a QL'-model that makes-true every sentence in Γ^*)
We construct this model, calling it " \mathfrak{M}^* " (\approx book's " \mathbf{I}^* ")
- ▶ Proof idea: since Γ^* is M-QND-C, for any sentence \mathcal{P} , either $\mathcal{P} \in \Gamma^*$ or $\sim \mathcal{P} \in \Gamma^*$ (you're either in the club or your '**nemesis**' is!)

Stage 3: The Maximal Consistency Lemma (\approx book's 11.4.7)

- ▶ \exists -C Maximal Consistency Lemma: every QL' set that is maximally-QND-consistent and \exists -complete is satisfiable
- ▶ (there exists a QL'-model that makes-true every sentence in Γ^*)
We construct this model, calling it " \mathfrak{M}^* " (\approx book's " \mathbf{I}^* ")
- ▶ Proof idea: since Γ^* is M-QND-C, for any sentence \mathcal{P} , either $\mathcal{P} \in \Gamma^*$ or $\sim \mathcal{P} \in \Gamma^*$ (you're either in the club or your '**nemesis**' is!)
This holds in particular for each QL'-atomic sentence

Stage 3: The Maximal Consistency Lemma (\approx book's 11.4.7)

- ▶ **\exists -C Maximal Consistency Lemma**: every QL' set that is maximally-QND-consistent and \exists -complete is satisfiable
- ▶ (there exists a QL' -model that makes-true every sentence in Γ^*)
We construct this model, calling it " \mathfrak{M}^* " (\approx book's " \mathbf{I}^* ")
- ▶ Proof idea: since Γ^* is M-QND-C, for any sentence \mathcal{P} , either $\mathcal{P} \in \Gamma^*$ or $\sim \mathcal{P} \in \Gamma^*$ (you're either in the club or your '**nemesis**' is!)
This holds in particular for each QL' -atomic sentence
- ▶ Construct a QL' -model \mathfrak{M}^* such that for each atomic QL' -sentence \mathcal{A} , $\mathfrak{M}^* \models \mathcal{A}$ iff $\mathcal{A} \in \Gamma^*$

Stage 3: The Maximal Consistency Lemma (\approx book's 11.4.7)

- ▶ **\exists -C Maximal Consistency Lemma**: every QL' set that is maximally-QND-consistent and \exists -complete is satisfiable
- ▶ (there exists a QL' -model that makes-true every sentence in Γ^*)
We construct this model, calling it " \mathfrak{M}^* " (\approx book's " \mathbf{I}^* ")
- ▶ Proof idea: since Γ^* is M-QND-C, for any sentence \mathcal{P} , either $\mathcal{P} \in \Gamma^*$ or $\sim \mathcal{P} \in \Gamma^*$ (you're either in the club or your '**nemesis**' is!)
This holds in particular for each QL' -atomic sentence
- ▶ Construct a QL' -model \mathfrak{M}^* such that for each atomic QL' -sentence \mathcal{A} , $\mathfrak{M}^* \models \mathcal{A}$ iff $\mathcal{A} \in \Gamma^*$
- ▶ Then by the recursive structure of QL' sentences, $\mathfrak{M}^* \models \mathcal{P}$ iff $\mathcal{P} \in \Gamma^*$

Stage 3 (i): the Membership Lemma (book's 11.4.6)

- To induct on QL' , we first constrain Γ^* membership

Stage 3 (i): the Membership Lemma (book's 11.4.6)

- ▶ To induct on QL' , we first constrain Γ^* membership
- ▶ Basically, Γ^* is *THE* club with the BADDEST MOTHAF**kin' bouncer you've eva seen, who enforces maximal consistency.

Stage 3 (i): the Membership Lemma (book's 11.4.6)

- ▶ To induct on QL' , we first constrain Γ^* membership
- ▶ Basically, Γ^* is *THE* club with the BADDEST MOTHAF**kin' bouncer you've eva seen, who enforces maximal consistency. Before this ma\$\$-hole lets a sentence into Γ^* ,

Stage 3 (i): the Membership Lemma (book's 11.4.6)

- ▶ To induct on QL' , we first constrain Γ^* membership
- ▶ Basically, Γ^* is *THE* club with the BADDEST MOTHAF**kin' bouncer you've eva seen, who enforces maximal consistency. Before this ma\$\$-hole lets a sentence into Γ^* , he checks who else is GOOD. You hear?

Stage 3 (i): the Membership Lemma (book's 11.4.6)

- ▶ To induct on QL' , we first constrain Γ^* membership
- ▶ Basically, Γ^* is *THE* club with the BADDEST MOTHAF**kin' bouncer you've eva seen, who enforces maximal consistency. Before this ma\$\$-hole lets a sentence into Γ^* , he checks who else is GOOD. You hear?
- ▶ **Membership Lemma** for club: if \mathcal{P} and \mathcal{Q} are QL' sentences, then:

Stage 3 (i): the Membership Lemma (book's 11.4.6)

- ▶ To induct on QL' , we first constrain Γ^* membership
- ▶ Basically, Γ^* is *THE* club with the BADDEST MOTHAF**kin' bouncer you've eva seen, who enforces maximal consistency. Before this ma\$\$-hole lets a sentence into Γ^* , he checks who else is GOOD. You hear?
- ▶ **Membership Lemma** for club: if \mathcal{P} and \mathcal{Q} are QL' sentences, then:
 - a.) $\sim\mathcal{P} \in \Gamma^*$ if and only if $\mathcal{P} \notin \Gamma^*$

Stage 3 (i): the Membership Lemma (book's 11.4.6)

- ▶ To induct on QL' , we first constrain Γ^* membership
- ▶ Basically, Γ^* is *THE* club with the BADDEST MOTHAF**kin' bouncer you've eva seen, who enforces maximal consistency. Before this ma\$\$-hole lets a sentence into Γ^* , he checks who else is GOOD. You hear?
- ▶ **Membership Lemma** for club: if \mathcal{P} and \mathcal{Q} are QL' sentences, then:
 - a.) $\sim\mathcal{P} \in \Gamma^*$ if and only if $\mathcal{P} \notin \Gamma^*$
 - b.) $\mathcal{P} \& \mathcal{Q} \in \Gamma^*$ if and only if both $\mathcal{P} \in \Gamma^*$ and $\mathcal{Q} \in \Gamma^*$

Stage 3 (i): the Membership Lemma (book's 11.4.6)

- ▶ To induct on QL' , we first constrain Γ^* membership
- ▶ Basically, Γ^* is *THE* club with the BADDEST MOTHAF**kin' bouncer you've eva seen, who enforces maximal consistency. Before this ma\$\$-hole lets a sentence into Γ^* , he checks who else is GOOD. You hear?
- ▶ **Membership Lemma** for club: if \mathcal{P} and \mathcal{Q} are QL' sentences, then:
 - a.) $\sim\mathcal{P} \in \Gamma^*$ if and only if $\mathcal{P} \notin \Gamma^*$
 - b.) $\mathcal{P} \& \mathcal{Q} \in \Gamma^*$ if and only if both $\mathcal{P} \in \Gamma^*$ and $\mathcal{Q} \in \Gamma^*$
 - c.) $\mathcal{P} \vee \mathcal{Q} \in \Gamma^*$ if and only if either $\mathcal{P} \in \Gamma^*$ or $\mathcal{Q} \in \Gamma^*$

Stage 3 (i): the Membership Lemma (book's 11.4.6)

- ▶ To induct on QL' , we first constrain Γ^* membership
- ▶ Basically, Γ^* is *THE* club with the BADDEST MOTHAF**kin' bouncer you've eva seen, who enforces maximal consistency. Before this ma\$\$-hole lets a sentence into Γ^* , he checks who else is GOOD. You hear?
- ▶ **Membership Lemma** for club: if \mathcal{P} and \mathcal{Q} are QL' sentences, then:
 - a.) $\sim\mathcal{P} \in \Gamma^*$ if and only if $\mathcal{P} \notin \Gamma^*$
 - b.) $\mathcal{P} \& \mathcal{Q} \in \Gamma^*$ if and only if both $\mathcal{P} \in \Gamma^*$ and $\mathcal{Q} \in \Gamma^*$
 - c.) $\mathcal{P} \vee \mathcal{Q} \in \Gamma^*$ if and only if either $\mathcal{P} \in \Gamma^*$ or $\mathcal{Q} \in \Gamma^*$
 - d.) $\mathcal{P} \supset \mathcal{Q} \in \Gamma^*$ if and only if either $\mathcal{P} \notin \Gamma^*$ or $\mathcal{Q} \in \Gamma^*$

Stage 3 (i): the Membership Lemma (book's 11.4.6)

- ▶ To induct on QL' , we first constrain Γ^* membership
- ▶ Basically, Γ^* is *THE* club with the BADDEST MOTHAF**kin' bouncer you've eva seen, who enforces maximal consistency. Before this ma\$\$-hole lets a sentence into Γ^* , he checks who else is GOOD. You hear?
- ▶ **Membership Lemma** for club: if \mathcal{P} and \mathcal{Q} are QL' sentences, then:
 - a.) $\sim \mathcal{P} \in \Gamma^*$ if and only if $\mathcal{P} \notin \Gamma^*$
 - b.) $\mathcal{P} \& \mathcal{Q} \in \Gamma^*$ if and only if both $\mathcal{P} \in \Gamma^*$ and $\mathcal{Q} \in \Gamma^*$
 - c.) $\mathcal{P} \vee \mathcal{Q} \in \Gamma^*$ if and only if either $\mathcal{P} \in \Gamma^*$ or $\mathcal{Q} \in \Gamma^*$
 - d.) $\mathcal{P} \supset \mathcal{Q} \in \Gamma^*$ if and only if either $\mathcal{P} \notin \Gamma^*$ or $\mathcal{Q} \in \Gamma^*$
 - e.) $\mathcal{P} \equiv \mathcal{Q} \in \Gamma^*$ iff either (i) $\mathcal{P} \in \Gamma^*$ and $\mathcal{Q} \in \Gamma^*$ or (ii) $\mathcal{P} \notin \Gamma^*$ and $\mathcal{Q} \notin \Gamma^*$

Stage 3 (i): the Membership Lemma (book's 11.4.6)

- ▶ To induct on QL' , we first constrain Γ^* membership
- ▶ Basically, Γ^* is *THE* club with the BADDEST MOTHAF**kin' bouncer you've ever seen, who enforces maximal consistency. Before this maff-hole lets a sentence into Γ^* , he checks who else is GOOD. You hear?
- ▶ **Membership Lemma** for club: if \mathcal{P} and \mathcal{Q} are QL' sentences, then:
 - a.) $\sim \mathcal{P} \in \Gamma^*$ if and only if $\mathcal{P} \notin \Gamma^*$
 - b.) $\mathcal{P} \& \mathcal{Q} \in \Gamma^*$ if and only if both $\mathcal{P} \in \Gamma^*$ and $\mathcal{Q} \in \Gamma^*$
 - c.) $\mathcal{P} \vee \mathcal{Q} \in \Gamma^*$ if and only if either $\mathcal{P} \in \Gamma^*$ or $\mathcal{Q} \in \Gamma^*$
 - d.) $\mathcal{P} \supset \mathcal{Q} \in \Gamma^*$ if and only if either $\mathcal{P} \notin \Gamma^*$ or $\mathcal{Q} \in \Gamma^*$
 - e.) $\mathcal{P} \equiv \mathcal{Q} \in \Gamma^*$ iff either (i) $\mathcal{P} \in \Gamma^*$ and $\mathcal{Q} \in \Gamma^*$ or (ii) $\mathcal{P} \notin \Gamma^*$ and $\mathcal{Q} \notin \Gamma^*$
 - f.) $(\forall \chi) \mathcal{P} \in \Gamma^*$ iff for each constant c , $\mathcal{P}[c/\chi] \in \Gamma^*$

Stage 3 (i): the Membership Lemma (book's 11.4.6)

- ▶ To induct on QL' , we first constrain Γ^* membership
- ▶ Basically, Γ^* is *THE* club with the BADDEST MOTHAF**kin' bouncer you've ever seen, who enforces maximal consistency. Before this maff-hole lets a sentence into Γ^* , he checks who else is GOOD. You hear?
- ▶ **Membership Lemma** for club: if \mathcal{P} and \mathcal{Q} are QL' sentences, then:
 - a.) $\sim \mathcal{P} \in \Gamma^*$ if and only if $\mathcal{P} \notin \Gamma^*$
 - b.) $\mathcal{P} \& \mathcal{Q} \in \Gamma^*$ if and only if both $\mathcal{P} \in \Gamma^*$ and $\mathcal{Q} \in \Gamma^*$
 - c.) $\mathcal{P} \vee \mathcal{Q} \in \Gamma^*$ if and only if either $\mathcal{P} \in \Gamma^*$ or $\mathcal{Q} \in \Gamma^*$
 - d.) $\mathcal{P} \supset \mathcal{Q} \in \Gamma^*$ if and only if either $\mathcal{P} \notin \Gamma^*$ or $\mathcal{Q} \in \Gamma^*$
 - e.) $\mathcal{P} \equiv \mathcal{Q} \in \Gamma^*$ iff either (i) $\mathcal{P} \in \Gamma^*$ and $\mathcal{Q} \in \Gamma^*$ or (ii) $\mathcal{P} \notin \Gamma^*$ and $\mathcal{Q} \notin \Gamma^*$
 - f.) $(\forall \chi) \mathcal{P} \in \Gamma^*$ iff for each constant c , $\mathcal{P}[c/\chi] \in \Gamma^*$
 - g.) $(\exists \chi) \mathcal{P} \in \Gamma^*$ iff for at least one constant c , $\mathcal{P}[c/\chi] \in \Gamma^*$

Stage 3 (i): **The Stairway** to heaven (book's 11.4.5)

- To prove the membership lemma's cases (a)-(g), we'll use another lemma (*NB*: and she's buying, a lemma, to

Stage 3 (i): **The Stairway** to heaven (book's 11.4.5)

- To prove the membership lemma's cases (a)-(g), we'll use another lemma (*NB*: and she's buying, a lemma, to heavennnnnnnnnn!):

Stage 3 (i): **The Stairway** to heaven (book's 11.4.5)

- ▶ To prove the membership lemma's cases (a)-(g), we'll use another lemma (*NB*: and she's buying, a lemma, to heavennnnnnnnnn!):
- ▶ **The Stairway**: if $\Gamma \vdash P$, and Γ^* is a maximally QND-consistent superset of Γ , then $P \in \Gamma^*$
(mnemonic: " $\Gamma \vdash P$ " pushes P up to QL'-heaven!)

Stage 3 (i): **The Stairway** to heaven (book's 11.4.5)

- ▶ To prove the membership lemma's cases (a)-(g), we'll use another lemma (*NB*: and she's buying, a lemma, to heavennnnnnnnnn!):
- ▶ **The Stairway**: if $\Gamma \vdash P$, and Γ^* is a maximally QND-consistent superset of Γ , then $P \in \Gamma^*$
(mnemonic: " $\Gamma \vdash P$ " pushes P up to QL'-heaven!)
- ▶ Proof: first, assume that $\Gamma \vdash P$ (we'll use this fact below)

Stage 3 (i): **The Stairway** to heaven (book's 11.4.5)

- ▶ To prove the membership lemma's cases (a)-(g), we'll use another lemma (*NB*: and she's buying, a lemma, to heavennnnnnnnnnn!):
- ▶ **The Stairway**: if $\Gamma \vdash P$, and Γ^* is a maximally QND-consistent superset of Γ , then $P \in \Gamma^*$
(mnemonic: " $\Gamma \vdash P$ " pushes P up to QL'-heaven!)
- ▶ Proof: first, assume that $\Gamma \vdash P$ (we'll use this fact below)
 - Next, assume for *reductio* that $P \notin \Gamma^*$. Then since Γ^* is maximally QND-consistent, $\Gamma^* \cup \{P\}$ must be **inconsistent in QND**.

Stage 3 (i): **The Stairway** to heaven (book's 11.4.5)

- ▶ To prove the membership lemma's cases (a)-(g), we'll use another lemma (*NB*: and she's buying, a lemma, to heavennnnnnnnnnn!):
- ▶ **The Stairway**: if $\Gamma \vdash P$, and Γ^* is a maximally QND-consistent superset of Γ , then $P \in \Gamma^*$
(mnemonic: " $\Gamma \vdash P$ " pushes P up to QL'-heaven!)
- ▶ Proof: first, assume that $\Gamma \vdash P$ (we'll use this fact below)
 - Next, assume for *reductio* that $P \notin \Gamma^*$. Then since Γ^* is maximally QND-consistent, $\Gamma^* \cup \{P\}$ must be **inconsistent in QND**.
 - Hence, by negation introduction, $\Gamma^* \vdash \sim P$

Stage 3 (i): **The Stairway** to heaven (book's 11.4.5)

- ▶ To prove the membership lemma's cases (a)–(g), we'll use another lemma (*NB*: and she's buying, a lemma, to heavennnnnnnnnn!):
- ▶ **The Stairway**: if $\Gamma \vdash P$, and Γ^* is a maximally QND-consistent superset of Γ , then $P \in \Gamma^*$
(mnemonic: “ $\Gamma \vdash P$ ” pushes P up to QL'–heaven!)
- ▶ Proof: first, assume that $\Gamma \vdash P$ (we'll use this fact below)
 - Next, assume for *reductio* that $P \notin \Gamma^*$. Then since Γ^* is maximally QND-consistent, $\Gamma^* \cup \{P\}$ must be **inconsistent in QND**.
 - Hence, by negation introduction, $\Gamma^* \vdash \sim P$
 - By assumption, $\Gamma \vdash P$, so also $\Gamma^* \vdash P$, since $\Gamma \subseteq \Gamma^*$

Stage 3 (i): **The Stairway** to heaven (book's 11.4.5)

- ▶ To prove the membership lemma's cases (a)–(g), we'll use another lemma (*NB*: and she's buying, a lemma, to heavennnnnnnnnnn!):
- ▶ **The Stairway**: if $\Gamma \vdash P$, and Γ^* is a maximally QND-consistent superset of Γ , then $P \in \Gamma^*$
(mnemonic: “ $\Gamma \vdash P$ ” pushes P up to QL'–heaven!)
- ▶ Proof: first, assume that $\Gamma \vdash P$ (we'll use this fact below)
 - Next, assume for *reductio* that $P \notin \Gamma^*$. Then since Γ^* is maximally QND-consistent, $\Gamma^* \cup \{P\}$ must be **inconsistent in QND**.
 - Hence, by negation introduction, $\Gamma^* \vdash \sim P$
 - By assumption, $\Gamma \vdash P$, so also $\Gamma^* \vdash P$, since $\Gamma \subseteq \Gamma^*$
 - So Γ^* derives both P and $\sim P$. *Reductio*! (since Γ^* is M-QND-C)

Stage 3 (i): **The Stairway** to heaven (book's 11.4.5)

- ▶ To prove the membership lemma's cases (a)–(g), we'll use another lemma (*NB*: and she's buying, a lemma, to heavennnnnnnnnnn!):
- ▶ **The Stairway**: if $\Gamma \vdash P$, and Γ^* is a maximally QND-consistent superset of Γ , then $P \in \Gamma^*$
(mnemonic: “ $\Gamma \vdash P$ ” pushes P up to QL'–heaven!)
- ▶ Proof: first, assume that $\Gamma \vdash P$ (we'll use this fact below)
 - Next, assume for *reductio* that $P \notin \Gamma^*$. Then since Γ^* is maximally QND-consistent, $\Gamma^* \cup \{P\}$ must be **inconsistent in QND**.
 - Hence, by negation introduction, $\Gamma^* \vdash \sim P$
 - By assumption, $\Gamma \vdash P$, so also $\Gamma^* \vdash P$, since $\Gamma \subseteq \Gamma^*$
 - So Γ^* derives both P and $\sim P$. *Reductio*! (since Γ^* is M-QND-C)
 - Hence, if $\Gamma \vdash P$ and $\Gamma \subseteq \Gamma^*$, then P must belong to Γ^*

Membership Lemma: Cases (a)–(e)

- ▶ I have a feeling that...

Membership Lemma: Cases (a)–(e)

- ▶ I have a feeling that...
- ▶ WE'VE SEEN THIS SH** BEFORE! (for SL)

Membership Lemma: Cases (a)–(e)

- ▶ I have a feeling that...
- ▶ WE'VE SEEN THIS SH** BEFORE! (for SL)
- ▶ see the next slide for a refresher

Membership Lemma: Cases (a)–(e)

- ▶ I have a feeling that...
- ▶ WE'VE SEEN THIS SH** BEFORE! (for SL)
- ▶ see the next slide for a refresher
- ▶ *Long story short*: There's a feeling I get

Membership Lemma: Cases (a)–(e)

- ▶ I have a feeling that...
- ▶ WE'VE SEEN THIS SH** BEFORE! (for SL)
- ▶ see the next slide for a refresher
- ▶ *Long story short*: There's a feeling I get

When I look to the west

Membership Lemma: Cases (a)–(e)

- ▶ I have a feeling that...
- ▶ WE'VE SEEN THIS SH** BEFORE! (for SL)
- ▶ see the next slide for a refresher
- ▶ *Long story short*: There's a feeling I get

When I look to the west

And my spirit is crying for leaving

Membership Lemma: Case (a)

- ▶ **Case (a):** $\sim\mathcal{P} \in \Gamma^*$ if and only if $\mathcal{P} \notin \Gamma^*$
- ▶ Two directions to prove:
 - \Rightarrow : Assume $\sim\mathcal{P} \in \Gamma^*$. Then if \mathcal{P} were in Γ^* , we could derive contradictory sentences.
So since Γ^* is QND-consistent, we must have $\mathcal{P} \notin \Gamma^*$
 - \Leftarrow : Assume $\mathcal{P} \notin \Gamma^*$. Then adding \mathcal{P} to Γ^* results in an QND-inconsistent set. Hence, there is some finite subset $\Delta \subset \Gamma^*$ s.t. $\Delta \cup \{\mathcal{P}\}$ is QND-inconsistent (i.e. derives contradictory sentence pair).
- ▶ So by negation introduction, $\Delta \vdash \sim\mathcal{P}$
- ▶ So by **The Stairway**, $\sim\mathcal{P} \in \Gamma^*$

Membership Lemma: Case (f) (something Universally new)

- **Case (f):** $(\forall \chi) \mathcal{P} \in \Gamma^*$ iff for each constant c , $\mathcal{P}[c/\chi] \in \Gamma^*$

Membership Lemma: Case (f) (something Universally new)

- ▶ **Case (f):** $(\forall \chi) \mathcal{P} \in \Gamma^*$ iff for each constant c , $\mathcal{P}[c/\chi] \in \Gamma^*$
- ▶ Two directions to prove:

Membership Lemma: Case (f) (something Universally new)

- ▶ **Case (f):** $(\forall \chi) \mathcal{P} \in \Gamma^*$ iff for each constant c , $\mathcal{P}[c/\chi] \in \Gamma^*$
- ▶ Two directions to prove:
 - \Rightarrow : Assume $(\forall \chi) \mathcal{P} \in \Gamma^*$

Membership Lemma: Case (f) (something Universally new)

- ▶ **Case (f):** $(\forall \chi)\mathcal{P} \in \Gamma^*$ iff for each constant c , $\mathcal{P}[c/\chi] \in \Gamma^*$
- ▶ Two directions to prove:
 - \Rightarrow : Assume $(\forall \chi)\mathcal{P} \in \Gamma^*$
 - Then **for any** substitution instance $\mathcal{P}[c/\chi]$, we note that $(\forall \chi)\mathcal{P} \vdash_{QND} \mathcal{P}[c/\chi]$ by $\forall E$. So by the Stairway, $\mathcal{P}[c/\chi] \in \Gamma^*$

Membership Lemma: Case (f) (something Universally new)

- ▶ **Case (f):** $(\forall \chi) \mathcal{P} \in \Gamma^*$ iff for each constant c , $\mathcal{P}[c/\chi] \in \Gamma^*$
- ▶ Two directions to prove:
 - \Rightarrow : Assume $(\forall \chi) \mathcal{P} \in \Gamma^*$
 - Then **for any** substitution instance $\mathcal{P}[c/\chi]$, we note that $(\forall \chi) \mathcal{P} \vdash_{QND} \mathcal{P}[c/\chi]$ by $\forall E$. So by the Stairway, $\mathcal{P}[c/\chi] \in \Gamma^*$
 - \Leftarrow : Assume $(\forall \chi) \mathcal{P} \notin \Gamma^*$. Show that for some constant c , $\mathcal{P}[c/\chi] \notin \Gamma^*$

Membership Lemma: Case (f) (something Universally new)

- ▶ **Case (f):** $(\forall \chi)\mathcal{P} \in \Gamma^*$ iff for each constant c , $\mathcal{P}[c/\chi] \in \Gamma^*$
- ▶ Two directions to prove:
 - \Rightarrow : Assume $(\forall \chi)\mathcal{P} \in \Gamma^*$
 - Then **for any** substitution instance $\mathcal{P}[c/\chi]$, we note that $(\forall \chi)\mathcal{P} \vdash_{QND} \mathcal{P}[c/\chi]$ by $\forall E$. So by the Stairway, $\mathcal{P}[c/\chi] \in \Gamma^*$
 - \Leftarrow : Assume $(\forall \chi)\mathcal{P} \notin \Gamma^*$. Show that for some constant c , $\mathcal{P}[c/\chi] \notin \Gamma^*$
 - Then $\sim(\forall \chi)\mathcal{P} \in \Gamma^*$ by membership clause (a)

Membership Lemma: Case (f) (something Universally new)

- ▶ **Case (f):** $(\forall \chi) \mathcal{P} \in \Gamma^*$ iff for each constant c , $\mathcal{P}[c/\chi] \in \Gamma^*$
- ▶ Two directions to prove:
 - \Rightarrow : Assume $(\forall \chi) \mathcal{P} \in \Gamma^*$
 - Then **for any** substitution instance $\mathcal{P}[c/\chi]$, we note that $(\forall \chi) \mathcal{P} \vdash_{QND} \mathcal{P}[c/\chi]$ by $\forall E$. So by the Stairway, $\mathcal{P}[c/\chi] \in \Gamma^*$
 - \Leftarrow : Assume $(\forall \chi) \mathcal{P} \notin \Gamma^*$. Show that for some constant c , $\mathcal{P}[c/\chi] \notin \Gamma^*$
 - Then $\sim(\forall \chi) \mathcal{P} \in \Gamma^*$ by membership clause (a)
 - Then the derivation on p. 573 or—if I have no life—the derivation on the next slide, shows by the Stairway that $(\exists \chi) \sim \mathcal{P} \in \Gamma^*$, i.e. $\sim(\forall \chi) \mathcal{P} \vdash_{QND} (\exists \chi) \sim \mathcal{P}$

Membership Lemma: Case (f) (something Universally new)

- ▶ **Case (f):** $(\forall \chi) \mathcal{P} \in \Gamma^*$ iff for each constant c , $\mathcal{P}[c/\chi] \in \Gamma^*$
- ▶ Two directions to prove:
 - \Rightarrow : Assume $(\forall \chi) \mathcal{P} \in \Gamma^*$
 - Then **for any** substitution instance $\mathcal{P}[c/\chi]$, we note that $(\forall \chi) \mathcal{P} \vdash_{QND} \mathcal{P}[c/\chi]$ by $\forall E$. So by the Stairway, $\mathcal{P}[c/\chi] \in \Gamma^*$
 - \Leftarrow : Assume $(\forall \chi) \mathcal{P} \notin \Gamma^*$. Show that for some constant c , $\mathcal{P}[c/\chi] \notin \Gamma^*$
 - Then $\sim(\forall \chi) \mathcal{P} \in \Gamma^*$ by membership clause (a)
 - Then the derivation on p. 573 or—if I have no life—the derivation on the next slide, shows by the Stairway that $(\exists \chi) \sim \mathcal{P} \in \Gamma^*$, i.e. $\sim(\forall \chi) \mathcal{P} \vdash_{QND} (\exists \chi) \sim \mathcal{P}$
 - Then since Γ^* is \exists -complete, there is at least one substitution instance $\sim \mathcal{P}[b/\chi] \in \Gamma^*$. So by (a), $\mathcal{P}[b/\chi] \notin \Gamma^*$, which is what we needed to show.

Membership Lemma: Case (g) (it's getting existential)

- **Case (g):** $(\exists \chi) \mathcal{P} \in \Gamma^*$ iff for at least one constant c , $\mathcal{P}[c/\chi] \in \Gamma^*$

Membership Lemma: Case (g) (it's getting existential)

- ▶ **Case (g):** $(\exists \chi) \mathcal{P} \in \Gamma^*$ iff for at least one constant c , $\mathcal{P}[c/\chi] \in \Gamma^*$
- ▶ \Rightarrow : Assume $(\exists \chi) \mathcal{P} \in \Gamma^*$

Membership Lemma: Case (g) (it's getting existential)

- ▶ **Case (g):** $(\exists \chi) \mathcal{P} \in \Gamma^*$ iff for at least one constant c , $\mathcal{P}[c/\chi] \in \Gamma^*$
- ▶ \Rightarrow : Assume $(\exists \chi) \mathcal{P} \in \Gamma^*$
Then since Γ^* is \exists -complete, there is at least one substitution instance $\mathcal{P}[c/\chi] \in \Gamma^*$

Membership Lemma: Case (g) (it's getting existential)

- ▶ **Case (g):** $(\exists \chi) \mathcal{P} \in \Gamma^*$ iff for at least one constant c , $\mathcal{P}[c/\chi] \in \Gamma^*$
- ▶ \Rightarrow : Assume $(\exists \chi) \mathcal{P} \in \Gamma^*$
Then since Γ^* is \exists -complete, there is at least one substitution instance $\mathcal{P}[c/\chi] \in \Gamma^*$
- ▶ \Leftarrow : assume that $\mathcal{P}[c/\chi] \in \Gamma^*$.

Membership Lemma: Case (g) (it's getting existential)

- ▶ **Case (g):** $(\exists \chi)\mathcal{P} \in \Gamma^*$ iff for at least one constant c , $\mathcal{P}[c/\chi] \in \Gamma^*$
- ▶ \Rightarrow : Assume $(\exists \chi)\mathcal{P} \in \Gamma^*$
Then since Γ^* is \exists -complete, there is at least one substitution instance $\mathcal{P}[c/\chi] \in \Gamma^*$
- ▶ \Leftarrow : assume that $\mathcal{P}[c/\chi] \in \Gamma^*$.
Note that $\mathcal{P}[c/\chi] \vdash_{QND} (\exists \chi)\mathcal{P}$ by Existential introduction

Membership Lemma: Case (g) (it's getting existential)

► **Case (g):** $(\exists \chi)\mathcal{P} \in \Gamma^*$ iff for at least one constant c , $\mathcal{P}[c/\chi] \in \Gamma^*$

► \Rightarrow : Assume $(\exists \chi)\mathcal{P} \in \Gamma^*$

Then since Γ^* is \exists -complete, there is at least one substitution instance $\mathcal{P}[c/\chi] \in \Gamma^*$

► \Leftarrow : assume that $\mathcal{P}[c/\chi] \in \Gamma^*$.

Note that $\mathcal{P}[c/\chi] \vdash_{QND} (\exists \chi)\mathcal{P}$ by Existential introduction

So by the Stairway, $(\exists \chi)\mathcal{P} \in \Gamma^*$

Membership Lemma: Case (g) (it's getting existential)

- ▶ **Case (g):** $(\exists \chi)\mathcal{P} \in \Gamma^*$ iff for at least one constant c , $\mathcal{P}[c/\chi] \in \Gamma^*$
- ▶ \Rightarrow : Assume $(\exists \chi)\mathcal{P} \in \Gamma^*$
Then since Γ^* is \exists -complete, there is at least one substitution instance $\mathcal{P}[c/\chi] \in \Gamma^*$
- ▶ \Leftarrow : assume that $\mathcal{P}[c/\chi] \in \Gamma^*$.
Note that $\mathcal{P}[c/\chi] \vdash_{QND} (\exists \chi)\mathcal{P}$ by Existential introduction
So by the Stairway, $(\exists \chi)\mathcal{P} \in \Gamma^*$
- ▶ This completes the Membership Lemma, so we proceed to construct a model that satisfies Γ^* (in virtue of being maximally-QND-consistent and \exists -complete)!

Stage 3 (ii): Model construction (smart choices=lazy choices)

- ▶ A model's domain can be *any* set of objects. Note that, conveniently, symbols *are* objects (“words are labels on boxes”)

Stage 3 (ii): Model construction (smart choices=lazy choices)

- ▶ A model's domain can be *any* set of objects. Note that, conveniently, symbols *are* objects (“words are labels on boxes”)
- ▶ We define $\mathfrak{M}^* := (D, I^*)$ as follows:

Stage 3 (ii): Model construction (smart choices=lazy choices)

- ▶ A model's domain can be *any* set of objects. Note that, conveniently, symbols *are* objects (“words are labels on boxes”)
- ▶ We define $\mathfrak{M}^* := (D, I^*)$ as follows:
 1. Let D = the set of constant symbols in QL' , which includes all QL-constants (e.g. unprimed subscripted constants like j_{22})

Stage 3 (ii): Model construction (smart choices=lazy choices)

- ▶ A model's domain can be *any* set of objects. Note that, conveniently, symbols *are* objects (“words are labels on boxes”)
- ▶ We define $\mathfrak{M}^* := (D, I^*)$ as follows:
 1. Let D = the set of constant symbols in QL' , which includes all QL-constants (e.g. unprimed subscripted constants like j_{22})
 2. For the 0-th place predicates, i.e. the sentence letters B ,
 $I^*(B) = \text{true}$ iff $B \in \Gamma^*$

Stage 3 (ii): Model construction (smart choices=lazy choices)

- ▶ A model's domain can be *any* set of objects. Note that, conveniently, symbols *are* objects (“words are labels on boxes”)
- ▶ We define $\mathfrak{M}^* := (D, I^*)$ as follows:
 1. Let D = the set of constant symbols in QL' , which includes all QL -constants (e.g. unprimed subscripted constants like j_{22})
 2. For the 0-th place predicates, i.e. the sentence letters B ,
 $I^*(B) = \text{true}$ iff $B \in \Gamma^*$
 3. For each QL' -constant c , define $I^*(c) = c$ (each names itself)

Stage 3 (ii): Model construction (smart choices=lazy choices)

- ▶ A model's domain can be *any* set of objects. Note that, conveniently, symbols *are* objects (“words are labels on boxes”)
- ▶ We define $\mathfrak{M}^* := (D, I^*)$ as follows:
 1. Let D = the set of constant symbols in QL' , which includes all QL -constants (e.g. unprimed subscripted constants like j_{22})
 2. For the 0-th place predicates, i.e. the sentence letters B , $I^*(B) = \text{true}$ iff $B \in \Gamma^*$
 3. For each QL' -constant c , define $I^*(c) = c$ (each names itself)
 4. For each k -place predicate P , $I^*(P) := \text{Ext}(P)$ includes all and only those k -tuples $\langle c_1, \dots, c_k \rangle$ such that $Pc_1 \dots c_k \in \Gamma^*$

Some important properties of our Model \mathfrak{M}^*

- By condition 3, each individual constant refers to a *unique* member of the domain, namely ‘itself’ (now ‘objectified’ in D !)

Some important properties of our Model \mathfrak{M}^*

- ▶ By condition 3, each individual constant refers to a *unique* member of the domain, namely ‘itself’ (now ‘objectified’ in D !)
- ▶ For each atomic sentence \mathcal{A} of QL' , $\mathfrak{M}^* \models \mathcal{A}$ iff $\mathcal{A} \in \Gamma^*$ (follows from conditions 2–4)

Some important properties of our Model \mathfrak{M}^*

- ▶ By condition 3, each individual constant refers to a *unique* member of the domain, namely ‘itself’ (now ‘objectified’ in D !)
- ▶ For each atomic sentence \mathcal{A} of QL' , $\mathfrak{M}^* \models \mathcal{A}$ iff $\mathcal{A} \in \Gamma^*$ (follows from conditions 2–4)
- ▶ By condition 3, every member of the domain is named by a constant, namely itself

Some important properties of our Model \mathfrak{M}^*

- ▶ By condition 3, each individual constant refers to a *unique* member of the domain, namely ‘itself’ (now ‘objectified’ in D !)
- ▶ For each atomic sentence \mathcal{A} of QL' , $\mathfrak{M}^* \models \mathcal{A}$ iff $\mathcal{A} \in \Gamma^*$ (follows from conditions 2–4)
- ▶ By condition 3, every member of the domain is named by a constant, namely itself
- ▶ We will occasionally rely on these properties in our induction

Stage 3 (iii): Induction on QL' (i.e. we still be clubbin')

- Goal: construct a QL' -model \mathfrak{M}^* that satisfies the \exists -C M-QND-C set Γ^* , i.e. that makes true everything in Γ^* ($\mathfrak{M}^* \models \Gamma^*$)

Stage 3 (iii): Induction on QL' (i.e. we still be clubbin')

- Goal: construct a QL' -model \mathfrak{M}^* that satisfies the \exists -C M-QND-C set Γ^* , i.e. that makes true everything in Γ^* ($\mathfrak{M}^* \models \Gamma^*$)
 - Suffices to construct \mathfrak{M}^* s.t. $\mathfrak{M}^* \models \mathcal{P}$ iff $\mathcal{P} \in \Gamma^*$

Stage 3 (iii): Induction on QL' (i.e. we still be clubbin')

- Goal: construct a QL' -model \mathfrak{M}^* that satisfies the \exists -C M-QND-C set Γ^* , i.e. that makes true everything in Γ^* ($\mathfrak{M}^* \models \Gamma^*$)
 - Suffices to construct \mathfrak{M}^* s.t. $\mathfrak{M}^* \models \mathcal{P}$ iff $\mathcal{P} \in \Gamma^*$
- Say that a sentence is “**clubbin'**” whenever it meets this property

Stage 3 (iii): Induction on QL' (i.e. we still be clubbin')

- ▶ Goal: construct a QL' -model \mathfrak{M}^* that satisfies the \exists -C M-QND-C set Γ^* , i.e. that makes true everything in Γ^* ($\mathfrak{M}^* \models \Gamma^*$)
 - Suffices to construct \mathfrak{M}^* s.t. $\mathfrak{M}^* \models \mathcal{P}$ iff $\mathcal{P} \in \Gamma^*$Say that a sentence is “**clubbin'**” whenever it meets this property
- ▶ We induct on the number of logical operators in a QL' sentence: i.e. the five connectives and two quantifiers (“conquans”)

Stage 3 (iii): Induction on QL' (i.e. we still be clubbin')

- ▶ Goal: construct a QL' -model \mathfrak{M}^* that satisfies the \exists -C M-QND-C set Γ^* , i.e. that makes true everything in Γ^* ($\mathfrak{M}^* \models \Gamma^*$)
 - Suffices to construct \mathfrak{M}^* s.t. $\mathfrak{M}^* \models \mathcal{P}$ iff $\mathcal{P} \in \Gamma^*$Say that a sentence is “**clubbin'**” whenever it meets this property
- ▶ We induct on the number of logical operators in a QL' sentence: i.e. the five connectives and two quantifiers (“conquans”)
- ▶ **Base case**: show that each QL' -sentence with **zero** logical operators is clubbin' (i.e. the QL' -atomics be clubbin')

Stage 3 (iii): Induction on QL' (i.e. we still be clubbin')

- ▶ Goal: construct a QL' -model \mathfrak{M}^* that satisfies the \exists -C M-QND-C set Γ^* , i.e. that makes true everything in Γ^* ($\mathfrak{M}^* \models \Gamma^*$)
 - Suffices to construct \mathfrak{M}^* s.t. $\mathfrak{M}^* \models \mathcal{P}$ iff $\mathcal{P} \in \Gamma^*$Say that a sentence is “**clubbin'**” whenever it meets this property
- ▶ We induct on the number of logical operators in a QL' sentence: i.e. the five connectives and two quantifiers (“conquans”)
- ▶ **Base case**: show that each QL' -sentence with **zero** logical operators is clubbin' (i.e. the QL' -atomics be clubbin')
- ▶ (Strong) **Induction hypothesis**: assume every QL' sentence with 1 to k -many operators is clubbin'

Stage 3 (iii): Induction on QL' (i.e. we still be clubbin')

- ▶ Goal: construct a QL' -model \mathfrak{M}^* that satisfies the \exists -C M-QND-C set Γ^* , i.e. that makes true everything in Γ^* ($\mathfrak{M}^* \models \Gamma^*$)
 - Suffices to construct \mathfrak{M}^* s.t. $\mathfrak{M}^* \models \mathcal{P}$ iff $\mathcal{P} \in \Gamma^*$Say that a sentence is “**clubbin'**” whenever it meets this property
- ▶ We induct on the number of logical operators in a QL' sentence: i.e. the five connectives and two quantifiers (“conquans”)
- ▶ **Base case**: show that each QL' -sentence with **zero** logical operators is clubbin' (i.e. the QL' -atomics be clubbin')
- ▶ (Strong) **Induction hypothesis**: assume every QL' sentence with 1 to k -many operators is clubbin'
- ▶ Induction step: show that an arbitrary QL' sentence with $k+1$ -many operators is clubbin'

Base Case (true by construction)

- Consider an arbitrary QL' -sentence \mathcal{A} that has zero logical operators. (Two directions to show! “iff”)

Base Case (true by construction)

- ▶ Consider an arbitrary QL' -sentence \mathcal{A} that has zero logical operators. (Two directions to show! “iff”)
- ▶ Then \mathcal{A} is either an atomic sentence letter B or of the form $Pc_1 \dots c_n$ for n -place predicate P .

Base Case (true by construction)

- ▶ Consider an arbitrary QL' -sentence \mathcal{A} that has zero logical operators. (Two directions to show! “iff”)
- ▶ Then \mathcal{A} is either an atomic sentence letter B or of the form $Pc_1 \dots c_n$ for n -place predicate P .
- ▶ If a sentence letter, then by part 2 of our definition of \mathfrak{M}^* , $I^*(B) = \text{true}$ iff $B \in \Gamma^*$ (i.e. $\mathfrak{M}^* \models B$ iff $B \in \Gamma^*$)

Base Case (true by construction)

- ▶ Consider an arbitrary QL'-sentence \mathcal{A} that has zero logical operators. (Two directions to show! “iff”)
- ▶ Then \mathcal{A} is either an atomic sentence letter B or of the form $Pc_1 \dots c_n$ for n -place predicate P .
- ▶ If a sentence letter, then by part 2 of our definition of \mathfrak{M}^* , $I^*(B) = \text{true}$ iff $B \in \Gamma^*$ (i.e. $\mathfrak{M}^* \models B$ iff $B \in \Gamma^*$)
- ▶ If \mathcal{A} is of form $Pc_1 \dots c_n$, then by definition $\mathfrak{M}^* \models Pc_1 \dots c_n$ iff $\langle c_1^D, \dots, c_n^D \rangle \in \text{Ext}(P)$.

Base Case (true by construction)

- ▶ Consider an arbitrary QL' -sentence \mathcal{A} that has zero logical operators. (Two directions to show! “iff”)
- ▶ Then \mathcal{A} is either an atomic sentence letter B or of the form $Pc_1 \dots c_n$ for n -place predicate P .
- ▶ If a sentence letter, then by part 2 of our definition of \mathfrak{M}^* , $I^*(B) = \text{true}$ iff $B \in \Gamma^*$ (i.e. $\mathfrak{M}^* \models B$ iff $B \in \Gamma^*$)
- ▶ If \mathcal{A} is of form $Pc_1 \dots c_n$, then by definition $\mathfrak{M}^* \models Pc_1 \dots c_n$ iff $\langle c_1^D, \dots, c_n^D \rangle \in \text{Ext}(P)$.
By part 4, $\langle c_1^D, \dots, c_n^D \rangle = \langle c_1, \dots, c_n \rangle \in \text{Ext}(P)$ iff $Pc_1 \dots c_n \in \Gamma^*$

Base Case (true by construction)

- ▶ Consider an arbitrary QL'-sentence \mathcal{A} that has zero logical operators. (Two directions to show! “iff”)
- ▶ Then \mathcal{A} is either an atomic sentence letter B or of the form $Pc_1 \dots c_n$ for n -place predicate P .
- ▶ If a sentence letter, then by part 2 of our definition of \mathfrak{M}^* , $I^*(B) = \text{true}$ iff $B \in \Gamma^*$ (i.e. $\mathfrak{M}^* \models B$ iff $B \in \Gamma^*$)
- ▶ If \mathcal{A} is of form $Pc_1 \dots c_n$, then by definition $\mathfrak{M}^* \models Pc_1 \dots c_n$ iff $\langle c_1^D, \dots, c_n^D \rangle \in \text{Ext}(P)$.
By part 4, $\langle c_1^D, \dots, c_n^D \rangle = \langle c_1, \dots, c_n \rangle \in \text{Ext}(P)$ iff $Pc_1 \dots c_n \in \Gamma^*$
- ▶ We proceed to do induction using our QL' induction schema: an arbitrary sentence \mathcal{P} with $k+1$ -many connectives has one of seven forms, coming from our seven operators

Induction on QL': Cases 1–5

- ▶ Cases 1–5 are just like what did to prove the completeness of SND
- ▶ See the next slide for a refresher (*mutatis mutandis*)!
- ▶ Need to show: \mathcal{P} be clubbin', i.e. \mathcal{P} is true on \mathfrak{M}^* iff $\mathcal{P} \in \Gamma^*$, where \mathcal{P} is arbitrary QL' sentence with $k+1$ -many operators
- ▶ **Induction hypothesis**: assume every QL sentence with 1 to k -many operators is clubbin'
- ▶ Case 1: \mathcal{P} has the form $\sim Q$
- ▶ Case 2: \mathcal{P} has the form $Q \& \mathcal{R}$
- ▶ Case 3: \mathcal{P} has the form $Q \vee \mathcal{R}$
- ▶ Case 4: \mathcal{P} has the form $Q \supset \mathcal{R}$
- ▶ Case 5: \mathcal{P} has the form $Q \equiv \mathcal{R}$

Induction on QL' : Case 1

- **Case 1:** \mathcal{P} has the form $\sim Q$, where since Q has k -operators, it is clubbin by the IH (i.e. $\mathfrak{M}^* \models Q$ if and only if $Q \in \Gamma^*$)

Induction on QL' : Case 1

- ▶ **Case 1:** \mathcal{P} has the form $\sim Q$, where since Q has k -operators, it is clubbin by the IH (i.e. $\mathfrak{M}^* \models Q$ if and only if $Q \in \Gamma^*$)
- ▶ NTS: (i) (the \Rightarrow direction) if $\mathfrak{M}^* \models \mathcal{P}$ then $\mathcal{P} \in \Gamma^*$ and
(ii) (the \Leftarrow direction) if $\mathcal{P} \in \Gamma^*$, then $\mathfrak{M}^* \models \mathcal{P}$

Induction on QL' : Case 1

- ▶ **Case 1:** \mathcal{P} has the form $\sim Q$, where since Q has k -operators, it is clubbin by the IH (i.e. $\mathfrak{M}^* \models Q$ if and only if $Q \in \Gamma^*$)
- ▶ NTS: (i) (the \Rightarrow direction) if $\mathfrak{M}^* \models \mathcal{P}$ then $\mathcal{P} \in \Gamma^*$ and
(ii) (the \Leftarrow direction) if $\mathcal{P} \in \Gamma^*$, then $\mathfrak{M}^* \models \mathcal{P}$
(Alternative (ii): show contrapositive: if $\mathfrak{M}^* \not\models \mathcal{P}$, then $\mathcal{P} \notin \Gamma^*$)

Induction on QL' : Case 1

- ▶ **Case 1:** \mathcal{P} has the form $\sim \mathcal{Q}$, where since \mathcal{Q} has k -operators, it is clubbin by the IH (i.e. $\mathfrak{M}^* \models \mathcal{Q}$ if and only if $\mathcal{Q} \in \Gamma^*$)
 - ▶ NTS: (i) (the \Rightarrow direction) if $\mathfrak{M}^* \models \mathcal{P}$ then $\mathcal{P} \in \Gamma^*$ and
(ii) (the \Leftarrow direction) if $\mathcal{P} \in \Gamma^*$, then $\mathfrak{M}^* \models \mathcal{P}$
(*Alternative (ii):* show contrapositive: if $\mathfrak{M}^* \not\models \mathcal{P}$, then $\mathcal{P} \notin \Gamma^*$)
- \Rightarrow if $\mathfrak{M}^* \models \mathcal{P}$, then $\mathfrak{M}^* \not\models \mathcal{Q}$. Since \mathcal{Q} is clubbin', we have $\mathcal{Q} \notin \Gamma^*$

Induction on QL' : Case 1

- ▶ **Case 1:** \mathcal{P} has the form $\sim Q$, where since Q has k -operators, it is clubbin by the IH (i.e. $\mathfrak{M}^* \models Q$ if and only if $Q \in \Gamma^*$)
 - ▶ NTS: (i) (the \Rightarrow direction) if $\mathfrak{M}^* \models \mathcal{P}$ then $\mathcal{P} \in \Gamma^*$ and
(ii) (the \Leftarrow direction) if $\mathcal{P} \in \Gamma^*$, then $\mathfrak{M}^* \models \mathcal{P}$
(Alternative (ii): show contrapositive: if $\mathfrak{M}^* \not\models \mathcal{P}$, then $\mathcal{P} \notin \Gamma^*$)
- \Rightarrow if $\mathfrak{M}^* \models \mathcal{P}$, then $\mathfrak{M}^* \not\models Q$. Since Q is clubbin', we have $Q \notin \Gamma^*$
By Membership lemma (a), $\sim Q \in \Gamma^*$, so $\mathcal{P} \in \Gamma^*$

Induction on QL' : Case 1

- ▶ **Case 1:** \mathcal{P} has the form $\sim Q$, where since Q has k -operators, it is clubbin by the IH (i.e. $\mathfrak{M}^* \models Q$ if and only if $Q \in \Gamma^*$)
 - ▶ NTS: (i) (the \Rightarrow direction) if $\mathfrak{M}^* \models \mathcal{P}$ then $\mathcal{P} \in \Gamma^*$ and
(ii) (the \Leftarrow direction) if $\mathcal{P} \in \Gamma^*$, then $\mathfrak{M}^* \models \mathcal{P}$
(*Alternative (ii):* show contrapositive: if $\mathfrak{M}^* \not\models \mathcal{P}$, then $\mathcal{P} \notin \Gamma^*$)
- \Rightarrow if $\mathfrak{M}^* \models \mathcal{P}$, then $\mathfrak{M}^* \not\models Q$. Since Q is clubbin', we have $Q \notin \Gamma^*$
By Membership lemma (a), $\sim Q \in \Gamma^*$, so $\mathcal{P} \in \Gamma^*$
- \Leftarrow if $\mathcal{P} \in \Gamma^*$, then $\sim Q \in \Gamma^*$. So by Membership lemma (a), $Q \notin \Gamma^*$.

Induction on QL' : Case 1

- ▶ **Case 1:** \mathcal{P} has the form $\sim Q$, where since Q has k -operators, it is clubbin by the IH (i.e. $\mathfrak{M}^* \models Q$ if and only if $Q \in \Gamma^*$)
 - ▶ NTS: (i) (the \Rightarrow direction) if $\mathfrak{M}^* \models \mathcal{P}$ then $\mathcal{P} \in \Gamma^*$ and
(ii) (the \Leftarrow direction) if $\mathcal{P} \in \Gamma^*$, then $\mathfrak{M}^* \models \mathcal{P}$
(Alternative (ii): show contrapositive: if $\mathfrak{M}^* \not\models \mathcal{P}$, then $\mathcal{P} \notin \Gamma^*$)
- \Rightarrow if $\mathfrak{M}^* \models \mathcal{P}$, then $\mathfrak{M}^* \not\models Q$. Since Q is clubbin', we have $Q \notin \Gamma^*$
By Membership lemma (a), $\sim Q \in \Gamma^*$, so $\mathcal{P} \in \Gamma^*$
- \Leftarrow if $\mathcal{P} \in \Gamma^*$, then $\sim Q \in \Gamma^*$. So by Membership lemma (a), $Q \notin \Gamma^*$.
Since Q is clubbin', we have $\mathfrak{M}^* \not\models Q$. (i.e. Q is false in \mathfrak{M}^*)

Induction on QL' : Case 1

- ▶ **Case 1:** \mathcal{P} has the form $\sim Q$, where since Q has k -operators, it is clubbin by the IH (i.e. $\mathfrak{M}^* \models Q$ if and only if $Q \in \Gamma^*$)
- ▶ NTS: (i) (the \Rightarrow direction) if $\mathfrak{M}^* \models \mathcal{P}$ then $\mathcal{P} \in \Gamma^*$ and
(ii) (the \Leftarrow direction) if $\mathcal{P} \in \Gamma^*$, then $\mathfrak{M}^* \models \mathcal{P}$
(Alternative (ii): show contrapositive: if $\mathfrak{M}^* \not\models \mathcal{P}$, then $\mathcal{P} \notin \Gamma^*$)
 \Rightarrow if $\mathfrak{M}^* \models \mathcal{P}$, then $\mathfrak{M}^* \not\models Q$. Since Q is clubbin', we have $Q \notin \Gamma^*$
By Membership lemma (a), $\sim Q \in \Gamma^*$, so $\mathcal{P} \in \Gamma^*$
 \Leftarrow if $\mathcal{P} \in \Gamma^*$, then $\sim Q \in \Gamma^*$. So by Membership lemma (a), $Q \notin \Gamma^*$.
Since Q is clubbin', we have $\mathfrak{M}^* \not\models Q$. (i.e. Q is false in \mathfrak{M}^*)
So by the truth conditions for negation, \mathcal{P} is true in \mathfrak{M}^* , i.e. $\mathfrak{M}^* \models \mathcal{P}$

Induction on QL' : Case 7 (existential quantifier)

- **Case 7:** \mathcal{P} has the form $(\exists x)Q$
(warning: “ Q ” is not a sentence, so sadly it can’t be clubbin’)

Induction on QL' : Case 7 (existential quantifier)

- ▶ **Case 7:** \mathcal{P} has the form $(\exists \chi)Q$
(warning: “ Q ” is not a sentence, so sadly it can’t be clubbin’)
- ▶ We will use Membership Lemma **Case (g)**:
 $(\exists \chi)Q \in \Gamma^*$ iff for at least one constant c , $Q[c/\chi] \in \Gamma^*$

Induction on QL' : Case 7 (existential quantifier)

- ▶ **Case 7:** \mathcal{P} has the form $(\exists \chi)Q$
(warning: “ Q ” is not a sentence, so sadly it can’t be clubbin’)
 - ▶ We will use Membership Lemma **Case (g)**:
 $(\exists \chi)Q \in \Gamma^*$ iff for at least one constant c , $Q[c/\chi] \in \Gamma^*$
- \Rightarrow Assume $\mathfrak{M}^* \models (\exists \chi)Q$. (need to show that $(\exists \chi)Q \in \Gamma^*$)

Induction on QL' : Case 7 (existential quantifier)

- ▶ **Case 7:** \mathcal{P} has the form $(\exists x)Q$
(warning: “ Q ” is not a sentence, so sadly it can’t be clubbin’)
 - ▶ We will use Membership Lemma **Case (g)**:
 $(\exists x)Q \in \Gamma^*$ iff for at least one constant c , $Q[c/x] \in \Gamma^*$
- \Rightarrow Assume $\mathfrak{M}^* \models (\exists x)Q$. (need to show that $(\exists x)Q \in \Gamma^*$)
- Then by the truth-conditions for existential, there is some object $r \in D$ that satisfies Q .

Induction on QL' : Case 7 (existential quantifier)

- ▶ **Case 7:** \mathcal{P} has the form $(\exists \chi)Q$
(warning: “ Q ” is not a sentence, so sadly it can’t be clubbin’)
 - ▶ We will use Membership Lemma **Case (g)**:
 $(\exists \chi)Q \in \Gamma^*$ iff for at least one constant c , $Q[c/\chi] \in \Gamma^*$
- \Rightarrow Assume $\mathfrak{M}^* \models (\exists \chi)Q$. (need to show that $(\exists \chi)Q \in \Gamma^*$)
- Then by the truth-conditions for existential, there is some object $r \in D$ that satisfies Q .
 - ‘ r ’ names object r , so substitution instance $Q[r/\chi]$ is true in \mathfrak{M}^*

Induction on QL' : Case 7 (existential quantifier)

- ▶ **Case 7:** \mathcal{P} has the form $(\exists \chi)Q$
(warning: “ Q ” is not a sentence, so sadly it can’t be clubbin’)
 - ▶ We will use Membership Lemma **Case (g)**:
 $(\exists \chi)Q \in \Gamma^*$ iff for at least one constant c , $Q[c/\chi] \in \Gamma^*$
- \Rightarrow Assume $\mathfrak{M}^* \models (\exists \chi)Q$. (need to show that $(\exists \chi)Q \in \Gamma^*$)
- Then by the truth-conditions for existential, there is some object $r \in D$ that satisfies Q .
 - ‘ r ’ names object r , so substitution instance $Q[r/\chi]$ is true in \mathfrak{M}^*
 - This substitution instance has less than $k + 1$ -operators, so it is clubbin’. Hence, by the IH, $Q[r/\chi] \in \Gamma^*$ (since $\mathfrak{M}^* \models Q[r/\chi]$)

Induction on QL' : Case 7 (existential quantifier)

- ▶ **Case 7:** \mathcal{P} has the form $(\exists \chi)Q$
(warning: “ Q ” is not a sentence, so sadly it can’t be clubbin’)
 - ▶ We will use Membership Lemma **Case (g)**:
 $(\exists \chi)Q \in \Gamma^*$ iff for at least one constant c , $Q[c/\chi] \in \Gamma^*$
- \Rightarrow Assume $\mathfrak{M}^* \models (\exists \chi)Q$. (need to show that $(\exists \chi)Q \in \Gamma^*$)
- Then by the truth-conditions for existential, there is some object $r \in D$ that satisfies Q .
 - ‘ r ’ names object r , so substitution instance $Q[r/\chi]$ is true in \mathfrak{M}^*
 - This substitution instance has less than $k + 1$ -operators, so it is clubbin’. Hence, by the IH, $Q[r/\chi] \in \Gamma^*$ (since $\mathfrak{M}^* \models Q[r/\chi]$)
 - So by membership case (g), $(\exists \chi)Q \in \Gamma^*$

Induction on QL' : Case 7 backwards direction

- **Case 7:** \mathcal{P} has the form $(\exists x)Q$

Induction on QL' : Case 7 backwards direction

- ▶ **Case 7:** \mathcal{P} has the form $(\exists \chi)Q$
- ▶ Use Membership Lemma **Case (g)**:
 $(\exists \chi)Q \in \Gamma^*$ iff for at least one constant c , $Q[c/\chi] \in \Gamma^*$

Induction on QL' : Case 7 backwards direction

- ▶ **Case 7:** \mathcal{P} has the form $(\exists \chi)Q$
 - ▶ Use Membership Lemma **Case (g)**:
 $(\exists \chi)Q \in \Gamma^*$ iff for at least one constant c , $Q[c/\chi] \in \Gamma^*$
- \Leftarrow Assume $(\exists \chi)Q \in \Gamma^*$. Show that $\mathfrak{M}^* \models (\exists \chi)Q$

Induction on QL' : Case 7 backwards direction

- ▶ **Case 7:** \mathcal{P} has the form $(\exists \chi)Q$
- ▶ Use Membership Lemma **Case (g)**:
 $(\exists \chi)Q \in \Gamma^*$ iff for at least one constant c , $Q[c/\chi] \in \Gamma^*$
- \Leftarrow Assume $(\exists \chi)Q \in \Gamma^*$. Show that $\mathfrak{M}^* \models (\exists \chi)Q$
 - Then by membership case (g), there is at least one substitution instance $Q[c/\chi] \in \Gamma^*$, for some constant c

Induction on QL' : Case 7 backwards direction

- ▶ **Case 7:** \mathcal{P} has the form $(\exists \chi) \mathcal{Q}$
- ▶ Use Membership Lemma **Case (g)**:
 $(\exists \chi) \mathcal{Q} \in \Gamma^*$ iff for at least one constant c , $\mathcal{Q}[c/\chi] \in \Gamma^*$
- \Leftarrow Assume $(\exists \chi) \mathcal{Q} \in \Gamma^*$. Show that $\mathfrak{M}^* \models (\exists \chi) \mathcal{Q}$
 - Then by membership case (g), there is at least one substitution instance $\mathcal{Q}[c/\chi] \in \Gamma^*$, for some constant c
 - Since $\mathcal{Q}[c/\chi]$ has fewer than $k + 1$ -operators, it is clubbin'.

Induction on QL' : Case 7 backwards direction

- ▶ **Case 7:** \mathcal{P} has the form $(\exists \chi)Q$
- ▶ Use Membership Lemma **Case (g)**:
 $(\exists \chi)Q \in \Gamma^*$ iff for at least one constant c , $Q[c/\chi] \in \Gamma^*$
- \Leftarrow Assume $(\exists \chi)Q \in \Gamma^*$. Show that $\mathfrak{M}^* \models (\exists \chi)Q$
 - Then by membership case (g), there is at least one substitution instance $Q[c/\chi] \in \Gamma^*$, for some constant c
 - Since $Q[c/\chi]$ has fewer than $k + 1$ -operators, it is clubbin'.
 - So by the Induction Hypothesis, $\mathfrak{M}^* \models Q[c/\chi]$.

Induction on QL' : Case 7 backwards direction

- ▶ **Case 7:** \mathcal{P} has the form $(\exists \chi)Q$
- ▶ Use Membership Lemma **Case (g)**:
 $(\exists \chi)Q \in \Gamma^*$ iff for at least one constant c , $Q[c/\chi] \in \Gamma^*$

\Leftarrow Assume $(\exists \chi)Q \in \Gamma^*$. Show that $\mathfrak{M}^* \models (\exists \chi)Q$

- Then by membership case (g), there is at least one substitution instance $Q[c/\chi] \in \Gamma^*$, for some constant c
- Since $Q[c/\chi]$ has fewer than $k + 1$ -operators, it is clubbin'.
- So by the Induction Hypothesis, $\mathfrak{M}^* \models Q[c/\chi]$.
- Since ' c ' names object c , we see that c satisfies Q in \mathfrak{M}^*

Induction on QL': Case 7 backwards direction

- ▶ **Case 7:** \mathcal{P} has the form $(\exists \chi)Q$
- ▶ Use Membership Lemma **Case (g)**:
 $(\exists \chi)Q \in \Gamma^*$ iff for at least one constant c , $Q[c/\chi] \in \Gamma^*$
- \Leftarrow Assume $(\exists \chi)Q \in \Gamma^*$. Show that $\mathfrak{M}^* \models (\exists \chi)Q$
 - Then by membership case (g), there is at least one substitution instance $Q[c/\chi] \in \Gamma^*$, for some constant c
 - Since $Q[c/\chi]$ has fewer than $k + 1$ -operators, it is clubbin'.
 - So by the Induction Hypothesis, $\mathfrak{M}^* \models Q[c/\chi]$.
 - Since ' c ' names object c , we see that c satisfies Q in \mathfrak{M}^*
 - So by the truth-conditions for existentials, $(\exists \chi)Q$ is true in \mathfrak{M}^*

Induction on QL' : Case 6 (universal quantifier)

- **Case 6:** \mathcal{P} has the form $(\forall x)Q$
(warning: “ Q ” is not a sentence, so it can’t be clubbin’)

Induction on QL' : Case 6 (universal quantifier)

- ▶ **Case 6:** \mathcal{P} has the form $(\forall \chi)Q$
(warning: “ Q ” is not a sentence, so it can’t be clubbin’)
- ▶ We will use Membership Lemma **Case (f)**:
 $(\forall \chi)Q \in \Gamma^*$ iff for each constant c , $Q[c/\chi] \in \Gamma^*$

Induction on QL' : Case 6 (universal quantifier)

- ▶ **Case 6:** \mathcal{P} has the form $(\forall \chi)Q$
(warning: “ Q ” is not a sentence, so it can’t be clubbin’)
 - ▶ We will use Membership Lemma **Case (f)**:
 $(\forall \chi)Q \in \Gamma^*$ iff for each constant c , $Q[c/\chi] \in \Gamma^*$
- \Rightarrow Assume $\mathfrak{M}^* \models (\forall \chi)Q$. Show that $(\forall \chi)Q \in \Gamma^*$

Induction on QL' : Case 6 (universal quantifier)

- ▶ **Case 6:** \mathcal{P} has the form $(\forall \chi) \mathcal{Q}$
(warning: “ \mathcal{Q} ” is not a sentence, so it can’t be clubbin’)
 - ▶ We will use Membership Lemma **Case (f)**:
 $(\forall \chi) \mathcal{Q} \in \Gamma^*$ iff for each constant c , $\mathcal{Q}[c/\chi] \in \Gamma^*$
- \Rightarrow Assume $\mathfrak{M}^* \models (\forall \chi) \mathcal{Q}$. Show that $(\forall \chi) \mathcal{Q} \in \Gamma^*$
- Then every object satisfies \mathcal{Q} , so every substitution instance for every constant is true in \mathfrak{M}^* (since each object is named by itself)

Induction on QL' : Case 6 (universal quantifier)

- ▶ **Case 6:** \mathcal{P} has the form $(\forall \chi)Q$
(warning: “ Q ” is not a sentence, so it can’t be clubbin’)
 - ▶ We will use Membership Lemma **Case (f)**:
 $(\forall \chi)Q \in \Gamma^*$ iff for each constant c , $Q[c/\chi] \in \Gamma^*$
- \Rightarrow Assume $\mathfrak{M}^* \models (\forall \chi)Q$. Show that $(\forall \chi)Q \in \Gamma^*$
- Then every object satisfies Q , so every substitution instance for every constant is true in \mathfrak{M}^* (since each object is named by itself)
 - These $Q[c/\chi]$ are clubbin’ by the IH, so they all belong to Γ^* .
So then by Membership Lemma case (f), $(\forall \chi)Q \in \Gamma^*$

Induction on QL' : Case 6 (universal quantifier)

- ▶ **Case 6:** \mathcal{P} has the form $(\forall \chi)Q$
(warning: “ Q ” is not a sentence, so it can’t be clubbin’)
 - ▶ We will use Membership Lemma **Case (f)**:
 $(\forall \chi)Q \in \Gamma^*$ iff for each constant c , $Q[c/\chi] \in \Gamma^*$
- \Rightarrow Assume $\mathfrak{M}^* \models (\forall \chi)Q$. Show that $(\forall \chi)Q \in \Gamma^*$
- Then every object satisfies Q , so every substitution instance for every constant is true in \mathfrak{M}^* (since each object is named by itself)
 - These $Q[c/\chi]$ are clubbin’ by the IH, so they all belong to Γ^* .
So then by Membership Lemma case (f), $(\forall \chi)Q \in \Gamma^*$
- \Leftarrow Assume $(\forall \chi)Q \in \Gamma^*$. Show that $\mathfrak{M}^* \models (\forall \chi)Q$

Induction on QL' : Case 6 (universal quantifier)

- ▶ **Case 6:** \mathcal{P} has the form $(\forall \chi)Q$
(warning: “ Q ” is not a sentence, so it can’t be clubbin’)
 - ▶ We will use Membership Lemma **Case (f)**:
 $(\forall \chi)Q \in \Gamma^*$ iff for each constant c , $Q[c/\chi] \in \Gamma^*$
- \Rightarrow Assume $\mathfrak{M}^* \models (\forall \chi)Q$. Show that $(\forall \chi)Q \in \Gamma^*$
- Then every object satisfies Q , so every substitution instance for every constant is true in \mathfrak{M}^* (since each object is named by itself)
 - These $Q[c/\chi]$ are clubbin’ by the IH, so they all belong to Γ^* .
So then by Membership Lemma case (f), $(\forall \chi)Q \in \Gamma^*$
- \Leftarrow Assume $(\forall \chi)Q \in \Gamma^*$. Show that $\mathfrak{M}^* \models (\forall \chi)Q$
- Practice this yourself!

Upshots of our Induction

- ▶ Having handled every case (in spirit), we conclude that every sentence of QL' is clubbin':

Upshots of our Induction

- ▶ Having handled every case (in spirit), we conclude that every sentence of QL' is clubbin':
- ▶ For all QL' -sentences \mathcal{P} , $\mathfrak{M}^* \models \mathcal{P}$ iff $\mathcal{P} \in \Gamma^*$

Upshots of our Induction

- ▶ Having handled every case (in spirit), we conclude that every sentence of QL' is clubbin':
- ▶ For all QL' -sentences \mathcal{P} , $\mathfrak{M}^* \models \mathcal{P}$ iff $\mathcal{P} \in \Gamma^*$
- ▶ Hence, the QL' -model \mathfrak{M}^* makes-true every sentence in Γ^* , showing that this set is satisfiable

Upshots of our Induction

- ▶ Having handled every case (in spirit), we conclude that every sentence of QL' is clubbin':
- ▶ For all QL' -sentences \mathcal{P} , $\mathfrak{M}^* \models \mathcal{P}$ iff $\mathcal{P} \in \Gamma^*$
- ▶ Hence, the QL' -model \mathfrak{M}^* makes-true every sentence in Γ^* , showing that this set is satisfiable
- ▶ Hence, we have proven the **\exists -C Maximal Consistency Lemma**: every QL' set that is maximally-QND-consistent and \exists -complete is satisfiable in QL'

Upshots of our Induction

- ▶ Having handled every case (in spirit), we conclude that every sentence of QL' is clubbin':
- ▶ For all QL' -sentences \mathcal{P} , $\mathfrak{M}^* \models \mathcal{P}$ iff $\mathcal{P} \in \Gamma^*$
- ▶ Hence, the QL' -model \mathfrak{M}^* makes-true every sentence in Γ^* , showing that this set is satisfiable
- ▶ Hence, we have proven the **\exists -C Maximal Consistency Lemma**: every QL' set that is maximally-QND-consistent and \exists -complete is satisfiable in QL'
- ▶ It remains to prove the **Consistency Lemma**, i.e. that any QND-consistent QL -set (like our O.G. Γ) is satisfiable **in QL !**

From satisfiability of Γ^* to satisfiability of Γ

- We have shown that the maximally-QND-consistent and existentially complete Γ^* is satisfiable in QL'

From satisfiability of Γ^* to satisfiability of Γ

- ▶ We have shown that the maximally-QND-consistent and existentially complete Γ^* is satisfiable in QL'
- ▶ It remains to show that QND-consistent Γ is satisfiable **in QL**

From satisfiability of Γ^* to satisfiability of Γ

- ▶ We have shown that the maximally-QND-consistent and existentially complete Γ^* is satisfiable in QL'
- ▶ It remains to show that QND-consistent Γ is satisfiable **in QL**
- ▶ i.e. we need a QL -model \mathfrak{M} s.t. $\mathfrak{M} \models \Gamma$

From satisfiability of Γ^* to satisfiability of Γ

- ▶ We have shown that the maximally-QND-consistent and existentially complete Γ^* is satisfiable in QL'
- ▶ It remains to show that QND-consistent Γ is satisfiable **in QL**
- ▶ i.e. we need a QL-model \mathfrak{M} s.t. $\mathfrak{M} \models \Gamma$
- ▶ **Hopes and dreams:** by construction $\Gamma \subset \Gamma^*$, so $\mathfrak{M}^* \models \Gamma$ in QL' .
But how are we to get a QL-model for Γ from this?????

From satisfiability of Γ^* to satisfiability of Γ

- ▶ We have shown that the maximally-QND-consistent and existentially complete Γ^* is satisfiable in QL'
- ▶ It remains to show that QND-consistent Γ is satisfiable **in QL**
- ▶ i.e. we need a QL-model \mathfrak{M} s.t. $\mathfrak{M} \models \Gamma$
- ▶ **Hopes and dreams**: by construction $\Gamma \subset \Gamma^*$, so $\mathfrak{M}^* \models \Gamma$ in QL' . But how are we to get a QL-model for Γ from this?????
- ▶ **Salvation**: note that the model \mathfrak{M}^* we constructed is *not only* a QL' model for Γ^* *BUT ALSO* a QL-model for Γ !

From satisfiability of Γ^* to satisfiability of Γ

- ▶ We have shown that the maximally-QND-consistent and existentially complete Γ^* is satisfiable in QL'
- ▶ It remains to show that QND-consistent Γ is satisfiable **in QL**
- ▶ i.e. we need a QL-model \mathfrak{M} s.t. $\mathfrak{M} \models \Gamma$
- ▶ **Hopes and dreams**: by construction $\Gamma \subset \Gamma^*$, so $\mathfrak{M}^* \models \Gamma$ in QL' .
But how are we to get a QL-model for Γ from this?????
- ▶ **Salvation**: note that the model \mathfrak{M}^* we constructed is *not only* a QL' model for Γ^* *BUT ALSO* a QL-model for Γ !
- ▶ Since the language of QL is contained in QL' , $\mathfrak{M}^* := (D, I^*)$ maps all symbols of QL to objects in D

From satisfiability of Γ^* to satisfiability of Γ

- ▶ We have shown that the maximally-QND-consistent and existentially complete Γ^* is satisfiable in QL'
- ▶ It remains to show that QND-consistent Γ is satisfiable **in QL**
- ▶ i.e. we need a QL-model \mathfrak{M} s.t. $\mathfrak{M} \models \Gamma$
- ▶ **Hopes and dreams:** by construction $\Gamma \subset \Gamma^*$, so $\mathfrak{M}^* \models \Gamma$ in QL' . But how are we to get a QL-model for Γ from this?????
- ▶ **Salvation:** note that the model \mathfrak{M}^* we constructed is *not only* a QL' model for Γ^* *BUT ALSO* a QL-model for Γ !
- ▶ Since the language of QL is contained in QL' , $\mathfrak{M}^* := (D, I^*)$ maps all symbols of QL to objects in D
- ▶ If you like, you can define a QL-model $\mathfrak{M} := (D, I)$ s.t. I is the restriction of I^* to unprimed constants in QL. Then $\mathfrak{M} \models \Gamma$.

From satisfiability of Γ^* to satisfiability of Γ

- ▶ We have shown that the maximally-QND-consistent and existentially complete Γ^* is satisfiable in QL'
- ▶ It remains to show that QND-consistent Γ is satisfiable **in QL**
- ▶ i.e. we need a QL-model \mathfrak{M} s.t. $\mathfrak{M} \models \Gamma$
- ▶ **Hopes and dreams:** by construction $\Gamma \subset \Gamma^*$, so $\mathfrak{M}^* \models \Gamma$ in QL' . But how are we to get a QL-model for Γ from this?????
- ▶ **Salvation:** note that the model \mathfrak{M}^* we constructed is *not only* a QL' model for Γ^* *BUT ALSO* a QL-model for Γ !
- ▶ Since the language of QL is contained in QL' , $\mathfrak{M}^* := (D, I^*)$ maps all symbols of QL to objects in D
- ▶ If you like, you can define a QL-model $\mathfrak{M} := (D, I)$ s.t. I is the restriction of I^* to unprimed constants in QL. Then $\mathfrak{M} \models \Gamma$.
- ▶ \square Q.E.D. motha f***ers!!! (i.e. *quod erat demonstrandum*)