

Oneway

Notes

Output Created		28-FEB-2022 22:38:54
Comments		
Input	Data	/Users/benjamin/Desktop/AP Research/21-22-PAS-AP-Research/Experiment 3/E3-Raw/E3-SA.csv
	Active Dataset	DataSet7
	Filter	<none>
	Weight	<none>
	Split File	<none>
	N of Rows in Working Data File	25
Missing Value Handling	Definition of Missing	User-defined missing values are treated as missing.
	Cases Used	Statistics for each analysis are based on cases with no missing data for any variable in the analysis.
Syntax		ONEWAY Difference BY ReferenceNum /ES=OVERALL /STATISTICS HOMOGENEITY /MISSING ANALYSIS /CRITERIA=CILEVEL (0.95) /POSTHOC=TUKEY ALPHA(0.05).
Resources	Processor Time	00:00:00.02
	Elapsed Time	00:00:00.00

Tests of Homogeneity of Variances

		Levene Statistic	df1	df2	Sig.
Difference	Based on Mean	2.554	4	20	.071
	Based on Median	1.679	4	20	.194
	Based on Median and with adjusted df	1.679	4	14.134	.210
	Based on trimmed mean	2.545	4	20	.071

ANOVA

Difference

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	.014	4	.004	28049.532	<.001
Within Groups	.000	20	.000		
Total	.014	24			

ANOVA Effect Sizes^a

		Point Estimate	95% Confidence Interval	
			Lower	Upper
Difference	Eta-squared	1.000	1.000	1.000
	Epsilon-squared	1.000	.999	1.000
	Omega-squared Fixed-effect	1.000	.999	1.000
	Omega-squared Random-effect	.999	.998	.999

a. Eta-squared and Epsilon-squared are estimated based on the fixed-effect model.

Post Hoc Tests

Multiple Comparisons

Dependent Variable: Difference

Tukey HSD

(I) ReferenceNum	(J) ReferenceNum	Mean Difference (I-J)	Std. Error	Sig.	95% ... Lower Bound
1	2	-.0458800 *	.0002252	<.001	-.046554
	3	-.0387400 *	.0002252	<.001	-.039414
	4	-.0460200 *	.0002252	<.001	-.046694
	5	.0090000 *	.0002252	<.001	.008326
2	1	.0458800 *	.0002252	<.001	.045206
	3	.0071400 *	.0002252	<.001	.006466
	4	-.0001400	.0002252	.970	-.000814
	5	.0548800 *	.0002252	<.001	.054206
3	1	.0387400 *	.0002252	<.001	.038066
	2	-.0071400 *	.0002252	<.001	-.007814
	4	-.0072800 *	.0002252	<.001	-.007954
	5	.0477400 *	.0002252	<.001	.047066
4	1	.0460200 *	.0002252	<.001	.045346
	2	.0001400	.0002252	.970	-.000534

Multiple Comparisons

Dependent Variable: Difference

Tukey HSD

		95% ...
(I) ReferenceNum	(J) ReferenceNum	Upper Bound
1	2	-.045206
	3	-.038066
	4	-.045346
	5	.009674
2	1	.046554
	3	.007814
	4	.000534
	5	.055554
3	1	.039414
	2	-.006466
	4	-.006606
	5	.048414
4	1	.046694
	2	.000814

Multiple Comparisons

Dependent Variable: Difference

Tukey HSD

		Mean			95% ...
(I) ReferenceNum	(J) ReferenceNum	Difference (I-J)	Std. Error	Sig.	Lower Bound
	3	.0072800 *	.0002252	<.001	.006606
	5	.0550200 *	.0002252	<.001	.054346
5	1	-.0090000 *	.0002252	<.001	-.009674
	2	-.0548800 *	.0002252	<.001	-.055554
	3	-.0477400 *	.0002252	<.001	-.048414
	4	-.0550200 *	.0002252	<.001	-.055694

Multiple Comparisons

Dependent Variable: Difference

Tukey HSD

95% ...

(I) ReferenceNum	(J) ReferenceNum	Upper Bound
5	3	.007954
	5	.055694
	1	-.008326
	2	-.054206
	3	-.047066
	4	-.054346

*. The mean difference is significant at the 0.05 level.

Homogeneous Subsets

Difference

Tukey HSD^a

ReferenceNum	N	Subset for alpha = 0.05			
		1	2	3	4
5	5	-.069820			
1	5		-.060820		
3	5			-.022080	
2	5				-.014940
4	5				-.014800
Sig.		1.000	1.000	1.000	.970

Means for groups in homogeneous subsets are displayed.

a. Uses Harmonic Mean Sample Size = 5.000.