## This function takes three inputs

x - a set of parameters t - the number of time-steps you wish to simulate data - actual data that you are attempting to fit

```
function f = sliroutput_big(x,t,data)
% set up transmission constants
k_{infections} = x(1);
k_{fatality} = x(2);
k_recover = x(3);
k lockdown = x(4);
k_vac = .3;
% set up initial conditions
ic\_susc = x(5);
ic_{lockdown} = x(6);
ic inf = x(7);
ic\_rec = x(8);
ic_fatality = x(9);
% Set up SIRD within-population transmission matrix
A = [1-(k infections + k lockdown + k vac) 0.4 0 0 0 0 0 0 0 0 0 0 0 0;
   k_lockdown 0.6 0.5 0 0 0 0 0 0 0 0 0 0;
   k infections 0 1-(0.5+k fatality + k recover) 0 0 0 0 0 0 0 0 0 0 0;
   k_vac 0 k_recover 1 0 0 0 0 0 0 0 0 0 0;
    0 0 k_fatality 0 1 0 0 0 0 0 0 0 0 0;
    0 0 0 0 1-(k_infections + k_lockdown + k_vac) 0.4 0 0 0 0 0 0 0;
    0 0 0 0 0 k lockdown 0.6 0.5 0 0 0 0 0 0;
    0 0 0 0 0 k_infections 0 1-(0.5+k_fatality + k_recover) 0 0 0 0 0 0 0;
    0 0 0 0 0 k_vac 0 k_recover 1 0 0 0 0 0;
    0 0 0 0 0 0 0 k_fatality 0 1 0 0 0 0;
    0 0 0 0 0 0 0 0 0 0 1-(k_infections + k_lockdown + k_vac) 0.4 0 0 0;
    0 0 0 0 0 0 0 0 0 0 k lockdown 0.6 0.5 0 0;
    0 0 0 0 0 0 0 0 0 0 k_infections 0 1-(0.5+k_fatality + k_recover) 0 0;
    0 0 0 0 0 0 0 0 0 0 k vac 0 k recover 1 0;
    0 0 0 0 0 0 0 0 0 0 0 0 k_fatality 0 1];
B = zeros(15,1);
% Set up the vector of initial conditions
x0 = [ic_susc, ic_lockdown, ic_inf, ic_rec, ic_fatality ic_susc, ic_lockdown,
 ic_inf, ic_rec, ic_fatality ic_susc, ic_lockdown, ic_inf, ic_rec,
 ic_fatality];
% simulate the SIRD model for t time-steps
sys sir base = ss(A,B,eye(15),zeros(15,1),1);
y = lsim(sys\_sir\_base, zeros(t,1), linspace(0,t-1,t), x0);
% return a "cost". This is the quantitity that you want your model to
% minimize. Basically, this should encapsulate the difference between your
% modeled data and the true data. Norms and distances will be useful here.
% Hint: This is a central part of this case study! choices here will have
```

```
% a big impact!

y2 = y(: , [4,5]);
y3 = cumsum(y2);
f = norm(data - y3); % norm of distance.

end

Not enough input arguments.

Error in sliroutput_big (line 9)
k_infections = x(1);
```

Published with MATLAB® R2021b