

Orthogonal complement

In the mathematical fields of linear algebra and functional analysis, the **orthogonal complement** of a subspace *W* of a vector space *V* equipped with a bilinear form *B* is the set *W*[⊥] of all vectors in *V* that are orthogonal to every vector in *W*. Informally, it is called the **perp**, short for **perpendicular complement**. It is a subspace of *V*.

Contents

- Example
- General bilinear forms
 - Properties
- Inner product spaces
 - Properties
 - Finite dimensions
- Banach spaces
- Applications
- See also
- References
- External links

Example

In the case that *W* is the subspace of *V* = ℝ⁵ (with the usual dot product) spanned by the rows of the next matrix,

$$\left(\begin{array}{cc|cc} 1 & 0 & 2 & 3 & 5 \\ 0 & 1 & 6 & 9 & 3 \end{array}\right)$$

its orthogonal complement *W*[⊥] is spanned by the three row-vectors of

$$\left(\begin{array}{cc|cc} -2 & -6 & 1 & 0 & 0 \\ -3 & -9 & 0 & 1 & 0 \\ -5 & -3 & 0 & 0 & 1 \end{array}\right).$$

The fact that every vector on the first list is orthogonal to every vector on the second list can be checked by direct computation. The fact that the spans of these vectors are orthogonal then follows by bilinearity of the dot product. Finally, the fact that these spaces are orthogonal complements follows from the dimension relationships given below.

General bilinear forms

Let V be a vector space over a field F equipped with a bilinear form B . We define u to be left-orthogonal to v , and v to be right-orthogonal to u , when $B(u, v) = 0$. For a subset W of V we define the left orthogonal complement W^\perp to be

$$W^\perp = \{x \in V : B(x, y) = 0 \text{ for all } y \in W\}.$$

There is a corresponding definition of right orthogonal complement. For a reflexive bilinear form, where $B(u, v) = 0$ implies $B(v, u) = 0$ for all u and v in V , the left and right complements coincide. This will be the case if B is a symmetric or an alternating form.

The definition extends to a bilinear form on a free module over a commutative ring, and to a sesquilinear form extended to include any free module over a commutative ring with conjugation.^[1]

Properties

- An orthogonal complement is a subspace of V ;
- If $X \subset Y$ then $X^\perp \supset Y^\perp$;
- The radical V^\perp of V is a subspace of every orthogonal complement;
- $W \subset (W^\perp)^\perp$;
- If B is non-degenerate and V is finite-dimensional, then $\dim(W) + \dim(W^\perp) = \dim V$.
- If L_1, L_2, \dots, L_r are subspaces of a finite-dimensional space V and $L_* = L_1 \cap L_2 \cap \dots \cap L_r$, then $L_*^\perp = L_1^\perp + L_2^\perp + \dots + L_r^\perp$.

Inner product spaces

This section considers orthogonal complements in inner product spaces.^[2]

Properties

The orthogonal complement is always closed in the metric topology. In finite-dimensional spaces, that is merely an instance of the fact that all subspaces of a vector space are closed. In infinite-dimensional Hilbert spaces, some subspaces are not closed, but all orthogonal complements are closed. In such spaces, the orthogonal complement of the orthogonal complement of W is the closure of W , i.e.,

$$(W^\perp)^\perp = \overline{W}.$$

Some other useful properties that always hold are the following. Let H be a Hilbert space and let X and Y be its linear subspaces. Then:

- $X^\perp = \overline{X}^\perp$;
- if $Y \subset X$, then $X^\perp \subset Y^\perp$;
- $X \cap X^\perp = \{0\}$;
- $X \subseteq (X^\perp)^\perp$;
- if X is a closed linear subspace of H , then $(X^\perp)^\perp = X$;
- if X is a closed linear subspace of H , then $H = X \oplus X^\perp$, the (inner) direct sum.

The orthogonal complement generalizes to the annihilator, and gives a Galois connection on subsets of the inner product space, with associated closure operator the topological closure of the span.

Finite dimensions

For a finite-dimensional inner product space of dimension *n*, the orthogonal complement of a *k*-dimensional subspace is an (*n* − *k*)-dimensional subspace, and the double orthogonal complement is the original subspace:

$$(W^\perp)^\perp = W.$$

If *A* is an *m* × *n* matrix, where Row *A*, Col *A*, and Null *A* refer to the row space, column space, and null space of *A* (respectively), we have

$$\begin{aligned}(\text{Row } A)^\perp &= \text{Null } A \\ (\text{Col } A)^\perp &= \text{Null } A^\mathsf{T}.\end{aligned}$$

Banach spaces

There is a natural analog of this notion in general Banach spaces. In this case one defines the orthogonal complement of *W* to be a subspace of the dual of *V* defined similarly as the annihilator

$$W^\perp = \{ \boldsymbol{x} \in V^* : \forall \boldsymbol{y} \in W, \boldsymbol{x}(\boldsymbol{y}) = 0 \}.$$

It is always a closed subspace of *V**. There is also an analog of the double complement property. *W*^{⊥⊥} is now a subspace of *V*** (which is not identical to *V*). However, the reflexive spaces have a natural isomorphism *i* between *V* and *V***. In this case we have

$$i\overline{W} = W^{\perp\perp}.$$

This is a rather straightforward consequence of the Hahn–Banach theorem.

Applications

In special relativity the orthogonal complement is used to determine the simultaneous hyperplane at a point of a world line. The bilinear form η used in Minkowski space determines a pseudo-Euclidean space of events. The origin and all events on the light cone are self-orthogonal. When a time event and a space event evaluate to zero under the bilinear form, then they are hyperbolic-orthogonal. This terminology stems from the use of two conjugate hyperbolas in the pseudo-Euclidean plane: conjugate diameters of these hyperbolas are hyperbolic-orthogonal.

See also

- Complemented lattice

References

1. Adkins & Weintraub (1992) p.359
 2. Adkins&Weintraub (1992) p.272
- Adkins, William A.; Weintraub, Steven H. (1992), *Algebra: An Approach via Module Theory*, Graduate Texts in Mathematics, **136**, Springer-Verlag, ISBN 3-540-97839-9, Zbl 0768.00003 (<https://zbmath.org/?format=complete&q=an:0768.00003>)
 - Halmos, Paul R. (1974), *Finite-dimensional vector spaces*, Undergraduate Texts in Mathematics, Berlin, New York: Springer-Verlag, ISBN 978-0-387-90093-3, Zbl 0288.15002 (<https://zbmath.org/?format=complete&q=an:0288.15002>)
 - Milnor, J.; Husemoller, D. (1973), *Symmetric Bilinear Forms*, Ergebnisse der Mathematik und ihrer Grenzgebiete, **73**, Springer-Verlag, ISBN 3-540-06009-X, Zbl 0292.10016 (<https://zbmath.org/?format=complete&q=an:0292.10016>)

External links

- Instructional video describing orthogonal complements (Khan Academy) (<http://www.khanacademy.org/math/linear-algebra/alternate-bases/othogonal-complements/v/linear-algebra-orthogonal-complements>)
-

Retrieved from "https://en.wikipedia.org/w/index.php?title=Orthogonal_complement&oldid=844368577"

This page was last edited on 4 June 2018, at 13:27.

Text is available under the Creative Commons Attribution-ShareAlike License; additional terms may apply. By using this site, you agree to the Terms of Use and Privacy Policy. Wikipedia® is a registered trademark of the Wikimedia Foundation, Inc., a non-profit organization.