

An in Depth Look at Burnside's Lemma and Its Applications

Ben Drummond

1018642

Friday April 2, 2021

Introduction

In this exposition we will be exploring Burnside's lemma along with some related theories and consequences of Burnside's lemma. We will also investigate some applications of Burnside's lemma and solve some examples.

Despite its name Burnside's lemma was not discovered by Burnside, and was known by Cauchy in 1945. Burnside included the formula and a subsequent proof in his book "Theory of Groups of Finite Order" and attributed it to Frobenius, as a result the lemma is sometimes referred to as "the lemma that is not Burnside's".

Burnside's lemma is a method of counting the number of orbits of a group acting on a finite set, and in plain English it states that the number of orbits is equal to the average number of fixed points of the group. In mathematical notation Burnside's lemma is defined by the equation.

$$|X/G| = \frac{1}{|G|} \sum_{g \in G} |X^g|$$

Where G is a group acting on a finite set X , with $|X/G|$ denoting the number of orbits and X^g denoting the set of points in X that are fixed by g .

You'll see in the examples later on that by using Burnside's lemma we are able to account for symmetry when counting objects. This is because any two elements that

are symmetrical are contained in the same orbit and thus the number of orbits is equal to the number of rotationally unique elements.

Before we can dive into the proof of Burnside's lemma and its applications we will first look at the orbit-stabilizer theorem as it will be used in the proof of Burnside's lemma and in the our examples.

The Orbit-Stabilizer Theorem

The orbit-stabilizer theorem is a useful way to determine the order of group G acting on a finite set X by choosing some element $x \in X$ and by examining its orbit and stabilizer. The orbit-stabilizer theorem states that if G is a group acting on a finite set X , then for any $x \in X$ the following must be true:

$$|G| = |Orb(x)||G_x|$$


Since this may or may not be the expected outcome I will provide an example to perhaps make this result a bit more intuitive.

Example 1: How many symmetries are there on a regular pentagon?

Another way of wording this question is "if you were given a pentagon, how many different orientations could you place it down in so that it occupies the same space?"

We will start off this question by numbering our pentagon, we can either number the

vertices or edges but in the diagram below I have chosen to number the vertices.



Now with some intuition from Gowers's [Gow11] we pick a vertex on the pentagon and choose where we want to place it. Note that we have 5 options for where we can place down our vertex, in this case lets choose to place our vertex in the 1 position. Now that our vertex is fixed notice that the pentagon can still be reflected about a line through 1 and the mid point of the edge between 4 and 5, this gives us 2 possible options for each vertex for a total of $5 \cdot 2 = 10$ ways to orient the pentagon.

In this example the 5 options we get from fixing the vertex is equivalent to the orbit of x and the 2 additional options on each vertex is equivalent to the stabilizer of x .

Proof of Burnside's Lemma

In this section we will be looking at a proof of Burnside's lemma; the proof presented closely follows the proof found on the Wolfram page on Burnside's lemma [Wei].

Let G be a finite group that acts on the set X . Earlier we mentioned that X^g and G_x are almost complementary, and there is a good reason for this. We know that :

$$\begin{aligned}\sum_{g \in G} |X^g| &= \sum_{g \in G} |\{x \in X : gx = x\}| \\ \sum_{x \in X} |G_x| &= \sum_{x \in X} |\{g \in G : gx = x\}| \end{aligned}$$

It's not hard to see that these are the same sums over different indices which means

$$\frac{1}{|G|} \sum_{g \in G} |X^g| = \frac{1}{|G|} \sum_{x \in X} |G_x|$$

Using the orbit stabilizer theorem we get

$$\begin{aligned}\frac{1}{|G|} \sum_{g \in G} |X^g| &= \frac{1}{|G|} \sum_{x \in X} \frac{|G|}{|Orb(x)|} \\ &= \sum_{x \in X} \frac{1}{|Orb(x)|} \end{aligned}$$

Since no $x \in X$ can be in more than one orbit, and the orbit of x can't be empty we find that X is the disjoint union of all its orbits denoted X/G . With this in mind we can break the sum into the following:

$$\begin{aligned}
\sum_{x \in X} \frac{1}{|Orb(x)|} &= \sum_{Orb(x) \in X/G} \left(\sum_{x \in Orb(x)} \frac{1}{|Orb(x)|} \right) \\
&= \sum_{Orb(x) \in X/G} 1 \\
&= |X/G|
\end{aligned}$$


We've arrived at Burnside's lemma.

$$\frac{1}{|G|} \sum_{g \in G} |X^g| = |X/G|$$

Examples

Example 2: Given two different colours of beads, how many unique bracelets of size 8 can you make?

The first step in solving this problem is determining the order of the group acting on the bracelet. We will use the graph on 8 vertices below to represent the bracelet because its vertices are connected in the same way as a bracelet with 8 beads.




Let X be the set of all possible bracelets with two colours of size 8, and let G be the group of symmetries that act on X . The first thing we must do is calculate the order of G using the orbit-stabilizer theorem.

Since G is the group of symmetries; rather than acting on the set of all possible colourings we can instead act on the set of vertices which we will denote Y . Looking at our diagram it's clear that any vertex $y \in Y$ can be rotated to any other vertex, and the only two symmetries that fix y are the identity element and a reflection on the line through y and its opposite vertex. So it follows that $Orb(y) = 8$ and $G_y = 2$, and as a corollary the elements of G can be generated by a rotation and a reflection. Applying the orbit-stabilizer theorem we find that $|G| = 8 \cdot 2 = 16$.

Next we need to find X^g for each $g \in G$, to specify actions in G we will write them in terms of a $\frac{1}{8}$ th clockwise rotation which we will call r and a reflection on the line

in the diagram below which we will call t .



Recall that a group G acting on X is a group homomorphism from G to the set of possible permutations on X , so it follows that $\phi(g) \in S(X)$. This means that r and t can be expressed as the following permutations of X .

$$r = (1\ 2\ 3\ 4\ 5\ 6\ 7\ 8)$$

$$t = (1\ 2)(3\ 8)(4\ 7)(5\ 6)$$

By expressing a group action in terms of permutation cycles, the elements of X that

are fixed by the group action become much more clear.

Take for example X^r , we can see that r takes the first vertex to the second vertex, this means that vertex 1 and 2 must be the same colour. Continuing on we see that r takes the second vertex to the third vertex, and so forth. Eventually we find that every vertex must have the same colour and since there are only 2 colours this means that $|X^r| = 2$.

Investigating further we find that if $x \in X^g$ then x must be coloured in such a way that each cycle in $\phi(g)$ contains vertices of the same colour. Now lets calculate the cycles for the elements in G .

g	Cycle	X^g	g	Cycle	X^g
e	(1)(2)(3)(4)(5)(6)(7)(8)	2^8	t	(1 2)(3 8)(4 7)(5 6)	2^4
r	(1 2 3 4 5 6 7 8)	2^1	rt	(1)(5)(2 8)(3 7)(4 6)	2^5
r^2	(1 3 5 7)(2 4 6 8)	2^2	r^2t	(1 8)(2 7)(3 6)(4 5)	2^4
r^3	(1 4 7 2 5 8 3 6)	2^1	r^3t	(4)(8)(1 7)(2 6)(3 5)	2^5
r^4	(1 5)(2 6)(3 7)(4 8)	2^4	r^4t	(1 6)(2 5)(3 4)(7 8)	2^4
r^5	(1 6 3 8 5 2 7 4)	2^1	r^5t	(7)(3)(1 5)(2 4)(6 8)	2^5
r^6	(1 7 5 3)(2 8 6 4)	2^2	r^6t	(1 4)(2 3)(5 8)(6 7)	2^4
r^7	(1 8 7 6 5 4 3 2)	2^1	r^7t	(2)(6)(1 3)(4 8)(5 7)	2^5

Plugging the values from the table below into Burnside's lemma we get the following

equation

$$\begin{aligned}|X/G| &= \frac{2^8 + 4 \cdot 2^1 + 2 \cdot 2^2 + 5 \cdot 2^4 + 4 \cdot 2^5}{16} \\ &= 30\end{aligned}$$

So with 8 beads of two colours there are a 30 possible bracelets that are unique.

Example 3: What if in the previous example we had 3 colours rather than 2 how many unique bracelets could you make?

In this problem the only difference is that the vertices in each cycle can be 3 three possible colours rather than 2. So the formula for the number of orbits is

$$\begin{aligned}|X/G| &= \frac{3^8 + 4 \cdot 3^1 + 2 \cdot 3^2 + 5 \cdot 3^4 + 4 \cdot 3^5}{16} \\ &= 498\end{aligned}$$

Example 4: Given two different colours of beads, how many unique necklaces of size 15 can you make?

This problem is similar to the previous problems, the only difference is that necklaces aren't symmetric under reflection. This problem is much easier to solve, especially if the number of beads is prime but we will get to that in the next example.

Since the necklace only has 2D rotational symmetry and 15 vertices, by inspection


we can determine that $|G| = 15$ and $G = \langle r \rangle$ where r is a $\frac{1}{15}$ th rotation.

Because the group G is generated by a single element it must be a cyclic group, an isomorphic to \mathbb{Z}_{15} . Note that 3 and 5 are prime divisors of $|G|$, which means that G must also have cyclic subgroups of prime order.

Suppose G is acting on a regular polygon with a set X of 15 vertices.

For any given rotation r^n we know that if n is co-prime to 15 then the $\text{LCM}(n, 15) = n \cdot 15$ which means that r^n is only equal to the identity after 15 permutations and must have a cycle of 15. This is true for $n \in \{1, 2, 4, 7, 8, 11, 13, 14\}$.

Now if $n \in \{3, 6, 9, 12\}$ then r^n contains three 5-cycles. As you can see in the figures below r^3 and r^6



This isn't much of a surprise as $3 \cdot 5 = 6 \cdot 5 = 9 \cdot 5 = 12 \cdot 5 = 0$ under addition mod 15.

Following the same logic as before, if $n \in \{5, 10\}$ then r^n contains five 3-cycles.

Making sure we don't forget the identity e with 15 fixed points were ready to move on.

For two different colours of beads, we end up with 2 choices per cycle in each symmetry of G . Plugging into Burnside's lemma we get

$$\begin{aligned} |X/G| &= \frac{8 \cdot 2^1 + 4 \cdot 2^3 + 2 \cdot 2^5 + 1 \cdot 2^{15}}{15} \\ &= 2192 \end{aligned}$$

Example 4: Given two different colours of beads, how many unique necklaces of size 31 can you make?

Using the same logic as the previous question we come to the conclusion that G is a cyclic group acting on a set of 31 vertices X . However in this case we find that $|G| = 31$ which is prime. This means that G has no non-trivial subgroups and as a consequence every possible permutation in G aside from the identity has a cycle of

size 31. This leaves us with the equation

$$\begin{aligned}|X/G| &= \frac{30 \cdot 2^1 + 1 \cdot 2^{31}}{31} \\ &= 69273668\end{aligned}$$

References

- [Gow11] Timothy Gowers. Group actions ii: The orbit-stabilizer theorem. <https://gowers.wordpress.com/2011/11/09/group-actions-ii-the-orbit-stabilizer-theorem>, Nov 2011.
- [Wei] Eric W Weisstein. Cauchy-frobenius lemma. <https://mathworld.wolfram.com/Cauchy-FrobeniusLemma.html>.