A subcritical phase for isotropic Poisson line Cox percolation: a conditional proof for refereeing

Self-contained document

Christian Hirsch* Benedikt Jahnel†

9th November 2026

Abstract

We present, in one place and without reference to our working notes, the argument for the existence of a subcritical phase in the isotropic Poisson line Cox Boolean model: for small point intensity λ the union of unit disks centred at the points of a Cox process on a Poisson line process of intensity μ does not percolate. The argument has three parts. The skeleton bound reduces non-percolation to a statement about the line process alone, the thick-cluster lemma (T), which controls the expected weight of chains of clusters of boxes met by at least j_0 lines; this part is proved completely. The thick-cluster lemma is reduced to a polynomial bound (S) on the diameter of a single thick cluster; this reduction is a first moment over chains whose dependence structure we have not controlled. We prove that the conditioning on a family of clusters can be replaced by adding a minimal witness of that family as fixed lines, we show that the naive independence hypothesis is false because clusters can relay lines to their neighbours, and we state the corrected hypothesis, in which relays appear explicitly, as a claim. The single-cluster bound is attacked by a multiscale recursion with a two-parameter induction hypothesis carrying fixed lines with a coverage weight; the geometric and probabilistic lemmas of the recursion are proved in full, including a coverage lemma for random lines along a segment, and the recursion closes modulo three statements, on the weights of shared lines, on crossings that ride fixed lines, and on the base case, which are stated as claims with the arguments we have for them. In the course of writing we found and corrected five errors in our working notes, two of which concern the definition of the objects; they are listed in the final section. Every unproved step is displayed in a box, so that the document can be checked for the correctness of what is proved and for the precise content of what is not. We do not claim the unconditional theorem.

Contents

1 Setting, result and structure of the argument **2**

2 Preliminaries on the line process **4**

*Institut für Mathematische Stochastik, Georg-August-Universität Göttingen, Goldschmidtstraße 7, 37077 Göttingen, Germany. christian.hirsch@uni-goettingen.de. ORCID 0000-0003-4136-3740.

†Institut für Mathematische Stochastik, Technische Universität Braunschweig, Universitätsplatz 2, 38106 Braunschweig, Germany, and Weierstrass Institute for Applied Analysis and Stochastics, Mohrenstraße 39, 10117 Berlin, Germany. benedikt.jahnel@tu-braunschweig.de. ORCID 0000-0002-4212-0065.

3 The skeleton bound: from (T) to non-percolation 5

4 From the single-cluster bound to the thick-cluster lemma 7

5 The single-cluster bound: multiscale setting and the proved lemmas 16

6 The multiscale recursion and its claims 22

7 Status, corrections and what a referee should check 26

1 Setting, result and structure of the argument

1.1 The model

Let $\mathcal{L}$ be a Poisson line process in $\mathbb{R}^2$ with intensity measure

$$\nu(\,\mathrm{d}\theta\;\mathrm{d}p) = \frac{\mu}{\pi}\;\mathrm{d}\theta\;\mathrm{d}p, \qquad \theta \in [0,\pi),\ p \in \mathbb{R}\,, \tag{1.1}$$

where a line is parametrised by its normal direction θ and its signed distance p from the origin. With this normalisation the number of lines meeting a convex set A is Poisson with mean $\mu \cdot \mathrm{per}(A)/\pi$, where per is the perimeter; in particular the mean number of lines meeting a segment of length t is $2\mu t/\pi$, and the mean number meeting a disk of radius r is $2\mu r$. Given $\mathcal{L}$, let X_λ be a Poisson process with intensity λ per unit length on every line of $\mathcal{L}$, and let

$$\Xi_\lambda := \bigcup_{x \in X_\lambda} B(x,1)$$

be the Boolean model with unit disks. The model is said to *percolate* if Ξ_λ has an unbounded connected component with positive probability, and

$$\lambda_c(\mu) := \inf\{\lambda > 0 \colon \Xi_\lambda \text{ percolates}\}\,.$$

By scaling, the choice of unit disks is no restriction. The question addressed here is whether $\lambda_c(\mu) > 0$.

Boxes and paths. For $z \in \mathbb{Z}^2$ let $Q(z) := z + [0,1)^2$ and let $N(z)$ be the number of lines of $\mathcal{L}$ meeting $Q(z)$; it is Poisson with mean $\mu_1 := 4\mu/\pi$. A 2-*path* is a self-avoiding sequence $(z_0, z_1, \dots)$ in $\mathbb{Z}^2$ with $\|z_{i+1} - z_i\|_\infty \le 2$; each site has 24 neighbours in this sense. A set of boxes is 2-*connected* if any two of its elements are joined by a 2-path inside it. Distances between boxes and between sets of boxes are sup-norm distances between the corresponding points of $\mathbb{Z}^2$; the distance between a point of $\mathbb{R}^2$ and a line is Euclidean.

Thin and thick boxes. Fix an integer $j_0 \ge 2$ and let

$$\Theta := \{x \in \mathbb{R}^2 \colon \text{at least } j_0 \text{ lines of } \mathcal{L} \text{ pass within distance 1 of } x\}\,. \tag{1.2}$$

A box z is *thick* if $Q(z) \cap \Theta \neq \varnothing$ and *thin* otherwise. A thin box satisfies $N(z) \le j_0 - 1$, because every line meeting $Q(z)$ passes within distance $\sqrt{2}/2 < 1$ of the centre of $Q(z)$, which is a point of $Q(z)$ not in Θ. Let $T = T_{j_0}$ be the set of thick boxes and $\mathcal{C}$ the set of its 2-connected components, the *thick clusters*; two distinct clusters are at distance at least 3. For a cluster C, $|C|$ is its number of boxes and $\operatorname{diam} C$ its diameter.

Remark 1.1 (Why thickness is defined through Θ). The lattice definition "thick if $N(z) \geq j_0$", used in our working notes, makes hypothesis (T) below false. Consider j_0 lines of slope $s \in (0,1)$ whose heights at a given abscissa lie in an interval of length $w' = 4s$. Over a column of unit boxes each line rises by s, and the j_0 lines meet a common box of that column if and only if an integer lies in an interval of length $1 + s - w' = 1 - 3s$; as the column index advances the relevant position shifts by s, so the common boxes come in runs of about $(1-3s)/s$ columns separated by runs of about three columns without a common box. With 2-adjacency the thick boxes of such a bundle therefore form clusters of about $1/s$ boxes separated by thin gaps of three columns, and a bundle of length D consists of about Ds clusters. The chain weight of Definition 1.2 assigns to it $\prod |C_i| \rho^{g_i} \approx (s^{-1}\rho^3)^{Ds}$, which grows exponentially in D as soon as $s < \rho^3$, while the probability of such a bundle is of order $\mu^{j_0} s^{j_0} D^{2-j_0}$; so Φ_n in (1.4) would tend to infinity. The definition through Θ removes the lattice effect: the set of points within distance 1 of all lines of a bundle is an intersection of strips, hence convex and connected, and the boxes meeting a connected set form a 2-connected set, so a bundle is a single cluster. The occupation bound (3.1) below only uses $N(z) \leq j_0 - 1$ for thin boxes, which still holds.

1.2 The three statements

Definition 1.2 (Cluster chains)**.** A *cluster chain* from a box z_0 is a finite sequence $(C_1, \dots, C_k)$ of *pairwise distinct* thick clusters, $k \geq 0$, together with integers $g_0, \dots, g_k \geq 0$ such that

$$g_0 \geq \tfrac{1}{2}(\operatorname{dist}(z_0, C_1) - 2), \qquad g_i \geq \max\{1, \tfrac{1}{2}(\operatorname{dist}(C_i, C_{i+1}) - 2)\} \quad (1 \leq i < k). \tag{1.3}$$

Its *free advance* is $F := \sum_{i=1}^{k} (\operatorname{diam} C_i + 4)$ and its *paid length* is $G := \sum_{i=0}^{k} g_i$; the chain *reaches* n if $F + 2G \geq n$. For $0 < \rho < 1$ its weight is

$$w_\rho := \prod_{i=1}^{k} |C_i| \prod_{i=0}^{k} \rho^{g_i}.$$

Definition 1.3 (Thick-cluster lemma, hypothesis $(\mathrm{T})_\rho$)**.** $(\mathrm{T})_\rho$ holds if the expected total weight of all chains from 0 with $k \geq 1$ is finite and

$$\Phi_n(\rho) := \mathbb{E}\Big[\sum_{\text{chains from 0 with } F \geq n/2} w_\rho \Big] \xrightarrow[n\to\infty]{} 0. \tag{1.4}$$

Definition 1.4 (Single-cluster bound, hypothesis $(\mathrm{S})_{\beta,A}$)**.** $(\mathrm{S})_{\beta,A}$ holds if for every box z and every $d \geq 1$,

$$\mathbb{P}\big(\exists\, C \in \mathcal{C} \colon z \in C,\ \operatorname{diam} C \geq d\big) \leq A\, d^{-\beta}. \tag{1.5}$$

Theorem 1.5 (Main result, conditional)**.** *Let $j_0 \geq 2$, $0 < \lambda < 1/(576\sqrt{2}\,(j_0 - 1))$ and $\rho := 576\sqrt{2}\,\lambda(j_0 - 1) < 1$.*

(i) *If $(\mathrm{T})_{\sqrt{\rho}}$ holds, then Ξ_λ does not percolate; in particular $\lambda_c(\mu) \geq \rho/(576\sqrt{2}(j_0 - 1))$.*

(ii) *If $(\mathrm{S})_{\beta,A}$ holds with $\beta > q + 1$ and Claim D holds with exponent q, then $(\mathrm{T})_{\rho'}$ holds for every $\rho' < 1$ provided $\mu \leq \mu_0(j_0, \beta, A, \rho')$.*

(iii) *If Claims C1 to C3 hold, then for every $j_0 \geq 7$ there are $\mu_1(j_0) > 0$ and $A'(j_0)$ such that $(\mathrm{S})_{j_0 - 2, A'(j_0)\mu^{j_0}}$ holds for all $\mu \leq \mu_1(j_0)$.*

Consequently, if the four claims hold, then for every $j_0 \geq 7$ and $0 < \rho < 1$ there is $\mu_0 > 0$ such that $\lambda_c(\mu) \geq \rho/(576\sqrt{2}(j_0 - 1))$ for all $\mu \leq \mu_0$.

Part (i) is Theorem 3.1, proved in Section 3. Part (ii) is the content of Section 4: Proposition 4.4 carries out the first moment over chains under an independence hypothesis, which is shown to be false in Proposition 4.7, and Claim D is the corrected hypothesis, whose second half is precisely the statement that the first moment converges. Part (iii) is Theorem 6.1 in Sections 5 and 6. Monotonicity in μ holds because the line process of intensity $\mu' < \mu$ is a thinning of that of intensity μ and the Boolean model is monotone in the lines.

1.3 How to read this document

Everything stated as a lemma, proposition or theorem is proved in full here, with the exception that a theorem may be conditional on claims, and then says so in its statement. The claims are displayed in boxes. There are four of them:

- Claim D (Section 4): the dependence between thick clusters sharing lines is controlled by relay terms. What we have is the independent-case computation in full, a lemma that replaces the conditioning on a family of clusters by fixed lines (a minimal witness), the proof that the naive independence hypothesis is false, a dichotomy lemma for the geometry of witnesses, and a reduction of the claim to a weighted form of the multiscale hypothesis plus one statement about the conditional law of witnesses (Section 4.6).
- Claims C1 to C3 (Section 6): the three places where the multiscale recursion is not closed. Claim C1 is a first-moment computation for the weights of shared lines that we have carried out for one sub-box but not for the product over several; Claim C3 is the base case, where the argument is complete except for the bookkeeping of short tubes; Claim C2, the treatment of crossings that ride the fixed lines, is the substantial open point, and we describe its three components.

The reader who wants to check only what is proved may read Sections 2 and 3 and the lemmas of Sections 5 and 6; the reader who wants to attack the open points will find in each box the precise statement needed and, after it, the state of our knowledge.

2 Preliminaries on the line process

Lemma 2.1 (Measures of sets of lines)**.** *Let ν be as in (1.1).*

(a) *The lines meeting a set of diameter r have measure at most $2\mu r$.*

(b) *Let A_1, A_2 be sets of diameter at most δ whose centres are at distance $D \geq 4\delta$. The lines meeting both have measure at most $4\mu\delta^2/D$, and any two of them make an angle at most $2\delta/D$.*

(c) *Let ℓ_0 be a line and $I \subseteq \ell_0$ a segment of length $t \geq 2$. The lines that stay within distance 1 of ℓ_0 along all of I have measure at most $8\mu/t$. The lines that stay within distance 1 of ℓ_0 along I and in addition meet a set A_2 of diameter at most δ at distance $D \geq 4\delta$ from I have measure at most $8\mu\delta/D$.*

Proof. (a) A set of diameter r is contained in a disk of radius r, and the lines meeting a disk of radius r have measure $(\mu/\pi) \cdot \pi \cdot 2r = 2\mu r$.

(b) Let c_1, c_2 be the centres. A line meeting both sets passes within distance $\delta/2$ of both centres, so if φ is its angle with the segment c_1c_2, then $D \sin\varphi \leq \delta$ and $\varphi \leq (\pi/2)\sin\varphi \leq \pi\delta/(2D)$. Two such lines therefore make an angle at most $\pi\delta/D \leq 4\delta/D$. The normal direction θ of such a line

ranges over an interval of length at most $\pi\delta/D$, and for each θ the offset p ranges over an interval of length at most δ, since the line meets A_1, whose projection on any direction has length at most δ. The measure is at most $(\mu/\pi)\cdot(\pi\delta/D)\cdot\delta = \mu\delta^2/D \le 4\mu\delta^2/D$.

(c) A line within distance 1 of ℓ_0 at both endpoints of I makes an angle φ with ℓ_0 satisfying $t\sin\varphi \le 2$, so $\varphi \le \pi/t$; its normal direction lies in an interval of length $2\pi/t$, and given the direction its offset lies in an interval of length 2, since it passes within distance 1 of a fixed endpoint of I; the measure is at most $(\mu/\pi)\cdot(2\pi/t)\cdot 2 = 4\mu/t \le 8\mu/t$. For the second statement apply the argument of (b) to the disk I' of radius 1 around the endpoint of I nearest to A_2 and to A_2: the normal direction lies in an interval of length at most $\pi\max\{\delta,2\}/D \le 2\pi\delta/D$ for $\delta \ge 1$, the offset in an interval of length at most 2, and the measure is at most $(\mu/\pi)\cdot(2\pi\delta/D)\cdot 2 = 4\mu\delta/D \le 8\mu\delta/D$. □

Lemma 2.2 (Poisson tails)**.** *Let M be Poisson with mean m. For $k \ge m$, $\mathbb{P}(M \ge k) \le (\mathrm{e}m/k)^k$, and $\mathbb{P}(M \ge 2\mathrm{e}m) \le \mathrm{e}^{-m}$.*

Proof. For $s > 0$, Markov's inequality gives $\mathbb{P}(M \ge k) \le \mathrm{e}^{-sk}\mathbb{E}\mathrm{e}^{sM} = \exp\{-sk + m(\mathrm{e}^s - 1)\}$, and $s = \log(k/m) \ge 0$ yields $\mathbb{P}(M \ge k) \le \mathrm{e}^{-m}(\mathrm{e}m/k)^k \le (\mathrm{e}m/k)^k$. For $k = 2\mathrm{e}m$ this is $\mathrm{e}^{-m}2^{-2\mathrm{e}m} \le \mathrm{e}^{-m}$. □

Lemma 2.3 (Multivariate Mecke formula)**.** *For a measurable function $f \ge 0$ of a finite set of lines and $n \ge 1$,*

$$\mathbb{E}\Big[\sum_{\substack{\Lambda\subseteq\mathcal{L}\\ |\Lambda|=n}} f(\Lambda)\Big] = \frac{1}{n!}\int f(\{\ell_1,\dots,\ell_n\})\,\nu^{\otimes n}(\,\mathrm{d}\ell_1,\dots,\mathrm{d}\ell_n)\,.$$

In particular, for a monotone increasing event E of the line process, $\mathbb{P}(E) \le \sum_{n\ge1} M_n/n!$, where M_n is the $\nu^{\otimes n}$-measure of the n-tuples $(\ell_1,\dots,\ell_n)$ such that $\{\ell_1,\dots,\ell_n\}$ is a minimal set realising E, i.e. realises E and no proper subset does.

Proof. The formula is [LP17, Theorem 4.4] for the Poisson process $\mathcal{L}$ with intensity measure ν. On E there is a finite set $\Lambda_0 \subseteq \mathcal{L}$ realising E (the event is determined by the lines meeting a bounded region in all our applications, and that set is almost surely finite); by monotonicity and finiteness Λ_0 contains a minimal realising set. Hence $\mathbf{1}_E \le \#\{\Lambda \subseteq \mathcal{L} \text{ minimal realising}\}$, and the expectation of the right-hand side is $\sum_n M_n/n!$ by the formula. □

Lemma 2.4 (Harris inequality)**.** *Increasing events of the Poisson line process are positively correlated: $\mathbb{P}(E\cap E') \ge \mathbb{P}(E)\mathbb{P}(E')$.*

Proof. [LP17, Theorem 20.4]. □

3 The skeleton bound: from (T) to non-percolation

Theorem 3.1 (Skeleton bound)**.** *Let $j_0 \ge 2$ and $0 < \lambda < 1/(576\sqrt{2}\,(j_0-1))$, and put $\rho := 576\sqrt{2}\,\lambda(j_0-1) < 1$. If $(\mathrm{T})_{\sqrt{\rho}}$ holds, then Ξ_λ does not percolate almost surely.*

Conditionally on $\mathcal{L}$, call a box z *occupied* if $X_\lambda(Q(z)) \ge 1$. Occupation events of distinct boxes are conditionally independent, since the Cox process restricted to disjoint sets consists of independent Poisson processes given $\mathcal{L}$. Each line meets $Q(z)$ in a segment of length at most $\sqrt{2}$, and a thin box has $N(z) \le j_0 - 1$, so

$$\mathbb{P}(z \text{ occupied} \,|\, \mathcal{L}) \le \min\{1, \sqrt{2}\,\lambda N(z)\} \le \begin{cases} \sqrt{2}\,\lambda(j_0-1) = \rho/576, & z \text{ thin},\\ 1, & z \text{ thick}. \end{cases} \tag{3.1}$$

Conditionally on $\mathcal{L}$, call a box *open* if it is thick or occupied. Thick boxes are open by definition, whether or not they are occupied; this is the only place where we give something away, and it is what makes the bookkeeping below independent of the paths inside clusters. For a 2-path γ let $\tau(\gamma)$ be its number of thin boxes.

Lemma 3.2 (Open paths visiting each cluster once)**.** *If some disk of an unbounded component of Ξ_λ has its centre in $Q(z_0)$, then for every $n \geq 1$ there is a 2-path $\gamma = (z_0, \dots, z_m)$ of open boxes with $\|z_m - z_0\|_\infty \geq n$ such that for every thick cluster C the boxes of γ in C form a contiguous stretch of γ.*

Proof. Let O be the set of boxes containing a centre of a disk of the component; its elements are occupied, hence open, and $z_0 \in O$. Two unit disks intersect only if their centres are at distance less than 2, and then their boxes are at sup-distance at most 2; since any two disks of the component are joined by a finite chain of pairwise intersecting disks of the component, O is 2-connected. It is infinite because the component is unbounded. Hence there is a 2-path in O from z_0 to a box z_m with $\|z_m - z_0\|_\infty \geq n$. Among all 2-paths of open boxes from z_0 to z_m choose one, γ, with $\tau(\gamma)$ minimal. Suppose γ visits a cluster C in two stretches, and let u be the last box of the first stretch and v the first box of the second. The part of γ strictly between u and v contains a thin box, since consecutive boxes of γ are at distance at most 2 while two clusters are at distance at least 3 and boxes within distance 2 of C outside C are thin. Replace this part by a 2-path inside C from u to v, which exists because C is 2-connected, and erase loops from the resulting sequence to make it self-avoiding; loop erasure deletes boxes only. The new path consists of open boxes, runs from z_0 to z_m and has fewer thin boxes than γ, a contradiction. □

Lemma 3.3 (From open paths to chains)**.** *Let $\gamma = (z_0, \dots, z_m)$ be a 2-path of open boxes with $\|z_m - z_0\|_\infty \geq n$ that visits every cluster in at most one contiguous stretch. Decompose γ into maximal thin stretches and maximal thick stretches. The thick stretches lie in pairwise distinct clusters $C_1, \dots, C_k$, listed in the order of their visits, and the thin stretches have lengths $g_0, \dots, g_k$, where g_0 is the length of the initial thin stretch and g_k that of the final one, both possibly zero. Then $(C_1, \dots, C_k; g_0, \dots, g_k)$ is a cluster chain from z_0 that reaches n.*

Proof. A maximal thick stretch is 2-connected and consists of thick boxes, so it lies in one cluster, and by assumption different thick stretches lie in different clusters. Maximal stretches alternate, so $g_i \geq 1$ for $1 \leq i < k$. The i-th thin stretch starts within distance 2 of a box of C_i and ends within distance 2 of a box of C_{i+1}, and its consecutive boxes are within distance 2, so $\operatorname{dist}(C_i, C_{i+1}) \leq 2 + 2(g_i - 1) + 2 = 2g_i + 2$, which is (1.3); the same argument gives the condition on g_0. The displacement of γ is at most the sum of the displacements of its stretches and of the at most $2k$ transitions between stretches, each a step of sup-norm at most 2: a thick stretch in C_i contributes at most $\operatorname{diam} C_i$, a thin stretch of g_i boxes at most $2g_i$, and the transitions at most $4k$; hence $n \leq F + 2G$. □

Proof of Theorem 3.1. Let $E(z_0)$ be the event that a disk of an unbounded component has its centre in $Q(z_0)$. Fix $\mathcal{L}$. By Lemmas 3.2 and 3.3 and the union bound, $\mathbb{P}(E(z_0) \mid \mathcal{L})$ is at most the sum, over cluster chains from z_0 that reach n, of the number of paths realising the chain times a bound on the conditional probability that the thin boxes of a given realisation are all occupied. We count realisations by their thin stretches and by the entry and exit boxes of each cluster visit: a thin stretch of g boxes has at most 24^g realisations given the box it starts from; the entry box of a cluster is within distance 2 of the last thin box, at most 24 choices; the exit box has at most $|C_i|$ choices; the path inside the cluster is not counted, since thick boxes are open regardless of

occupation. By (3.1) and conditional independence, all thin boxes of a realisation are occupied with conditional probability at most $(\rho/576)^G$. Since interior thin stretches have $g_i \geq 1$, $k \leq G+1$ and $24^k \leq 24 \cdot 24^G$; therefore

$$\mathbb{P}(E(z_0) \,|\, \mathcal{L}) \leq \sum_{\text{chains}} 24^{k+G} \Big(\frac{\rho}{576}\Big)^G \prod_{i=1}^{k} |C_i| \leq 24 \sum_{\text{chains}} w_\rho \,, \tag{3.2}$$

where both sums run over the chains from z_0 reaching n. A chain reaching n has $F \geq n/2$ or $G \geq n/4$. For the latter, $\rho^G \leq \rho^{n/8} \rho^{G/2}$ and $w_\rho \leq \rho^{n/8} w_{\sqrt{\rho}}$. Taking expectations and using stationarity,

$$\mathbb{P}(E(0)) \leq 24\, \rho^{n/8}\, \mathbb{E}\Big[\sum_{\text{all chains from } 0} w_{\sqrt{\rho}} \Big] + 24\, \Phi_n(\sqrt{\rho}) \,.$$

The expectation is finite by $(\mathrm{T})_{\sqrt{\rho}}$: chains with $k = 0$ contribute $\sum_{g \geq 0} \rho^{g/2} < \infty$, and chains with $k \geq 1$ have finite expected total weight by assumption. The second term tends to zero by (1.4). Letting $n \to \infty$, the origin box contains no centre of an unbounded component almost surely, and a union over $z_0 \in \mathbb{Z}^2$ shows that there is no unbounded component almost surely. □

Remark 3.4. The bound (3.2) contains no sum over paths inside thick clusters: a cluster is entered, traversed at cost 1, and left, with the exit box as the only choice. Thin boxes have $N < j_0$, so their cost is uniform and the annealed moments of the line counts, which are superexponential, play no role. The constant $576 = 24^2$ is not optimised.

Remark 3.5 (Distinct clusters). The requirement in Definition 1.2 that the clusters of a chain be pairwise distinct is essential and corrects our working note [HJ26a], where repetitions were allowed. With repetitions, the chains $(C, C, \ldots, C)$ with $g_i = 1$ have weight $\rho^{k-1}|C|^k$, and under a polynomial size tail of clusters, which is the true behaviour, the expected total weight is infinite, so that (T) as stated there could not hold. Lemma 3.2 removes the repetitions by choosing the open path with the fewest thin boxes.

4 From the single-cluster bound to the thick-cluster lemma

This section carries out the first moment over cluster chains and isolates the dependence step. The first moment is complete under an independence hypothesis (D_0) (Proposition 4.4); we then prove that the conditioning on a family of clusters can be replaced by fixed lines (Lemma 4.6), show that (D_0) is false because of relays (Proposition 4.7), and state the corrected hypothesis (Claim D).

4.1 Single-cluster moments

For a box z let $C(z)$ be the thick cluster containing z if z is thick, and $C(z) = \varnothing$ otherwise, with $\operatorname{diam} \varnothing = -1$. Put

$$p_0 := \mathbb{P}(z \in T) = \mathbb{P}(N(z) \geq j_0) \leq \Big(\frac{\mathrm{e}\mu_1}{j_0}\Big)^{j_0} \qquad (\text{Lemma 2.2, for } j_0 \geq \mu_1) \,, \tag{4.1}$$

and, for $d \geq 0$,

$$A_3(d) := p_0 \mathbf{1}\{d = 0\} + \sum_{d' \geq \max\{d,1\}} (d'+1)^3 A\, d'^{-\beta}, \qquad A_3 := A_3(0) \,. \tag{4.2}$$

Under $(\mathrm{S})_{\beta,A}$ with $\beta > 4$ these are finite, and for $d \geq 1$,

$$A_3(d) \leq 8A \sum_{d' \geq d} {d'}^{3-\beta} \leq \frac{16A}{\beta - 4}\, d^{4-\beta}\,. \tag{4.3}$$

Lemma 4.1 (Third moment of one cluster)**.** *Assume* $(\mathrm{S})_{\beta,A}$ *with* $\beta > 4$*. For every box* z *and* $d \geq 0$*,*

$$\mathbb{E}\big[|C(z)|(\operatorname{diam} C(z) + 1)\mathbf{1}\{\operatorname{diam} C(z) \geq d\}\big] \leq A_3(d)\,.$$

Proof. A set of sup-norm diameter d' is contained in a square of $(d'+1)^2$ lattice points, so $|C(z)|(\operatorname{diam} C(z)+1) \leq (\operatorname{diam} C(z)+1)^3$. Hence the expectation is at most $\sum_{d' \geq d}(d'+1)^3 \mathbb{P}(\operatorname{diam} C(z) = d')$, where $\mathbb{P}(\operatorname{diam} C(z) = 0) \leq p_0$ and $\mathbb{P}(\operatorname{diam} C(z) = d') \leq \mathbb{P}(\exists C \ni z,\ \operatorname{diam} C \geq d') \leq A d'^{-\beta}$ for $d' \geq 1$ by (1.5). □

4.2 Counting boxes near a cluster

Lemma 4.2 (Lattice counts)**.** *Let* C *be a set of boxes of sup-norm diameter* d*. For* $r \geq 1$*, the number of boxes at sup-distance exactly* r *from* C *is at most* $8(d+1)(r+1)$*, and the number of boxes at sup-distance exactly* r *from a single box is at most* $8r$*.*

Proof. C lies in a square of side d (in lattice units), and the boxes at distance exactly r from C lie in the square of side $d+2r$ concentric with it and outside the square of side $d+2r-2$; these have $(d+2r+1)^2 - (d+2r-1)^2 = 4(d+2r) \leq 8(d+1)(r+1)$ lattice points. For a single box, $d = 0$ and the count is $8r$. □

For $0 < \rho < 1$ put

$$S_\rho := \frac{1}{1-\rho}\Big(1 + \sum_{r \geq 1} 8r\, \rho^{(r-2)^+/2}\Big), \qquad S'_\rho := \frac{1}{1-\rho} \sum_{r \geq 3} 8(r+1)\, \rho^{(r-2)/2}\,. \tag{4.4}$$

4.3 The independence hypothesis, and why it fails

For a finite family $\mathcal{F} = \{C^1, \dots, C^i\}$ of finite, 2-connected, pairwise disjoint sets of boxes, pairwise at distance at least 3, let $\{\mathcal{F} \subseteq \mathcal{C}\}$ be the event that every C^j is a thick cluster. The first moment over chains, carried out in Proposition 4.4 below, needs a bound on the size of the cluster of a box z conditionally on this event. The simplest conceivable bound is the following.

Definition 4.3 (Hypothesis (D_0))**.** There is $\kappa = \kappa(j_0) \geq 1$ such that for every family $\mathcal{F}$ as above with $\mathbb{P}(\mathcal{F} \subseteq \mathcal{C}) > 0$, every box z at distance at least 3 from all members of $\mathcal{F}$, and every $d \geq 0$,

$$\mathbb{E}\big[|C(z)|(\operatorname{diam} C(z) + 1)\mathbf{1}\{\operatorname{diam} C(z) \geq d\} \;\big|\; \mathcal{F} \subseteq \mathcal{C}\big] \leq \kappa\, A_3(d)\,. \tag{4.5}$$

If the thick clusters were an independent family, (4.5) would be Lemma 4.1 with $\kappa = 1$. We show in Proposition 4.7 that (D_0) is false, and in Section 4.5 what replaces it. We first record the first moment over chains under (D_0), because any correct replacement will have to be fed into the same arithmetic.

4.4 The first moment over chains

Proposition 4.4 (From (S) to (T) under (D_0))**.** *Assume* $(S)_{\beta,A}$ *with* $\beta > 4$ *and* (D_0)*. Let* $0 < \rho < 1$ *and suppose*

$$\eta := \kappa A_3 S'_\rho < 1\,. \tag{4.6}$$

Then $(T)_\rho$ *holds. More precisely, the expected total weight of all chains from* 0 *with* $k \geq 1$ *is at most* $\kappa A_3 S_\rho/((1-\rho)(1-\eta))$*, and* $\Phi_n(\rho) = O(n^{4-\beta})$*.*

Condition (4.6) is a smallness condition on A_3, i.e. on p_0 and A. With $A = A'(j_0)\mu^{j_0}$ as in Theorem 6.1 and $p_0 \leq (\mathrm{e}\mu_1/j_0)^{j_0}$, (4.6) holds for $\mu \leq \mu_0(j_0,\rho)$; the dependence on ρ is through $S'_\rho \leq C(1-\rho)^{-3}$.

Proof. Step 1: summing out the gaps. For a chain $(C_1,\dots,C_k;g_0,\dots,g_k)$ the gaps are constrained only by (1.3). Summing ρ^{g_i} over the admissible g_i gives, by (1.3),

$$\sum_{g_0}\rho^{g_0} \leq \frac{\rho^{(\mathrm{dist}(0,C_1)-2)^+/2}}{1-\rho}, \quad \sum_{g_i}\rho^{g_i} \leq \frac{\rho^{\max\{1,(\mathrm{dist}(C_i,C_{i+1})-2)/2\}}}{1-\rho} \quad (1\leq i<k), \quad \sum_{g_k}\rho^{g_k} \leq \frac{1}{1-\rho}\,.$$

Hence the total weight of the chains from 0 with clusters $(C_1,\dots,C_k)$ is at most

$$\frac{\rho^{(\mathrm{dist}(0,C_1)-2)^+/2}}{(1-\rho)^{k+1}} \prod_{i=1}^{k-1}\rho^{\max\{1,(\mathrm{dist}(C_i,C_{i+1})-2)/2\}} \prod_{i=1}^{k}|C_i|\,. \tag{4.7}$$

Step 2: from clusters to boxes. For a cluster C and a set C' of boxes with $\mathrm{dist}(C,C') \geq 3$, let $z(C,C')$ be the lexicographically first box of C at minimal distance from C'; similarly $z(C,0)$. Then the sum over clusters C distinct from given clusters $C_1,\dots,C_i$ of $|C|f(\mathrm{dist}(C_i,C))$, for a decreasing $f \geq 0$, is at most the sum over boxes z at distance at least 3 from C_i of $|C(z)|f(\mathrm{dist}(z,C_i))$, since every such cluster is $C(z(C,C_i))$ and $\mathrm{dist}(C,C_i) = \mathrm{dist}(z(C,C_i),C_i)$. Note that a box z with $\mathrm{dist}(z,C_i) \leq 2$ is thin, so that the sum over all boxes with $C(z) \neq \varnothing$ and $C(z) \neq C_j$ for $j \leq i$ is the same as the sum over boxes at distance at least 3 from $C_1,\dots,C_i$.

Step 3: the last cluster. Fix distinct clusters $C_1,\dots,C_{k-1}$ with $k \geq 2$, that is, fix a deterministic family $\mathcal{F} = \{C_1,\dots,C_{k-1}\}$ and work on $\{\mathcal{F} \subseteq \mathcal{C}\}$. By Step 2, Lemma 4.2 and (4.5) with $d = 0$,

$$\begin{aligned}
\mathbb{E}\Big[\sum_{C_k}|C_k|\rho^{\max\{1,(\mathrm{dist}(C_{k-1},C_k)-2)/2\}} \;\Big|\; \mathcal{F}\subseteq\mathcal{C}\Big] &\leq \sum_{r\geq 3}\sum_{z:\ \mathrm{dist}(z,C_{k-1})=r}\rho^{(r-2)/2}\,\mathbb{E}\big[|C(z)|\,\big|\,\mathcal{F}\subseteq\mathcal{C}\big] \\
&\leq \sum_{r\geq 3} 8(\mathrm{diam}\,C_{k-1}+1)(r+1)\rho^{(r-2)/2}\,\kappa A_3 \\
&= (1-\rho)S'_\rho\,\kappa A_3\,(\mathrm{diam}\,C_{k-1}+1)\,,
\end{aligned}$$

where we used $|C(z)| \leq |C(z)|(\mathrm{diam}\,C(z)+1)$. The factor $(\mathrm{diam}\,C_{k-1}+1)$ is attached to the cluster C_{k-1}, whose weight in (4.7) becomes $|C_{k-1}|(\mathrm{diam}\,C_{k-1}+1)$; this is why the third moment A_3 appears.

Step 4: iteration. Write Ψ_k for the expected total weight of chains from 0 with exactly k clusters. Expanding the expectation of (4.7) summed over $(C_1,\dots,C_k)$ as a sum over deterministic families of the indicator $\mathbf{1}\{\{C_1,\dots,C_k\}\subseteq\mathcal{C}\}$, taking first the expectation over C_k given $\{C_1,\dots,C_{k-1}\}\subseteq\mathcal{C}$ as in Step 3, then over C_{k-1} given $\{C_1,\dots,C_{k-2}\}\subseteq\mathcal{C}$ with the weight $|C_{k-1}|(\mathrm{diam}\,C_{k-1}+1)$ in place of $|C_{k-1}|$, and so on down to C_1, for which the unconditional Lemma 4.1 is used with $\sum_{r\geq 0}\#\{z\colon \|z\|_\infty = r\}\rho^{(r-2)^+/2} \leq (1-\rho)S_\rho$ by Lemma 4.2, we obtain

$$\Psi_k \leq \frac{(1-\rho)^k S_\rho\,(S'_\rho)^{k-1}\,(\kappa A_3)^k}{(1-\rho)^{k+1}} = \frac{S_\rho\,\kappa A_3}{1-\rho}\,\eta^{k-1} \qquad (k\geq 1)\,, \tag{4.8}$$

where the k factors $(1-\rho)$ in the numerator come from the k cluster sums and the denominator from Step 1. Summing over $k \geq 1$ gives the first assertion.

Step 5: the free advance. A chain with k clusters and $F \geq n/2$ has a cluster C_i with $\operatorname{diam} C_i + 4 \geq n/(2k)$. Repeating Steps 3 and 4 with the indicator $\mathbf{1}\{\operatorname{diam} C_i \geq n/(2k) - 4\}$ inserted at the i-th cluster replaces the factor κA_3 at that cluster by $\kappa A_3(n/(2k) - 4)$, by (4.5) with $d = \lceil n/(2k) - 4 \rceil$ (or by Lemma 4.1 if $i = 1$). Hence, with (4.3),

$$\Phi_n(\rho) \leq \sum_{k \geq 1} k\, \Psi_k \, \frac{A_3(\lceil n/(2k) - 4 \rceil)}{A_3} \leq \sum_{k \leq n/16} k\, \Psi_k \, \frac{16A}{(\beta - 4)A_3} \Big(\frac{n}{4k}\Big)^{4-\beta} + \sum_{k > n/16} k\, \Psi_k \,,$$

where for $k \leq n/16$ we used $n/(2k) - 4 \geq n/(4k)$. By (4.8) the first sum is at most $C\, n^{4-\beta} \sum_k k^{\beta-3} \eta^{k-1} = O(n^{4-\beta})$ and the second is at most $(1-\rho)^{-1} S_\rho \kappa A_3 \sum_{k > n/16} k \eta^{k-1} = O(n \eta^{n/16})$. Both tend to zero. □

Remark 4.5. The only property of thick clusters used in the proof is the size tail (S) together with (4.5). The exponent $\beta > 4$ is what the third moment in Step 3 requires and is far below the exponent $\beta = j_0 - 2$ aimed at by Theorem 6.1 for large j_0.

4.5 The dependence step: witnesses and relays

We now remove the conditioning in (4.5) in favour of fixed lines, show that (D_0) is false, and state what has to replace it.

Witnesses. Let $N(\mathcal{F})$ be the set of boxes within distance 2 of a member of $\mathcal{F}$, let $U(\mathcal{F})$ be the 1-neighbourhood of $\bigcup_{z \in N(\mathcal{F})} Q(z)$, a bounded set, and let $\mathcal{L}_N$ be the set of lines of $\mathcal{L}$ meeting $U(\mathcal{F})$ and $\mathcal{L}' \coloneqq \mathcal{L} \setminus \mathcal{L}_N$; these are independent Poisson processes, and $\mathcal{L}_N$ is almost surely finite. Whether a box of $N(\mathcal{F})$ is thick is determined by $\mathcal{L}_N$, so the events

$$T_{\mathcal{F}} \coloneqq \{\text{all boxes of all members of } \mathcal{F} \text{ are thick}\},$$
$$\Theta_{\mathcal{F}} \coloneqq \{\text{all boxes of } N(\mathcal{F}) \text{ outside the members of } \mathcal{F} \text{ are thin}\}$$

belong to $\sigma(\mathcal{L}_N)$, $T_{\mathcal{F}}$ is increasing and $\Theta_{\mathcal{F}}$ decreasing in $\mathcal{L}_N$, and $\{\mathcal{F} \subseteq \mathcal{C}\} = T_{\mathcal{F}} \cap \Theta_{\mathcal{F}}$, since a 2-connected set of thick boxes whose 2-neighbourhood consists of thin boxes is a cluster. A finite set w of lines is a *witness* if all boxes of all members of $\mathcal{F}$ are thick in w, and a *minimal witness* if no proper subset of w is a witness. Fix a measurable total order on finite sets of lines (by cardinality, then lexicographically in the coordinates (θ, p) of the lines listed in increasing order). On $T_{\mathcal{F}}$ the finite set $\mathcal{L}_N$ is a witness and hence contains minimal witnesses, finitely many; let $W = W(\mathcal{L}_N)$ be the least of them.

For a box z at distance at least 3 from $\mathcal{F}$ and a locally finite set A of lines let $C^*_A(z)$ be the 2-connected component of z in the set of boxes outside $N(\mathcal{F})$ that are thick in A, with $C^*_A(z) = \varnothing$ if z is thin in A, and put

$$f^*_d(A) \coloneqq |C^*_A(z)|(\operatorname{diam} C^*_A(z) + 1)\mathbf{1}\{\operatorname{diam} C^*_A(z) \geq d\}\,. \tag{4.9}$$

The map $A \mapsto f^*_d(A)$ is increasing, and on $\{\mathcal{F} \subseteq \mathcal{C}\}$ one has $C^*_{\mathcal{L}}(z) = C(z)$, because $C(z)$ is a cluster distinct from the members of $\mathcal{F}$ and hence disjoint from $N(\mathcal{F})$.

Lemma 4.6 (Conditioning on a family of clusters is dominated by adding a witness)**.** *Let $g \geq 0$ be a measurable increasing functional of locally finite sets of lines, and let $G(w) := \mathbb{E}\, g(w \cup \mathcal{L})$ for finite sets w of lines. Then*

$$\mathbb{E}[g(\mathcal{L})\mathbf{1}\{\mathcal{F} \subseteq \mathcal{C}\}] \leq \mathbb{E}[G(W)\mathbf{1}\{\mathcal{F} \subseteq \mathcal{C}\}]\,, \tag{4.10}$$

and consequently $\mathbb{E}[g(\mathcal{L}) \mid \mathcal{F} \subseteq \mathcal{C}] \leq \sup\{G(w)\colon w$ a minimal witness with $\Theta_{\mathcal{F}}$ holding in $w\}$ whenever $\mathbb{P}(\mathcal{F} \subseteq \mathcal{C}) > 0$.

Proof. Put $\tilde{g}(A) := \mathbb{E}\, g(A \cup \mathcal{L}')$ for finite $A \subseteq \{$lines meeting $U(\mathcal{F})\}$, an increasing function, so that $\mathbb{E}[g(\mathcal{L})\mathbf{1}_{T\cap\Theta}] = \mathbb{E}[\tilde{g}(\mathcal{L}_N)\mathbf{1}_{T\cap\Theta}(\mathcal{L}_N)]$ by the independence of $\mathcal{L}_N$ and $\mathcal{L}'$, where we abbreviate $T = T_{\mathcal{F}}$, $\Theta = \Theta_{\mathcal{F}}$. On T exactly one subset of $\mathcal{L}_N$ equals $W(\mathcal{L}_N)$, so

$$\mathbb{E}[\tilde{g}(\mathcal{L}_N)\mathbf{1}_{T\cap\Theta}(\mathcal{L}_N)] = \sum_{n\geq 1} \mathbb{E}\Big[\sum_{w\subseteq\mathcal{L}_N,\ |w|=n} h(w, \mathcal{L}_N \setminus w)\Big]\,,$$

where $h(w, A) := \mathbf{1}\{w = W(w\cup A)\}\,\mathbf{1}_\Theta(w\cup A)\,\tilde{g}(w\cup A)$ and the inner sum runs over the subsets w of $\mathcal{L}_N$ with n elements. By the multivariate Mecke formula [LP17, Theorem 4.4], the n-th term equals

$$\frac{1}{n!}\int \mathbb{E}[h(w, \mathcal{L}_N)]\,\nu_N^{\otimes n}(\,\mathrm{d}w)\,,$$

where ν_N is ν restricted to the lines meeting $U(\mathcal{F})$. Fix w. The map $A \mapsto \mathbf{1}\{w = W(w\cup A)\}$ is decreasing: if $A \subseteq A'$ and w is the least minimal witness of $w \cup A'$, then w is a minimal witness contained in $w \cup A$, every minimal witness of $w \cup A$ is also one of $w \cup A'$ and hence not below w, so $w = W(w \cup A)$. The map $A \mapsto \mathbf{1}_\Theta(w \cup A)$ is decreasing and $A \mapsto \tilde{g}(w \cup A)$ is increasing. By the Harris inequality for the Poisson process $\mathcal{L}_N$ [LP17, Theorem 20.4], applied to the decreasing bounded function $\mathbf{1}\{w = W(w\cup A)\}\mathbf{1}_\Theta(w\cup A)$ and the increasing function $\tilde{g}(w\cup A)\wedge M$, and letting $M \to \infty$,

$$\mathbb{E}[h(w, \mathcal{L}_N)] \leq \mathbb{E}[\mathbf{1}\{w = W(w\cup\mathcal{L}_N)\}\mathbf{1}_\Theta(w\cup\mathcal{L}_N)]\cdot\mathbb{E}[\tilde{g}(w\cup\mathcal{L}_N)]\,,$$

and $\mathbb{E}[\tilde{g}(w\cup\mathcal{L}_N)] = \mathbb{E}\, g(w\cup\mathcal{L}_N\cup\mathcal{L}') = G(w)$. Multiplying by $\frac{1}{n!}$, integrating over w and applying the Mecke formula backwards gives

$$\mathbb{E}[g(\mathcal{L})\mathbf{1}_{T\cap\Theta}] \leq \sum_{n\geq 1}\mathbb{E}\Big[\sum_{w\subseteq\mathcal{L}_N,\ |w|=n} \mathbf{1}\{w = W(\mathcal{L}_N)\}\mathbf{1}_\Theta(\mathcal{L}_N)\,G(w)\Big] = \mathbb{E}[G(W)\mathbf{1}_{T\cap\Theta}]\,,$$

which is (4.10). For the consequence, divide by $\mathbb{P}(T\cap\Theta)$ and note that on $T\cap\Theta$ the witness W is minimal and Θ holds in $W \subseteq \mathcal{L}_N$, since Θ is decreasing. □

Applied to $g = f_d^*$, the lemma says that the left-hand side of (4.5) is at most $\sup_w G_d(w)$, where $G_d(w) = \mathbb{E}f_d^*(w\cup\mathcal{L})$ is the third moment of the cluster of z in the Poisson process with the finite set w added as fixed lines and the boxes of $N(\mathcal{F})$ removed. The dependence step is thereby reduced to a *fixed-line single-cluster bound*, in which the fixed lines are a minimal witness of a family of clusters. The lemma does not use anything about lines and holds for any Poisson process; what is specific to the problem is what minimal witnesses look like.

Proposition 4.7 ((D_0) is false)**.** *Let $j_0 \geq 3$ and $\mu \leq \mu_0(j_0)$. There is $c = c(j_0) > 0$ such that for every $d_1 \geq d_1(j_0)$ there are a family $\mathcal{F} = \{C_1\}$ consisting of one set of diameter d_1 with $\mathbb{P}(\mathcal{F}\subseteq\mathcal{C}) > 0$ and a box z at distance 3 from C_1 such that*

$$\mathbb{P}(\operatorname{diam} C(z) \geq d \mid \mathcal{F}\subseteq\mathcal{C}) \geq \frac{c\,\mu}{d} \qquad \text{for all } 1 \leq d \leq cd_1\,. \tag{4.11}$$

Consequently the left-hand side of (4.5) *with* $d = 0$ *is at least* $c'\mu d_1^2$*, and no constant* κ *satisfies* (D_0).

Proof sketch. Let C_1 be the set of boxes meeting the convex set $\{x\colon |x_2| \le 1/2,\ 0 \le x_1 \le d_1\}$, and $z = (d_1 + 4, 0)$. The typical realisation of $\{\mathcal{F} \subseteq \mathcal{C}\}$ consists of j_0 lines whose common 1-neighbourhood is a convex region ending near $x_1 = d_1$, with no further lines nearby; its probability is of order $(\mu/d_1)^{j_0}$ up to constants, by Lemma 2.1(c) for the lines and a bounded factor for the absence of further lines in the bounded region where they would spoil $\Theta_{\mathcal{F}}$. The directions of the j_0 lines differ by angles of order $1/d_1$, so the common 1-neighbourhood of $j_0 - 1$ of them, the region Θ_{j_0-1}, extends beyond $x_1 = d_1$ by a length of order d_1 with probability bounded below; this is the *release* of one line of the bundle. A further line of $\mathcal{L}$ that enters Θ_{j_0-1} at abscissa $d_1 + 4$ and stays within it over a length $2d$ makes the boxes along $[d_1 + 4, d_1 + 4 + d]$ thick, while the boxes in $[d_1 + 1, d_1 + 3]$ remain thin because the new line is at distance greater than 1 from Θ_{j_0-1} there and C_1's own j_0-th line has left; such a line has angle of order $1/d$ with the axis and offset in a window of bounded length, so its ν-measure is of order μ/d, and it is independent of the lines realising $\{\mathcal{F} \subseteq \mathcal{C}\}$ except for the requirement that it does not make a box of $N(\mathcal{F})$ thick, which costs a bounded factor. Hence the probability of $\{\mathcal{F} \subseteq \mathcal{C}\} \cap \{\operatorname{diam} C(z) \ge d\}$ is at least $c\,\mu/d$ times a lower bound for $\mathbb{P}(\mathcal{F} \subseteq \mathcal{C})$ of the same order as its typical realisation, which gives (4.11). The moment bound follows by summing $(d+1)^3\mathbb{P}(\operatorname{diam} C(z) = d)$, bounded below by $\sum_{d \le cd_1} d^2 \cdot c\mu/d \cdot (\text{a constant})$, i.e. by $c'\mu d_1^2$. We have not written out the constants, which requires an upper bound for $\mathbb{P}(\mathcal{F} \subseteq \mathcal{C})$ of the order of its typical realisation; this is a bounded-region estimate of the kind in Claim C3. □

The mechanism is a *relay*: the next cluster inherits $j_0 - 1$ lines from the previous one and buys the j_0-th at the price μ/d of a single line along a prescribed axis. Relays are harmless for the chain sum, since a relay of length d attached to a bundle of length d_1 is rarer than a single bundle of length $d_1 + d$ by a factor of order $\mu(d_1 + d)^{j_0-2}/(d\, d_1^{j_0-2})$, which is small; but they are not uniform in $\mathcal{F}$, and a correct dependence hypothesis must mention them.

4.6 An attempt at the dependence step

We now try to prove the dependence step in the form suggested by Lemma 4.6. The attempt does not succeed, but it clarifies the statement: it shows that the bound cannot be uniform in the witness, so that Claim D below has to be an average over the conditional law of the witness, and it reduces the claim to a weighted version of the multiscale hypothesis together with one statement about witnesses, which we isolate.

The geometry of witnesses. For a finite set S of lines let

$$\Theta(S) \coloneqq \{x \in \mathbb{R}^2\colon \text{every line of } S \text{ passes within distance 1 of } x\}\,,$$

the intersection of the closed strips of width 2 around the lines of S; it is convex. Write $Q(\mathcal{F})$ for the union of the boxes of the members of $\mathcal{F}$ and $Q(N(\mathcal{F}))$ for the union of the boxes of $N(\mathcal{F})$.

Lemma 4.8 (Dichotomy for witnesses)**.** *Let* w *be a finite set of lines in which* $\Theta_{\mathcal{F}}$ *holds, and let* $S \subseteq w$ *with* $|S| = j_0$*. Then either* $\Theta(S) \subseteq Q(C)$ *for one member* $C \in \mathcal{F}$*, or* $\Theta(S) \cap Q(N(\mathcal{F})) = \varnothing$*. In particular, if* w *is a witness with exactly* j_0 *lines and* $\mathcal{F} = \{C\}$*, then every point outside* $Q(N(\mathcal{F}))$ *has at most* $j_0 - 1$ *lines of* w *within distance* 1*; and in general a point outside* $Q(N(\mathcal{F}))$ *with at least* j_0 *lines of* w *within distance* 1 *lies in* $\Theta(S)$ *for an* external j_0*-subset* S*, one with* $\Theta(S) \cap Q(N(\mathcal{F})) = \varnothing$.

Proof. A point of $\Theta(S)$ in a box of $N(\mathcal{F})$ outside the members would make that box thick in w, contradicting $\Theta_{\mathcal{F}}$; so $\Theta(S)$ does not meet the boxes of $N(\mathcal{F}) \setminus \bigcup \mathcal{F}$. Suppose $\Theta(S)$ contains a point x in a box of a member C and a point y outside $Q(C)$. The segment $[x, y]$ lies in $\Theta(S)$. Along a segment, the boxes visited are consecutively adjacent (sharing an edge or a corner), so their sup-distances from C change by at most 1 from one box to the next; since the segment starts at sup-distance 0 and ends at sup-distance at least 1, it visits a box at sup-distance exactly 1 or 2 from C unless it ends in one, and these boxes belong to $N(\mathcal{F}) \setminus \bigcup \mathcal{F}$ because distinct members are at distance at least 3. This contradicts the first sentence. Hence $\Theta(S)$ meets the boxes of at most one member, and if it meets them it is contained in $Q(C)$. If w has exactly j_0 lines and is a witness for $\mathcal{F} = \{C\}$, then $\Theta(w)$ meets every box of C, so $\Theta(w) \subseteq Q(C)$, and a point outside $Q(N(\mathcal{F}))$ with j_0 lines of w within distance 1 would lie in $\Theta(w)$. □

The lemma is the precise form of the statement that a bundle cannot reconverge after its ring: on the event $\{\mathcal{F} \subseteq \mathcal{C}\}$, the j_0 lines carrying a single bundle cluster are never all within distance 1 of a point outside the neighbourhood of the cluster. This is why the relay of Proposition 4.7 needs one new line. The lemma also shows what else can happen: lines from two different members may converge outside $N(\mathcal{F})$ and form an external j_0-subset, and then the boxes of $\Theta(S)$ are thick in w alone.

Proposition 4.9 (No bound uniform in the witness)**.** *For $j_0 \geq 3$ there are families $\mathcal{F} = \{C_1, C_2\}$, minimal witnesses w of $T_{\mathcal{F}}$ in which $\Theta_{\mathcal{F}}$ holds, and boxes z at distance 3 from C_1, such that $C_w^*(z)$ is deterministic with diameter t of order $\min\{\operatorname{diam} C_1, \operatorname{diam} C_2\}$, so that $\mathbb{E} f_0^*(w \cup \mathcal{L}) \geq t^3$. Consequently no inequality of the form (4.12) below with a factor μ in the second term can hold for every minimal witness.*

Proof. Let C_1 be the set of boxes meeting the convex region $\{|x_2| \leq 1/2,\ 0 \leq x_1 \leq d_1\}$ and let w_1 consist of j_0 lines through this region whose common 1-neighbourhood is contained in it and ends near $x_1 = d_1$, with $j_0 - 1$ of them, S_1, having a common 1-neighbourhood that continues along the axis beyond $x_1 = d_1 + 4$ for a length t (for instance $j_0 - 1$ lines exactly parallel to the axis at mutual distance less than 1 and one line at a small angle leaving the strip at $x_1 = d_1$). Let C_2 be a second such set, with its own witness w_2 of j_0 lines, placed so that one line $\ell_2 \in w_2$ runs within distance 1 of the axis of C_1 along $[d_1 + 4, d_1 + 4 + t]$ after leaving C_2, and so that C_2, its ring and its other witness lines stay away from this region; this is possible with $\operatorname{diam} C_2$ of order t. Then $w = w_1 \cup w_2$ is a minimal witness in which $\Theta_{\mathcal{F}}$ holds, $S = S_1 \cup \{\ell_2\}$ is an external j_0-subset, and the boxes along $[d_1 + 4, d_1 + 4 + t]$ are thick in w, so $C_w^*(z)$ for $z = (d_1 + 4, 0)$ contains them whatever $\mathcal{L}$ is. The second assertion follows because t can be chosen of order $\min\{\operatorname{diam} C_1, \operatorname{diam} C_2\}$ while μ is arbitrary. □

Such witnesses have small probability under the conditional law given $\{\mathcal{F} \subseteq \mathcal{C}\}$, because the line ℓ_2 has to be aligned with the release region of C_1 to within an angle $1/t$. The right statement is therefore about the chosen witness W of Lemma 4.6 under the conditional law, and this is how Claim D is formulated. We write $G_{z,d}(w) := \mathbb{E} f_d^*(w \cup \mathcal{L})$ with f_d^* defined for the box z.

Claim D (Dependence step, relay form). There are $\kappa = \kappa(j_0)$ and $q = q(j_0)$ such that for $\mu \leq \mu_0(j_0)$, $0 < \rho < 1$, every family $\mathcal{F}$ as above with $\mathbb{P}(\mathcal{F} \subseteq \mathcal{C}) > 0$, every member $C \in \mathcal{F}$ and

every $d \geq 0$,

$$\mathbb{E}\Big[\sum_{r\geq 3}\sum_{z:\ \operatorname{dist}(z,C)=r}\rho^{(r-2)/2}\,G_{z,d}(W)\ \Big|\ \mathcal{F}\subseteq\mathcal{C}\Big] \leq \kappa A_3(d)(\operatorname{diam}C+1)+\kappa\mu\sum_{C'\in\mathcal{F}}(\operatorname{diam}C'+1)^q\,\rho^{(\operatorname{dist}(C,C')-2\operatorname{diam}C'-2)^+/2}. \tag{4.12}$$

Moreover, the first moment over chains converges under (4.12) for $\beta > q+1$ and $\mu \leq \mu_0(j_0,\rho,\beta,A)$, in the sense that the iteration of Step 4 of Proposition 4.4, in which a relay term charges the moment $(\operatorname{diam}C'+1)^q$ to an earlier cluster C', has a finite sum.

By Lemma 4.6, the left-hand side of (4.12) dominates the conditional expectation that Step 3 of Proposition 4.4 needs, so the first half of the claim is what replaces (4.5). The damping $\rho^{(\operatorname{dist}(C,C')-2\operatorname{diam}C'-2)^+/2}$ starts only beyond the release region of C', which extends to a distance of the order of $\operatorname{diam}C'$ from C'.

Reduction to a weighted multiscale hypothesis. The quantity $G_{z,d}(w)$ is a single-cluster moment in the Poisson process with the deterministic lines w added and the boxes of $N(\mathcal{F})$ removed. To bound it by the multiscale hypothesis we need the hypothesis in a form with an excluded set of boxes and with a weight that charges fixed lines only where they can actually make boxes thick. For a finite set G' of lines, a point p and a set E of boxes, let $c^{(1)}_{G'}(p)$ be the maximum, over points x with $|x-p|\leq 3$ not in the boxes of E, of the number of lines of G' within distance 1 of x, and define $n^E_L(\ell_0)$ and $W^E_L(G')$ as in Definition 5.3 with $c_{G'}$ replaced by $c^{(1)}_{G'}$ and the stretches I required to avoid the boxes of E. Let $\mathcal{H}^E_2(L)$ be the statement that for every box B of side L, all finite $G'\subseteq G$ and every set E of boxes, the probability that B contains a crossing path in $\mathcal{L}\cup G$ avoiding the boxes of E and having fewer than $\varepsilon_L L$ boxes within distance 1 of $G\setminus G'$ is at most $C_0\rho_L W^E_L(G')$. Since excluding boxes only shrinks the event, $\mathcal{H}^E_2$ differs from $\mathcal{H}_2$ in the weight alone; Remark 5.5(b) applies to $c^{(1)}$ as it does to c, and we expect the recursion of Section 6 to go through for it unchanged, but we have not verified this.

Proposition 4.10 (Claim D from the weighted hypothesis and a witness bound)**.** *Assume $\mathcal{H}^E_2(L)$ for all $L\geq 8$. Then for every family $\mathcal{F}$, every finite set w of lines, every box z outside $N(\mathcal{F})$ and every $d\geq 0$,*

$$G_{z,d}(w)\leq \mathbf{1}\{d<8\}\,8^4\,\mathbb{P}(z \text{ thick in } w\cup\mathcal{L})+C_0\sum_{d'\geq\max\{d,8\}}(d'+1)^3\rho_{d'}\,W^{N(\mathcal{F})}_{2\lfloor d'/2\rfloor}\big(w\cap B_{d'}(z)\big), \tag{4.13}$$

where $B_{d'}(z)$ is the box of side $2\lfloor d'/2\rfloor$ centred at z. Consequently the first half of Claim D follows from the witness weight bound

$$\mathbb{E}\Big[\sum_{r\geq 3}\sum_{z:\ \operatorname{dist}(z,C)=r}\rho^{(r-2)/2}\sum_{d'\geq\max\{d,8\}}(d'+1)^3\rho_{d'}\Big(W^{N(\mathcal{F})}_{2\lfloor d'/2\rfloor}(W\cap B_{d'}(z))-1\Big)\ \Big|\ \mathcal{F}\subseteq\mathcal{C}\Big] \leq \kappa'\mu\sum_{C'\in\mathcal{F}}(\operatorname{diam}C'+1)^q\rho^{(\operatorname{dist}(C,C')-2\operatorname{diam}C'-2)^+/2} \tag{4.14}$$

together with the analogous bound for the thickness term, $\mathbb{E}[\sum_z\rho^{(r-2)/2}\mathbb{P}(z \text{ thick in } W\cup\mathcal{L})\,|\,\mathcal{F}\subseteq\mathcal{C}]\leq\kappa' p_0(\operatorname{diam}C+1)+(\text{relay term})$.

Proof. If $\operatorname{diam} C_w^*(z) \geq d' \geq 8$, then, as in Lemma 5.2, the box $B_{d'}(z)$ contains a crossing path of boxes thick in $w \cup \mathcal{L}$ avoiding the boxes of $N(\mathcal{F})$, since $C_w^*(z)$ is 2-connected, contains z, avoids $N(\mathcal{F})$ and reaches the boundary layer; so $\mathcal{H}_2^E$ with $E = N(\mathcal{F})$, $G = G' = w \cap B_{d'}(z)$ and no avoidance requirement bounds its probability by $C_0 \rho_{d'} W^E(w \cap B_{d'}(z))$ up to the factor 2^{j_0-2} from $\rho_{2\lfloor d'/2 \rfloor} \leq 2^{j_0-2}\rho_{d'}$, which we absorb into C_0. Summing $(d'+1)^3 \mathbb{P}(\operatorname{diam} C_w^*(z) = d') \leq (d'+1)^3 \mathbb{P}(\operatorname{diam} C_w^*(z) \geq d')$ over $d' \geq 8$ gives the second term of (4.13), and the diameters $d' < 8$ contribute at most $\sum_{d'<8}(d'+1)^3 \mathbb{P}(C_w^*(z) \neq \varnothing) \leq 8^4\, \mathbb{P}(z \text{ thick in } w \cup \mathcal{L})$. For the consequence, insert (4.13) into the left-hand side of (4.12): the part of the sum with the weights replaced by 1 is at most $\sum_{r \geq 3} 8(\operatorname{diam} C + 1)(r+1)\rho^{(r-2)/2} \cdot C_0 \sum_{d' \geq d}(d'+1)^3 \rho_{d'}$ by Lemma 4.2, and $C_0 \sum_{d' \geq \max\{d,1\}}(d'+1)^3 \rho_{d'}$ is $A_3(d)$ for the constants $A = C_0 2^{j_0} \mu^{j_0}$ and $\beta = j_0 - 2$ of Theorem 6.1, up to a constant; the remaining part is (4.14). The thickness term is the analogous bound assumed in the statement. □

What the witness weight bound asks, and what we can say. The left-hand side of (4.14) is the conditional expectation of the *excess weight* of the witness near C: for each line ℓ_0 of W passing within distance d' of a box z near C, the term $(d'+1)^3 \rho_{d'} k_{d'}(m) = K^m \mu^{j_0-m}(d'+1)^3 d'^{-(j_0-m)}$ with $m = n_{d'}^{N(\mathcal{F})}(\ell_0)$ the number of witness lines within distance 1 of common points along a stretch of length $d'/4$ near z outside $N(\mathcal{F})$. Three mechanisms contribute, and we can prove something about the first only.

1. *Released lines of one member.* By Lemma 4.8, if the witness of a member C' consists of j_0 lines, at most $j_0 - 1$ of them are within distance 1 of a common point outside $N(\mathcal{F})$, so $m \leq j_0 - 1$ for stretches carried by these lines, and the term is at most $K^{j_0-1}\mu(d'+1)^3/d'$. This is the relay term. The stretch has to lie in the region where the $j_0 - 1$ lines are still within width 2, which extends to distance $O(\operatorname{diam} C')$ from C' for lines with mutual angles of order $1/\operatorname{diam} C'$, but is unbounded for parallel lines; under the conditional law the angles are of order $1/\operatorname{diam} C'$ with high probability, which is the first unproved statement. Summing $(d'+1)^3\mu/d'$ over $d' \leq c \operatorname{diam} C'$ gives $\mu(\operatorname{diam} C')^3$, i.e. $q = 3$ in (4.14) for this mechanism, and the position of z is one of $O(1)$ boxes at the ends of the release region, which is within distance $O(\operatorname{diam} C')$ of C'; this is the damping in (4.12).

2. *Internal relays.* A member whose witness has more than j_0 lines, because it is itself a chain of relays, may release $j_0 - 1$ lines at several places. The number of released groups is bounded by the number of internal relays, each of which costs a factor of order μ in the conditional law; we have not quantified this.

3. *External bundles.* Lines from different members converging outside $N(\mathcal{F})$ into an external j_0-subset give $m \geq j_0$ and a trivial bound on the stretch they cover. Proposition 4.9 shows that this cannot be excluded for individual witnesses; under the conditional law, an external j_0-subset of length t requires an alignment to within angle $1/t$ of a line of one member with the release region of another, of conditional probability of order $1/t$ times the number of candidate pairs, and the resulting contribution to (4.14) is of the order of the relay term with μ replaced by this probability. Bounding it requires the conditional law of the directions of the witness lines, which is the second unproved statement.

The conditional law of W is explicit in principle: by the proof of Lemma 4.6 it is proportional to $\mathbf{1}\{w \text{ minimal witness}\}\, \mathbb{E}[\mathbf{1}\{w = W(w \cup \mathcal{L}_N)\}\mathbf{1}_\Theta(w \cup \mathcal{L}_N)]\, \nu_N^{\otimes n}(\mathrm{d}w)/n!$, and the two unproved statements are statements about this measure. Both say that the lines of a cluster's witness behave, in direction and in number, like the j_0 long lines of a single bundle; a proof would have

to bound the alternative realisations of $\{\mathcal{F} \subseteq \mathcal{C}\}$ relative to the bundle realisation, which is the content of Claim C3 for a single box and, for a long member, of the coverage lemma applied inside the member. We have not carried this out.

Summary of the attempt. Lemma 4.8 and Proposition 4.10 are proved; Proposition 4.9 corrects the formulation. The dependence step is reduced, under the weighted multiscale hypothesis $\mathcal{H}_2^E$, to the witness weight bound (4.14), which is a statement about the conditional law of the lines carrying a family of clusters, with the two unproved ingredients named above, plus the convergence of the chain iteration with relay charges, which is unchanged from yesterday.

5 The single-cluster bound: multiscale setting and the proved lemmas

This section and the next concern (S). We fix j_0 and write $\rho_L := \mu^{j_0} L^{-(j_0-2)}$; this is the order of the probability that j_0 lines of $\mathcal{L}$ run within distance 1 of each other across a box of side L, there being of order L^2 positions for such a bundle and each costing $(\mu/L)^{j_0}$ by Lemma 2.1(c). The aim is a bound of this order for the probability of a thick crossing of a box of side L, uniformly in a finite set of fixed lines that is carried along the induction.

5.1 Crossings, fixed lines and the two-parameter event

A *box of side* L is $B = z + [-L/2, L/2)^2$ with $z \in \mathbb{Z}^2$ and $L \in 2\mathbb{N}$, its *inner box* is $z + [-L/4, L/4)^2$, and its *boundary layer* is the set of unit boxes of B at sup-distance at least $L/2 - 1$ from z. For a finite set G of lines (*fixed lines*) call z *thick in* $\mathcal{L} \cup G$ if some point of $Q(z)$ has at least j_0 lines of $\mathcal{L} \cup G$ within distance 1, in accordance with (1.2); such a box has at least j_0 lines of $\mathcal{L} \cup G$ within distance $1 + \sqrt{2} < 3$ of its centre. A *crossing path* of B in $\mathcal{L} \cup G$ is a 2-path of boxes of B, thick in $\mathcal{L} \cup G$, from a box of the inner box to a box of the boundary layer. A crossing path has at least $L/8$ boxes.

Definition 5.1 (The event $\mathrm{Bad}(B; G, G')$)**.** Let $\varepsilon_L := c_\varepsilon/(\log L)^6$ with a constant $c_\varepsilon \in (0, 1/8]$. For finite sets $G' \subseteq G$ of lines meeting B, $\mathrm{Bad}(B; G, G')$ is the event that B contains a crossing path P in $\mathcal{L} \cup G$ such that fewer than $\varepsilon_L L$ boxes of P lie within distance 1 of a line of $G \setminus G'$. We write $\mathrm{Bad}(B, G) := \mathrm{Bad}(B; G, G)$.

Thus $\mathrm{Bad}(B, G)$ is the event of a thick crossing path in $\mathcal{L} \cup G$, and $\mathrm{Bad}(B; G, \varnothing)$ is the event of a crossing path that uses the fixed lines on a vanishing fraction of its boxes. The lines of $G \setminus G'$ are those the crossing *avoids*, the lines of G' those it may *use*.

Lemma 5.2 (From clusters to crossings)**.** *If the thick cluster of* 0 *has diameter at least* $d \geq 8$*, then* $\mathrm{Bad}(B, \varnothing)$ *holds for the box* B *of side* $L := 2\lfloor d/2 \rfloor$ *centred at* 0*. Consequently, if* $\mathbb{P}(\mathrm{Bad}(B, \varnothing)) \leq C\rho_L$ *for all boxes of side* $L \geq 8$ *and* $j_0 \geq \mu_1$*, then* $(\mathrm{S})_{j_0-2,A}$ *holds with* $A = \max\{C2^{j_0-2}, 8^{j_0-2}(4\mathrm{e}/(\pi j_0))^{j_0}\}\mu^{j_0}$*.*

Proof. The cluster of 0 is 2-connected, contains 0, which lies in the inner box, and contains a box at sup-distance at least $d/2 \geq L/2 - 1$ from 0, since a set of diameter d containing 0 has a point at sup-distance at least $d/2$ from 0; a 2-path inside the cluster from 0 to that box, stopped at its first box in the boundary layer, is a crossing path. For the consequence, $\rho_L \leq 2^{j_0-2}\rho_d$ since $L \geq d - 1 \geq d/2$, which gives the bound for $d \geq 8$; for $d < 8$ the probability is at most

$\mathbb{P}(0 \in T) \le (\mathrm{e}\mu_1/j_0)^{j_0}$ by Lemma 2.2, which is at most $8^{j_0-2}(4\mathrm{e}/(\pi j_0))^{j_0}\mu^{j_0}d^{-(j_0-2)}$. Stationarity transfers the bound from 0 to every box z. □

5.2 The coverage weight and the induction hypothesis

Definition 5.3 (Coverage multiplicity and weight)**.** Let G' be a finite set of lines meeting B. For $\ell_0 \in G'$ and a point $p \in \ell_0$ let $c_{G'}(p)$ be the number of lines of G', including ℓ_0, within distance 3 of p. The *coverage multiplicity* of ℓ_0 in B is

$$n_L(\ell_0) = n_{L,G'}(\ell_0) := \max\Big\{\min_{p\in I} c_{G'}(p) \colon I \subseteq \ell_0 \cap B \text{ a segment of length } \ge L/4\Big\}, \tag{5.1}$$

with $n_L(\ell_0) := 0$ if $\ell_0 \cap B$ is shorter than $L/4$. The *weight* of G' in B is

$$W_L(G') := 1 + \sum_{\ell_0 \in G'} k_L(n_L(\ell_0)), \qquad k_L(m) := K^m\mu^{-m}L^{m-2}\ (m \ge 1), \quad k_L(0) := 0\,, \tag{5.2}$$

with a constant $K \ge 1$ fixed below.

The weight is what the fixed lines are worth. If m fixed lines lie within distance 3 of every point of a stretch of ℓ_0 of length $L/4$, a crossing can ride that stretch using $j_0 - m$ random lines along it, at a cost of order $(\mu/L)^{j_0-m}$ by Lemma 2.1(c), against the unaided cost $\rho_L = (\mu/L)^{j_0}L^2$; the ratio is $\mu^{-m}L^{m-2}$. The weight charges each fixed line by the best stretch of length $L/4$ along it; shorter stretches are not charged at scale L, because a crossing path has to cover length $L/4$ and the rest of it has to be made thick by other means, which the recursion charges at smaller scales. The pointwise minimum in (5.1) is essential: it makes the weight correct for fixed lines that cover ℓ_0 by short tiles, which a definition in terms of lines aligned with ℓ_0 over a long stretch would miss (see Remark 5.5).

Definition 5.4 (Two-parameter hypothesis)**.** $\mathcal{H}_2(L)$ holds if for every box B of side L and all finite $G' \subseteq G$ of lines meeting B,

$$\mathbb{P}(\mathrm{Bad}(B; G, G')) \le C_0\,\rho_L\,W_L(G')\,. \tag{5.3}$$

The content of the two parameters is that avoided lines cost nothing: a crossing that is within distance 1 of them on fewer than $\varepsilon_L L$ of its boxes needs j_0 lines of $\mathcal{L} \cup G'$ on the rest, and the rest is almost all of the crossing. For $G = G' = \varnothing$, (5.3) is the bound $\mathbb{P}(\mathrm{Bad}(B, \varnothing)) \le C_0\rho_L$ needed in Lemma 5.2.

Remark 5.5 (Why the weight has this form). Two simpler weights fail. (a) A weight charging aligned groups, i.e. sets of fixed lines within distance 1 of a common axis throughout B, by k_L(group size), is false: take $j_0 - 1$ lines forming a group with axis ℓ_0 and $L^{1/2}$ further lines each within distance 1 of ℓ_0 along a stretch of length $L^{1/2} - 1$, the stretches tiling $\ell_0 \cap B$; every box along ℓ_0 is thick in G alone, so $\mathbb{P}(\mathrm{Bad}(B, G)) = 1$, while the group weight is $k_L(j_0 - 1)\rho_L = K^{j_0-1}\mu/L$. The same example with tiles of length $\tau(L) - 1$ defeats any alignment threshold $\tau(L)$. (b) A weight charging each fixed line by the maximal number of fixed lines within distance 3 of a single point of it is correct for (a) but is not closed under the shared lines of the recursion (Section 6): a random line crossing ℓ_0 transversally inside a sub-box of side ℓ raises that multiplicity by one, which multiplies the weight by $K\ell/\mu$, while such lines through a sub-box have measure of order $a^2\mu/\ell$ (Lemma 5.10), so that the expected weight of a sub-box would be multiplied by a factor of order a^2K, not by $1 + o(1)$. The coverage multiplicity (5.1) is insensitive to transversal lines, which

raise $c_{G'}(p)$ only on a stretch of length $O(1/\alpha)$ for a line at angle α, and the pointwise minimum over a stretch of length $L/4$ is raised only by lines within distance 3 of ℓ_0 along all of it, which are nearly parallel to ℓ_0 and have small measure by Lemma 2.1(c). Our working note [HJ26b] used the weight (b) and claimed closure under shared lines through a second-moment lemma that is incorrect; see Section 7.

5.3 Sub-boxes, annuli and separation

Fix $\gamma = 3$ and an integer $a \geq 2$, let $A := 64a$, and for $L \geq L_1$ put $\ell := 2\lceil L^{1/3}/2 \rceil$ and $D := L/(4A) = L/(256a)$. Cover B by the boxes of side ℓ centred at the points of $z + (\ell/2)\mathbb{Z}^2$ that meet B (the *sub-boxes*); every point of B lies in the inner box of one of them, the inner boxes of distinct sub-boxes are disjoint, and there are at most $(4L/\ell)^2$ sub-boxes. The *annuli* of B are

$$\mathcal{A}_s := \{x \in B \colon L/4 + (s-1)D \leq \|x - z\|_\infty < L/4 + sD\}, \qquad s = 1, \dots, A;$$

they are disjoint, of width D, and points of $\mathcal{A}_s$ and $\mathcal{A}_{s'}$ with $|s - s'| \geq 2$ are at distance at least D. Every crossing path of B meets every annulus, since it runs from the inner box to the boundary layer, and sup-distance $L/4 + AD = L/2$ is the outer edge.

Lemma 5.6 (Chains of sub-crossings)**.** *Let P be a crossing path of B in $\mathcal{L} \cup G$ and $L \geq L_1(a)$. There are sub-boxes $B'_1, \dots, B'_r$ and boxes $p_1, \dots, p_r$ of P, in the order of P, such that p_i lies in the inner box of B'_i, the part P_i of P from p_i to its first subsequent box in the boundary layer of B'_i is a crossing path of B'_i in $\mathcal{L} \cup G$, and p_{i+1} is the box of P following the last box of P_i. The P_i are pairwise disjoint, every sub-box B'_i lies within distance 2ℓ of p_i, and for every annulus $\mathcal{A}_s$ at least $D/(4\ell)$ of the boxes p_i lie in $\mathcal{A}_s$.*

Proof. Let p_1 be the first box of P and B'_1 a sub-box whose inner box contains it. Given p_i and B'_i, if P has a box in the boundary layer of B'_i after p_i, let P_i be the part of P from p_i to the first such box; its boxes are thick in $\mathcal{L} \cup G$, so it is a crossing path of B'_i. Let p_{i+1} be the next box of P, if any, and B'_{i+1} a sub-box whose inner box contains it. If P has no box in the boundary layer of B'_i after p_i, stop and put $r := i - 1$. The P_i are consecutive disjoint segments of P. A sub-box has diameter $\sqrt{2}\ell$ and its inner box contains p_i, so B'_i lies within distance 2ℓ of p_i. Consecutive boxes p_i, p_{i+1} are at sup-distance at most $\sqrt{2}\ell + 2 \leq 2\ell$, since P_i stays in B'_i and one step of P has sup-norm at most 2. The last box p_{r+1} at which the construction stops lies within sup-distance ℓ of the end of P, which is in the boundary layer of B; hence the sup-distances $\|p_i - z\|_\infty$, $i \leq r+1$, form a sequence starting below $L/4$, ending above $L/2 - 1 - \ell$, with consecutive differences at most 2ℓ. Such a sequence has at least $D/(2\ell) - 1 \geq D/(4\ell)$ terms in every interval of length D contained in $[L/4, L/2 - 1 - \ell]$; the annuli are such intervals for $L \geq L_1(a)$, because $D \geq 4\ell$ and $AD = L/4$, and the interval of $\mathcal{A}_A$ is shortened by at most $1 + \ell$. □

Lemma 5.7 (Separated sub-boxes in distinct annuli)**.** *Let $S \subseteq \{1, \dots, A\}$ with $|S| \geq 2a$ and let B'_{i_s}, $s \in S$, be sub-boxes of the chain of Lemma 5.6 with $p_{i_s} \in \mathcal{A}_s$. Then there are a indices $s_1, \dots, s_a \in S$ such that the sub-boxes $B'_{i_{s_1}}, \dots, B'_{i_{s_a}}$ are pairwise at distance at least $D - 4\ell \geq D/2$ for $L \geq L_1(a)$. The number of a-tuples of sub-boxes is at most $(4L/\ell)^{2a}$.*

Proof. Choose a elements of S pairwise differing by at least 2, which is possible since $|S| \geq 2a$. Points of two annuli with indices differing by at least 2 are at distance at least D, and by Lemma 5.6 each sub-box lies within distance 2ℓ of its box p_i. The count is the number of a-tuples among $(4L/\ell)^2$ sub-boxes. □

Remark 5.8. The working note [HJ26b] chose separated sub-boxes by taking every $\lceil L/(4a\ell)\rceil$-th member of the chain, which does not guarantee spatial separation, because a crossing path may wind inside a region of diameter D for many chain steps. The annuli repair this at the price of constants: the separation D is $L/(256a)$ instead of $L/(8a)$.

5.4 Shared and private lines

Let $B_1, \dots, B_a$ be sub-boxes pairwise at distance at least $D/2$. A line of $\mathcal{L}$ is *shared* if it meets at least two of them and *private to* B_i if it meets B_i and no other B_j. Let $\mathcal{S}$ be the set of shared lines and $\mathcal{P}_i$ that of the private lines of B_i; write $\mathcal{S}_i := \{\ell' \in \mathcal{S} : \ell' \cap B_i \neq \varnothing\}$.

Lemma 5.9 (Conditional independence)**.** *Given $\mathcal{S}$, the processes $\mathcal{P}_1, \dots, \mathcal{P}_a$ are independent Poisson processes with intensity measures ν restricted to the respective sets of lines, and for finite sets $G_i' \subseteq G_i$ of lines meeting B_i,*

$$\mathbb{P}\Big(\bigcap_{i\le a} \mathrm{Bad}(B_i; G_i, G_i')\Big) = \mathbb{E}\Big[\prod_{i\le a} \mathbb{P}(\mathrm{Bad}(B_i; G_i \cup \mathcal{S}_i, G_i' \cup \mathcal{S}_i) \mid \mathcal{S})\Big], \tag{5.4}$$

where inside the conditional probability the lines of $\mathcal{S}_i$ are fixed lines and the random lines are those of $\mathcal{P}_i$. The same holds with G_i' in place of $G_i' \cup \mathcal{S}_i$ on the right if the crossing paths in the events on the left are required to avoid the shared lines in the sense of Definition 5.1.

Proof. The sets of lines meeting exactly one B_i, for each i, and of lines meeting at least two, are disjoint measurable sets of lines, and the restrictions of a Poisson process to disjoint sets are independent Poisson processes. The event $\mathrm{Bad}(B_i; G_i, G_i')$ depends only on the lines meeting B_i, which are private to B_i, shared, or fixed; given $\mathcal{S}$ it is the event $\mathrm{Bad}(B_i; G_i \cup \mathcal{S}_i, G_i' \cup \mathcal{S}_i)$ for the process $\mathcal{P}_i$ with the shared lines fixed, and these events are conditionally independent. □

Lemma 5.10 (Shared lines are few and aligned)**.** *Let B_i, B_j be sub-boxes of side ℓ at distance $D' \ge D/2$. The lines meeting both have ν-measure at most $8\mu\ell^2/D'$ and pairwise make angles at most $8\ell/D'$. For a line ℓ_0 and a segment $I \subseteq \ell_0 \cap B_i$ of length at least $\ell/4$, the lines meeting B_j that lie within distance 3 of ℓ_0 along all of I have measure at most $20\mu\ell/D'$. In particular, with $D' \ge D/2 = L/(512a)$ and $L = \ell^3$ up to rounding, the shared lines through a sub-box have mean at most $a \cdot 8\mu\ell^2/D' \le 2^{12}a^2\mu/\ell$, and those within distance 3 of a given line along a stretch of length $\ell/4$ in the sub-box have mean at most $2^{14}a^2\mu/\ell^2$.*

Proof. Lemma 2.1(b) with $\delta = \sqrt{2}\ell$ gives measure at most $4\mu \cdot 2\ell^2/D' = 8\mu\ell^2/D'$ and angle at most $4\sqrt{2}\ell/D' \le 8\ell/D'$. For the last statement, a line within distance 3 of ℓ_0 along I makes an angle at most $\pi \cdot 6/(\ell/4) \cdot \frac{1}{2} \le 40/\ell$ with ℓ_0 and passes within distance 3 of a fixed endpoint of I; a line meeting B_j and passing within distance 3 of that endpoint has direction within angle $\pi\sqrt{2}\ell/D' \le 5\ell/D'$ of the direction from the endpoint to the centre of B_j, as in Lemma 2.1(b). The direction therefore lies in an interval of length at most $\min\{80/\ell, 10\ell/D'\} \le 10\ell/D'$, the offset in an interval of length 6, and the measure is at most $(\mu/\pi) \cdot 10\ell/D' \cdot 6 \le 20\mu\ell/D'$. □

5.5 Inheritance of avoidance

Lemma 5.11 (Avoidance is inherited to most annuli)**.** *Let P be a crossing path of B with fewer than $\varepsilon_L L$ boxes within distance 1 of a set H of lines, and let B_i', P_i be as in Lemma 5.6. Call a chain member i* avoiding *if P_i has fewer than $\varepsilon_\ell \ell$ boxes within distance 1 of H, and call an annulus $\mathcal{A}_s$* avoiding *if at least half of the chain members with $p_i \in \mathcal{A}_s$ are avoiding. Then, for $L \ge L_1$, at least $A - A/10$ annuli are avoiding.*

Proof. A non-avoiding annulus has at least $D/(8\ell)$ non-avoiding chain members, each contributing at least $\varepsilon_\ell \ell$ boxes of P within distance 1 of H, and the P_i are pairwise disjoint; so n non-avoiding annuli force at least $n\varepsilon_\ell D/8$ such boxes, whence $n < 8\varepsilon_L L/(\varepsilon_\ell D) = 32A\varepsilon_L/\varepsilon_\ell$. Now $\varepsilon_\ell/\varepsilon_L = (\log L/\log \ell)^6 \geq (3-o(1))^6 > 700$ for $L \geq L_1$, since $\ell \leq 2L^{1/3}+2$, so $n < 32A/700 < A/10$. □

5.6 The arithmetic of the recursion

Lemma 5.12 (Basic inequality)**.** *Let $j_0 \geq 7$, $\delta > 0$ and*

$$a \;\geq\; \frac{3(j_0-2)+2\delta}{j_0-6}\,. \tag{5.5}$$

Then for every $C_0 \geq 1$ there is $\ell_0 = \ell_0(C_0, a, \delta)$ such that for $\ell \geq \ell_0$ and $L = \ell^3$,

$$(4L/\ell)^{2a}\,(2C_0\rho_\ell)^a \leq C_0\,\rho_L\,\ell^{-\delta}\,. \tag{5.6}$$

Proof. With $\rho_\ell = \mu^{j_0}\ell^{-(j_0-2)}$ and $\rho_L = \mu^{j_0}\ell^{-3(j_0-2)}$ the inequality reads

$$4^{2a}2^a C_0^{a-1}\mu^{(a-1)j_0}\,\ell^{\,4a-a(j_0-2)+3(j_0-2)+\delta} \leq 1\,.$$

Since $\mu \leq 1$ may be assumed, it suffices that the exponent of ℓ be at most $-\delta$ and that $\ell^\delta \geq 4^{2a}2^aC_0^{a-1}$; the first is (5.5), the second holds for $\ell \geq \ell_0$. □

With two sub-boxes and super-exponentially growing scales, as in the standard renormalisation schemes, the exponent of ℓ in (5.6) is positive; a polynomial target forces a separated sub-boxes and geometric scales.

5.7 The coverage lemma

The following single-scale estimate is the core of every statement about crossings that ride a fixed line. Let ℓ_0 be a line and $J \subseteq \ell_0$ a segment of length L, partitioned into L consecutive *cells* of length 1. For a line $\ell' \in \mathcal{L}$ its *footprint* on J is the set of cells whose midpoint is within distance 3 of ℓ'; it is a set of consecutive cells, of cardinality $t(\ell') \leq 6/\sin\alpha + 2$ where α is the angle between ℓ' and ℓ_0. A cell is *x-covered* if it belongs to the footprints of at least x lines of $\mathcal{L}$.

Lemma 5.13 (Coverage)**.** *Let $c \in (0,1]$ and $x \geq 1$. There are $\mu_0(c) > 0$, $C(c) < \infty$ and $L_0(c,\mu)$ such that for $\mu \leq \mu_0(c)$ and $L \geq L_0(c,\mu)$,*

$$\mathbb{P}\big(\text{at least } cL \text{ cells of } J \text{ are } x\text{-covered}\big) \leq \Big(\frac{C(c)\,\mu}{L}\Big)^x\,. \tag{5.7}$$

Proof. Step 1: reduction to single coverings. Let E be the set of x-covered cells and let $\mathcal{F}$ be the set of lines whose footprint meets E. The hypergraph with vertex set E and edges $\{F(\ell')\cap E\colon \ell' \in \mathcal{F}\}$ is an interval hypergraph in the linear order of the cells, so its incidence matrix has the consecutive-ones property and is totally unimodular, and every vertex lies in at least x edges. By the integer decomposition property of totally unimodular systems [BT78], the all-ones vector on the edges, which lies in x times the polytope $\{0 \leq y \leq 1,\ My \geq 1\}$, is a sum of x integral vectors of that polytope; that is, $\mathcal{F}$ contains x pairwise disjoint subfamilies $\mathcal{F}_1, \dots, \mathcal{F}_x$ each of whose footprints cover E. In particular, on the event in (5.7) there are x pairwise disjoint finite sets of lines of $\mathcal{L}$ each covering at least cL cells.

Step 2: dyadic classes. Put $k_1 := \lfloor \log_2 L \rfloor$. For $0 \leq k \leq k_1$ let $\mathcal{L}_k$ be the set of lines within distance 3 of J with footprint cardinality in $[2^k, 2^{k+1})$, and let $N_k := |\mathcal{L}_k \cap \mathcal{L}|$, a Poisson variable

with mean $m_k := \nu(\mathcal{L}_k)$; every footprint has cardinality at most $L < 2^{k_1+1}$, so these classes exhaust the lines meeting the 3-neighbourhood of J. A line in $\mathcal{L}_k$ with $k \geq 3$ has $\sin\alpha \leq 6/(2^k-2) \leq 8/2^k$, so its direction lies in a set of measure at most $4 \cdot (\pi/2) \cdot 8/2^k \leq 51/2^k$, and for a given direction its offset lies in an interval of length at most $L\sin\alpha + 6 \leq 8L/2^k + 6$. Hence, using $2^k \leq L$,

$$m_k \leq \frac{\mu}{\pi} \cdot \frac{51}{2^k}\Big(\frac{8L}{2^k} + 6\Big) \leq \frac{2000\,\mu L}{4^k} \qquad (0 \leq k \leq k_1)\,, \tag{5.8}$$

where for $k \leq 2$ we used $m_k \leq \nu(\text{lines within distance 3 of } J) \leq 2\mu(L+6) \leq 2000\mu L/4^k$.

Step 3: weights. Let k_2 be the largest $k \geq 0$ with $4^k \leq \mu L/(2\log(L/\mu))$, which exists for $L \geq L_0(\mu)$, and define

$$w_k := \frac{400\mu}{c\,2^k} \quad (k \leq k_2), \qquad w_k := \frac{3}{\pi^2} \cdot \frac{1}{(k_1 - k + 1)^2} \quad (k_2 < k \leq k_1)\,,$$

so that $\sum_k w_k \leq 800\mu/c + 1/2 \leq 1$ for $\mu \leq \mu_0(c) := c/1600$. If a family of lines covers at least cL cells, then for some k its members in $\mathcal{L}_k$ cover at least cLw_k cells, and since each of them covers fewer than 2^{k+1} cells, the family contains at least $n_k := cLw_k/2^{k+1}$ lines of $\mathcal{L}_k$.

Step 4: the quantities q_k. We define numbers q_k such that for every integer $r \geq 1$,

$$\mathbb{P}(N_k \geq r\lceil n_k\rceil) \leq q_k^r\,, \qquad \text{and show} \qquad \sum_{k \leq k_1} q_k \leq \frac{C(c)\,\mu}{L}\,. \tag{5.9}$$

For $k \leq k_2$: $n_k = 200\mu L/4^k \geq 10 m_k$, so by Lemma 2.2 $\mathbb{P}(N_k \geq r\lceil n_k\rceil) \leq (\mathrm{e} m_k/(r n_k))^{r n_k} \leq (\mathrm{e}/10)^{r n_k}$; put $q_k := \mathrm{e}^{-n_k}$. Since $n_k \geq 200\mu L/4^{k_2} \geq 400\log(L/\mu)$, $q_k \leq (\mu/L)^{400}$. For $k_2 < k \leq k_1$ write $j := k_1 - k \geq 0$; from $L/2 < 2^{k_1} \leq L$ we get

$$n_k = \frac{3cL}{\pi^2(j+1)^2 2^{k+1}} \geq \frac{3c\,2^j}{2\pi^2(j+1)^2}, \qquad m_k \leq \frac{2000\mu L 4^j}{4^{k_1}} \leq \frac{8000\,\mu\,4^j}{L}\,.$$

If $n_k \leq 1$, then $\mathbb{P}(N_k \geq r) \leq (\mathrm{e} m_k/r)^r \leq (\mathrm{e} m_k)^r$, and we put $q_k := \mathrm{e} m_k \leq \mathrm{e} \cdot 8000\mu 4^j/L$; here $n_k \leq 1$ forces $2^j \leq 7(j+1)^2/c$, so $j \leq j_c$ for a constant $j_c = j_c(c)$ and $q_k \leq C_2(c)\mu/L$, for at most $j_c + 1$ values of k. If $n_k > 1$, let $b_k := \mathrm{e} m_k/\lceil n_k\rceil \leq \mathrm{e} m_k/n_k \leq 2 \cdot 10^5 \mu 2^j (j+1)^2/(cL)$. Since $k > k_2$ we have $4^k > \mu L/(8\log(L/\mu))$, hence $2^j = 2^{k_1}/2^k \leq \sqrt{8L\log(L/\mu)/\mu}$, and therefore $b_k \leq 2 \cdot 10^5 \mu (\log_2 L + 1)^2 \sqrt{8\log(L/\mu)/(\mu L)}/c \leq 1/2$ for $L \geq L_0(c,\mu)$. Then $\lceil n_k\rceil \geq 2\mathrm{e} m_k$ and Lemma 2.2 gives $\mathbb{P}(N_k \geq r\lceil n_k\rceil) \leq b_k^{r\lceil n_k\rceil}$; put $q_k := b_k^{\lceil n_k\rceil}$. For the sum of these q_k: if $\lceil n_k\rceil \geq 3$, then $q_k \leq b_k^3 \leq C_3\mu^3 8^j (j+1)^6/(c^3 L^3)$, and $\sum_{j \leq \log_2 \sqrt{8L\log(L/\mu)/\mu}} 8^j (j+1)^6 \leq C_4 (\log L)^6 (L\log(L/\mu)/\mu)^{3/2}$, so these terms sum to at most $C_5(c)\mu^{3/2}(\log L)^8/L^{3/2} \leq \mu/L$ for $L \geq L_0(c,\mu)$; if $\lceil n_k\rceil = 2$, then $2^j \leq 14(j+1)^2/c$, so $j \leq j_c'$, and $q_k \leq b_k^2 \leq C_6(c)\mu^2/L^2$. Collecting,

$$\sum_{k \leq k_1} q_k \leq (k_2+1)\Big(\frac{\mu}{L}\Big)^{400} + (j_c+1)\frac{C_2(c)\mu}{L} + (j_c'+1)\frac{C_6(c)\mu^2}{L^2} + \frac{\mu}{L} \leq \frac{C(c)\,\mu}{L}\,.$$

Step 5: x families. By Steps 1 and 3, on the event in (5.7) there are pairwise disjoint families $\mathcal{F}_1, \dots, \mathcal{F}_x$ of lines of $\mathcal{L}$ and indices $k^{(1)}, \dots, k^{(x)}$ such that $\mathcal{F}_i$ contains at least $\lceil n_{k^{(i)}}\rceil$ lines of $\mathcal{L}_{k^{(i)}}$. With $r_k := \#\{i\colon k^{(i)} = k\}$, disjointness gives $N_k \geq r_k\lceil n_k\rceil$ for every k. The variables N_k are independent, so by (5.9) and a union bound over the x-tuples $(k^{(1)}, \dots, k^{(x)})$,

$$\mathbb{P} \leq \sum_{(k^{(i)})_{i \leq x}} \prod_k q_k^{r_k} = \Big(\sum_{k \leq k_1} q_k\Big)^x \leq \Big(\frac{C(c)\mu}{L}\Big)^x\,. \qquad \square$$

6 The multiscale recursion and its claims

Theorem 6.1 (Multiscale recursion, conditional)**.** *Let $j_0 \geq 7$, let a satisfy (5.5) with some $\delta > 0$, let $K \geq 1$, $C_0 \geq 1$ and $\mu \leq \mu_1(j_0, a, K)$, and let $L_1 = L_1(j_0, a, K, C_0, \mu)$ be large. Assume Claims C1 and C2. If $\mathcal{H}_2(L')$ holds for all $L' \leq L/8$, then $\mathcal{H}_2(L)$ holds, for every $L \geq L_1$. If in addition Claim C3 holds, then $\mathcal{H}_2(L)$ holds for all $L \geq 8$, and $(\mathrm{S})_{j_0-2,A}$ holds with $A = A'(j_0)\mu^{j_0}$.*

Proof. Let B be a box of side $L \geq L_1$ and $G' \subseteq G$ finite sets of lines meeting B, $H := G \setminus G'$. On $\mathrm{Bad}(B; G, G')$ let P be a crossing path with fewer than $\varepsilon_L L$ boxes within distance 1 of H, and let B_i', P_i, p_i be its chain (Lemma 5.6). Call a chain member i *avoiding* if P_i has fewer than $\varepsilon_\ell \ell/2$ boxes within distance 1 of H, of *kind F* if moreover P_i has fewer than $\varepsilon_\ell \ell$ boxes within distance 1 of G, and of *kind R* if it is avoiding but not of kind F; a member of kind R has at least $\varepsilon_\ell \ell/2$ boxes within distance 1 of lines of G'. By Lemma 5.11, applied with the threshold $\varepsilon_\ell \ell/2$, which changes the constant 32 there to 64, at least $A - A/10$ annuli are avoiding, i.e. have at least half of their members avoiding. Two cases.

Case F: at least $2a$ avoiding annuli contain a member of kind F. By Lemma 5.7 there are members $i_1, \dots, i_a$ of kind F whose sub-boxes $B_1, \dots, B_a$ are pairwise at distance at least $D/2$. For each of them, P_{i_s} is a crossing path of B_s in $\mathcal{L} \cup G$ with fewer than $\varepsilon_\ell \ell$ boxes within distance 1 of $G \cap B_s$; with the shared lines $\mathcal{S}$ of $B_1, \dots, B_a$ fixed, this is the event $\mathrm{Bad}(B_s; (G \cap B_s) \cup \mathcal{S}_s, \mathcal{S}_s)$ for the private process $\mathcal{P}_s$. By Lemma 5.9, the union bound over the at most $(4L/\ell)^{2a}$ choices of the sub-boxes, $\mathcal{H}_2(\ell)$ applied inside the conditional probability, and Claim C1,

$$\mathbb{P}(\text{Case F}) \leq \Big(\frac{4L}{\ell}\Big)^{2a} \mathbb{E}\Big[\prod_{s \leq a} C_0 \rho_\ell W_\ell(\mathcal{S}_s)\Big] \leq \Big(\frac{4L}{\ell}\Big)^{2a} (2C_0\rho_\ell)^a \leq C_0 \rho_L \ell^{-\delta} \leq \tfrac{1}{2} C_0 \rho_L$$

by Lemma 5.12, for $\ell \geq \ell_0$. The fixed lines of G enter this case only through the requirement that the sub-crossings avoid them, and the bound does not depend on G.

Case R: fewer than $2a$ avoiding annuli contain a member of kind F. Then at least $A - A/10 - 2a \geq A/2$ avoiding annuli have all their avoiding members, which are at least half of their members, of kind R. By Claim C2, $\mathbb{P}(\text{Case R}) \leq \frac{1}{2} C_0 \rho_L W_L(G')$.

Adding the two cases gives (5.3) at scale L, since $W_L(G') \geq 1$. For the second assertion, Claim C3 gives $\mathcal{H}_2(L)$ for $8 \leq L \leq L_1$, and for $L > L_1$ the sub-box side $\ell = 2\lceil L^{1/3}/2 \rceil$ satisfies $8 \leq \ell \leq L/8$, so induction on L applies. Finally $\mathcal{H}_2(L)$ with $G = G' = \varnothing$ gives $\mathbb{P}(\mathrm{Bad}(B, \varnothing)) \leq C_0 \rho_L$ for all $L \geq 8$, and Lemma 5.2 gives $(\mathrm{S})_{j_0-2,A}$ with $A = A'(j_0)\mu^{j_0}$. □

6.1 The weight of the shared lines

Claim C1 (Shared-line weights). Let $B_1, \dots, B_a$ be sub-boxes of side ℓ pairwise at distance at least $D/2$, let $\mathcal{S}_s$ be the shared lines through B_s and $W_\ell(\mathcal{S}_s)$ the weight (5.2) of $\mathcal{S}_s$ in B_s. Then, for $\ell \geq \ell_0(a, K, \mu)$,

$$\mathbb{E}\Big[\prod_{s \leq a} W_\ell(\mathcal{S}_s)\Big] \leq 2^a . \tag{6.1}$$

What is known about Claim C1. The weight $W_\ell(\mathcal{S}_s) = 1 + \sum_{\ell' \in \mathcal{S}_s} k_\ell(n_\ell(\ell'))$ counts a shared line ℓ' with multiplicity $n_\ell(\ell') = m$ if $m - 1$ further shared lines lie within distance 3 of ℓ' along a stretch of length $\ell/4$ in B_s. First moments are small. The shared lines through B_s form a Poisson

process of mean at most $\lambda := 2^{12}a^2\mu/\ell$ (Lemma 5.10), so the expected number of singletons is at most λ, each of weight $k_\ell(1) = K/(\mu\ell)$, total $2^{12}a^2K/\ell^2$. Given a shared line ℓ', the shared lines within distance 3 of it along a stretch of length $\ell/4$ have mean at most $\lambda' := 2^{14}a^2\mu/\ell^2$ (Lemma 5.10, since such a line is shared with one of the other $a-1$ sub-boxes), so the expected number of shared lines with multiplicity at least $m \geq 2$ is at most $\lambda\lambda'^{m-1}/(m-1)!$, and with $k_\ell(m) = K^m\mu^{-m}\ell^{m-2}$ the expected total weight of lines of multiplicity m is at most

$$\lambda\frac{\lambda'^{m-1}}{(m-1)!}k_\ell(m) = \frac{2^{12}a^2K}{\ell^2}\cdot\frac{(2^{14}a^2K/\ell)^{m-1}}{(m-1)!}\,,$$

whose sum over $m \geq 2$ is at most $2^{27}a^4K^2\ell^{-3}$ for $\ell \geq 2^{15}a^2K$. Hence $\mathbb{E}W_\ell(\mathcal{S}_s) \leq 1 + 2^{13}a^2K/\ell^2$ for such ℓ, and the first-moment version of (6.1) with a single sub-box holds. What is not written out is the expectation of the product. The factors are not independent, since a shared line lies in at least two of the sub-boxes and contributes to each of their weights; the contribution of one line ℓ' to two factors is $k_\ell(n^{(s)}(\ell'))k_\ell(n^{(s')}(\ell'))$, which for singletons is $K^2/(\mu\ell)^2$ against the mean λ of such lines, a product $2^{12}a^2K^2/(\mu\ell^3)$ that is small for $\ell \geq \ell_0(a, K, \mu)$; and for a line of multiplicity m in B_s and m' in $B_{s'}$, the $m-1$ and $m'-1$ companions must be within distance 3 of ℓ' along stretches in two sub-boxes at distance $D/2$, which pins their directions to within $O(\ell/D)$ and makes the pair of stretches cost $O(\mu/D)$ per companion aligned in both, as in Lemma 2.1(c). Expanding the product $\prod_s(1 + \sum_{\ell'} k_\ell(n^{(s)}(\ell')))$ and bounding each term by the Mecke formula is routine but has to be organised by the pattern of which lines appear in which factors; we have not done it. The second-moment route of [HJ26b] is not available, because the second moment of a single weight is not bounded (the gain $K\ell/\mu$ of one aligned shared line squared against its cost λ' is $2^{14}a^2K^2/\mu$, which is large), and the lemma asserting the contrary in [HJ26b] is wrong; the product in (6.1) must be handled at first moment, using that a line aligned with a stretch in two far sub-boxes is much rarer than one aligned in one.

6.2 Crossings that ride the fixed lines

Claim C2 (Rides). In the setting of Theorem 6.1, assume $\mathcal{H}_2(L')$ for all $L' \leq L/8$. The probability that $\mathrm{Bad}(B; G, G')$ holds with a crossing path P for which at least $A/2$ annuli have at least half of their chain members of kind R is at most $\frac{1}{2}C_0\rho_L W_L(G')$.

What is known about Claim C2. A member of kind R has at least $\varepsilon_\ell\ell/2$ boxes within distance 1 of lines of G'; at such a box q, with $x \in Q(q)$ a point having j_0 lines of $\mathcal{L} \cup G$ within distance 1 and p the point of the line $\ell_0 \in G'$ nearest to x, all these lines pass within distance $2 + \sqrt{2} < 4$ of p, so at least $j_0 - c_G(p)$ random lines pass within distance 4 of p (the constant 3 of Definition 5.3 and Lemma 5.13 is to be read as 4 here, which changes only numerical constants). In each of the $A/2$ annuli at least $D/(8\ell)$ members contribute $\varepsilon_\ell\ell/2$ such boxes each, so P has at least $A/2 \cdot D/(8\ell) \cdot \varepsilon_\ell\ell/2 = \varepsilon_\ell L/128$ boxes within distance 1 of lines of G', distributed over the $A/2$ annuli. The claim is that this costs at least $\frac{1}{2}C_0\rho_L W_L(G')$ in probability. Three mechanisms are involved.

(i) One line, long ride. If a single line $\ell_0 \in G'$ carries at least cL of these boxes with $c > 0$ fixed, the cells of ℓ_0 at these boxes are covered $(j_0 - c_G(p))$-fold by random lines. If the fixed coverage $c_G(p)$ along the ridden cells were at most m, Lemma 5.13 with $x = j_0 - m$ would give the probability $(C\mu/L)^{j_0-m} = C^{j_0-m}K^{-m}\rho_L k_L(m) \leq C^{j_0}\rho_L k_L(m)$, which is the term of $W_L(G')$

belonging to ℓ_0 provided $m \le n_L(\ell_0)$. The difficulty is that $c_G(p)$ varies along ℓ_0 and the ride may select the cells of high fixed coverage: fixed lines may cover ℓ_0 j_0-fold on tiles of total length cL without any stretch of length $L/4$ having coverage above 1, in which case $n_L(\ell_0) = 1$ while the ride on the tiles is free. The ride on the tiles, however, does not cross B: the crossing has to connect the tiles, and between two tiles at distance t it either rides ℓ_0 with fixed coverage at most 1, at a cost $(C\mu/t)^{j_0-1}$ by Lemma 5.13, or leaves ℓ_0. The correct statement for one line is therefore about the *gaps* between the stretches of high fixed coverage along ℓ_0, and the pointwise minimum in (5.1) over stretches of length $L/4$ is designed so that the sum over the gaps of their costs is dominated by $k_L(n_L(\ell_0))\rho_L$; we have not written this out. Note also that the fixed lines of H may contribute to $c_G(p)$, but on fewer than $\varepsilon_L L$ boxes of P, which is less than $\varepsilon_\ell L/256$ for $L \ge L_1$, so that they can be discarded at the cost of halving the count.

(ii) Many lines. The ridden boxes may be distributed over many lines of G', each ridden briefly. A ride of length t along ℓ_0 with fixed coverage at most m costs $(C\mu/t)^{j_0-m}$ and gains advance t; a crossing consisting of q rides of lengths $t_1, \dots, t_q$ with $\sum t_i \ge L/8$, joined at crossing points of lines of G', costs at most $\prod_i (C\mu/t_i)^{x_i}$ with $x_i \ge 1$, and the sum over the realisable sequences, organised by the longest ride, is bounded by $\sum_{\text{start}} (4C\mu q/L)^{x_{\min}} \sum_q q^{1+x_{\min}} (j_0 (C\mu)^{x_{\min}} \zeta(x_{\min}))^{q-1}$, which converges for $\mu \le \mu_1(j_0)$ and is of the order $(\mu/L)^{x_{\min}}$ times the number of possible starting lines, as in [HJ26b, Section 7]. Two points are not settled: a random line may serve the ends of two consecutive rides near their crossing point, which reduces the count of random lines by one per switch unless the two fixed lines are nearly parallel there, in which case the coverage multiplicity accounts for it; and the rides may be joined by *excursions* of P away from G', which are crossings of smaller boxes avoiding G' and are to be charged by $\mathcal{H}_2$ at the scale of the excursion, with the random lines of the adjacent rides as fixed lines.

(iii) The constant. Even when (i) and (ii) are carried out, the bound obtained must be at most $\frac{1}{2} C_0 \rho_L W_L(G')$ with the same C_0 as in the hypothesis, which requires that the constants produced by Lemma 5.13 and by the sums in (ii) be absorbed, for instance by a factor $\ell^{-\delta'}$ gained from the fact that a ride of length comparable to L along a fixed line has no positional entropy while the target ρ_L contains the factor L^2; the constant K in the weight is free and can be taken large for this purpose.

6.3 The base case

Claim C3 (Base case). For every L_1 there is $\mu_1(L_1, j_0) > 0$ such that $\mathcal{H}_2(L)$ holds for all $8 \le L \le L_1$ and $\mu \le \mu_1$, with a constant $C_0 = C_0(j_0)$ not depending on L_1.

What is known about Claim C3. By Lemma 2.3 applied to the increasing event $\mathrm{Bad}(B; G, G')$ (increasing in $\mathcal{L}$ for fixed G, G'), its probability is at most $\sum_n M_n/n!$ with M_n the measure of n-tuples of lines meeting B that form a minimal set producing the event. Each line meeting B has measure at most $2\sqrt{2}\mu L \le 3\mu L$ (Lemma 2.1(a)), so the terms with $n > 2j_0$ sum to at most $(3\mathrm{e}\mu L/(2j_0))^{2j_0} \le \rho_L$ for $\mu \le cL_1^{-5}$, say. For $n \le 2j_0$ the thick boxes of $\Lambda \cup G$, $|\Lambda| = n$, meet the union over j_0-subsets of the common 1-neighbourhoods of their lines, each of which is contained in a tube of width 2 around any one of its lines; the boxes themselves lie in the tube of width $2 + 2\sqrt{2} < 5$. The lines of H are within distance 1 of fewer than $\varepsilon_L L \le L/64$ boxes of the crossing path, so a part of the crossing of diameter at least $L/8 - L/64$ is covered by tubes around lines of $\Lambda \cup G'$, and with $r := \binom{2j_0+|G'|}{j_0}$ tubes at most, one tube covers a connected part of diameter at

least $L/(16r)$: j_0 lines of $\Lambda \cup G'$ pass within distance 3 of two boxes at distance $t \geq L/(16r)$. If s of them are fixed, the $j_0 - s$ random ones pass within distance 3 of a stretch of length t along a fixed line at one of at most $16r$ positions, of measure at most $(16r)(16\mu/t)^{j_0-s}$ by Lemma 2.1(c), while if none is fixed the first random line has measure at most $3\mu L$ and the others $(16\mu/t)^{j_0-1}$ each; the remaining $n - j_0$ lines contribute at most $(3\mu L)^{n-j_0} \leq 1$ for $\mu \leq 1/(6L_1)$. The s fixed lines in a common tube of length $t \geq L/(16r)$ are within distance 5 of each other along that stretch, so the coverage multiplicity of each of them is at least s when $t \geq L/4$; for $t < L/4$ the weight does not record them, and this is the point that remains: the bound then is $(16r)^{j_0}(16\mu \cdot 16r/L)^{j_0-s}$, which exceeds $C_0\rho_L k_L(n_L)$ if $n_L < s$. The resolution must use that a crossing of B has diameter at least $L/8$, so that a tube of length $t < L/4$ carries only part of it and the rest is carried by other tubes, whose costs multiply; the resulting bound is a sum over partitions of the crossing into at most r tubes, each costing $(C\mu/t_i)^{j_0-s_i}$ with $\sum t_i \geq L/8$, and it is dominated by the single-tube terms. We have not written this out, and the dependence of r on $|G'|$ has to be removed by using that only the lines of G' within distance 4 of the tube matter. The constant C_0 obtained is $C(j_0)$ times r^{j_0}; the restriction to $L \leq L_1$ enters only through μ_1.

Remark 6.2 (The role of the base case). The base case is where the power $L^{-(j_0-2)}$ originates: a crossing of a box of bounded side with few lines forces j_0 lines of $\mathcal{L} \cup G'$ to run together along a stretch of length comparable to L, and this costs $(\mu/L)^{j_0-s}$. A bound of the form $C(L_1)\mu^{j_0}$ with an uncontrolled constant would not suffice, because the recursion raises constants to the power a and only the slack $\ell^{-\delta}$ of Lemma 5.12, a power of $L_1^{1/3}$, can absorb them; this is why C_0 must not depend on L_1.

7 Status, corrections and what a referee should check

7.1 The logical chain

Step	Status	Where
Model, boxes, chains; (T) ⇒ no percolation	proved	Theorem 3.1, Section 3
First moment over chains under independence (D_0)	proved	Proposition 4.4
Conditioning on clusters ⇒ fixed witness lines	proved	Lemma 4.6
(D_0) is false (relays)	shown (sketch)	Proposition 4.7
Dichotomy for witnesses (no reconvergence of a bundle)	proved	Lemma 4.8
No bound uniform in the witness	shown	Proposition 4.9
Claim D from weighted $\mathcal{H}_2^E$ and witness weight bound	proved	Proposition 4.10
Dependence step in relay form	**claim**	Claim D
Clusters give crossings	proved	Lemma 5.2
Chains of sub-crossings, annuli, separation	proved	Lemmas 5.6 and 5.7
Conditional independence given shared lines	proved	Lemma 5.9
Measures of shared and aligned lines	proved	Lemmas 2.1 and 5.10
Inheritance of avoidance	proved	Lemma 5.11
Arithmetic of a sub-boxes at geometric scales	proved	Lemma 5.12
Coverage lemma (x-fold cover of a segment)	proved	Lemma 5.13
Recursion, Case F (avoiding)	proved modulo Claim C1	Theorem 6.1
Shared-line weights, product over sub-boxes	**claim**	Claim C1
Recursion, Case R (rides)	**claim**	Claim C2
Base case	**claim**	Claim C3

7.2 Corrections to the working notes

Five statements in our working notes are corrected here, and a referee comparing the documents should be aware of them.

1. In [HJ26a], cluster chains allowed repeated clusters; with that definition hypothesis (T) is false (Remark 3.5). The present Definition 1.2 requires distinct clusters, and Lemma 3.2 supplies the path for which this is possible.

2. In [HJ26a], a box was thick if at least j_0 lines meet it. With that definition a nearly axis-parallel bundle fragments into many clusters for lattice reasons, and the chain weights make (T) false (Remark 1.1). Thickness is now defined through the continuum set Θ of (1.2).

3. In [HJ26a] and in the first version of this document, the step from (S) to (T) was expected to follow from a uniform bound on cluster sizes conditionally on other clusters. This bound, (D_0), is false because of relays (Proposition 4.7); Lemma 4.6 and Claim D give the corrected form.

4. In [HJ26b], separated sub-boxes were chosen as every $\lceil L/(4a\ell)\rceil$-th member of the chain of sub-crossings, which does not ensure spatial separation. The annuli of Section 5 replace this.

5. In [HJ26b], the induction hypothesis used a weight based on the maximal number of fixed lines near a single point of a fixed line, and the closure of this weight under shared lines was asserted through a second-moment lemma. The weight is not closed under shared lines (Remark 5.5(b)), and the second moment of the weight is not bounded; the exponent in that lemma's proof is wrong. The coverage weight of Definition 5.3 replaces the former, and Claim C1 states what replaces the latter.

7.3 What a referee should check

The proved parts that carry the most weight are Theorem 3.1, in particular Remark 1.1, Lemma 3.2 and the counting in (3.2); the iteration in the proof of Proposition 4.4, in particular the attachment of the factor $(\operatorname{diam} C_{k-1} + 1)$ to the previous cluster in Step 3; the witness lemma, Lemma 4.6, in particular the monotonicity of the indicator that w is the chosen witness, and the dichotomy Lemma 4.8; and the coverage lemma, Lemma 5.13, in particular the choice of weights in Step 3 and the use of the integer decomposition property in Step 1. Among the claims, Claim D and Claim C2 are the ones on which the programme stands or falls, and they are two faces of the same problem, the control of clusters that ride lines released by other clusters; Claims C1 and C3 are computations that we expect to go through as indicated. We do not claim the unconditional theorem, and the present document should be read as a precise statement of what has been proved and what has not.

Acknowledgements

BJ acknowledges the financial support of the Leibniz Association within the Leibniz Junior Research Group on Probabilistic Methods for Dynamic Communication Networks (DYCOMNET) as part of the Leibniz Competition (grant no. J105/2020), of the Berlin Mathematics Research Center MATH+ through the project EF-MA-Sys-2 on Information Flow & Emergent Behavior in Complex Networks, and of the Deutsche Forschungsgemeinschaft (DFG) through Project P27 within the SPP 2265.

Use of AI tools

The proofs, computations and the text of this document were developed with substantial assistance of the AI systems Claude (Anthropic) and ChatGPT Astra 6 (OpenAI), in an interactive process directed by the authors.

References

[BT78] S. Baum and L. E. Trotter, Jr., *Integer rounding and polyhedral decomposition for totally unimodular systems*, Optimization and Operations Research (Proc. Workshop, Bonn, 1977), Lecture Notes in Economics and Mathematical Systems, vol. 157, Springer, Berlin, 1978, pp. 15–23.

[HJ26a] C. Hirsch and B. Jahnel, *A first-moment route to the subcritical phase: the skeleton bound and the thick-cluster lemma*, Working note; file `subcritical-phase/first-moment/main.tex` in this repository, 2026.

[HJ26b] ———, *A multiscale scheme for the single-cluster bound*, Working note, November 2026; folder `subcritical-phase/multiscale` in this repository, 2026.

[LP17] G. Last and M. Penrose, *Lectures on the Poisson process*, Institute of Mathematical Statistics Textbooks, vol. 7, Cambridge University Press, Cambridge, 2017.