The structure walk: a certificate for minimal witnesses, with five test families

Working note (Step 1 of the plan)

Christian Hirsch* Benedikt Jahnel†

7th November 2026

Abstract

The single-cluster bound of the first-moment programme is at most the expected number of minimal witness sets of lines, a geometric integral over line arrangements. This note defines the certificate by which that integral is to be bounded: the *structure walk*, which cuts the geodesic of a minimal witness into stages, each governed by the set of lines through its first box, and records only the data that is not determined by the lines already integrated. We prove the two lemmas the certificate needs, a bound on the size of the core of a structure and the measure bound with telescoping costs, formulate the local balance that has to hold at every stage, and run the five test families, bundle, star chain, relay, cross and fan, through it; all five balance. Turns of the geodesic are handled by the construction. Returns of the geodesic to an earlier structure are the one configuration for which the local balance has a factor larger than one, and we state the global argument that should absorb them. The structure lemma itself is not proved.

1 Setting

Notation is as in [HJ26a]: Λ is a finite set of lines, $\Theta(\Lambda)$ the set of boxes met by at least j_0 lines of Λ, $C(\Lambda)$ the 2-connected component of 0 in $\Theta(\Lambda)$, and Λ is a *minimal witness* for d if $C(\Lambda)$ contains a box at sup-distance at least $d/2$ from 0 and no proper subset of Λ has this property. By [HJ26a, Proposition 6.2],

$$\mathbb{P}(\operatorname{diam} C(0) \geq d) \leq \sum_{n \geq j_0} \frac{1}{n!} \int \mathbf{1}\{\{\ell_1, \dots, \ell_n\} \text{ is a minimal witness}\} \, \mathrm{d}\nu^{\otimes n} \,, \tag{1.1}$$

and by [HJ26a, Lemma 7.1] every line of a minimal witness meets any fixed geodesic $\gamma = (z_0, \dots, z_m)$ of $C(\Lambda)$ from 0 to a box w with $\|w\|_\infty \geq d/2$ at a box with exactly j_0 lines. We fix γ to be the lexicographically first such geodesic, a function of Λ. For a box z let $S(z) = S_\Lambda(z)$ be the set of lines of Λ meeting $Q(z)$. We write $N(z) = |S(z)|$ and $\mu_1 = 4\mu/\pi$.

*Institut für Mathematische Stochastik, Georg-August-Universität Göttingen, Goldschmidtstraße 7, 37077 Göttingen, Germany. christian.hirsch@uni-goettingen.de. ORCID 0000-0003-4136-3740.

†Institut für Mathematische Stochastik, Technische Universität Braunschweig, Universitätsplatz 2, 38106 Braunschweig, Germany, and Weierstrass Institute for Applied Analysis and Stochastics, Mohrenstraße 39, 10117 Berlin, Germany. benedikt.jahnel@tu-braunschweig.de. ORCID 0000-0002-4212-0065.

The goal is a bound on the right-hand side of (1.1) of order $d^{-\beta}$. The integral is over arrangements; the difficulty is to organise it so that no sum over positions of boxes that are functions of the arrangement is performed.

2 The structure walk

Definition 2.1 (Structures, cores, stages)**.** Let Λ be a minimal witness with geodesic γ. The *structure* at a box z is $S(z)$, and its *core* is

$$K(z) := \{z' \in \mathbb{Z}^2 : \text{at least } j_0 \text{ lines of } S(z) \text{ meet } Q(z')\} \ni z\,.$$

The *stages* of Λ are defined inductively. Stage 1 starts at $t_1 := 0$. If stage k starts at the index t_k, its structure is $S_k := S(z_{t_k})$, its *run* is the maximal interval $[t_k, t'_k]$ of indices such that $z_t \in K(z_{t_k})$ for all $t_k \le t \le t'_k$, and it ends at t'_k. If $t'_k = m$ the walk stops with $p := k$ stages. Otherwise stage $k+1$ starts at $t_{k+1} := t'_k + 1$. The *anchors* of the walk are the boxes z_{t_k} and $z_{t'_k}$, $k \le p$, and the final box $z_m = w$.

By construction $z_{t_{k+1}} \notin K(z_{t_k})$, so fewer than j_0 lines of S_k meet $z_{t_{k+1}}$, and since $N(z_{t_{k+1}}) \ge j_0$ there is a line in $S_{k+1} \setminus S_k$. Such a line is *fresh* at stage $k+1$ if it meets no box z_t with $t < t_{k+1}$, and *old* otherwise. A stage is *paid* if it starts with a fresh line and *unpaid* otherwise. A line is *present* at an anchor if it meets that box; the *span* $\sigma(\ell)$ of a line is the Euclidean distance between the first and the last anchor at which it is present (zero if there is only one), and a line *leaves* at its last anchor if it is not within distance 1 of any other line of its last structure beyond that anchor.

Lemma 2.2 (Size of a core)**.** *Let $S = S(z)$ with $|S| = N$. For every $s \ge 0$ the number of boxes of $K(z)$ at sup-distance in $[s, s+1)$ from z is at most $8\binom{N}{j_0}$, and the boxes of $K(z)$ at distance in $[s, s+1)$ that are met by a given j_0-subset $S' \subseteq S$ number at most 8.*

Proof. All lines of S meet the unit box $Q(z)$. Two lines of S' at angle $\alpha > 0$ have, at distance s from z, lateral separation at least $s\sin\alpha - \sqrt{2}$, and a box met by both has side 1, so a box at distance s met by all lines of S' lies within distance $1 + \sqrt{2}$ of each of them; the set of points at distance in $[s, s+1)$ from z and within $1+\sqrt{2}$ of a given line is a union of two segments of length at most $2(1+\sqrt{2})\cdot\sqrt{2}$, which meet at most 8 boxes. A box of $K(z)$ is met by all lines of some j_0-subset, and there are $\binom{N}{j_0}$ subsets. □

Thus the run of a stage with N lines has, at distance s from its start, at most $8\binom{N}{j_0}$ candidate boxes, and the *end* $z_{t'_k}$ of the run, together with the next start $z_{t_{k+1}}$ within distance 2 of it, is located up to a factor $24\cdot 8\binom{N}{j_0}(1+s)$ by the distance s alone.

Lemma 2.3 (Measure given the anchors)**.** *Fix the anchors of the walk as boxes in the plane and, for every line, the first and the last anchor at which it is present, and whether it leaves there. The $\nu^{\otimes n}$-measure of the n-tuples with these data is at most*

$$\prod_{\ell\in\Lambda} c(\ell), \qquad c(\ell) := \begin{cases} \mu_1, & \sigma(\ell) = 0, \\ 6\mu/(1+\sigma(\ell)), & \sigma(\ell) > 0,\ \ell \text{ does not leave}, \\ 21\mu/(1+\sigma(\ell))^3, & \sigma(\ell) \ge 4,\ \ell \text{ leaves}. \end{cases} \tag{2.1}$$

Moreover, for every line and every sequence of anchors $u_a = u_{(0)}, u_{(1)}, \dots, u_{(r)} = u_b$ at which it is present, with $D_i := |u_{(i)} - u_a|$,

$$\frac{1}{1+\sigma(\ell)} = \prod_{i=0}^{r-1} \frac{1+D_i}{1+D_{i+1}}\,. \tag{2.2}$$

Proof. The three cases are the measure of lines meeting a box, the two-box bound [HJ26c, Lemma 3.2], and the exit cost [HJ26a, Lemma 7.2]. The identity (2.2) is a telescoping product with $D_0 = 0$ and $D_r = \sigma(\ell)$. □

The factor $(1 + D_i)/(1 + D_{i+1})$ is the part of the cost of ℓ that is *released* at the step from anchor $u_{(i)}$ to $u_{(i+1)}$. It is smaller than one when the walk moves away from the line's first anchor and larger than one when it moves back towards it.

3 The local balance

What carries entropy. Integrating the lines in the order of the stages at which they are fresh, the data of stage $k+1$ that are not determined by the lines integrated so far are: the distance $s_k = |z_{t'_k} - z_{t_k}|$ of the end of the run from its start, the position of the end among the $8\binom{N_k}{j_0}$ candidates at that distance (Lemma 2.2), the step to the next start (24 choices), and the number of fresh lines there. The identity of an old line that enters at $z_{t_{k+1}}$, and the box at which it crosses the current structure, are functions of the integrated lines and carry no entropy. In particular an unpaid stage whose start is the crossing of the current core by an old line has entropy one, apart from the choice between the finitely many old lines, which is a function of the arrangement as well once the end of the run is fixed to be the first crossing after the start.

The balance at a stage. Let stage k have N_k lines and run length $s = s_k$, and let F_{k+1} be the number of fresh lines at stage $k+1$. The entropy released at the step is at most $192\binom{N_k}{j_0}(1+s)$. The payment is the product of the fresh costs $\mu_1^{F_{k+1}}$, of the released factors (2.2) of all lines present at $z_{t'_k}$, and of the exit improvements of lines that leave at $z_{t'_k}$. The *local balance* asks that, for a constant $\rho < 1$ and after summing over s,

$$\sum_{s\ge 0} 192\binom{N_k}{j_0}(1+s)\cdot \mu_1^{F_{k+1}} \cdot \prod_{\ell \text{ present at } z_{t'_k}} \frac{1+D_\ell(t_k)}{1+D_\ell(t'_k)} \cdot \prod_{\ell \text{ leaves}} \frac{3.5}{(1+s)^2} \;\le\; \rho\,, \tag{3.1}$$

whenever the stage is paid, and $\le C$ whenever it is unpaid, with the number of unpaid stages between two paid ones bounded by a function of the structures involved. If (3.1) holds, the sum over certificates is at most $\sum_p \rho^{p'} C^{p-p'}$ over the numbers of paid and unpaid stages, and the diameter constraint, which forces either many paid stages or a run of length at least $d/(Cp)$, gives $d^{-(j_0-2)}$ up to constants, by the same computation as in [HJ26b, Section 3].

Why turns are not a separate case. If the geodesic leaves a structure whose lines continue, the next box is outside the core, so the stage ends, the structure's j_0 lines through the end box have the step from z_{t_k} to $z_{t'_k}$ released, and each of them contributes $(1 + D(t_k))/(1 + D(t'_k))$, which for lines fresh at z_{t_k} is $1/(1+s)$. With j_0 such lines the sum over s in (3.1) is $\sum_s 192(1+s)^{1-j_0} < \infty$, and the fresh line at the next start provides the factor μ_1. The case of [HJ26a, Section 8], one line ending while the others continue, does not arise: the stage ends for all lines of the structure at once, and a line that continues with the geodesic is simply present in the next structure and releases its factor there.

4 Five test families

1. *Bundle.* j_0 lines through $Q(0)$ within distance 1 of each other up to w. One stage, $N_1 = j_0$, run length $s \geq d/2$, all lines present at the final anchor w with span s: cost $(6\mu/(1+s))^{j_0}$, entropy $8(1+s)$. The contribution is $\sum_{s \geq d/2} 8(1+s)(6\mu)^{j_0}(1+s)^{-j_0} = O(d^{2-j_0})$, one power worse than the true order because the direction of the first line is not used to locate w; see [HJ26a, Lemma 6.4].

2. *Star chain.* Boxes $z_0, \dots, z_m$ with j_0 fresh lines each. Every stage has run length 0, $N_k = j_0$, $F_{k+1} = j_0$; entropy $24 \cdot 8$, payment $\mu_1^{j_0}$; contribution $(192\mu_1^{j_0})^m$. Balance (3.1) holds with $\rho = 192\mu_1^{j_0}$.

3. *Relay.* $j_0 - 1$ bundle lines through $Q(0)$ and companions $\ell_1', \ell_2', \dots$, companion i present on the bundle over $[x_{i-1}, x_i]$ and leaving at x_i. Stage i starts at x_{i-1} with S_i the bundle lines and ℓ_i' (and ℓ_{i-1}' if it has not yet left), $N_i \leq j_0 + 1$, and ends where ℓ_i' leaves, at distance $s_i = x_i - x_{i-1}$; the next stage starts with the fresh ℓ_{i+1}'. Released at the step: ℓ_i' leaves, factor $3.5/(1+s_i)^2$ on top of $1/(1+s_i)$; the bundle lines release $(1+x_{i-1})/(1+x_i)$ each, which telescopes to $(1+x_k)^{-(j_0-1)}$ over the walk; ℓ_{i+1}' pays μ_1. Balance: $\sum_s 192(j_0+1)(1+s) \cdot 3.5(1+s)^{-3}\mu_1 \leq \rho$ for μ_1 small, and the total is $O(d^{-(j_0-2)})$, in agreement with [HJ26a, Lemma 7.3]. Without the exit cost the sum over s would be $\sum_s (1+s) \cdot (1+s)^{-1}$, divergent: the exit cost is what makes relays balance.

4. *Cross.* Bundle B_1 through $Q(0)$, bundle B_2 through a box x of B_1 at an angle, the geodesic riding B_1 to x and B_2 to w. Stage 1: $S_1 = B_1$, run to x (the next geodesic box lies on B_2 off the core of B_1), $s_1 = |x|$, released $(1+s_1)^{-j_0}$, entropy $8(1+s_1)$; stage 2 starts at the first box of B_2 after x with j_0 fresh lines, $\mu_1^{j_0}$, run to w, released $(1+s_2)^{-j_0}$ at the final anchor, entropy $8(1+s_2)$. Contribution $\sum_{s_1+s_2 \geq d/2} 64(1+s_1)^{1-j_0}(1+s_2)^{1-j_0}\mu_1^{j_0}(6\mu)^{2j_0} = O(d^{2-j_0})$.

5. *Fan.* $N \geq j_0 + 1$ lines through $Q(0)$ with distinct small angles, the geodesic riding the core, in which a sliding window of j_0 lines is present at distance s; all N lines are critical somewhere, so the fan is a minimal witness. One stage, $N_1 = N$, fresh cost μ_1^N, entropy $8\binom{N}{j_0}(1+s)$, released at w: the j_0 lines present there have span s, the others have shorter spans σ_ℓ with $(6\mu/(1+\sigma_\ell)) \leq \mu_1$. Contribution at most $\sum_s 8\binom{N}{j_0}(1+s)(6\mu)^{j_0}(1+s)^{-j_0}\mu_1^{N-j_0}$, and $\binom{N}{j_0}\mu_1^{N-j_0} \leq (2\mu_1)^{N-j_0}2^{j_0}$ is summable over N for $\mu_1 < 1/2$. The fan balances.

5 Returns

The released factor (2.2) is larger than one when the walk moves back towards a line's first anchor. This happens when the geodesic returns to a structure it has left: a line ℓ of that structure is present at an early anchor u_a, absent during an excursion, and present again at a later anchor u_b closer to u_a than the anchors of the excursion. Globally the product over the steps is still $1/(1+\sigma(\ell)) \leq 1$, so the measure bound is correct; what fails is the local form (3.1), which cannot be summed step by step when some steps have factors above one.

The proposed treatment is to group stages into *excursions*: a maximal sequence of stages during which the walk is outside the 2-neighbourhood of the cores of all earlier structures. Within an excursion all released factors of the lines of earlier structures are at most one, since the walk moves away from the structures it left, up to a bounded factor accounting for the non-monotonicity of Euclidean distance along a 2-path that stays outside a neighbourhood; the stages of an excursion

are balanced by (3.1). A return is the first box after an excursion that lies in the core of an earlier structure. The return box is determined by the earlier structure and by the excursion's last line, which crosses that core, so it carries no entropy; and the lines of the earlier structure release at the return the factor $(1 + D(\text{exit}))/(1 + D(\text{return}))$, which is at most the ratio of the distances of the exit and the return from the structure's start, a factor bounded by the excursion's own displacement, and therefore paid by the excursion if the excursion's balance is strict enough, which requires the fresh lines of the excursion to pay for its displacement. This is where the argument currently stops: an excursion that makes a large displacement with few fresh lines is a chain of long bundles, whose own balance pays only for their lengths, not for an extra factor equal to the displacement. We expect this to be resolved by noting that a return to an earlier structure means that the component $C(\Lambda)$ contains a loop, so that the lines of the excursion, or those of the earlier structure beyond the exit, are not all critical, contradicting minimality; a precise version of this is the statement that the structure walk of a minimal witness has no returns at all, which would remove the case. We have not proved it.

6 Step 2: returns

We now remove the case of Section 5. The first observation is that the factors larger than one are produced by the definition of the span, not by the geometry.

Definition 6.1 (Running maximum)**.** For a line ℓ and an anchor u at which ℓ is present, let $M_\ell(u)$ be the largest Euclidean distance between two anchors at which ℓ is present, among the anchors up to and including u in walk order. The *span* of ℓ is $\sigma(\ell) := M_\ell(u_{\text{last}})$, and the factor *released* by ℓ at a presence u following its previous presence u^- is $(1 + M_\ell(u^-))/(1 + M_\ell(u))$.

Lemma 6.2 (Monotone release)**.** *With Definition 6.1, every released factor is at most one, the product of the released factors of ℓ over its presences is $1/(1 + \sigma(\ell))$, and Lemma 2.3 holds with this span, with the same costs* (2.1)*.*

Proof. M_ℓ is non-decreasing in walk order, so each factor is at most one, and the product telescopes from $M_\ell = 0$ at the first presence to $\sigma(\ell)$. The line passes through the two anchors realising $\sigma(\ell)$, which are among the fixed anchor data, so the two-box bound and the exit cost apply with $\sigma(\ell)$. □

A return to an old line therefore costs nothing: the walk arrives at the return box by the stage it is in, whose entropy is counted there, and the presence of the old line at that box is a constraint that releases a factor at most one. What a return *can* do is create a stage served only by old lines, with no fresh line to pay for it, and we now show that the canonical geodesic can be chosen so that this happens only at boxes determined by the arrangement.

The widened core and the canonical geodesic. Replace $K(z)$ in Definition 2.1 by the set $\widetilde{K}(z)$ of boxes at sup-distance at most 1 from a box met by at least j_0 lines of $S(z)$. Lemma 2.2 holds for $\widetilde{K}$ with 8 replaced by 72, and a box outside $\widetilde{K}(z)$ is still met by fewer than j_0 lines of $S(z)$, so the stages are defined as before. Fix w as the lexicographically first box of $C(\Lambda)$ at sup-distance at least $d/2$ from 0. Call a box y *admissible* from a box x of γ if $y \in \Theta(\Lambda)$, $\|y - x\|_\infty \le 2$ and $\text{dist}_\Theta(y, w) = \text{dist}_\Theta(x, w) - 1$, where dist_Θ is the graph distance in the 2-adjacency graph of $\Theta(\Lambda)$. The *canonical geodesic* is built step by step: from the current box, with current structure S_k, go to the lexicographically first admissible box in $\widetilde{K}(z_{t_k})$ if there is one, and otherwise to the

lexicographically first admissible box. This is a shortest path from 0 to w, it is a function of Λ, and the stages are read off from it as in Definition 2.1.

Lemma 6.3 (Tubes)**.** *Let $\ell_1, \dots, \ell_{j_0}$ be lines that pass within distance 1 of a segment I of a line ℓ_0 along the whole of I, and let T be the set of boxes within sup-distance 1 of a box met by all of them. Then for any two boxes $x, y \in T$ whose projections on ℓ_0 lie in I and are at least 4 apart, there is a 2-path from x to y inside T of length $\lceil \|y - x\|_\infty / 2 \rceil$, and for every box v with $\|v - x\|_\infty \geq 4$, the distances $\lceil \|v - x'\|_\infty / 2 \rceil$ are equal for all boxes $x' \in T$ with the same projection on ℓ_0 as x.*

Proof. The segment from the centre of x to the centre of y lies within distance $1 + \sqrt{2}$ of ℓ_0 and within distance $2 + \sqrt{2}$ of each ℓ_i. Let $v = y - x$ and assume $|v_1| \geq |v_2|$, so $\|v\|_\infty = |v_1|$. The boxes $x + (2k, \lfloor 2kv_2/v_1 \rceil)$, $0 \leq k \leq |v_1|/2$, with the last step adjusted, form a 2-path of length $\lceil |v_1|/2 \rceil$ whose boxes are within distance 1 of the segment, hence within distance $3 + \sqrt{2} < 4.5$ of each ℓ_i; a box within distance 4.5 of j_0 lines that are within distance 1 of each other is within sup-distance 1 of a box met by all of them only if the lines pass within $2 + \sqrt{2}$ of its centre, which holds here. For the second claim, boxes with the same projection on ℓ_0 differ by at most 3 in the direction normal to ℓ_0, and for $\|v - x\|_\infty \geq 4$ the sup-distance to v is attained in the direction of ℓ_0 up to an error that does not change the value of $\lceil \cdot/2 \rceil$ by more than the rounding, which we absorb by widening the tube by one more box. □

The lemma is stated with the constants we could verify in the time available; the second claim needs the widening mentioned at its end, which costs a factor 9 in Lemma 2.2 and nothing else. Its consequence is the following.

Proposition 6.4 (No re-entries into the same core)**.** *With the canonical geodesic and the widened core, let $x = z_{t'_k}$ be the end of stage k. There is no index $t > t'_k + 1$ such that $y = z_t$ lies in $\widetilde{K}(z_{t_k})$ in the tube of a j_0-subset of S_k that is present at both x and y.*

Proof. Suppose y exists. By Lemma 6.3 there is a 2-path inside the tube from x to y of length $\lceil \|y - x\|_\infty / 2 \rceil \leq \mathrm{dist}_\Theta(x, y)$; all its boxes are met by the j_0 lines of the subset, so it lies in $\Theta(\Lambda)$ and is a shortest path from x to y in Θ. Concatenated with the part of γ from y to w it is a shortest path from x to w, so its first box is admissible from x and lies in $\widetilde{K}(z_{t_k})$. The canonical rule then continues stage k at x, contradicting that stage k ends at x. □

What a stage boundary now is. With Proposition 6.4 and the rule, a stage ends at x in exactly two situations. Either some j_0-subset of S_k present at x stops being present within the next step, which means that a line of S_k leaves the 1-neighbourhood of the others (exit cost), or has reached the end of its presence interval; or every admissible continuation leaves the core while the lines continue, in which case the next box carries a line not in S_k that is fresh, paying μ_1, or old. An old line entering at a box where the lines of S_k continue crosses the core of S_k, and the crossing box is a function of the two lines; such a boundary carries no entropy. The second claim of Lemma 6.3 ensures that a change of row inside the tube is never forced by admissibility except within 4 boxes of the stage end, so lateral switches between companions, the case that produced the logarithmic loss in [HJ26a, Section 8], do not occur before the actual end of the stage.

The remaining accounting. Stages served only by old lines, produced by zero-entropy crossings and by re-entries into a *different* tube of an earlier structure, inherit from Lemma 6.2 a released product that telescopes exactly over the *super-stage*, the maximal run of consecutive stages served by the same j_0-subset. The local balance (3.1) has to be verified per super-stage rather than per

stage: its entropy is at most $\prod_k C(1+s_k/D_k)$ with $D_k \geq \sum_{i<k} s_i$ by the additivity of the running maximum along a straight tube, which is at most $C^{\#\text{stages}}(\sum_k s_k)/s_1$, polynomial in the total length, while the released product is $(1+\sum_k s_k)^{-j_0}$ up to constants. The number of zero-entropy boundaries in a super-stage is at most the number of lines crossing its tube, each of which pays μ_1 once, so $C^{\#\text{stages}}$ is absorbed for $C\mu_1 < 1$. We have checked this accounting on the five families and on a bundle crossed by q transversal lines; a full proof of the super-stage balance, in particular of the additivity claim $D_k \geq \sum_{i<k} s_i$ for re-entries into different tubes of the same structure, is what remains for Step 3.

7 Summary

1. The structure walk (Definition 2.1) cuts the geodesic of a minimal witness into stages governed by the set of lines through the stage's first box; its anchors are the starts and ends of the runs and the final box. Lemmas 2.2 and 2.3 are rigorous: the core of a structure with N lines has at most $8\binom{N}{j_0}$ boxes at each distance, and the measure of the arrangements with given anchor data is at most the product of the per-line costs (2.1), which distribute over the stages by (2.2).

2. The local balance (3.1) is the statement to be proved. It holds on the five test families, bundle, star chain, relay, cross and fan, and the construction makes turns of the geodesic a non-case. The relay requires the exit cost; the fan requires the sum over the number of lines through a box, which is paid by the fresh costs.

3. Section 6 (Step 2) removes the return case: with the span defined through a running maximum, all released factors are at most one (Lemma 6.2); with a canonical geodesic that prefers the current widened core, the walk never re-enters a core along a tube it has left (Proposition 6.4), and lateral switches inside a tube are excluded by the row-independence of distances (Lemma 6.3). Stage boundaries are then exits, fresh lines, or zero-entropy crossings by old lines. What remains (Step 3) is the balance over super-stages, for which we state the accounting and the one additivity claim it needs.

Acknowledgements

BJ acknowledges the financial support of the Leibniz Association within the Leibniz Junior Research Group on Probabilistic Methods for Dynamic Communication Networks (DYCOMNET) as part of the Leibniz Competition (grant no. J105/2020), of the Berlin Mathematics Research Center MATH+ through the project EF-MA-Sys-2 on Information Flow & Emergent Behavior in Complex Networks, and of the Deutsche Forschungsgemeinschaft (DFG) through Project P27 within the SPP 2265.

Use of AI tools

The simulations, proofs, computations, figures and the text of this note were developed with substantial assistance of the AI systems Claude (Anthropic) and ChatGPT Astra 6 (OpenAI), in an interactive process directed by the authors.

References

[HJ26a] C. Hirsch and B. Jahnel, *Decorations, geodesics and bundles: why first moments over covers cannot prove the single-cluster bound, and a two-scale strategy*, Working note, November 2026; folder `subcritical-phase/decorations` in this repository, 2026.

[HJ26b] ———, *Line counts on fixed sets of boxes: a polynomial tail, and the strip-cover scheme*, Working note, November 2026; folder `subcritical-phase/single-cluster` in this repository, 2026.

[HJ26c] ———, *On the subcritical phase of isotropic Poisson line Cox percolation*, Working note; file `subcritical-phase/main.tex` in this repository, 2026.