

## Section 13.8 Extrema of Functions of Two Variables

MATH211 Calculus III

Instructor: Ben Huang




DEPARTMENT OF  
COMPUTING, MATHEMATICS  
AND PHYSICS

# Relative Extrema

## Section 13.8


B.H.



# Relative Extrema

## Section 13.8

B.H.







# Critical points

## Section 13.8

B.H.

Where are the critical points?







# Critical points

## Section 13.8

B.H.


Where are the critical points?




# Second Derivative Test From Calculus I

## Section 13.8

B.H.



$f'(c) = 0$  and  $f''(c) > 0$   
 $f(c)$  is a local minimum




$f'(c) = 0$  and  $f''(c) < 0$   
 $f(c)$  is a local maximum


# Classification of Critical Points

## Section 13.8

B.H.



$f_{xx} < 0$  &  $f_{yy} < 0$   
at the critical point  
local maximum




$f_{xx} > 0$  &  $f_{yy} > 0$   
at the critical point  
local minimum

# Classification of Critical Points

## Section 13.8

B.H.




$f_{xx} > 0$  &  $f_{yy} < 0$  at the critical point  
saddle point



# Classification of Critical Points

## Section 13.8

B.H.



$f_{xx} > 0$  &  $f_{yy} > 0$  at the critical point  
saddle point

# Failure of the Second Partial Test

## Section 13.8

B.H.

When  $d(x, y) = f_{xx}f_{yy} - f_{xy}^2 = 0$  at the critical point in question, the Second Partial Test is **inconclusive**.




Figure 1:  $z = y^2$




Figure 2:  $z = -x^2$




Figure 3:  $z = \frac{1}{50}x^3$