

Model of the spread of a disease in a population of mobile agents

NIDDAM Benjamin

November 30, 2021

Contents

A	Introduction	2
B	Model presentation	2
C	Experiments, results	4
	C.1 Population density	4
	C.2 Insatisfaction rate	5
D	Conclusion	6

A Introduction

The modeling of spatial segregation proposed in the 1970s by Thomas C. Schelling made an impression because of the perverse effect she suggested: a strong segregation could be the collective result of individual decisions that are not aimed at such segregation. We would be dealing with an almost spontaneous phenomenon. The problem is that such a conclusion outright contradicts the entropy principle. An exam more attentive to the model makes it possible here to identify no less than four biases which condition the results. We will study here the impacts that population density and the dissatisfaction rate can have on segregation.

B Model presentation

Schelling's model is a model of spatial segregation. It takes the form of a matrix of $n * n$ boxes. Each cell is represented by an integer between -1, 0 and 1. Empty cells are represented by 0 while cells occupied by an individual are represented by a -1 or 1. We therefore obtain a stylized representation of two populations which share an urban area.

Initially, the agents are placed at random to simulate the mix of populations. We then add a rule of movement of agents: individuals will change boxes if in their immediate neighborhood, there is not at least one rate "n" agents of the same type as them. When an agent is not satisfied, we generate a random place among the empty cells and we place the agent in this box; which frees up its old place. This operation is carried out until all the agents are satisfied, or until the defined maximum time is reached. Under these conditions, we obtain groupings of populations which appeared without being intentionally provoked.




Figure 1: Graphic representations of the initial and segregated populations with Schelling's model

C Experiments, results

With this model, we were able to study the effects of population density and dissatisfaction rate on movements agents.

C.1 Population density

As said, we have run several simulations with different population density values. In other words, we wanted to study whether the number of free sites played a role in the separation of populations.

We varied the number of free cells from 15 to 24 and obtained the following results:




Figure 2: Graphic representations of the initial and segregated states of Schelling's model with 24 free cells over 400 at an dissatisfaction rate of 0.5





Figure 3: Graphic representations of the initial and segregated states of Schelling's model with 24 free cells over 400 at an insatisfaction rate of 0.5

On observe que dans les deux cas, les agents se sont séparés en deux groupes distincts en fonction de leur couleur avant la fin du temps imparti. Nous avons donc représenté le temps mit par nos agent pour arriver à une situation stable en fonction de la densité de population.



(a) Population segregation after simulation

Figure 4: Graphic representations of the initial and segregated states of Schelling's model with 24 free cells over 400 at an insatisfaction rate of 0.5

C.2 Insatisfaction rate

D Conclusion