

# Modelling the Human Immunodeficiency Virus

## Lecture 4

**Ago Merico**

**28 January 2020**

# More reasons for modelling

Another motivation for learning mathematical biology is that equations typically "say" more than a prose; being able to "read" equations is very important both to verify the logical construct of a system and to evaluate the limitations of the assumptions;

# More reasons for modelling

Another motivation for learning mathematical biology is that equations typically "say" more than a prose; being able to "read" equations is very important both to verify the logical construct of a system and to evaluate the limitations of the assumptions;

In this lecture we will focus on the biological insights obtained from a mathematical model developed to investigate the ecology and epidemiology of the Human Immunodeficiency Virus (HIV);

# More reasons for modelling

Another motivation for learning mathematical biology is that equations typically "say" more than a prose; being able to "read" equations is very important both to verify the logical construct of a system and to evaluate the limitations of the assumptions;

In this lecture we will focus on the biological insights obtained from a mathematical model developed to investigate the ecology and epidemiology of the Human Immunodeficiency Virus (HIV);

Mathematical models (1) have allowed biologists to understand otherwise hidden aspects of HIV, (2) have produced testable predictions about how HIV replicates and spreads, and (3) have generated forecasts that improve the efficacy of prevention and health care programmes.

# Acquired Immunodeficiency Syndrome (AIDS)

On June 5, 1981, the Centers for Disease Control in Los Angeles reported the deaths of five males, all of whom had died from pneumocystis, a form of pneumonia that rarely causes death in individuals with healthy immune systems;

# Acquired Immunodeficiency Syndrome (AIDS)

On June 5, 1981, the Centers for Disease Control in Los Angeles reported the deaths of five males, all of whom had died from pneumocystis, a form of pneumonia that rarely causes death in individuals with healthy immune systems;

Since this first report, Acquired Immunodeficiency Syndrome (AIDS), as the disease has come to be known, has reached epidemic proportions, having caused more than 20 million deaths worldwide;

# Acquired Immunodeficiency Syndrome (AIDS)


On June 5, 1981, the Centers for Disease Control in Los Angeles reported the deaths of five males, all of whom had died from pneumocystis, a form of pneumonia that rarely causes death in individuals with healthy immune systems;

Since this first report, Acquired Immunodeficiency Syndrome (AIDS), as the disease has come to be known, has reached epidemic proportions, having caused more than 20 million deaths worldwide;

AIDS results from the deterioration of the immune system, which then fails to ward off various cancers (e.g. Kaposi's sarcoma) and infectious agents (e.g. the protozoa that cause pneumocystis, the viruses that cause retinitis, and the bacteria that cause tuberculosis).

# Human Immunodeficiency Virus (HIV)

The collapse of the immune system is caused by infection with HIV, which is transmitted from infected to susceptible individuals by the exchange of bodily fluids, primarily through sexual intercourse without condoms, sharing of unsterilised needles, or transfusion with infected blood supplies;




Scanning electron micrograph of HIV (in green), grown in cultured lymphocytes. A lymphocyte is one of the subtypes of white blood cell in a vertebrate's immune system.



# Human Immunodeficiency Virus (HIV)

The collapse of the immune system is caused by infection with HIV, which is transmitted from infected to susceptible individuals by the exchange of bodily fluids, primarily through sexual intercourse without condoms, sharing of unsterilised needles, or transfusion with infected blood supplies;

Once inside the body, HIV particles infect white blood cells ( $CD4^+$  T cells, or CD4 cells);




Scanning electron micrograph of HIV (in green), grown in cultured lymphocytes. A lymphocyte is one of the subtypes of white blood cell in a vertebrate's immune system.

# Human Immunodeficiency Virus (HIV)

The collapse of the immune system is caused by infection with HIV, which is transmitted from infected to susceptible individuals by the exchange of bodily fluids, primarily through sexual intercourse without condoms, sharing of unsterilised needles, or transfusion with infected blood supplies;

Once inside the body, HIV particles infect white blood cells ( $CD4^+$  T cells, or CD4 cells);

The genome of the virus, made of RNA, enters these cells and is reverse transcribed (hence also called retrovirus) into DNA, which is subsequently incorporated into the genome of the host.



Scanning electron micrograph of HIV (in green), grown in cultured lymphocytes. A lymphocyte is one of the subtypes of white blood cell in a vertebrate's immune system.

# Human Immunodeficiency Virus (HIV)

The virus may remain latent within the genome of the host cell or become activated;




Diagram of HIV.

# Human Immunodeficiency Virus (HIV)

The virus may remain latent within the genome of the host cell or become activated;

When actively replicating, HIV can produce hundreds of daughter viruses per day per host cell, often killing the host cell in the process;




Diagram of HIV.

# Human Immunodeficiency Virus (HIV)

The virus may remain latent within the genome of the host cell or become activated;

When actively replicating, HIV can produce hundreds of daughter viruses per day per host cell, often killing the host cell in the process;

These virus particles (or virions) then go on to infect other cells.





Diagram of HIV.

# Human Immunodeficiency Virus (HIV)

Eventually, without treatment, the population of the infected cells declines dramatically from about 1000 cells  $\text{mm}^{-3}$  of blood to about 200 cells  $\text{mm}^{-3}$ , signaling the onset of AIDS.



The time evolution of HIV infection within an individual; viral loads and infected CD4 cells are plotted over time since infection.

# Human Immunodeficiency Virus (HIV)

What makes HIV particularly harmful to the immune system is that the virus preferentially attacks and destroys cells that recognize and fight other infections.

# Human Immunodeficiency Virus (HIV)

What makes HIV particularly harmful to the immune system is that the virus preferentially attacks and destroys cells that recognize and fight other infections.

Early on in the epidemic, the median period between infection with HIV and the onset of AIDS was about ten years; the median survival time following the onset of AIDS was just under one year.



# Human Immunodeficiency Virus (HIV)

What makes HIV particularly harmful to the immune system is that the virus preferentially attacks and destroys cells that recognize and fight other infections.

Early on in the epidemic, the median period between infection with HIV and the onset of AIDS was about ten years; the median survival time following the onset of AIDS was just under one year.

Survival statistics have improved dramatically with the development of effective antiretroviral therapies (ART).

# Human Immunodeficiency Virus (HIV)

What makes HIV particularly harmful to the immune system is that the virus preferentially attacks and destroys cells that recognize and fight other infections.

Early on in the epidemic, the median period between infection with HIV and the onset of AIDS was about ten years; the median survival time following the onset of AIDS was just under one year.

Survival statistics have improved dramatically with the development of effective antiretroviral therapies (ART).

The median survival after diagnosis with an AIDS rose from 17 months between 1990 and 1994 to 59 months between 1995 and 1998.

# Human Immunodeficiency Virus (HIV)

What makes HIV particularly harmful to the immune system is that the virus preferentially attacks and destroys cells that recognize and fight other infections.

Early on in the epidemic, the median period between infection with HIV and the onset of AIDS was about ten years; the median survival time following the onset of AIDS was just under one year.

Survival statistics have improved dramatically with the development of effective antiretroviral therapies (ART).

The median survival after diagnosis with an AIDS rose from 17 months between 1990 and 1994 to 59 months between 1995 and 1998.

Unfortunately, modern drug therapies are extremely expensive and cannot be afforded by the majority of individuals infected with HIV worldwide.

# Human Immunodeficiency Virus (HIV)

Until effective therapy or vaccines become freely available, HIV will continue to take a devastating toll...



Number of individuals living with HIV (source: Joint United Nations Programme on HIV/AIDS, 2004).

Mathematical modeling has been a very important tool in HIV/AIDS research;

# Models of HIV / AIDS

Mathematical modeling has been a very important tool in HIV/AIDS research;

Every aspect of the natural history, treatment, and prevention of HIV has been the subject of mathematical models, from the thermodynamic characteristics of HIV to its replication rate both within and among individuals;

# Models of HIV / AIDS


Mathematical modeling has been a very important tool in HIV/AIDS research;

Every aspect of the natural history, treatment, and prevention of HIV has been the subject of mathematical models, from the thermodynamic characteristics of HIV to its replication rate both within and among individuals;

In the following, [we will explore one](#) of these [models](#) in detail. This model was chosen because of its implication for our understanding of HIV.

# Model – Dynamics of HIV after initial infection

After an individual is infected by HIV, the number of virions within the bloodstream skyrockets and then drops down again; this period is known as the acute phase, lasts for about 100 days, and leads to flu-like symptoms;




The time evolution of HIV infection within an individual; viral loads and infected CD4 cells are plotted over time since infection.



# Model – Dynamics of HIV after initial infection

After an individual is infected by HIV, the number of virions within the bloodstream skyrockets and then drops down again; this period is known as the acute phase, lasts for about 100 days, and leads to flu-like symptoms;

The rapid rise in virions reflects the infection of CD4 cells and the replication of HIV within the infected host cells;




The time evolution of HIV infection within an individual; viral loads and infected CD4 cells are plotted over time since infection.

# Model – Dynamics of HIV after initial infection

After an individual is infected by HIV, the number of virions within the bloodstream skyrockets and then drops down again; this period is known as the acute phase, lasts for about 100 days, and leads to flu-like symptoms;

The rapid rise in virions reflects the infection of CD4 cells and the replication of HIV within the infected host cells;

But what cause the decline in virions?



The time evolution of HIV infection within an individual; viral loads and infected CD4 cells are plotted over time since infection.


# Model – Dynamics of HIV after initial infection

After an individual is infected by HIV, the number of virions within the bloodstream skyrockets and then drops down again; this period is known as the acute phase, lasts for about 100 days, and leads to flu-like symptoms;

The rapid rise in virions reflects the infection of CD4 cells and the replication of HIV within the infected host cells;

**But what cause the decline in virions?**

The most obvious answer is that the immune system acts to recognise and suppress the viral infection...



The time evolution of HIV infection within an individual; viral loads and infected CD4 cells are plotted over time since infection.

# Model – Dynamics of HIV after initial infection

Phillips (1996), however, suggested an alternative explanation:

# Model – Dynamics of HIV after initial infection

Phillips (1996), however, suggested an alternative explanation:

The number of virions might decline because most of the susceptible CD4 cells have already been infected and thus there are fewer host cells to infect.

# Model – Dynamics of HIV after initial infection

Phillips (1996), however, suggested an alternative explanation:

The number of virions might decline because most of the susceptible CD4 cells have already been infected and thus there are fewer host cells to infect.

Phillips developed a model to assess whether this alternative explanation could mimic the observed rise and fall of virions in the blood stream over the right time frame. The model contains 4 *variables*:

# Model – Dynamics of HIV after initial infection

Phillips (1996), however, suggested an alternative explanation:

The number of virions might decline because most of the susceptible CD4 cells have already been infected and thus there are fewer host cells to infect.

Phillips developed a model to assess whether this alternative explanation could mimic the observed rise and fall of virions in the blood stream over the right time frame. The model contains 4 *variables*:

- ▶  $R$ : number of activated but uninfected CD4 cells;

# Model – Dynamics of HIV after initial infection

Phillips (1996), however, suggested an alternative explanation:

The number of virions might decline because most of the susceptible CD4 cells have already been infected and thus there are fewer host cells to infect.

Phillips developed a model to assess whether this alternative explanation could mimic the observed rise and fall of virions in the blood stream over the right time frame. The model contains 4 *variables*:

- ▶  $R$ : number of activated but uninfected CD4 cells;
- ▶  $L$ : number of latently infected CD4 cells;



# Model – Dynamics of HIV after initial infection

Phillips (1996), however, suggested an alternative explanation:

The number of virions might decline because most of the susceptible CD4 cells have already been infected and thus there are fewer host cells to infect.

Phillips developed a model to assess whether this alternative explanation could mimic the observed rise and fall of virions in the blood stream over the right time frame. The model contains 4 *variables*:

- ▶  $R$ : number of activated but uninfected CD4 cells;
- ▶  $L$ : number of latently infected CD4 cells;
- ▶  $E$ : number of actively infected CD4 cells;

# Model – Dynamics of HIV after initial infection

Phillips (1996), however, suggested an alternative explanation:

The number of virions might decline because most of the susceptible CD4 cells have already been infected and thus there are fewer host cells to infect.

Phillips developed a model to assess whether this alternative explanation could mimic the observed rise and fall of virions in the blood stream over the right time frame. The model contains 4 *variables*:

- ▶  $R$ : number of activated but uninfected CD4 cells;
- ▶  $L$ : number of latently infected CD4 cells;
- ▶  $E$ : number of actively infected CD4 cells;
- ▶  $V$ : number of virions in the blood stream.

# Model – Dynamics of HIV after initial infection

Phillips (1996), however, suggested an alternative explanation:

The number of virions might decline because most of the susceptible CD4 cells have already been infected and thus there are fewer host cells to infect.

Phillips developed a model to assess whether this alternative explanation could mimic the observed rise and fall of virions in the blood stream over the right time frame. The model contains 4 *variables*:

- ▶  $R$ : number of activated but uninfected CD4 cells;
- ▶  $L$ : number of latently infected CD4 cells;
- ▶  $E$ : number of actively infected CD4 cells;
- ▶  $V$ : number of virions in the blood stream.

The *dynamics* of each variable depends on the values of the remaining variables.

# Model – Dynamics of HIV after initial infection

Phillips' model contains several *parameters*, such as the death rate of actively infected cells ( $\delta$ ) and the death rate of viruses ( $\sigma$ );

# Model – Dynamics of HIV after initial infection

Phillips' model contains several *parameters*, such as the death rate of actively infected cells ( $\delta$ ) and the death rate of viruses ( $\sigma$ );

Phillips built into his model the crucial assumption that the body does not get better at eliminating infected cells or virions over time, under the null hypothesis that the immune system does not mount a defense against HIV during the acute phase;

# Model – Dynamics of HIV after initial infection

Phillips' model contains several *parameters*, such as the death rate of actively infected cells ( $\delta$ ) and the death rate of viruses ( $\sigma$ );

Phillips built into his model the crucial assumption that the body does not get better at eliminating infected cells or virions over time, under the null hypothesis that the immune system does not mount a defense against HIV during the acute phase;

To model the progression of HIV within the body, Phillips then needed values for each of the parameters in the model; unfortunately, few data existed at the time for many of them;

# Model – Dynamics of HIV after initial infection

Phillips' model contains several *parameters*, such as the death rate of actively infected cells ( $\delta$ ) and the death rate of viruses ( $\sigma$ );


Phillips built into his model the crucial assumption that the body does not get better at eliminating infected cells or virions over time, under the null hypothesis that the immune system does not mount a defense against HIV during the acute phase;

To model the progression of HIV within the body, Phillips then needed values for each of the parameters in the model; unfortunately, few data existed at the time for many of them;

To proceed, Phillips chose plausible values for each parameter and numerically ran the model;

# Model – Dynamics of HIV after initial infection

Phillip observed that the number of virus particles typically rose and then fell by several orders of magnitude over a period of a few days to weeks:




Number of virions in the blood stream ( $V$ ), based on Phillips' model, as a function of the number of days since primary infection.




# Model – Dynamics of HIV after initial infection

Phillip observed that the number of virus particles typically rose and then fell by several orders of magnitude over a period of a few days to weeks:




Number of virions in the blood stream ( $V$ ), based on Phillips' model, as a function of the number of days since primary infection.




The time evolution of HIV infection within an individual; viral loads and infected CD4 cells are plotted over time since infection.

# Model – Dynamics of HIV after initial infection

Phillip observed that the number of virus particles typically rose and then fell by several orders of magnitude over a period of a few days to weeks:



Number of virions in the blood stream ( $V$ ), based on Phillips' model, as a function of the number of days since primary infection.



The time evolution of HIV infection within an individual; viral loads and infected CD4 cells are plotted over time since infection.

Phillips thus came to the counter-intuitive conclusion that *"the reduction in virions during acute infection may not reflect the ability of the HIV-specific immune response to control the virus replication"*.

# Model – Dynamics of HIV after initial infection

Phillips' model accomplished a number of important things:

# Model – Dynamics of HIV after initial infection

Phillips' model accomplished a number of important things:

1. It changed our view of what was possible: without such a model, it would seem unlikely that a dramatic viral peak and decline could be caused by the dynamics of CD4 cells;

# Model – Dynamics of HIV after initial infection

Phillips' model accomplished a number of important things:

1. It changed our view of what was possible: without such a model, it would seem unlikely that a dramatic viral peak and decline could be caused by the dynamics of CD4 cells;
2. It produced testable predictions: one being that the viral peak and decline should be observed even in individuals that do not mount an immune response (indeed, this prediction has been confirmed in several patients);

# Model – Dynamics of HIV after initial infection

Phillips' model accomplished a number of important things:

1. It changed our view of what was possible: without such a model, it would seem unlikely that a dramatic viral peak and decline could be caused by the dynamics of CD4 cells;
2. It produced testable predictions: one being that the viral peak and decline should be observed even in individuals that do not mount an immune response (indeed, this prediction has been confirmed in several patients);
3. Phillips' model generated a useful null hypothesis: viral dynamics do not reflect an immune response; this null hypothesis might be wrong, but at least it can be tested.


# Phillips' model

We illustrate Phillips' model in the form of a flow diagram.

# Phillips' model


We illustrate Phillips' model in the form of a flow diagram.

The circles represent (1) the number of susceptible CD4 cells,  $R(t)$ , (2) the number of latently infected CD4 cells,  $L(t)$ , (3) the number of actively infected CD4 cells,  $E(t)$ , and (4) the number of virions in the blood stream,  $V(t)$ .






# Phillips' HIV model – flow diagram



# Phillips' HIV model – flow diagram


The immune systems produces new CD4 cells at a rate  $\Gamma$ , of which a fraction  $\tau$  becomes susceptible of being attacked by HIV;



# Phillips' HIV model – flow diagram

The immune systems produces new CD4 cells at a rate  $\Gamma$ , of which a fraction  $\tau$  becomes susceptible of being attacked by HIV;

Even without HIV infection, CD4 cells die at a rate  $\mu$  per cell per day, leading to a total flow out of the circle of  $\mu \cdot R(t)$  per day;




# Phillips' HIV model – flow diagram

The immune system produces new CD4 cells at a rate  $\Gamma$ , of which a fraction  $\tau$  becomes susceptible of being attacked by HIV;

Even without HIV infection, CD4 cells die at a rate  $\mu$  per cell per day, leading to a total flow out of the circle of  $\mu \cdot R(t)$  per day;

CD4 cells become infected if they encounter a virus; new infections are assumed to occur at a rate  $\beta \cdot V(t)$  per cell per day, leading to a total flow of  $\beta \cdot V(t) \cdot R(t)$  per day;




# Phillips' HIV model – flow diagram

The immune systems produces new CD4 cells at a rate  $\Gamma$ , of which a fraction  $\tau$  becomes susceptible of being attacked by HIV;

Even without HIV infection, CD4 cells die at a rate  $\mu$  per cell per day, leading to a total flow out of the circle of  $\mu \cdot R(t)$  per day;

CD4 cells become infected if they encounter a virus; new infections are assumed to occur at a rate  $\beta \cdot V(t)$  per cell per day, leading to a total flow of  $\beta \cdot V(t) \cdot R(t)$  per day;

Once infected, CD4 cells may host HIV in a latent (non-replicating) state (a proportion  $p$ ) or in its actively replicating state ( $1 - p$ );



# Phillips' HIV model – flow diagram


The immune systems produces new CD4 cells at a rate  $\Gamma$ , of which a fraction  $\tau$  becomes susceptible of being attacked by HIV;

Even without HIV infection, CD4 cells die at a rate  $\mu$  per cell per day, leading to a total flow out of the circle of  $\mu \cdot R(t)$  per day;

CD4 cells become infected if they encounter a virus; new infections are assumed to occur at a rate  $\beta \cdot V(t)$  per cell per day, leading to a total flow of  $\beta \cdot V(t) \cdot R(t)$  per day;

Once infected, CD4 cells may host HIV in a latent (non-replicating) state (a proportion  $p$ ) or in its actively replicating state ( $1 - p$ );

Since HIV is hidden in the genome of latently infected CD4 cells, these cells die at the same rate as uninfected cells ( $\mu$  per cell per day);



# Phillips' HIV model – flow diagram

The immune systems produces new CD4 cells at a rate  $\Gamma$ , of which a fraction  $\tau$  becomes susceptible of being attacked by HIV;


Even without HIV infection, CD4 cells die at a rate  $\mu$  per cell per day, leading to a total flow out of the circle of  $\mu \cdot R(t)$  per day;

CD4 cells become infected if they encounter a virus; new infections are assumed to occur at a rate  $\beta \cdot V(t)$  per cell per day, leading to a total flow of  $\beta \cdot V(t) \cdot R(t)$  per day;

Once infected, CD4 cells may host HIV in a latent (non-replicating) state (a proportion  $p$ ) or in its actively replicating state ( $1 - p$ );

Since HIV is hidden in the genome of latently infected CD4 cells, these cells die at the same rate as uninfected cells ( $\mu$  per cell per day);

Latently infected cells may be activated at a rate  $\alpha$  per cell per day; actively infected cells are thus produced by two means: by the infection of susceptible cells at a rate  $(1 - p) \cdot \beta \cdot V(t) \cdot R(t)$  or by the conversion of latently infected cells at a rate  $\alpha \cdot L(t)$ ;



# Phillips' HIV model – flow diagram

The immune systems produces new CD4 cells at a rate  $\Gamma$ , of which a fraction  $\tau$  becomes susceptible of being attacked by HIV;

Even without HIV infection, CD4 cells die at a rate  $\mu$  per cell per day, leading to a total flow out of the circle of  $\mu \cdot R(t)$  per day;


CD4 cells become infected if they encounter a virus; new infections are assumed to occur at a rate  $\beta \cdot V(t)$  per cell per day, leading to a total flow of  $\beta \cdot V(t) \cdot R(t)$  per day;

Once infected, CD4 cells may host HIV in a latent (non-replicating) state (a proportion  $p$ ) or in its actively replicating state ( $1 - p$ );

Since HIV is hidden in the genome of latently infected CD4 cells, these cells die at the same rate as uninfected cells ( $\mu$  per cell per day);

Latently infected cells may be activated at a rate  $\alpha$  per cell per day; actively infected cells are thus produced by two means: by the infection of susceptible cells at a rate  $(1 - p) \cdot \beta \cdot V(t) \cdot R(t)$  or by the conversion of latently infected cells at a rate  $\alpha \cdot L(t)$ ;

Actively infected cells die at a fast rate  $\delta$  per cell per day, due to the continual budding of virus particles at a rate  $\pi$  per infected cell per day; a dashed arrow between actively infected cells and viruses indicates that viral production by budding,  $\pi \cdot E(t)$ , does not directly eliminate an infected cell;





# Phillips' HIV model – flow diagram

The immune systems produces new CD4 cells at a rate  $\Gamma$ , of which a fraction  $\tau$  becomes susceptible of being attacked by HIV;

Even without HIV infection, CD4 cells die at a rate  $\mu$  per cell per day, leading to a total flow out of the circle of  $\mu \cdot R(t)$  per day;

CD4 cells become infected if they encounter a virus; new infections are assumed to occur at a rate  $\beta \cdot V(t)$  per cell per day, leading to a total flow of  $\beta \cdot V(t) \cdot R(t)$  per day;


Once infected, CD4 cells may host HIV in a latent (non-replicating) state (a proportion  $p$ ) or in its actively replicating state ( $1 - p$ );

Since HIV is hidden in the genome of latently infected CD4 cells, these cells die at the same rate as uninfected cells ( $\mu$  per cell per day);


Latently infected cells may be activated at a rate  $\alpha$  per cell per day; actively infected cells are thus produced by two means: by the infection of susceptible cells at a rate  $(1 - p) \cdot \beta \cdot V(t) \cdot R(t)$  or by the conversion of latently infected cells at a rate  $\alpha \cdot L(t)$ ;

Actively infected cells die at a fast rate  $\delta$  per cell per day, due to the continual budding of virus particles at a rate  $\pi$  per infected cell per day; a dashed arrow between actively infected cells and viruses indicates that viral production by budding,  $\pi \cdot E(t)$ , does not directly eliminate an infected cell;

Finally, virus particles degrade or are eliminated from the body at a rate  $\sigma$  per virion per day.



# Phillips' HIV model – flow diagram



Flow diagram of Phillips' HIV model, which describes the number of viruses in the blood stream after HIV infection.

# Phillips' HIV model – equations

# Phillips' HIV model – equations

From the flow diagram we can write down the differential equations describing the rate of change of each variable over time (e.g.  $dV(t)/dt$  for the rate of change of virus particles). Each variable represented by a circle in the flow diagram changes at a rate equal to the sum of all of the arrows entering the circle minus all of the arrows exiting the circle:

# Phillips' HIV model – equations

From the flow diagram we can write down the differential equations describing the rate of change of each variable over time (e.g.  $dV(t)/dt$  for the rate of change of virus particles). Each variable represented by a circle in the flow diagram changes at a rate equal to the sum of all of the arrows entering the circle minus all of the arrows exiting the circle:

$$\frac{dR(t)}{dt} = \Gamma \tau - \mu R(t) - \beta V(t) R(t),$$

$$\frac{dL(t)}{dt} = p \beta V(t) R(t) - \mu L(t) - \alpha L(t),$$

$$\frac{dE(t)}{dt} = (1 - p) \beta V(t) R(t) + \alpha L(t) - \delta E(t),$$

$$\frac{dV(t)}{dt} = \pi E(t) - \sigma V(t).$$

# Phillips' HIV model – equations

From the flow diagram we can write down the differential equations describing the rate of change of each variable over time (e.g.  $dV(t)/dt$  for the rate of change of virus particles). Each variable represented by a circle in the flow diagram changes at a rate equal to the sum of all of the arrows entering the circle minus all of the arrows exiting the circle:

$$\frac{dR(t)}{dt} = \Gamma \tau - \mu R(t) - \beta V(t) R(t),$$

$$\frac{dL(t)}{dt} = p \beta V(t) R(t) - \mu L(t) - \alpha L(t),$$

$$\frac{dE(t)}{dt} = (1 - p) \beta V(t) R(t) + \alpha L(t) - \delta E(t),$$

$$\frac{dV(t)}{dt} = \pi E(t) - \sigma V(t).$$

Technically, the rate at which virus particles infect susceptible cells,  $\beta V(t) R(t)$ , should also be subtracted off from  $dV(t)/dt$ , but this rate is assumed small relative to the large number of virus particles in the bloodstream.


# Phillips' HIV model – parameter values

Symbol	Description	Value (per day)	Comments
$\Gamma$	Rate of CD4 cell production	1.36	Equivalent to $3.4 \times 10^8$ cells per day in whole body
$\mu$	HIV-independent death rate of susceptible CD4 cells	$1.36 \times 10^{-3}$	Average life span of uninfected CD4 cells: 2 years
$\tau$	Fraction of cells becoming susceptible of HIV attack	0.2	
$\beta$	Rate of infection of cells per virion	0.00027	Cannot be estimated directly
$\rho$	Proportion of cells becoming latently infected upon infection	0.1	
$\alpha$	Activation rate of latently infected cells	$3.6 \times 10^{-2}$	
$\sigma$	removal rate of cell-free virus	2.0	Average life span: 1/2 day
$\delta$	Removal (death) rate of actively infected cells	0.33	Average life span: 3 days
$\pi$	Reate of production of virions by an actively infected cell	100.0	

- ▶ The total number of CD4 cells is:  $1000 \times (1 - \tau) + R(t) + L(t) + E(t)$
- ▶ Initial conditions are:  $R(t = 0) = 200$ ,  $L(t = 0) = 0$ ,  $E(t = 0) = 0$ ,  $V(t = 0) = 4.0 \times 10^{-7}$

# Phillips' HIV model – results

Check the original article by Phillips (1996) [available via CampusNet]:



**Fig. 1.** Temporal changes in the number of (A) total CD4 lymphocytes [ $1000(1 - \tau) + R + L + E$ ] and free virions ( $V$ ) and (B) activated, uninfected CD4 lymphocytes ( $R$ ), latently infected cells ( $L$ ), and actively infected cells ( $E$ ) in the first 120 days after HIV infection as predicted by the model on the basis of a simulation with iterations of step length 1 hour and 20 min. The total (whole-body) inoculum of HIV was assumed to consist of 100 virions (and no infected cells); thus, there were initially  $4 \times 10^{-7}$  virions in the quantity being considered.



# Plotting suggestions

## plot using a single y-axis

```
import numpy as np
import matplotlib.pyplot as plt

t=np.linspace(0,15,100)
y=np.sin(t)
z=np.cos(t)

plt.plot(t,y, '-r', linewidth=3, label='sin(x)')
plt.plot(t,z, '-b', linewidth=3, label='cos(x)')
plt.xlim(0,15)
plt.ylim(-2,2)
plt.xlabel('x', fontsize=20)
plt.ylabel('Trigonometric function', fontsize=20)
plt.legend(loc='best')

plt.show()
```

# Plotting suggestions

## plot using both y-axes (with the `twinx` method)

```
import numpy as np
import matplotlib.pyplot as plt

t=np.linspace(0,15,100)

y=np.sin(t)
z=np.cos(t)


ax1 = plt.subplot(1,1,1) # create a first y-axis object
ax1.plot(t, y, '-r', linewidth=3)
ax1.set_xlim(0,15)
ax1.set_ylim(-2,2)
ax1.set_xlabel('x', fontsize=20)
ax1.set_ylabel('sin(x)', color='red', fontsize=20)
ax1.tick_params(axis='y', colors='red')

ax2 = ax1.twinx() # create a twin y-axis object sharing the x-axis
ax2.plot(t, z, '-b', linewidth=3)
ax2.set_ylabel('cos(x)', color='blue', fontsize=20)
ax2.tick_params(axis='y', colors='blue')
ax2.set_ylim(-4,4)
ax2.set_xlim(0,15)


plt.show()
```

# Plotting suggestions

plot using a single y-axis



plot using both y-axes



# Further reading



Sarah P. Otto & Troy Day 2007

*A Biologist's Guide to Mathematical Modelling in Ecology and Evolution*  
Princeton University Press



A. N. Phillips

Reduction of HIV concentration during acute infection: Independence from a specific immune response

*Science*, 271:497–499, 1996