

Divergent evolutionary trajectories of two young, homomorphic, and closely related sex chromosome systems

Benjamin L. S. Furman*, and Ben J. Evans*

*McMaster University, Hamilton, Ontario, Canada

*Corresponding authors: E-mail: furmanbl@mcmaster.ca, evansb@mcmaster.ca.

Data Deposition: Sequence data for this paper have been deposited in the NCBI SRA, BioProject accession numbers: PRJNA421148 (*X. laevis* RADSeq), PRJNA421481 (*X. borealis* WGS).

Abstract

There exists extraordinary variation among species in the degree and nature of sex chromosome divergence. However, much of our knowledge about sex chromosomes is based on comparisons between deeply diverged species with different ancestral sex chromosomes, making it difficult to establish how fast and why sex chromosomes acquire variable levels of divergence. To address this problem, we studied sex chromosome evolution in two species of African clawed frog (*Xenopus*), both of whom acquired novel systems for sex determination from a recent common ancestor, and both of whom have female (ZW/ZZ) heterogamy. Derived sex chromosomes of one species, *X. laevis*, have a small region of suppressed recombination that surrounds the sex determining locus, and have remained this way for millions of years. In the other species, *X. borealis*, a younger sex chromosome system exists on a different pair of chromosomes, but the region of suppressed recombination surrounding an unidentified sex determining gene is vast, spanning almost half of the sex chromosomes. Differences between these sex chromosome systems are also apparent in the extent of nucleotide divergence between the sex chromosomes carried by females. Our analyses also indicate that in autosomes of both of these species, recombination during oogenesis occurs more frequently and in different genomic locations than during spermatogenesis. These results demonstrate that new sex chromosomes can assume radically different evolutionary trajectories, with far-reaching genomic consequences. They also suggest that in some instances the origin of new triggers for sex determination may be coupled with rapid evolution sex chromosomes, including recombination suppression of large genomic regions.

Key words: sex chromosomes, linkage map, recombination suppression, differentiation, amphibian, *Xenopus*

Introduction

Sex chromosomes originate when an autosome acquires a mutation that triggers development of one sex or the other. Recombination between sex chromosomes (the X and Y or Z and W) can be suppressed in regions that include and flank the sex determining mutation, which causes sex-specific inheritance of a sex determining trigger (Charlesworth et al., 1991). Portions of sex chromosomes that lack recombination (e.g., the sex specific portions of the Y or W) and portions that have a reduced level of recombination compared to the autosomes (e.g., the non-pseudoautosomal regions of the X or Z) are subject to distinct population genetic phenomena from autosomes. These genomic regions generally have a lower effective population size than autosomes and thus experience weaker purifying selection (Rice et al., 1994). Portions of each sex chromosome that have a sex-biased mode of inheritance may also have distinct mutation rates (e.g., Makova and Li, 2002)

and generation times (Amster and Sella, 2016). Differences in the variance of reproductive success between each sex can further contribute to the disparity in extent of genetic drift (the effective population size) of these regions (Charlesworth, 2009).

A lack of recombination causes portions of the two sex chromosomes to diverge from one another in nucleotide sequence, gene content, and the abundance and distribution of transposable and other repetitive elements (Charlesworth and Charlesworth, 2000; Bachtrog, 2013). Additionally, the non-recombining region may expand due to accumulation of sexually antagonistic genes, because sex-biased inheritance can mitigate sexual antagonism (Rice, 1987; Wright et al., 2017). Over time, these factors can lead to cytological distinctions between the sex chromosomes, a condition known as sex chromosome heteromorphy. In various taxa (e.g., some mammals, birds, and plants), divergence of sex chromosomes occurred incrementally along the length of the sex chromosomes due to se-

quential inversions or natural selection on recombination modifiers, expanding the non-recombining regions in a stepwise fashion (Bergero and Charlesworth, 2009; Coop and Przeworski, 2007; Vicoso et al., 2013).

Interestingly and perhaps counterintuitively, the age of the sex chromosomes does not seem to be tightly correlated with whether or not sex chromosomes are cytologically distinct (heteromorphic) or indistinct (homomorphic) (reviewed in Wright et al., 2016). In some old sex chromosomes, for example those of neoaves (> 100 MY; Zhou et al. 2014) and therian mammals (~150 MY; Graves 2006), and also some young sex chromosomes, such as those of *Drosophila miranda* (~1 MY; Bachtrog and Charlesworth, 2002) and *Silene latifolia* (10–20 MY; Bergero et al., 2007), divergence between the sex chromosomes is pronounced. In contrast, in the old sex chromosomes of ratite birds (> 100 MY; Zhou et al. 2014), recombination is suppressed over large regions of the sex chromosomes, but accompanied at the nucleotide level by relatively modest differentiation between the sex chromosomes and minimal cytological differentiation (Vicoso et al., 2013; Yazdi and Ellegren, 2014). An extreme case of homomorphy exists in the young sex chromosomes of tiger pufferfish, where a single mutation appears to control sexual differentiation and there is no evidence of suppressed recombination (Kamiya et al., 2012). In the young sex chromosomes of hylid tree frogs (~5 my old) and Palearctic green toads (~3.3 MY old), recombination appears to be low or absent in heterogametic males, but there is not substantial nucleotide divergence (Stöck et al., 2011, 2013). Why sex chromosomes of some species are homomorphic whereas those of others are heteromorphic, and why some heteromorphic sex chromosomes are more cytologically diverged than others remains enigmatic (Wright et al., 2016).

Sex chromosomes evolved multiple times in *Xenopus*

Insights into the origin of variation among species in sex chromosome divergence may be gained by examining whether, to what extent, why, and for how long recombination is suppressed in genomic regions flanking the sex determining locus in multiple species. For this reason, we quantified and compared recombination on the sex chromosomes of the African clawed frog, *Xenopus laevis*, and the Marsabit clawed frog, *Xenopus borealis*. The most recent common ancestor of these two species experienced allotetraploidization about 18–34 MY ago (Evans et al., 2015;

Session et al., 2016). These and other allotetraploid species in subgenus *Xenopus* have $2n=4s=36$ chromosomes, where n refers to the number of chromosomes in a haploid gamete and s refers to the number of chromosomes in an ancestral gamete prior to genome duplication. Chromosomes in tetraploids in subgenus *Xenopus* are numbered 1–18 followed by an L or an S, indicating from which of two diploid ancestors each chromosome was derived (Matsuda et al., 2015).

Species in genus *Xenopus* have homomorphic sex chromosomes (Tymowska and Fischberg, 1973; Tymowska, 1991), and three non-homologous sex determining systems have been identified in this group. One is on chromosome 2L of the allotetraploid species *X. laevis* (Yoshimoto et al., 2008) and also several other allopolyploid *Xenopus* species (Bewick et al., 2011). In these species, the W chromosome carries a gene called *DM-W* that triggers female sexual differentiation (Yoshimoto et al., 2008). *DM-W* originated after the whole genome duplication event ancestral to subgenus *Xenopus* species (Bewick et al., 2011). A second sex determination system in *Xenopus* is located on chromosome 8L in the allotetraploid species *X. borealis* (Furman and Evans, 2016). This sex determination system evolved in *X. borealis* from an ancestor that carried *DM-W* (Furman and Evans, 2016). A third sex determination system in *Xenopus* is located on chromosome 7 in the diploid species *Xenopus silurana tropicalis* (Olmstead et al., 2010; Evans et al., 2015). In *X. tropicalis*, Z, W, and Y chromosomes segregate (Roco et al., 2015). Overall then, of the three sets of sex chromosomes in *Xenopus*, at least two – those of *X. laevis* and *X. borealis* are newly evolved, and the system of *X. borealis* is proposed to be derived with respect to (i.e., younger than) the system of *X. laevis* (Fig. 1; Furman and Evans, 2016).

This variation in sex chromosomes among *Xenopus* species presents an opportunity to compare the evolutionary trajectories of two newly established sex chromosome systems (i.e., the sex chromosomes of *X. borealis* and *X. laevis*). Some differences between the W and Z chromosomes of *X. laevis* have been detected, including differences in gene content, insertion-deletion mutations, and nucleotide divergence, but this limited to only a few hundred Kb (< 1% of the chromosome length; Mawaribuchi et al., 2016). However, in general, in *X. laevis* and most other *Xenopus* species little is known about fundamental evolutionary genomic characteristics of sex and recombination, such as sex chromosome-wide levels of divergence, the extent of

sex-linkage of genes on sex chromosomes, genome-wide variation in rates of recombination, or sex differences in rates of recombination. We therefore used reduced genome sequencing of parents and offspring of each species to assess sex-linkage of SNPs and construct sex specific linkage maps for both species. We found that these two systems differ greatly in the extent of sex chromosome recombination suppression during oogenesis, with the younger system in *X. borealis* exhibiting a substantially larger region than the older system of *X. laevis*. Whole genome sequence data indicate that the non-recombining portions of the *X. borealis* sex chromosomes have a modest, but detectable, level of nucleotide divergence. Finally, linkage mapping in both species demonstrates that females have higher rates of recombination than males of both species, and that the location of crossovers is distinctive between females and males in both species, but similar in same sex comparisons across species. These findings demonstrate that newly evolved sex chromosomes in different species may rapidly assume radically different evolutionary trajectories.

Methods

Reduced representation genome sequences from *X. laevis* and *X. borealis* families

To assess genome wide sex-linkage we used reduced genome sequencing (genotype by sequencing (GBS) (Elshire et al., 2011) and restriction site associated DNA sequencing (RADSeq) (Baird et al., 2008)) on parents and offspring of an *X. borealis* family and an *X. laevis* family, respectively. For the *X. borealis* family, we used GBS data that we previously reported (Furman and Evans, 2016), with a female and male obtained from *XenopusExpress* (Brooksville, FL, USA). These GBS data included mother, father, 24 daughters, and 23 sons (22 and 17 individuals, respectively, after filtering, see Supplement S1.1), with offspring sex determined by dissection after euthanasia. The GBS data were 100 base pairs (bp) single-end sequences; library preparation and sequencing was performed at Cornell University Institute of Biotechnology Genome Diversity Facility on an Illumina HiSeq 2500; other details about these data available in Furman and Evans (2016). For the *X. laevis* family, we obtained female and male individuals from Boreal Science (St. Catharines, ON, Canada). We induced breeding with injection of human chorionic gonadotropin and determined the sex of tadpoles using primers for *DM-W*, which amplifies only in females, and sex-shared primers for *DMRT1* as a positive control

(Yoshimoto et al., 2008). The RADSeq library was generated by Floragenex (Portland, Oregon, USA) on both *X. laevis* parents, 17 daughters, and 20 sons and 150 bp single-end sequencing was performed at the University of Oregon using an Illumina HiSeq 2500 machine. Though slightly different procedures were used to generate reduced representation genome sequences from each species, the nature of the data is essentially the same – both methods produced sequence data from many homologous regions in most or all individuals from each family.

GBS or RADSeq data from each *X. borealis* or *X. laevis* individual were de-multiplexed, trimmed, and aligned to the *X. laevis* genome version 9.1 (www.xenbase.org) followed by genotyping and filtering steps that are described in the Supplement S1.1. This yielded a panel of SNPs for each family that were used to study recombination as described next. We discuss the potential impacts that the differences in the datasets of *X. borealis* and *X. laevis* may have on our study in Supplement S1.1, Fig. S4.

Sex-linked genomic regions

In *X. laevis* and *X. borealis*, females are the heterogametic sex (Yoshimoto et al., 2008; Furman and Evans, 2016). Using the filtered data for both families, we thus calculated maternal genotype association with the phenotypic sex (male or female) of each individual SNP following Goudet et al. (1996). Significance was assessed using a false discovery rate correction on the *P*-value of association with sex ($\alpha = 0.05$, using R; R Core Team 2016) and we discarded from this analysis maternal SNPs that were also heterozygous in the father. In order make inferences discussed below about the region of suppressed recombination that flanks the trigger for sex determination, for each maternal SNP we also determined the frequency of the most common genotype in daughters and then the frequency of this same genotype in sons. We refer to this frequency as the “major daughter genotype frequency”. At a completely sex-linked site that was heterozygous in the mother and homozygous in the father, we expected offspring genotypes to be homozygous in one sex and heterozygous in the other (which sex is heterozygous depends on whether the SNP was on the maternal Z or W). Thus, the major daughter genotype frequency at a completely sex-linked site would be 1.0 for daughters, and 0.0 for sons. Conversely, at an autosomal site the major daughter genotype frequency in daughters should be about 50% (but always $\geq 50\%$ because we excluded from this

analysis positions with more than two variants). In sons, the major daughter genotype frequency should also be about 50% at autosomal sites, but could be lower or higher than this value.

Linkage maps

We set out to evaluate rates and locations of recombination events in the mother and the father of our laboratory crosses. To accomplish this, we used the R package *OneMap* (Margarido et al., 2007) to construct linkage groups based on variable sites from the *X. borealis* and *X. laevis* families that mapped to each of the 18 *X. laevis* chromosomes in the reference genome. For each *X. laevis* chromosome and separately for each species, linkage groups were constructed with a maximum recombination fraction of 0.4 and a LOD threshold of five. With perfect synteny between the *X. laevis* and *X. borealis* and an even genomic distribution of genotyped SNPs, there should be one linkage group per *X. laevis* chromosome. However, we frequently identified several linkage groups per *X. laevis* chromosome in each species cross and we suspect that this was a consequence of genotyping and mapping errors (see below) and regions with sparse SNPs due to poor mapping of *X. borealis* reads to the *X. laevis* reference genome. For the *X. borealis* family, rearrangements between *X. borealis* and *X. laevis* could also break up a chromosome-specific linkage group. For either species, genome assembly errors could also prevent assembly of one linkage group for a chromosome. We note that our linkage maps did not include a particularly large number of offspring (39 in *X. borealis* and 37 in *X. laevis*), and this contributed to a lack of statistical power to form whole-chromosome linkage groups. However, this was not a concern for (or an objective of) our analyses, which focus on genomic regions for which assembly of linkage groups was possible.

In order to evaluate rates of recombination in the mother and father of each species, we selected the largest linkage group from each chromosome and divided the markers in each linkage group into those that were heterozygous in the mother, in the father, or in both parents. Then, using each of the maternal and paternal sets of markers from each of the largest linkage groups per chromosome, we re-computed recombination fractions between the sets of sex-specific markers and constrained marker order to match the mapping position in the v.9.1 *X. laevis* genome. For the *X. borealis* family, some chromosomes had very few or no double heterozygous sites (sites that were heterozygous

in both parents), which is a consequence of the lower overall amount of data for this cross compared to the *X. laevis* cross (due to mapping of *X. borealis* but not *X. laevis* data to a diverged reference genome, and the lower overall coverage we obtained from the GBS data compared to the RADSeq data). This meant that the recombination fractions between male and female markers were unable to be estimated for some chromosomes, and thus the first step of creating a joint linkage group could not be performed. For these chromosomes, we instead selected the largest female-specific and largest male-specific linkage group for each chromosome independently to estimate sex-specific linkage maps. Thus for these chromosomes, the male and female linkage groups do not span identical genomic regions.

Error correction and haplotype estimation

Genotyping errors create genotypes resembling recombined haplotypes that distort linkage maps and lead to inflated map lengths (Hackett and Broadfoot, 2003). Although we filtered incompatible parent-offspring genotypes (Supplement S1.1), under-calling of heterozygous sites can also produce incorrect homozygous genotypes in offspring that are nonetheless compatible with parental genotypes. To deal with this problem, we identified putative genotype errors based on phased offspring haplotypes. Each parent has two haplotypes per chromosome, and sites inherited by offspring can be assigned to one or the other haplotype for each parent. Recombination during gametogenesis creates new combinations of the two parental haplotypes within an offspring, with the ‘phase’ referring to which parental haplotype an offspring site comes from (see Fig. S1 for a visual explanation). Genotyping errors appears as a change in phase for a single SNPs (or a few SNPs in a row) when compared to surrounding SNPs. This pattern at one or few sites can also arise biologically from a double recombination (a crossover on either side of a variable position). However, double recombination events in small genomic windows are considered to be unlikely because of recombination interference (reviewed in Zickler and Kleckner, 2016).

To identify putative genotype errors, we used the parental phase estimated during linkage map construction (using *OneMap*; see Wu et al. 2002 for details on phase estimation of outcross maps) to estimate the parental haplotypes inherited by each offspring individual, for each chromosome-specific linkage map (Fig. S1a,b). Under the assumption that double recombination events are rare in small genomic windows, we

set to missing data any single genotype supporting a phase change in an individual at just that site (i.e., sites whose flanking genotypes were consistent double recombination event around a single genotyped site). As well, any genotypes in an individual that indicated a double recombination event that only encompassed a small genomic window of < 5 Mb were set to missing data (i.e., a series of sites within 5 Mb who were in an alternate phase compared to adjacent sites). For the *X. laevis* cross, which involved substantially more markers than the *X. borealis* cross, there were more of these potential genotyping errors (4% of all genotyped sites in individuals indicated a double recombination at either a single site or phase changes encompassing < 5MB in the *X. laevis* maps, compared to <0.5% for either in the *X. borealis* map; Table S1). Over 90% of the putative genotyping errors that were identified based on double recombination like phase changes in the *X. laevis* maps were homozygous, which is consistent with the bulk of these errors having been generated by under-called heterozygous positions (Table S1). After setting these genotypes to missing data in the affected individuals, we re-estimated linkage maps for each chromosome, for each parent for each species. Map distances were then calculated using the Kosambi function (Kosambi, 1943).

To quantify recombination events across all maps, we counted all phase changes in each linkage map for each individual based on haplotypes that were constructed from phased SNPs in each offspring. The location of recombination events was approximated as half the distance between the two markers bordering a recombination event in the *X. laevis* reference genome. We assessed the relationship between linkage map length and the amount of bp covered (on the *X. laevis* genome) by each map using a linear model, fitting an interaction between sex and species, along with a three-way interaction between sex, species, and the Mb covered by a linkage map (after scaling and centering Mb) using R. This strategy allowed us to assess for each sex and species slopes for the relationship between cM and Mb. We then used the `confint` function to compute confidence intervals on the estimates.

Divergence between the W and Z chromosomes of *X. borealis*

As discussed below, our analysis identified a large region of the *X. borealis* sex chromosomes that had sex-linked inheritance. If recombination has been suppressed in this region for a protracted period of evolu-

tionary time, we expected molecular polymorphism in the mother to be higher than the homologous region of the father due to the accumulation of diverged sites between the W and Z. For this reason, we also predicted that polymorphism in this region of the maternal sex chromosomes would be higher than other recombining portions of the maternal genome.

To explore the effects of this lack of recombination at the nucleotide level, we performed whole genome sequencing on the parents of our *X. borealis* family using the Illumina HiSeqX platform at The Center for Applied Genomics (Toronto, Canada), with both individuals multiplexed across two lanes. We trimmed the data, mapped it to the *X. laevis* reference genome, and genotyped and filtered the data as described in the Supplement S1.4. Mapping to a diverged reference genome could lead to a bias of more conserved sequences mapping, than sequences that have evolved quickly. With sex chromosomes, faster-Z (i.e., rapid evolution of Z-linked genes) or degeneration of the W sequences could lead to an underrepresentation of rapidly evolved sequences, leading to an underestimation of divergence. Contrary to this expectation, however, the number of reads mapped to the sex linked region of chromosome 8L in the female (10.2 million) was similar to other identically sized regions of other chromosomes (range 7.8–11 million).

One concern in the quantification of divergence in the non-recombining portion of the sex chromosomes is that intergenic regions may have many mapping errors due to repetitive sequences. For this reason, we focused our calculation of nucleotide diversity on genomic regions that are within and flank genes, because these areas contain less repetitive DNA (at least in *X. tropicalis*; Shen et al. 2013). We used the *X. laevis* genome annotation (version 9.1 primary gene models gff file; www.xenbase.org) to separately calculate nucleotide diversity (π) in each parent for coding sequence of genes (hereafter CDS), introns, 5' and 3' untranslated regions (hereafter UTR), 5,000 bp upstream of the 5' UTR, and 5,000 bp downstream of the 3' UTR for genes on all chromosomes. We considered only estimates that were generated from at least 200 bp of contiguous data from both *X. borealis* individuals. Overall, we measured π in 30,876 CDS regions, 3,092 5' UTRs, 14,954 3' UTRs, 119,420 introns, 30,326 upstream regions, and 30,270 downstream regions (for a total of 230,016 genomic regions) in the female and the male *X. borealis* individuals.

To test whether the W and Z chromosomes were

more diverged in the mother than the homologous Z region in the father, we used a linear mixed model implemented by the `lme4` package in R (Bates et al., 2015). We set as fixed effects sex (female or male) and sex-linkage (defined as sex-linked if between bp 4,605,306 and 51,708,524 (corresponding to 100% sex linked tags; Fig. 1) of chromosome 8L as defined by the analysis of sex-linked GBS tags discussed below). The six categories of gene regions (CDS, 5' & 3' UTRs, introns, up/down-stream) were set as a random effects. The model also included an interaction between the two fixed effects (sex and sex-linkage). We then used likelihood profiles (using the `profile` command in `lme4`) to calculate confidence intervals on the estimated coefficients.

To visualize and test for differences in divergence of the sex-linked region, we calculated median π for the mother and father in 1 Mb windows of chromosome 8L, using the π estimates from each of the genomic regions (intragenic, 5'UTR, 3'UTR, introns, 5,000 bp upstream of genes, and 5,000 bp downstream of genes). Because the mother and father had different levels of polymorphism, we needed to control for this difference in our comparisons between genomic regions of each individual. We therefore first calculated the median π value of all 1 Mb windows across chromosome 8L for each individual. We then standardized the maternal and paternal estimates of π by dividing by their corresponding chromosome-wide median. In order to compare these standardized values of diversity, we then divided the standardized estimates of π measured in each 1 Mb window of the mother by the standardized estimates of π measured in the homologous window of the father. With no difference in level of divergence between alleles we expected this ratio to be equal to one; if the W and Z chromosome were more diverged from each other in the mother than the Z and Z chromosome were from each other in the father, this ratio should be greater than one. We tested for a difference between the sex-linked and non-sex-linked portions using a Wilcoxon rank sum test and the measured disparity between parents of each 1 Mb estimates of standardized π . We also explored whether there was a higher rate of synonymous and nonsynonymous substitutions in genes on the non-recombining portion of the sex chromosomes to the rest of the genome using the WGS sequence data as described in detail in the Supplement S1.5. Finally, we explored the possibility of an accumulation of deletions and/or insertions on the sex chromosomes. Further details of these analyses are presented in the Supplement S1.6.

Validation of *X. borealis* sex chromosomes and recombination suppression

To explore whether the expansive region of suppressed recombination in *X. borealis* was limited to our lab raised family, we raised a second family of *X. borealis* using different parents. We then sequenced two genes (*SOX3* and *NR5A-1* (alternatively, *SF-1*)) located 25 Mb apart within the sex linked region (according to placement in the *X. laevis* genome v9.1) to look at co-inheritance of alleles from parents to offspring. We also surveyed a panel of adults that were not used in either cross from both sexes to assess linkage of alleles at these two genes. Further details of these assessments are in the Supplement S1.3.

Results

Diverse evolutionary fates of newly evolved sex chromosomes

Our analysis of the sex chromosomes of *X. borealis* and *X. laevis* identified a far larger region of sex-linked SNPs in *X. borealis* (Fig. 1). In *X. borealis*, 40 maternal SNPs spanning ~52 Mb (43%) of the sex chromosome (8L) had a significant association with the phenotypic sex of offspring (positions 4,605,306 – 56,690,925 of a total chromosome length of ~120 Mb in the *X. laevis* genome assembly; $P < 0.05$ after FDR correction; Fig. 1, S2). Within this region, daughters had identical genotypes at 34 of the 40 SNPs, with only one daughter differing for the last seven in the region (see below). Similarly in most sons, maternally inherited molecular variation in this genomic region was also almost entirely sex-linked, with exceptions discussed below. Across the entire genome after filtering, the SNP dataset consisted of 1,813 variable positions and there were more heterozygous SNPs in the mother than the father (1,103 and 644 SNPs in the mother and father, respectively, and 66 positions were heterozygous in both parents, with 15–133 SNPs per chromosome, and a mean of 61.8 maternal SNPs per chromosome). For maternal heterozygous positions used for assessing sex linkage in *X. borealis*, daughters had a median depth of 68 and genotype quality of 99 (maximum possible value), sons had a depth of 31 and a genotype quality of 99 (Fig. S4). Aligning to the diverged *X. laevis* genome substantially reduced the number of SNPs recovered to about 10% of the *de novo* SNP discovery method that did not involve mapping to the *X. laevis* genome (Furman and Evans, 2016)).

In sharp contrast, on the *X. laevis* sex chromo-




FIG. 1.—Sex-linkage of SNPs on sex chromosomes of *X. borealis* and *X. laevis*. In each graph, the x-axis is the position on the sex chromosome using the coordinates of the *X. laevis* reference genome and the y-axis is the major daughter genotype frequency in sons and daughters (see Methods for details) with colors as defined in the key indicating whether or not a SNP is significantly associated with sex (FDR corrected $P < 0.05$). For each species a diagram of a chromosome is shaded darker in the region of suppressed recombination. The inset phylogeny is from (Furman and Evans, 2016); *DM-W* is carried by female *X. clivii*, but its presence on chr2L has not been confirmed.

somes (2L) significant sex-linkage was only detected at only 6 maternal SNPs spanning 2 Mb (1%; positions 178,144,865 to 180,779,644, and possibly to the end of the chromosome at ~181,296,000; $P < 0.05$ after FDR correction; Fig. 1,S5). In *X. laevis*, SNPs immediately adjacent to the statically associated SNPs also had a strongly sex-biased pattern of inheritance, which is consistent with recombination suppression (Fig. 1). A lack of a statistically significant sex-linkage of some SNPs in this small genomic region may be a consequence of under-called heterozygous positions (Table S1 and see Methods). Across the entire genome, there were 7,779 SNPs, and in this family the father was more polymorphic (1,618 and 4,547 in mother and father, respectively, and 1,614 positions were heterozygous in both parents). For maternal heterozygous positions used in the sex linkage analysis of *X. laevis*, daughters had a median depth of 67, and a genotype quality of 99, sons had a depth of 61 and a genotype quality of 99 (Fig. S4).

Within the sex-linked region of *X. borealis*, there was a section with no recombination, and an adjacent section with reduced recombination between positions 51,708,524 – 56,690,925 of chromosome 8L (Fig. 1). Seven consecutive SNPs on the end of this region indicated recombination between the W and Z in one daughter, who had the same genotype as the sons at these positions (Fig. 1). Additionally, by inspecting changes in parental phase in the offspring (see below), another maternal recombination event was observed immediately adjacent to the region of completely suppressed recombination in one of the sons (Fig. S3). We note that additional information from more offspring or other families could potentially identify further recombinational activity within the genomic region where we did not observe recombination.

The genomic locations of several SNPs in the *X. borealis* family suggested genotyping or mapping error (Fig. S2). For a few sites within the otherwise completely sex-linked region of chromosome 8L, different individual sons had the same genotype as their sisters (Fig. 1). If this were due to a real recombination event, we would expect these sons to have the same genotype as their sisters at adjacent SNPs as well. Although this pattern could arise from independent double recombination events around these single sites in different sons, a more plausible explanation is that these are genotyping errors.

We observed three SNPs that mapped to the middle of the sex-linked region of chromosome 8L that were

not associated with sex ($P > 0.05$, following FDR, two sites are overlapping on the plot; Fig. 1), and we also found five SNPs that were completely sex-linked that mapped chromosome 8S. These genotypes are best explained by mapping error between *X. borealis* sequence reads and the *X. laevis* genome, or perhaps assembly error in the *X. laevis* genome wherein homeologous portions of the 8L and 8S chromosomes are intermingled in the assembly. It is also possible that sections of homeologous sequences of *X. laevis* and *X. borealis* were lost in an asymmetric fashion after whole genome duplication, such that chromosome 8L in *X. laevis* is missing portions that were not lost in *X. borealis*. This could cause reads from *X. borealis* to map to homeologous sequence in the *X. laevis* genome, instead of the missing orthologous sequence in *X. laevis*.

We also identified a sex-linked site in *X. borealis* that mapped to *X. laevis* chromosome 5S (Fig. S2). We blasted sequence from the GBS tag that contained this SNP to a *de novo* assembly of the maternal *X. borealis* HiSeqX data that was assembled using SOAPdenovo v.2.04, with a *kmer* = 23, and default parameters. We then blasted the top hit scaffold back to the *X. laevis* genome and found that its best matches were chromosomes 8S and 8L with similar affinities. This suggests that that this site could be a translocation between *X. borealis* and *X. laevis*, an assembly error in the *X. laevis* genome, or a mapping error due to the short sequence length (< 100 bp) of each GBS tag.

Recombination is higher in females of both species

Sex differences in the linkage maps revealed higher recombination rates in females of both species. The female linkage maps of both species were longer (*X. laevis* = 1,572 cM; *X. borealis* = 719 cM) than the same-species male linkage maps (*X. laevis* = 1,275 cM; *X. borealis* = 165 cM; Fig. 2). Longer female maps were recovered despite female markers spanning fewer base pairs of the *X. laevis* genome in both species (*X. laevis* female = 1.76 giga base pairs (Gb), male = 2.28 Gb; *X. borealis* female = 0.96 Gb, male = 1.72 Gb; Fig. 2). Consistent with this, the number of crossovers higher in oogenesis than spermatogenesis in both species (*X. laevis*: oogenesis = 558 total; 15.1/offspring, spermatogenesis = 467 total; 12.6/offspring; *X. borealis*: oogenesis = 270 total, 7.3/offspring; spermatogenesis = 62 total; 1.6/offspring).

Also of note is that the locations of crossovers were distinctive in females and males of both species. Female crossovers were more concentrated in the middle of




FIG. 2.—Linkage map length (in cM) is positively correlated with the number of bp spanned by the map (based on the *X. laevis* genome) for maternal but not paternal linkage maps. Black “sex chr” dots indicate the linkage map of the sex chromosome of each species (chromosome 8L in *X. borealis*, chromosome 2L in *X. laevis*). Lines reflect linear model relationships; gray shading indicates the 95% confidence interval of this relationship. Additionally, chromosome 8S is highlighted for *X. borealis*, because it is the homeolog of the sex chromosome 8L (see Results for details).

the chromosomes, whereas male crossovers occurred more often at the ends of chromosomes (Fig. 3). Possibly related to this (see Discussion), the length in cM of female linkage maps of both species was positively correlated with the number of bp covered by a map, but this relationship was not found in the male linkage maps from either species (linear model slope estimates, 95% confidence intervals: *X. borealis* female = 36.96, 24.78–49.13, male = -0.50 -14.96–13.95, *X. laevis* female = 40.80, 30.66–50.94, male = 5.40, -8.04–18.83; Fig. 2). Similar results were recovered when total length of chromosome was used instead of the number of bp covered by the linkage map, or when the number of crossover events was used instead of total cM (results not shown).

For the *X. borealis* family, the largest female linkage group on chromosome 8L (the sex chromosome, which includes both the Z and the W chromosomes) was formed from markers that mapped to the sex-linked portion (Fig. 1), and did not include markers from the non-sex-linked portion (see Methods for possible explanations). This region spanned 52 Mb (43% of the total *X. laevis* chromosome 8L) and was only 5 cM in length. That this recombination probability is not 0 cM is attributable to two recombination events at the end of the region, each of which is illustrated in plots of offspring haplotype assignment (Fig. S3). The female

linkage map of chromosome 8L was much shorter in recombination probability (cM) than other female and male linkage maps that spanned similar numbers of bp on other chromosomes (Fig. 2). The male map of chromosome 8L in the *X. borealis* family, which corresponds to a pair of Z chromosomes, spanned almost the entire chromosome, and had a length of 13 cM, which is similar to other chromosomes (Fig. 2). In the father, we detected five recombination events within the portion of chromosome 8L (i.e., between two Z chromosomes) that had suppressed recombination in the mother (i.e., the region where there was almost no recombination between the W and Z chromosomes; Fig. S3).

Interestingly, even though it is not a sex chromosome, the maternal linkage map of the *X. borealis* chromosome that is homeologous to the sex chromosome – chromosome 8S – was also substantially shorter in cM than other linkage maps spanning a similar amount of megabases (it was below the best fit line; Fig. 2). This suggests that recombination is less frequent on this homeologous chromosome than other autosomes, even though it is not sex-linked.

The *X. laevis* female linkage map of chromosome 2L did not include the last 20 Mb, which is where *DM-W* resides (Session et al., 2016), and where we detected sex-linked SNPs (Fig. 1). Therefore, we did not detect




FIG. 3.—Density plots of recombination events with respect to the relative position along chromosomes (chromosome length scaled to be between 0 and 1) in the maternal and paternal linkage maps of *X. borealis* and *X. laevis*.

any restricted recombination in this map, and the size (in cM) of the linkage map of this chromosome was similar to the size of the linkage maps for other chromosomes spanning similar amounts of Mbp (Fig. 2).

Divergence between the sex-linked portions of the W and Z chromosomes of *X. borealis*

We analyzed genotypes inferred from whole genome sequencing (WGS) data from the mother and the father to test whether we could detect evidence of sex chromosome divergence between sex-linked portions of the W and Z sex chromosomes. Compared to the pseudoautosomal portion of chromosome 8L and also to the autosomes, the sex-linked portion of chromosome 8L had the highest median nucleotide diversity in the female (pairwise nucleotide diversity (π) = 0.012; Fig. 4a). In this female genome, diversity within the non-sex-linked (pseudoautosomal) portion of chromosome 8L was similar to that of other chromosomes (π = 0.009; Fig. 4a). In the male genome, diversity of each portion of chromosome 8L fell within the range of estimates from other chromosomes from this genome (sex linked: π = 0.0072; non-sex linked: π = 0.009; Fig. 4a). The nucleotide diversity measured for these chromosomes is far less than the 7% divergence of homeologous sequences (Evans and Kwon, 2015); the considerably lower π estimates reported here suggest that cross map-

ping of reads across subgenomes was relatively rare.

Analyses of nucleotide diversity in and around genes (divided into 6 categories; see Methods), which used a linear mixed model, recovered a significant interaction between sex and sex-linkage, indicating that the mother had a higher π than the father in the sex-linked portion of chromosome 8L compared to the rest of the genome, and after controlling for differences in polymorphism between these individuals (estimate of female diversity in the sex linked region = 0.0018, 0.0009 – 0.0027 95% CI, t-stat = 4.09; Fig. 4b). For this analysis, we discarded the first 4 million bp of chromosome 8L because we lack information on whether this region is also sex-linked (Fig. 1).

We note that nucleotide diversity in the sex-linked portion of the female sex chromosomes includes fixed differences between the W and Z chromosomes and also positions that are segregating on the Z chromosome. Thus, this measurement is influenced by demographic differences between the female and male (the female genome is more polymorphic; Fig. 4). However, we found that standardizing the estimates of nucleotide diversity by the genome-wide average for each individual (by dividing diversity estimates from the male or female genome by the corresponding genome-wide mean for each genome) did not affect the results of the linear mixed model (see Results and S1.4). In the analysis

of nucleotide diversity, the sex linked portion of chromosome 8L stood out as the most polymorphic region in the female genome, supporting the existence of fixed divergent sites between the W and Z chromosomes.

The disparity between the female and male in nucleotide diversity along chromosome 8L was greater in the sex-linked portion than the pseudoautosomal portion of chromosome 8L (Wilcoxon rank sum test: $P < 0.001$; Fig. 4c). This result is consistent with the results of the linear mixed model (above). There was also a peak of divergence near end of the chromosome in the non-sex-linked region (Fig. 4c), that overlapped with a region where *X. borealis* daughters were mostly inheriting the same allele, suggesting partial sex-linkage (Fig. 1). This could be due to an inversion, although we did not explore this possibility in our data.

Within coding regions, dN and dS were very slightly, but significantly (statistically) elevated in the sex-linked region compared to the rest of the genome for both the female and male, but dN/dS was not (based on a permutation test; see Supplement 1.5). But, unlike the analysis of all SNPs (above), which included more data, the sex linked region was not the highest for any value (dN , dS , or dN/dS) compared individually to the other chromosomes. This emphasizes the subtlety of the divergence in the sex linked region and indicates that the time since recombination suppression is recent. We did not recover evidence of substantial differences in coverage between the female and male on the sex chromosomes (see Supplement S1.6).

Discussion

More expansive recombination suppression on younger sex chromosomes

The homomorphic sex chromosomes of *X. borealis* and *X. laevis* experienced distinctive evolutionary histories since they originated. In *X. laevis*, the sex-linked region is restricted to a small portion on the end of a chromosome (2L). In *X. borealis*, however, the sex-linked region encompasses almost half of a chromosome (8L; Fig. 1), even though this sex chromosome system is thought to be derived with respect to the ancestral sex determination system of *X. laevis* (Furman and Evans, 2016). Within the region of suppressed recombination of both of these species, there is evidence of sex chromosome divergence at the molecular level (*X. borealis*: Fig. 4a-c, Supplement S1.5; *X. laevis*: Mawaribuchi et al. 2016). Although the magnitude of sex chromosome divergence in the large sex-linked re-

gion of the *X. borealis* sex chromosomes is modest, it appears that recombination has been suppressed over sufficient evolutionary time for these differences to be detectable, presumably for many thousands of generations or more. Supporting this, our second family of lab-reared *X. borealis* and the surveyed panel of adults also had completely suppressed recombination in this large region (there were some sex linked female heterozygous sites that appeared in both families and others that were unique to one family or the other, see Supplement S1.3). Together, these findings are consistent with observations made in other, more diverged species that the extent of recombination suppression need not be more expansive in older than younger sex chromosomes (reviewed in Wright et al., 2016). They further demonstrate that newly established sex chromosomes may assume radically different evolutionary trajectories.

We infer here that the younger sex chromosomes of *X. borealis* have a larger region of suppressed recombination than the older sex chromosomes of *X. laevis*. One possibility is that this is due to a large scale genomic change, such as an inversion or deletion leading to widespread recombination suppression (Charlesworth et al., 2005). We were unable to characterize rearrangements in the sex chromosomes of *X. borealis* here due to the nature of our WGS data (short reads and relatively low coverage). However, there were two crossover events detected in the sex linked region (Fig. 1, S3). As well, the level of divergence between the W and Z was lower in the last 1/3 of the sex linked region, consistent with a more recent cessation of recombination (and possibly indicating the presence of genomic regions – strata – with different levels of divergence). These results suggest that a single large scale inversion encompassing the entire sex-linked region is not likely the reason for suppressed recombination. We cannot rule out the possibility that there are smaller inversions within the sex linked region that causes recombination suppression in flanking regions. In some sex chromosome systems, inversions are not thought to be the driver of recombination suppression. For example, in the plant *S. latifolia*, inversions in the non-recombining portion of the sex chromosomes may have occurred after recombination suppression evolved (Bergero et al., 2008). We did not recover any evidence of major coverage differences between the sequenced female and male *X. borealis* (Supplement S1.4), suggesting a lack of deletions or insertion differences between the Z and W. However, our inference is limited by a lack of a con-specific reference genome, because




FIG. 4.—Nucleotide diversity (π) in *X. borealis* based on WGS data mapped to the *X. laevis* reference genome. (a) Median π by chromosome as measured in the six genomic categories; error bars indicate 95% CI bootstrap estimates (for further information on differences see Supplement S1.4). The 8L_NL category refers to the diversity measured on chromosome 8L in the non-sex-linked region (57 Mb – 120 Mb). (b) Box and whisker plot of π across six genomic categories (described in Methods); the y-axis is truncated at 0.05 for clarity. (c) Standardized nucleotide diversity of the female divided by the standardized nucleotide diversity of male in 1 Mb windows across chr8L; the completely sex-linked region is highlighted in dark purple, and the significantly sex-linked region with suppressed recombination in light purple (see Fig. 1).

unique or rapidly evolving sequences on the sex chromosomes of *X. borealis* may not map to the homologous portion of or be present in the *X. laevis* reference genome.

Alternatively, modifiers of recombination can be favored by natural selection to suppress recombination (Charlesworth et al., 2005; Coop and Przeworski, 2007). These genetic factors control chiasmata formation during meiosis, possibly by modifying chromosome structure, or via the action of genes or repetitive elements (Ji et al., 1999; Otto and Lenormand, 2002). Curiously, chromosome 8S in *X. borealis* also had a lower recombination rate than other chromosome linkage maps of similar size (Fig. 2). This chromosome is homeologous (i.e., related by genome duplication) to the sex chromosomes 8L (Session et al., 2016). This result offers the intriguing possibility that whatever is acting to suppress recombination on the sex chromosome may also influence recombination of homeologous sequence on chromosome 8S (genome-wide, the L and S nucleotide divergence is about 6%; (Session et al., 2016)). This is unlikely to be an artifact of mapping errors because linkage groups would not form from markers that were a mix of chromosome 8L and 8S, because SNPs on different chromosomes should have a recombination fraction of about 0.5 (above our threshold; Methods).

Sex-linkage with minimal divergence (similar to our observations in *X. borealis*) has also been found in other species. For instance, the Japan sea population of stickleback fish have a recently evolved set of sex chromosomes, which were generated by a fusion of the ancestral sex chromosome and an autosome (Kitano et al., 2009). In this system, recombination suppression spread from the point of sex chromosome fusion to an ancestral autosome along a large fraction of the neo-sex chromosome (Natri et al., 2013). Sex-linked genomic regions with variable levels of divergence suggest that the boundaries of recombination suppression evolve over time, and may encompass areas that are not yet diverged. As such, recombination may occasionally happen in these regions until a hard recombination boundary is established (Bergero and Charlesworth, 2009). In some other amphibians, periodic recombination may prevent divergence of the sex chromosomes (Perrin, 2009; Stöck et al., 2011; Dufresnes et al., 2014). Though recombination was not detected in this region for either family of *X. borealis*, it is possible that over long timescales the sex chromosomes of *X. borealis* may occasionally recombine.

However, the divergence detected here between the Z and W, though modest, indicates that recombination is not happening frequently enough to completely prevent divergence (Fig. 4a-b).

The relative ages of the sex chromosomes of *X. laevis* and *X. borealis*

Our inference that recombination suppression expanded more quickly in *X. borealis* than *X. laevis* is based on (i) the inferred origin of *DM-W* in subgenus after the whole genome duplication event shared by all extant subgenus *Xenopus* species (Bewick et al., 2011) and (ii) inferred phylogenetic relationships within subgenus *Xenopus* (Furman and Evans, 2016), which indicates that the *DM-W* based sex determination system is ancestral to the system of *X. borealis* (Fig. 1). If this phylogenetic inference were erroneous and instead the sex determining system of *X. borealis* were ancestral to the *DM-W* based system of *X. laevis*, the rate that recombination suppression expanded over the sex chromosomes of *X. borealis* could be slower than it seems here.

However, there are several lines of evidence that argue against *X. borealis* having the older sex chromosomes than *X. laevis*. First, the strongest phylogenetic signal found using 1,585 genes supports a paraphyletic clade of *DM-W* possessing species (Fig. 1; Furman and Evans, 2016). More specifically, the alternate hypothesis of monophyly of *DM-W*-possessing species is supported by substantially fewer genes than the hypothesis of paraphyly of *DM-W*-possessing species with a sister relationship between *DM-W*-possessing *X. clivii* and *X. borealis* (as presented in Fig. 1; Furman and Evans 2016). In fact, the hypothesis of monophyly of *DM-W*-possessing species has an equal support to another paraphyletic relationship among *DM-W*-possessing species where *X. borealis* is more closely related to *X. laevis* than *X. clivii* is to *X. laevis* (Furman and Evans, 2016).

Additional evidence against the possibility of older sex chromosomes in *X. borealis* is provided by divergence of orthologous autosomal genes of *X. borealis* and *X. laevis* (e.g., divergence of synonymous site of ~14%; Chain et al. 2008) that is substantially greater than that observed between the non-recombining regions of the *X. borealis* sex chromosomes (Fig. 4). Likewise, homeologous coding sequences (including nonsynonymous and synonymous sites) also have higher divergence (~7%; Evans and Kwon, 2015) than the non-recombining region of the *X. borealis* sex chromosomes. These genomic patterns are consistent with

the proposal that suppressed recombination in the sex chromosomes of *X. borealis* occurred after allotetraploidization. Thus, even if previous phylogenetic inferences (Furman and Evans, 2016) are incorrect, the level of divergence between these sex chromosomes still argues that the expansion of the non-recombining region occurred after the origin of *DM-W* (i.e., post-whole genome duplication in subgenus *Xenopus*) after or at least within a similar time frame (post-whole genome duplication).

More recombination in females than males, and in different genomic regions

Heterochiasmy refers to differences in sex-specific rates of recombination. Here, in two independently derived sex chromosome systems with female heterogamy, we observed heterochiasmy with females having a higher rate of recombination than males. In some species of bird and crab with female heterogamy, recombination rates appear to be similar between the sexes (Groenen et al., 2009; Backström et al., 2010; Nietlisbach et al., 2015; Cui et al., 2015). But in some fish and other bird species the rate of recombination is higher in heterogametic females (Ruan et al., 2010; Hansson et al., 2010), or higher in homogametic males (Kawakami et al., 2014). In vertebrates with male heterogamy, the rate of recombination is often higher in females, particularly in XY mammals (e.g., Wong et al., 2010; Ottolini et al., 2015), though exceptions are known where rates are similar between the sexes, or higher in males (Mank, 2009; Johnston et al., 2016, respectively).

In several other frog species with male heterogamy, heterochiasmy has been observed with a higher recombination rate in females (Berset-Brändli et al., 2008; Brelsford et al., 2016). This was interpreted to be consistent with the Haldane-Huxely Rule (Haldane, 1922; Huxley, 1928) which postulates that when one sex does not recombine (i.e., when one sex is achiasmatic), that sex is the heterogametic sex (Berset-Brändli et al., 2008; Brelsford et al., 2016). Our results suggest instead that in species with heterochiasmy, the sex with lower recombination is not strongly linked to which sex is heterogametic (Lenormand and Dutheil, 2005). Heterochiasmy may be more prominently influenced by haploid selection (Lenormand and Dutheil, 2005), sexual antagonism (Mank, 2009), or other explanations.

The locations of recombination events was sex-biased in both species of *Xenopus* investigated, with recombination most frequent in the center of chromo-

somes in females, versus the ends of chromosomes in males (Fig. 3). Sex specific differences in crossover location have been observed in other taxa, including, for example, frogs, dogs, and primates (Wong et al., 2010; Venn et al., 2014; Ottolini et al., 2015; Brelsford et al., 2016). Female linkage map length (in cM) and the number of crossover events was positively correlated with the amount of bp covered by the map and the total length of a chromosome, whereas in males this relationship was not observed (Fig. 2). A similar disparity between the sexes in the relationship of cM and Mb spanned by linkage maps has been observed in the frog *Hyla arborea* (Brelsford et al., 2016) and in humans (Ottolini et al., 2015). This sex specific difference could be due to the differences in recombination location. In females, because recombination is spread out across the middle of chromosomes, longer chromosomes may permit more recombination events to occur without crossover interference. In males, where recombination occurs mostly on the tips of chromosomes, crossover interference is less likely to vary among chromosomes with different lengths. Similar findings have been recovered in soay sheep, where male recombination is mostly biased to the last 18 Mb of each of the chromosome tips, with chromosomes ranging in size from ~50–200 Mb (Johnston et al., 2016), encompassing the chromosome length variation of *Xenopus* (Session et al., 2016). Why females and males have differences in recombination locations is potentially due to differences in meiosis. During spermatogenesis there appears to be more control over formation and number of crossover events compared to oogenesis, with crossovers stopping in the presence of errors and more often restricted to one per arm (Hunt and Hassold, 2002; Hassold et al., 2004; Coop and Przeworski, 2007). As well, maintenance of favorable allelic combination by haploid selection, which is generally stronger in males, may limit the breadth of possible crossover locations to genomic regions, such as chromosome tips, that have low gene density (Lenormand and Dutheil, 2005).

One possible caveat our conclusions on sex specific differences in recombination rate is that in some cases maternal and paternal linkage groups spanned non-overlapping genomic regions, which themselves may vary in the local rate of recombination (Groenen et al., 2009; Kawakami et al., 2014; Ottolini et al., 2015). Since male recombination rate is biased towards tips of chromosomes (Fig. 3), it is possible that crossover events were not accounted for in these linkage maps if tags do not span to the ends of chromo-

somes. Kawakami et al. (2014) also noted that RAD based studies in birds may also underestimate linkage map lengths if they also underrepresent microchromosomes and ends of chromosomes. In this study, the disparity between female and male linkage map lengths in *X. laevis* (1.2:1 ratio of map length) is much less than *X. borealis* (4.4:1). The total map lengths in *X. laevis* (females: 1,572 cM and males: 1,275 cM) was not far from a total map length of 1,800 cM, which is the expected length if there were an obligate rate of one crossover per chromosome arm. This suggests our estimate of recombination in *X. laevis* is not unreasonably low. As well, the female to male map length ratio in *X. laevis* of 1.2:1 is within the range of a wide variety of other species (1.4:1 for a fish, Ruan et al. 2010; 1.2:1 for a mammal, Wong et al. 2010; 1.1:1 for a bird, Kawakami et al. 2014). Thus, the sex specific differences detected in *X. laevis* are likely genuine. We note that the magnitude of the sex difference in recombination rate for *X. borealis* (females: 719 cM and males: 165 cM) may be exaggerated due to lower genomic coverage in the *X. borealis* family (though large differences in recombination between closely related species is known Kawakami et al. 2014). Furthermore, our linkage maps are not capturing all recombination events in either species because the per gamete rates of recombination are much less than the expectation of one event per chromosome of 18 (Results). As such, caution should be used when interpreting linkage maps from reduced genome sequencing technologies (e.g., RAD-seq, GBS), especially when a closely related reference genome is lacking to assess marker distribution across chromosomes.

Drivers of sex chromosome evolution and stasis

Information from a diversity of organisms suggest that the age of sex chromosomes is not a strong predictor of the amount divergence between sex chromosomes within a species (Wright et al., 2016). Our findings from the sex chromosomes of *X. borealis* and *X. laevis* support this inference. One possible explanation for these observations is that the genomic context in which a new sex chromosome system is established plays a large role in determining the extent of divergence a newly established will experience. For example, the ability to cope with dosage imbalances or the potential for dosage compensation mechanisms to evolve could strongly influence whether sex chromosomes become heteromorphic or not (Batada and Hurst 2007, but see Mank 2009). If, for instance, the sex chromosomes of *X. laevis* (chromosome 2L), con-

tains more dosage sensitive genes than the sex chromosomes of *X. borealis* (chromosome 8L), this could hinder the expansion of recombination suppression in *X. laevis* but not *X. borealis*. In ratites, for example, an inability to accommodate dosage imbalances may prevent sex chromosome divergence beyond the limited regions thought to no longer recombine (Adolfsson and Ellegren, 2013; Vicoso et al., 2013; Yazdi and Ellegren, 2014). As well, the life history or ecological context of a population can influence the fate of sex chromosomes. Guppies, which similar to *X. borealis* have a large sex linked region without extensive degeneration, show variability in the extent of sex linkage on the chromosomes depending on an interplay between the strength of sexual antagonism and predation pressures in the population (Wright et al., 2017). A compelling direction for further inquiry is to explore factors that govern sex chromosome divergence and stasis in African clawed frogs, including the role of natural selection (e.g., favoring balanced gene dosage between the sexes, sexually antagonistic selection, haploid selection (Rice et al., 1994; Lenormand, 2003; Adolfsson and Ellegren, 2013)), and non-selective events (e.g., recombination in sex reversed individuals (Perrin, 2009), or large scale inversions).

Supplementary Material

Supplementary figures and tables are available online.

Acknowledgements

We thank Brian Golding for providing computational resources. We also thank NSERC for funding support (CGSD3-475567-2015 to BLSF, RGPIN/283102-2012 and RGPIN-2017-05770 to BJE).

Literature Cited

- Adolfsson S, Ellegren H. 2013. Lack of dosage compensation accompanies the arrested stage of sex chromosome evolution in ostriches. *Molecular Biology and Evolution*. 30(4):806–810.
- Amster G, Sella G. 2016. Life history effects on the molecular clock of autosomes and sex chromosomes. *Proceedings of the National Academy of Sciences*. 113(6):1588–1593.
- Bachtrog D. 2013. Y chromosome evolution: emerging insights into processes of y chromosome degeneration. *Nature Reviews Genetics*. 14(2):113.
- Bachtrog D, Charlesworth B. 2002. Reduced adaptation of a non-recombining neo-y chromosome. *Nature*. 416(6878):323–326.
- Backström N, Forstmeier W, Schielzeth H, Mellenius H, Nam K, Bolund E, Webster MT, Öst T, Schneider M, Kempnaers B, et al. 2010. The recombination landscape of the zebra finch

- Taeniopygia guttata* genome. *Genome Research*. 20(4):485–495.
- Baird NA, Etter PD, Atwood TS, Currey MC, Shiver AL, Lewis ZA, Selker EU, Cresko WA, Johnson EA. 2008. Rapid SNP discovery and genetic mapping using sequenced rad markers. *PLoS One*. 3(10):e3376.
- Batada NN, Hurst LD. 2007. Evolution of chromosome organization driven by selection for reduced gene expression noise. *Nature Genetics*. 39(8):945–949.
- Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*. 67(1):1–48.
- Bergero R, Charlesworth D. 2009. The evolution of restricted recombination in sex chromosomes. *Trends in Ecology & Evolution*. 24(2):94–102.
- Bergero R, Charlesworth D, Filatov DA, Moore RC. 2008. Defining regions and rearrangements of the *Silene latifolia* y chromosome. *Genetics*. 178(4):2045–2053.
- Bergero R, Forrest A, Kamau E, Charlesworth D. 2007. Evolutionary strata on the X chromosomes of the dioecious plant *Silene latifolia*: evidence from new sex-linked genes. *Genetics*. 175(4):1945–1954.
- Berset-Brändli L, Jaquière J, Broquet T, Ulrich Y, Perrin N. 2008. Extreme heterochiasmy and nascent sex chromosomes in european tree frogs. *Proceedings of the Royal Society of London B: Biological Sciences*. 275(1642):1577–1585.
- Bewick AJ, Anderson DW, Evans BJ. 2011. Evolution of hte closely related, sex-related genes *DM-W* and *dmrt1* in african clawed frogs (*Xenopus*). *Evolution*. 65(3):698–712.
- Brelsford A, Dufresnes C, Perrin N. 2016a. High-density sex-specific linkage maps of a european tree frog (*Hyla arborea*) identify the sex chromosome without information on offspring sex. *Heredity*. 116(2):177–181.
- Brelsford A, Rodrigues N, Perrin N. 2016b. High-density linkage maps fail to detect any genetic component to sex determination in a *Rana temporaria* family. *Journal of Evolutionary Biology*. 29(1):220–225.
- Chain FJ, Ilieva D, Evans BJ. 2008. Duplicate gene evolution and expression in the wake of vertebrate allopolyploidization. *BMC Evolutionary Biology*. 8(1):43.
- Charlesworth B. 2009. Fundamental concepts in genetics: effective population size and patterns of molecular evolution and variation. *Nature Reviews Genetics*. 10(3):195.
- Charlesworth B, et al. 1991. The evolution of sex chromosomes. *Science*. 251(4997):1030–1033.
- Charlesworth B, Charlesworth D. 2000. The degeneration of y chromosomes. *Philosophical Transactions of the Royal Society of London B: Biological Sciences*. 355(1403):1563–1572.
- Charlesworth D, Charlesworth B, Marais G. 2005. Steps in the evolution of heteromorphic sex chromosomes. *Heredity*. 95(2):118–128.
- Coop G, Przeworski M. 2007. An evolutionary view of human recombination. *Nature Reviews Genetics*. 8:23–34.
- Cui Z, Hui M, Liu Y, Song C, Li X, Li Y, Liu L, Shi G, Wang S, Li F, et al. 2015. High-density linkage mapping aided by transcriptomics documents ZW sex determination system in the chinese mitten crab *Eriocheir sinensis*. *Heredity*. 115(3):206.
- Dufresnes C, Bertholet Y, Wassef J, Ghali K, Savary R, Pasteur B, Brelsford A, Rozenblut-Kościsty B, Ogielska M, Stöck M, et al. 2014. Sex-chromosome differentiation parallels post-glacial range expansion in european tree frogs (*Hyla arborea*). *Evolution*. 68(12):3445–3456.
- Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K, Buckler ES, Mitchell SE. 2011. A robust, simple genotyping-by-sequencing (GBS) approach for high diversity species. *PLoS One*. 6(5):e19379.
- Evans BJ, Carter TF, Greenbaum E, Gvoždík V, Kelley DB, McLaughlin PJ, Pauwels OS, Portik DM, Stanley EL, Tinsley RC, et al. 2015. Genetics, morphology, advertisement calls, and historical records distinguish six new polyploid species of african clawed frog (*Xenopus*, pipidae) from west and central africa. *PLoS one*. 10(12):e0142823.
- Evans BJ, Kwon T. 2015. Molecular polymorphism and divergence of duplicated genes in tetraploid african clawed frogs (*Xenopus*). *Cytogenetic and genome research*. 145(3–4):243–252.
- Furman BLS, Evans BJ. 2016. Sequential turnovers of sex chromosomes in african clawed frogs (*Xenopus*) suggest some genomic regions are good at sex determination. *G3: Genes/Genomes/Genetics*. 6(11):3625–3633.
- Goudet J, Raymond M, de Meelis T, Rousset F. 1996. Testing differentiation in diploid populations. *Genetics*. 144(4):1933–1940.
- Graves JAM. 2006. Sex chromosome specialization and degeneration in mammals. *Cell*. 124(5):901–914.
- Groenen MA, Wahlberg P, Foglio M, Cheng HH, Megens HJ, Crooijmans RP, Besnier F, Lathrop M, Muir WM, Wong GKS, et al. 2009. A high-density snp-based linkage map of the chicken genome reveals sequence features correlated with recombination rate. *Genome Research*. 19(3):510–519.
- Hackett C, Broadfoot L. 2003. Effects of genotyping errors, missing values and segregation distortion in molecular marker data on the construction of linkage maps. *Heredity*. 90(1):33.
- Haldane JB. 1922. Sex ratio and unisexual sterility in hybrid animals. *Journal of Genetics*. 12(2):101–109.
- Hansson B, Jungqvist M, Dawson D, Mueller J, Olano-Marin J, Ellegren H, Nilsson JÅ. 2010. Avian genome evolution: insights from a linkage map of the blue tit (*Cyanistes caeruleus*). *Heredity*. 104(1):67.
- Hassold T, Judis L, Chan E, Schwartz S, Seftel A, Lynn A. 2004. Cytological studies of meiotic recombination in human males. *Cytogenetic and Genome Research*. 107(3–4):249–255.
- Hunt PA, Hassold TJ. 2002. Sex matters in meiosis. *Science*. 296(5576):2181–2183.
- Huxley J. 1928. Sexual difference of linkage in *Gammarus chevreuxi*. *Journal of Genetics*. 20(2):145–156.
- Ji Y, Stelly DM, De Donato M, Goodman MM, Williams CG. 1999. A candidate recombination modifier gene for *Zea mays* l. *Genetics*. 151(2):821–830.
- Johnston SE, Bérénos C, Slate J, Pemberton JM. 2016. Conserved genetic architecture underlying individual recombination rate variation in a wild population of soay sheep (*Ovis aries*). *Genetics*. 203(1):583–598.
- Kamiya T, Kai W, Tasumi S, Oka A, Matsunaga T, Mizuno N, Fujita M, Suetake H, Suzuki S, Hosoya S, et al. 2012. A trans-species missense snp in *Amhr2* is associated with sex determination in the tiger pufferfish, *Takifugu rubripes* (fugu). *PLoS Genetics*. 8(7):e1002798.
- Kawakami T, Smeds L, Backström N, Husby A, Qvarnström A, Mugal CF, Olason P, Ellegren H. 2014. A high-density linkage

- map enables a second-generation collared flycatcher genome assembly and reveals the patterns of avian recombination rate variation and chromosomal evolution. *Molecular Ecology*. 23(16):4035–4058.
- Kitano J, Ross JA, Mori S, Kume M, Jones FC, Chan YF, Absher DM, Grimwood J, Schmutz J, Myers RM, et al. 2009. A role for a neo-sex chromosome in Stickleback speciation. *Nature*. 461(7267):1079–1083.
- Kosambi D. 1943. The estimation of map distances from recombination values. *Annals of Human Genetics*. 12(1):172–175.
- Lenormand T. 2003. The evolution of sex dimorphism in recombination. *Genetics*. 163(2):811–822.
- Lenormand T, Dutheil J. 2005. Recombination difference between sexes: a role for haploid selection. *PLoS Biology*. 3(3):e63.
- Makova KD, Li WH. 2002. Strong male-driven evolution of dna sequences in humans and apes. *Nature*. 416(6881):624–626.
- Mank JE. 2009a. The evolution of heterochiasmy: the role of sexual selection and sperm competition in determining sex-specific recombination rates in eutherian mammals. *Genetics Research*. 91(5):355–363.
- Mank JE. 2009b. The W, X, Y and Z of sex-chromosome dosage compensation. *Trends in Genetics*. 25(5):226–233.
- Margarido GRA, de Souza AP, Garcia AAF. 2007. Onemap: software for genetic mapping in outcrossing species. *Hereditas*. 144:78–79.
- Matsuda Y, Uno Y, Kondo M, Gilchrist MJ, Zorn AM, Rokhsar DS, Schmid M, Taira M. 2015. A new nomenclature of *Xenopus laevis* chromosomes based on the phylogenetic relationship to *Silurana/Xenopus tropicalis*. *Cytogenetic and Genome Research*. 145(3-4):187–191.
- Mawaribuchi S, Takahashi S, Wada M, Uno Y, Matsuda Y, Kondo M, Fukui A, Takamatsu N, Taira M, Ito M. 2016. Sex chromosome differentiation and the W-and Z-specific loci in *Xenopus laevis*. *Developmental Biology*. 426(2):393–400.
- Natri HM, Shikano T, Merilä J. 2013. Progressive recombination suppression and differentiation in recently evolved neo-sex chromosomes. *Molecular Biology and Evolution*. 30(5):1131–1144.
- Nietlisbach P, Camenisch G, Bucher T, Slate J, Keller LF, Postma E. 2015. A microsatellite-based linkage map for song sparrows (*Melospiza melodia*). *Molecular Ecology Resources*. 15(6):1486–1496.
- Olmstead AW, Lindberg-Livingston A, Degitz SJ. 2010. Genotyping sex in the amphibian, *Xenopus (Silurana) tropicalis*, for endocrine disruptor bioassays. *Aquatic Toxicology*. 98(1):60–66.
- Otto SP, Lenormand T. 2002. Resolving the paradox of sex and recombination. *Nature Reviews Genetics*. 3(4):252.
- Ottolini CS, Newnham LJ, Capalbo A, Natesan SA, Joshi HA, Cimadomo D, Griffin DK, Sage K, Summers MC, Thornhill AR, et al. 2015. Genome-wide maps of recombination and chromosome segregation in human oocytes and embryos show selection for maternal recombination rates. *Nature Genetics*. 47(7):727–735.
- Perrin N. 2009. Sex reversal: a fountain of youth for sex chromosomes? *Evolution*. 63(12):3043–3049.
- R Core Team. 2016. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.
- Rice WR. 1987. The accumulation of sexually antagonistic genes as a selective agent promoting the evolution of reduced recombination between primitive sex chromosomes. *Evolution*. 41(4):911–914.
- Rice WR, et al. 1994. Degeneration of a nonrecombining chromosome. *Science*. 263(5144):230–231.
- Roco AS, Olmstead AW, Degitz SJ, Amano T, Zimmerman LB, Bullejos M. 2015. Coexistence of Y, W, and Z sex chromosomes in *Xenopus tropicalis*. *Proceedings of the National Academy of Sciences*. 112(34):E4752–E4761.
- Ruan X, Wang W, Kong J, Yu F, Huang X. 2010. Genetic linkage mapping of turbot (*Scophthalmus maximus L.*) using microsatellite markers and its application in qtl analysis. *Aquaculture*. 308(3):89–100.
- Session AM, Uno Y, Kwon T, Chapman JA, Toyoda A, Takahashi S, Fukui A, Hikosaka A, Suzuki A, Kondo M, et al. 2016. Genome evolution in the allotetraploid frog *Xenopus laevis*. *Nature*. 538(7625):336–343.
- Shen JJ, Dushoff J, Bewick AJ, Chain FJ, Evans BJ. 2013. Genomic dynamics of transposable elements in the western clawed frog (*Silurana tropicalis*). *Genome Biology and Evolution*. 5(5):998–1009.
- Stöck M, Horn A, Grossen C, Lindtke D, Sermier R, Betto-Colliard C, Dufresnes C, Bonjour E, Dumas Z, Luquet E, et al. 2011. Ever-young sex chromosomes in european tree frogs. *PLoS Biology*. 9(5):e1001062.
- Stöck M, Savary R, Betto-Colliard C, Biollay S, Jourdan-Pineau H, Perrin N. 2013. Low rates of X-Y recombination, not turnovers, account for homomorphic sex chromosomes in several diploid species of palearctic green toads (*Bufo viridis* subgroup). *Journal of Evolutionary Biology*. 26(3):674–682.
- Tymowska J. 1991. Polyploidy and cytogenetic variation in frogs of the genus *Xenopus*. In: Green DM, Sessions SK, editors. *Amphibian Cytogenetics and Evolution*. Academic Press San Diego, CA. p. 259–297.
- Tymowska J, Fischberg M. 1973. Chromosome complements of the genus *Xenopus*. *Chromosoma*. 44(3):335–342.
- Venn O, Turner I, Mathieson I, de Groot N, Bontrop R, McVean G. 2014. Strong male bias drives germline mutation in chimpanzees. *Science*. 344(6189):1272–1275.
- Vicoso B, Kaiser VB, Bachtrog D. 2013. Sex-biased gene expression at homomorphic sex chromosomes in emus and its implication for sex chromosome evolution. *Proceedings of the National Academy of Sciences*. 110(16):6453–6458.
- Wong AK, Ruhe AL, Dumont BL, Robertson KR, Guerrero G, Shull SM, Ziegler JS, Millon LV, Broman KW, Payseur BA, et al. 2010. A comprehensive linkage map of the dog genome. *Genetics*. 184(2):595–605.
- Wright AE, Darolti I, Bloch NI, Oostra V, Sandkam B, Buechel SD, Kolm N, Breden F, Vicoso B, Mank JE. 2017. Convergent recombination suppression suggests role of sexual selection in guppy sex chromosome formation. *Nature Communications*. 8:14251.
- Wright AE, Dean R, Zimmer F, Mank JE. 2016. How to make a sex chromosome. *Nature Communications*. 7:12087.
- Wu R, Ma CX, Painter I, Zeng ZB. 2002. Simultaneous maximum likelihood estimation of linkage and linkage phases in outcrossing species. *Theoretical Population Biology*. 61(3):349–363.
- Yazdi HP, Ellegren H. 2014. Old but not (so) degenerated—slow evolution of largely homomorphic sex chromosomes in ratites.

Molecular Biology and Evolution.:msu101.

- Yoshimoto S, Okada E, Umemoto H, Tamura K, Uno Y, Nishida-Umehara C, Matsuda Y, Takamatsu N, Shiba T, Ito M. 2008. A w-linked dm-domain gene, dm-w, participates in primary ovary development in *Xenopus laevis*. Proceedings of the National Academy of Sciences. 105(7):2469–2474.
- Zhou Q, Zhang J, Bachtrog D, An N, Huang Q, Jarvis ED, Gilbert MTP, Zhang G. 2014. Complex evolutionary trajectories of sex chromosomes across bird taxa. Science. 346(6215):1246338.
- Zickler D, Kleckner N. 2016. A few of our favorite things: Pairing, the bouquet, crossover interference and evolution of meiosis. In: Seminars in Cell & Developmental Biology, Volume 54. Elsevier. p. 135–148.