

Supplemental material for ‘Early high rates and disparity in the evolution of fish-shaped reptiles’

Benjamin C. Moon

Thomas L. Stubbs

27 September 2019 o.2.965

Contents

Supplemental methods	1
Supplemental results	2
Supplemental figures	3
Supplementary tables	64
Supplemental code	71

Supplemental methods

Comparison of time-scaling methods. To assess the effects of variation in the timing of ichthyosaur evolution on discrete evolutionary rates, we further used the minimum branch length (MBL) tree-scaling method^{1,2}. This scales the tree according to occurrence dates, but ensures that each branch length is greater than a given value, rescaling ancestral branches as necessary to ensure this minimum length. Here, we used a MBL of 1 Ma as a reasonable minimum between speciation events and to avoid forcing excessive branch lengths where speciation may occur rapidly. We used the same sample of 120 phylogenetic trees as the main analysis from the Bayesian phylogenetic posterior distribution of Moon [3]. Trees were time-scaled in R⁴ using the function `timePaleoPhy` in the package `paleotree`¹ with point ages sampled from a uniform distribution between their first and last occurrences. Each tree was resampled 10 times to account for the occurrence ranges for each taxon (120 tree topologies × 10 samples = 1200 time-scaled trees total). These MBL time-scaled trees were then used for a further set of discrete character evolutionary rates analyses using function `DiscreteCharacterRate` of R package `Claddis`⁵. The results of this were used to produce ‘spaghetti’ plots for epoch-length bins and equal-length bins using modified scripts from Close *et al.* [6]. Code for all these analyses is included in Supplemental Code s1.

Additional disparity metrics. Our main results present ichthyosauriform disparity using per-bin pairwise differences between taxa from a distance matrix calculated using maximum observed rescaled distances⁵. Additionally, we compared different distance conversion and disparity metrics.

Claddis provides four distance metrics for discrete character data⁵: raw Euclidean distances (RAW), generalized Euclidean distances (GED)⁷, Gower’s coefficient (GOW)⁸, and maximum observable rescaled distances (MAX)⁵. All four distance metrics were run through the same disparity work flow. Recent

studies have shown that GED as implemented in Claddis is susceptible to the completeness of the original data matrix, which may have a strong effect on the resulting disparity^{9,10}; therefore we prefer MAX.

Similarly, several different disparity metrics have been developed, each with varying properties. Our main results present mean and weighted mean pairwise distances on MAX as this comes directly from the original data matrix, but we also calculated the pairwise distances for RAW, GED, and GOW distances matrices (fig. 1). We ordinated the data using Principal Coordinates Analysis (PCA), both with and without applying a correction to negative eigenvalues¹¹ and compared the correlation of the PCA data with the original distance matrix.

From the PCA data we used all the resultant axes to calculate per-bin sum of variances, sum of ranges, and centroid distances. These metrics have been used extensively in previous analyses^{9,12,13}, so we considered it pertinent to compare them. Binning, bootstrap resampling with 10 000 replicates, and complete rarefaction were completed using the functions `custom.subsets` and `boot.matrix`, and disparity calculations used the function `dispRity`, all from package `dispRity`¹⁴ in R. Code for this is included in Supplemental Code s1.

Supplemental results

Pairwise disparity. Broadly speaking, trends in disparity across all four distance matrices are similar: disparity peaks in the Late Triassic then declines through the Jurassic and Cretaceous (fig. 1). The bins that preserve the most completely coded taxa (fig. 1 CHAR: Early Jurassic; 201.3 Ma to 171.3 Ma) also show relatively increased disparity in RAW and GED distance matrices compared to GOW and MAX. Indeed, the earliest Jurassic bins are the most disparate for the RAW distance metric with both binning schemes, and for GED the earliest Jurassic bins have relatively higher disparity than GOW and MAX distance matrices. This is most likely a further effect of incompleteness degrading the disparity signal by averaging the difference between taxa^{9,10}, therefore we prefer the results given by GOW and MAX distance matrices.

Correlation of ordinated data. Negative eigenvalue correction notably decreased the variance described by the first few principal coordinate axes. The highest correlations between the original and ordinated data were found when including all ordinated axes (??). Without negative eigenvalue correction RAW and GED had the highest correlation, whereas GOW and MAX were reduced to ~0.8. With negative eigenvalue correction the pattern of correlations with increasing number of axes was more complex: RAW gradually increased whereas GED strongly decreased, but both rapidly increased to 1.0 with the last axes; GOW and MAX correlations both immediately decreased, increased to a peak at ~axis 60, then rapidly increased again when including the last axes.

Disparity of ordinated data. Wills [12] asserted that variance based disparity metrics are more suited to measuring overall dissimilarity whereas range-based metrics are appropriate for disparity as they are affected by occurrence and thus show the diversification of morphology. In this context, our results support our conclusions that ichthyosaurs represent an early burst of evolution: both of these metrics show initial high disparity from all distance matrices (fig. 2). Sum of variances also has a marked increase between the Early to Middle Triassic and a substantial decline in disparity between the Late Triassic–Early Jurassic in the combination of GOW/MAX distance matrix and uncorrected PCO; otherwise all curves follow similar trends. Sum of variances proves more resilient to sample size in rarefaction than either sum of ranges or centroid distance (fig. 3).

All sum of ranges curves display the same trends in disparity, differing only in the magnitude. Similarly, we find early high disparity and an increase between the Early–Middle Triassic (fig. 2). Disparity decreases substantially through the later Triassic, but broadly recovers in the Early Jurassic before more log-term

decline through to the extinction of the ichthyosaurs. Particularly low disparity (e.g. Middle Jurassic; 171.3 Ma to 161.3 Ma) are those represented by few taxa and relative incompleteness.

In the case of centroid distance, although this has been shown to be especially susceptible to issues of centroid slippage^{9,10}, our results show the same trends as for sum of variances: high early disparity that is sustained through to the Late Jurassic/Early Cretaceous before decline, with dips that are most likely related to incompleteness of specimens (fig. 2).

Morphospace occupation of ordinated data. Morphospace occupation between Triassic and post-Triassic Ichthyosauriformes is separated in almost all cases (fig. 4; except RAW and GED distances). Late Triassic taxa are also separated from earlier Triassic taxa in GOW and MAX distance without negative eigenvalue correction, and are consistently positioned more closely towards the Early Jurassic taxa. The variation in Jurassic and Cretaceous taxa is markedly increased in RAW and GED distances relative to GOW and MAX. Differences within Jurassic and Cretaceous taxa are more represented in PCo axis 2 than axis 3 in the RAW and GED morphospace plots, but in a combination of PCo axes 1 and 3 in GOW and MAX. All RAW and GED morphospace plots show more points towards the origins of the plots than GOW and MAX, a results of 'centroid slippage'^{9,10}; in particular these represent the least complete taxa.

Time-scaling and rates. Using the MBL time-scaling method created trees with a root age of 253.8 Ma to 268.5 Ma; older than the corresponding root ages from the Hedman scaling method. Rates of discrete character evolution are relatively lower for during the Early–Middle Triassic, but these earlier bins nonetheless show significantly higher rates of evolution than subsequent bins (fig. 5). Trends across the whole of ichthyosaur evolution remain similar, although there are increased peaks in the later Early Jurassic and the Late Cretaceous bins. Significantly low rates of discrete character evolution are reached in the Early Jurassic (epoch bins) or Late Triassic (10 Ma bins).

Supplemental figures

1	Per-bin discrete skeletal disparity of Ichthyosauriformes though the Mesozoic	3
2	Per-bin discrete skeletal disparity of Ichthyosauriformes through the Mesozoic from ordinated data	5
3	Per-bin rarefaction curves for each disparity-time curve shown in fig. 2	9
4	Morphospace occupation of Ichthyosauriformes through the Mesozoic	58
5	Rates of discrete skeletal character evolution in Ichthyosauriformes	61
6	Rates of skull size evolution in Ichthyosauriformes	63

Figure 1: (following page) **Per-bin discrete skeletal disparity of Ichthyosauriformes though the Mesozoic.** Pairwise and weighted pairwise dissimilarity measured from raw Euclidean (RAW), generalised Euclidean (GED), Gower (GOW), and maximum observed rescaled (MAX) distances between taxa in the cladistic dataset of Moon [3] binned into epochs and equal 10-million-year bins. Also, pairwise number of comparable characters between taxa (CHAR) indicating the variation in completeness and comparability in each bin. Mean values and 95% confidence intervals are shown from 10,000 bootstrap replicates.

Ichthyosaur disparity (mean pairwise dissimilarity) through time






Figure 2: (following pages) **Per-bin discrete skeletal disparity of Ichthyosauriformes through the Mesozoic from ordinated data.** Ichthyosaur disparity represented by mean sum of variances, mean sum of ranges, and mean centroid distance from each of eight PCA (four distance matrices: RAW, GED, GOW, MAX; with and without negative eigenvalue correction) on the cladistic matrix of Moon [3]. Error bars show 95% confidence intervals from 10,000 bootstrap replicates.

Ichthyosaur disparity (mean sum of variances) through time



Ichthyosaur disparity (mean sum of ranges) through time



Ichthyosaur disparity (mean centroid distance) through time






Figure 3: (following pages) **Per-bin rarefaction curves for each disparity-time curve shown in fig. 2**
Disparity for each bin is sequentially rarefied on taxon occurrences. Error polygon gives 95% confidence interval from 10,000 bootstrap replicates.


Rarefaction curves: mean sum of variances of uncorrected RAW distance matrix in epoch-length bins




Rarefaction curves: mean sum of variances of uncorrected GED distance matrix in epoch-length bins




Rarefaction curves: mean sum of variances of uncorrected GOW distance matrix in epoch-length bins




Rarefaction curves: mean sum of variances of uncorrected MAX distance matrix in epoch-length bins




Rarefaction curves: mean sum of variances of Cailleuz-corrected RAW distance matrix in epoch-length bins




Rarefaction curves: mean sum of variances of Cailleuz-corrected GED distance matrix in epoch-length bins




Rarefaction curves: mean sum of variances of Cailleuz-corrected GOW distance matrix in epoch-length bins




Rarefaction curves: mean sum of variances of Cailleuz-corrected MAX distance matrix in epoch-length bins




Rarefaction curves: mean sum of variances of uncorrected RAW distance matrix in 10 Ma bins




Rarefaction curves: mean sum of variances of uncorrected GED distance matrix in 10 Ma bins




Rarefaction curves: mean sum of variances of uncorrected GOW distance matrix in 10 Ma bins




Rarefaction curves: mean sum of variances of uncorrected MAX distance matrix in 10 Ma bins




Rarefaction curves: mean sum of variances of Cailleuz-corrected RAW distance matrix in 10 Ma bins




Rarefaction curves: mean sum of variances of Cailleuz-corrected GED distance matrix in 10 Ma bins




Rarefaction curves: mean sum of variances of Cailleuz-corrected GOW distance matrix in 10 Ma bins




Rarefaction curves: mean sum of variances of Cailleuz-corrected MAX distance matrix in 10 Ma bins




Rarefaction curves: mean sum of ranges of uncorrected RAW distance matrix in epoch-length bins




Rarefaction curves: mean sum of ranges of uncorrected GED distance matrix in epoch-length bins




Rarefaction curves: mean sum of ranges of uncorrected GOW distance matrix in epoch-length bins




Rarefaction curves: mean sum of ranges of uncorrected MAX distance matrix in epoch-length bins




Rarefaction curves: mean sum of ranges of Cailleuz-corrected RAW distance matrix in epoch-length bins




Rarefaction curves: mean sum of ranges of Cailleuz-corrected GED distance matrix in epoch-length bins




Rarefaction curves: mean sum of ranges of Cailleuz-corrected GOW distance matrix in epoch-length bins




Rarefaction curves: mean sum of ranges of Cailleuz-corrected MAX distance matrix in epoch-length bins




Rarefaction curves: mean sum of ranges of uncorrected RAW distance matrix in 10 Ma bins




Rarefaction curves: mean sum of ranges of uncorrected GED distance matrix in 10 Ma bins




Rarefaction curves: mean sum of ranges of uncorrected GOW distance matrix in 10 Ma bins




Rarefaction curves: mean sum of ranges of uncorrected MAX distance matrix in 10 Ma bins




Rarefaction curves: mean sum of ranges of Cailleuz-corrected RAW distance matrix in 10 Ma bins




Rarefaction curves: mean sum of ranges of Cailleuz-corrected GED distance matrix in 10 Ma bins




Rarefaction curves: mean sum of ranges of Cailleuz-corrected GOW distance matrix in 10 Ma bins




Rarefaction curves: mean sum of ranges of Cailleuz-corrected MAX distance matrix in 10 Ma bins




Rarefaction curves: mean centroid distance of uncorrected RAW distance matrix in epoch-length bins




Rarefaction curves: mean centroid distance of uncorrected GED distance matrix in epoch-length bins




Rarefaction curves: mean centroid distance of uncorrected GOW distance matrix in epoch-length bins




Rarefaction curves: mean centroid distance of uncorrected MAX distance matrix in epoch-length bins




Rarefaction curves: mean centroid distance of Cailleuz-corrected RAW distance matrix in epoch-length bins




Rarefaction curves: mean centroid distance of Cailleuz-corrected GED distance matrix in epoch-length bins




Rarefaction curves: mean centroid distance of Cailleuz-corrected GOW distance matrix in epoch-length bins




Rarefaction curves: mean centroid distance of Cailleuz-corrected MAX distance matrix in epoch-length bins




Rarefaction curves: mean centroid distance of uncorrected RAW distance matrix in 10 Ma bins




Rarefaction curves: mean centroid distance of uncorrected GED distance matrix in 10 Ma bins




Rarefaction curves: mean centroid distance of uncorrected GOW distance matrix in 10 Ma bins




Rarefaction curves: mean centroid distance of uncorrected MAX distance matrix in 10 Ma bins




Rarefaction curves: mean centroid distance of Cailleuz-corrected RAW distance matrix in 10 Ma bins



Rarefaction curves: mean centroid distance of Cailleuz-corrected GED distance matrix in 10 Ma bins



Rarefaction curves: mean centroid distance of Cailleuz-corrected GOW distance matrix in 10 Ma bins



Rarefaction curves: mean centroid distance of Cailleuz-corrected MAX distance matrix in 10 Ma bins







Figure 4: (following pages) **Morphospace occupation of Ichthyosauriformes through the Mesozoic.** Principal coordinate axis 1 against axes 2 (top row) and 3 (bottom row) from each of eight PCA (four distance matrices: RAW, GED, GOW, MAX; with and without negative eigenvalue correction) on the cladistic matrix of Moon [3] binned into epochs.

PCo 1-3 scatter plots without negative eigenvalue correction



PCo 1-3 scatter plots with negative eigenvalue correction








Figure 6: (previous page) **Rates of skull size evolution in Ichthyosauriformes.** Evolutionary rate results from 100 Hedman-dated phylogenies. Branches are scaled and branches and taxon names coloured to the rate of skull size change on that branch.

Supplementary tables

1	Bin boundaries of 10 Ma bins used in this study	64
2	Occurrence dates of outgroup taxa used to date the tree of Ichthyosauriformes	64
3	Occurrence stratigraphy and dates of Ichthyosauriformes included in the analyses	65
4	Skull lengths of Ichthyosauriformes included in the analyses	70

Table 1: **Bin boundaries of 10 Ma bins used in this study.** Approximate age ranges are given as indicators.

Bin	Start (Ma)	End (Ma)	Approximate age range
1	251.3	241.3	Olenekian–Ladinian
2	241.3	231.3	Ladinian–Carnian
3	231.3	221.3	Carnian–Norian
4	221.3	211.3	Norian
5	211.3	201.3	Norian–Rhaetian
6	201.3	191.3	Hettangian–Sinemurian
7	191.3	181.3	Sinemurian–Toarcian
8	181.3	171.3	Toarcian–Aalenian
9	171.3	161.3	Aalenian–Oxfordian
10	161.3	151.3	Oxfordian–Tithonian
11	151.3	141.3	Tithonian–Berriasian
12	141.3	131.3	Berriasian–Hauterivian
13	131.3	121.3	Hauterivian–Aptian
14	121.3	111.3	Aptian–Albian
15	111.3	101.3	Albian
16	101.3	91.3	Albian–Turonian

Table 2: **Occurrence dates of outgroup taxa used to date the tree of Ichthyosauriformes.** Stratigraphic occurrence intervals are taken from the given references. Occurrences are converted to absolute ages using Gradstein *et al.* [15]. FAD, first appearance date; FAS, first appearance stratigraphy; LAD, last appearance date; LAS, last appearance stratigraphy.

Taxon	FAD (Ma)	LAD (Ma)	FAS	LAS	Reference
<i>Petrolacosaurus</i>	307.0	298.9	Upper Pennsylvanian	Upper Pennsylvanian	[16, 17]
<i>Hovasaurus</i>	259.8	251.2	Upper Permian	Induan	[18]
<i>Claudiosaurus</i>	253.2	252.17	Late Changsinghian	Late Changsinghian	[19]
<i>Thadeosaurus</i>	254.14	252.17	Changsinghian	Changsinghian	[19]
<i>Milleretta</i>	253.2	252.5	Changsinghian	Changsinghian	[20]
<i>Broomia</i>	265.1	260.5	Capitanian	Capitanian	[20]
<i>Mesosaurus</i>	290.1	286	Early Artinskian	Early Artinskian	[20]
<i>Captorhinus</i>	280	270.6	Leonardian	Leonardian	[21]

Table 3: Occurrence stratigraphy and dates of Ichthyosauriformes included in the analyses. Stratigraphical occurrences are given to the nearest ammonite or conodont biozone horizon where possible. Occurrences are converted to absolute ages using Gradstein *et al.* [15]. FAD, first appearance date; LAD, last appearance date.

Taxon	FAD stratigraphy	LAD stratigraphy	FAD (Ma)	LAD (Ma)	References
<i>Acamptonectes densus</i>	D2D horizon, Speeton Clay Formation, basal Hauterivian	<i>Sinikites concinnus/staffi</i> Biozone, upper Hauterivian, Lower Cretaceous	132.9	129.4	[22]
<i>Aegirosaurus leptospondylus</i>	Malm Zeta2b, early lower Tithonian, Upper Jurassic	Malm Zeta2b, early lower Tithonian, Upper Jurassic	153.96	149.87	[23]
<i>Arthropterygius chrisorum</i>	Oxfordian, Upper Jurassic	Kimmeridgian, Upper Jurassic	163.47	152.06	[24]
<i>Athabascasaurus bitumineus</i>	Lower Albian, Lower Cretaceous	Lower Albian, Lower Cretaceous	113	111.5	[25]
<i>Barracudasauroides paxaniensis</i>	Anisian, Middle Triassic	<i>Nicarella germanicus</i> Conodont Biozone, Anisian, Middle Triassic	244.94	243.99	[26]
<i>Besanosaurus leptorhynchus</i>	Nevadites Conodont Biozone, uppermost Anisian, Middle Triassic	Nevadites Conodont Biozone, uppermost Anisian, Middle Triassic	242.1	241.5	[27]
<i>Brachypterygius extremus</i>	<i>Pectinatites wheatleyensis</i> Ammonite Biozone, Thithonian, Upper Jurassic	<i>Pectinatites hudsoni</i> Ammonite Biozone, Thithonian, Upper Jurassic	151	150	[28]
<i>Brachypterygius pseudoscythica</i>	<i>Ilovaisya pseudoscythica</i> Ammonite Biozone, Thithonian, Upper Jurassic	<i>Ilovaisya pseudoscythica</i> Ammonite Biozone, Tithonian, Upper Jurassic	150.1	149.5	[29]
<i>Californosaurus perrini</i>	<i>Trachyceras</i> Beds, Hosselkus Limestone, Carnian	<i>Trachyceras</i> Beds, Hosselkus Limestone, Carnian	233.5	228.35	[30, 31]
<i>Callawayia neoscapularis</i>	<i>Epigondolella triangularis</i> Conodont Biozone, early Norian, Upper Triassic	<i>Epigondolella quadrata</i> Conodont Biozone, early Norian, Upper Triassic	221.5	217.5	[32]
<i>Cartorhynchus lenticarpus</i>	Subcolumnites Ammonite Biozone, Olenekian, Lower Triassic	Subcolumnites Ammonite Biozone, Olenekian, Lower Triassic	247.7	247.2	[33]
<i>Caypullisaurus bonapartei</i>	<i>Emileia giebelii</i> Ammonite Biozone, early Bajocian, Middle Jurassic	<i>Emileia giebelii</i> Ammonite Biozone, early Bajocian, Middle Jurassic	152.1	139.4	[34]
<i>Chacaicosaurus cayi</i>	<i>Procolumnites</i> Ammonite Biozone, Olenekian, Lower Triassic	<i>Subcolumnites</i> Ammonite Biozone, Olenekian, Lower Triassic	170.3	169.45	[35]
<i>Chaohusaurus chaoxianensis</i>	<i>Procolumnites</i> Ammonite Biozone, Olenekian, Lower Triassic	<i>Subcolumnites</i> Ammonite Biozone, Olenekian, Lower Triassic	247.9	247.2	[36, 37]
<i>Chaohusaurus geishanensis</i>	<i>Procolumnites</i> Ammonite Biozone, Olenekian, Lower Triassic	<i>Subcolumnites</i> Ammonite Biozone, Olenekian, Lower Triassic	247.9	247.2	[33, 37]
<i>Chaohusaurus zhangjiawanensis</i>	<i>Neospathodus homeri</i> Conodont Biozone, Spathian, Lower Triassic	<i>Neospathodus triangulus</i> Conodont Biozone, Spathian, Lower Triassic	247.9	247.2	[38]
<i>Contectopalatus atavus</i>	<i>Pelsonia</i> , Anisian, Middle Triassic	Illyrian, Anisian, Middle Triassic	244.94	241.5	[39]
<i>Cryptopterygius kielmae</i>	Middle Volgian, Lower Tithonian, Upper Jurassic	Middle Volgian, Lower Tithonian, Upper Jurassic	149	147	[40]
<i>Cryptopterygius kristiansenae</i>	<i>Dorsoplantis maximus</i> Ammonite biozone, Tithonian, Upper Jurassic	<i>Dorsoplantis ilovaiskyi</i> Ammonite biozone, Tithonian, Upper Jurassic	148.3	147.4	[41]
<i>Cymbospondylus buchseri</i>	Upper Anisian, Middle Triassic	lower Ladinian, Middle Triassic	242.57	240.3	[42]
<i>Cymbospondylus nicholisi</i>	<i>Kellnerites fesocevagensis</i> Ammonite Biozone, Anisian, Middle Triassic	<i>Kellnerites fesocevagensis</i> Ammonite Biozone, Anisian, Middle Triassic	243.99	243.05	[43]

Table 3 continued

Taxon	FAD stratigraphy	LAD stratigraphy	FAD (Ma)	LAD (Ma)	References
<i>Cymbospondylus petrinus</i>	<i>Paragoniodella ex gr. excelsa</i> Conodont Biozone, Anisian, Middle Triassic	<i>Paragoniodella ex gr. excelsa</i> Conodont Biozone, Anisian, Middle Triassic	243.99	241.5	[31]
<i>Cymbospondylus piscesmus</i>	<i>Pleydellia aalensis</i> Ammonite Biozone, Toarcian, Lower Jurassic	<i>Anisian, Middle Triassic</i>	247.2	241.5	[31]
<i>Dearcmhra schawcrossi</i>	<i>Dactylioceras tenuicostatum</i> Ammonite Biozone, Toarcian, Upper Jurassic	<i>Stephanoceras humphriesianum</i> Ammonite Biozone, Bajocian, Middle Jurassic	174.43	169.45	[44]
<i>Eurhinosaurus longirostris</i>	<i>Arietites bucklandii</i> Ammonite Biozone, lower Sinemurian, Lower Jurassic	<i>Harpoceras falciferum</i> Ammonite Biozone, Toarcian, Lower Sinemurian, Lower Jurassic	182.7	180.36	[45, 46]
<i>Excalibosaurus costini</i>	Calcareous facies Sacocoma Formation, Late Kimmeridgian, Upper Jurassic	<i>Arietites bucklandii</i> Ammonite Biozone, Middle Volgian, Lower Tithonian, Upper Jurassic	199.3	197.8	[46, 47]
<i>Gengasaurus nicosiae</i>	<i>Dorsoplanites panderi</i> Ammonite Biozone, Tithonian, Upper Jurassic	<i>Dorsoplanites panderi</i> Ammonite Biozone, Tithonian, Upper Jurassic	149.6	147.9	[49]
<i>Grendelius alekseevi</i>	Middle Volgian, Lower Tithonian, Upper Jurassic	Middle Volgian, Lower Tithonian, Upper Jurassic	149.6	147.9	[50]
<i>Grendelius zhuravlevi</i>	<i>Subcolumnites</i> Ammonite Biozone, Olenekian, Lower Triassic	<i>Subcolumnites</i> Ammonite Biozone, Olenekian, Lower Triassic	247.7	247.2	[51]
<i>Grippia longirostris</i>	Carnian, Upper Triassic	Carnian, Upper Triassic	233.5	228.35	[52]
<i>Guizhouichthyosaurus tangae</i>	Carnian, Upper Triassic	Carnian, Upper Triassic	233.5	228.35	[53]
<i>Guizhouichthyosaurus wologangense</i>	<i>Subcolumnites</i> Ammonite Biozone, Olenekian, Lower Triassic	<i>Subcolumnites</i> Ammonite Biozone, Olenekian, Lower Triassic	247.7	247.2	[54, 55]
<i>Guiosaurus helmi</i>	<i>Dactylioceras tenuicostatum</i> , Ammonite Biozone, Toarcian	<i>Harpoceras falciferum</i> Ammonite Biozone, Toarcian	182.7	181.25	[56–58]
<i>Hauffipteryx typicus</i>	Norian, Upper Triassic	Norian, Upper Triassic	228.4	209.5	[59]
<i>Himalayasaurus tibetensis</i>	<i>Epigondolella quadrata</i> Conodont Biozone, Norian, Upper Triassic	<i>Epigondolella quadrata</i> Conodont Biozone, Norian, Upper Triassic	226.5	221.25	[60]
<i>Hudsonelpidia brevirostris</i>	<i>Subcolumnites</i> Ammonite Biozone, Olenekian, Lower Triassic	<i>Subcolumnites</i> Ammonite Biozone, Olenekian, Lower Triassic	247.9	247.2	[61]
<i>Hupishuchus nanchangensis</i>	<i>Hildoceras bifrons</i> Ammonite Biozone, Toarcian, Lower Jurassic	<i>Dactylioceras communis</i> Ammonite Biozone, Toarcian, Lower Jurassic	180.36	175.6	[47]
<i>Ichthyosaurus acutirostris</i>	Asteroceras obtusum? Ammonite Biozone, Sinemurian, Lower Jurassic	<i>Uptonia jamesoni</i> Ammonite Biozone, Pliensbachian, Lower Jurassic	193.81	189.35	[62]
<i>Ichthyosaurus anningae</i>	<i>Schlotheimia angulata</i> Ammonite Biozone, Sinemurian, Lower Jurassic	<i>Artioceras semicostatum</i> Ammonite Biozone, Sinemurian, Lower Jurassic	200.1	196.31	[63]
<i>Ichthyosaurus breviceps</i>	Upper Rhaetian, Upper Triassic	<i>Artioceras semicostatum</i> Ammonite Biozone, Lower Jurassic	201.3	196.31	[63]
<i>Ichthyosaurus communis</i>	<i>Schlotheimia angulata</i> Ammonite Biozone, Hettangian	<i>Artioceras semicostatum</i> Ammonite Biozone, Sinemurian	200.1	196.31	[63]
<i>Ichthyosaurus conybeari</i>	Pre- <i>Planorbis</i> beds, Hettangian, Lower Jurassic	Pre- <i>Planorbis</i> beds, Hettangian, Lower Jurassic	201.3	200.85	[64]
<i>Ichthyosaurus larkini</i>	Pre- <i>Planorbis</i> beds, Hettangian, Lower Jurassic	Pre- <i>Planorbis</i> beds, Hettangian, Lower Jurassic	201.3	200.85	[64]
<i>Ichthyosaurus somersetensis</i>	<i>Subcolumnites</i> Ammonite Biozone, Olenekian, Lower Triassic	<i>Subcolumnites</i> Ammonite Biozone, Olenekian, Lower Triassic	247.7	247.2	[65]

Table 3 continued

Taxon	FAD stratigraphy	LAD stratigraphy	FAD (Ma)	LAD (Ma)	References
<i>Janusaurus lundi</i>	<i>Dorsoplantis maximus</i> Ammonite Biozone, Tithonian, Upper Jurassic	<i>Dorsoplantis ilovaiskyi</i> Ammonite Biozone, Tithonian, Upper Jurassic	148.3	147.4	[66]
<i>Keihauia nui</i>	Slottsmøya Member, Agardfjellet Formation, Berriasian, Lower Cretaceous	Slottsmøya Member, Agardfjellet Formation, Berriasian, Lower Cretaceous	145	143	[67]
<i>Leniniastellans</i>	<i>Deshayesites volgensis</i> Ammonite Biozone, Lower Aptian, Lower Cretaceous	<i>Deshayesites volgensis</i> Ammonite Biozone, Lower Aptian, Lower Cretaceous	126.3	123	[68]
<i>Leptonectes moorei</i>	Lower Pliensbachian, Lower Jurassic	Lower Pliensbachian, Lower Jurassic	190.82	187.56	[69]
<i>Leptonectes solei</i>	<i>Arnioceras semicostatum</i> Ammonite Biozone, Sinemurian, Lower Jurassic	<i>Asteroceras obustum</i> Ammonite Biozone, Sinemurian, Lower Jurassic	195.31	193.81	[70]
<i>Leptonectes tenuirostris</i>	Pre-Planorbis beds, Hettangian, Lower Jurassic	<i>Amaltheus marginatus</i> Ammonite Biozone, Pliensbachian, Lower Jurassic	201.3	190.8	[47, 71]
<i>Macgowania janiceps</i>	<i>Epigondolella matthewi</i> Conodont Biozone, middle Norian, Upper Triassic	<i>Epigondolella multidentata</i> Conodont Biozone, middle Norian, Upper Triassic	220	216.9	[72]
<i>Maiaspondylus lindoei</i>	Middle Albian, Lower Cretaceous	Middle Albian, Lower Cretaceous	111.27	110.22	[73]
<i>Malawania anachronus</i>	Late Hauterivian, Early Cretaceous	Barremian, Early Cretaceous	131	125	[74]
<i>Mikadocephalus gracilirostris</i>	Upper Illyrian, Anisian, Middle Triassic	lower Fassiniian, Ladinian, Middle Triassic	242.57	240.3	[75]
<i>Micosaurus cornalianus</i>	Upper Illyrian, Anisian, Middle Triassic	lower Fassiniian, Ladinian, Middle Triassic	242.57	240.3	[47, 76]
<i>Micosaurus kuhnschneidleri</i>	Upper Illyrian, Anisian, Middle Triassic	lower Fassiniian, Ladinian, Middle Triassic	242.57	240.3	[77]
<i>Micosaurus sindianensis</i>	<i>Nicorrella kockeli</i> Conodont Biozone, Pelsonian, Anisian, Middle Triassic	<i>Nicorrella kockeli</i> Conodont Biozone, Pelsonian, Anisian, Middle Triassic	244.94	241.5	[78]
<i>Mollesaurus perialis</i>	<i>Emilia giebelii</i> Ammonite Biozone, early Bajocian, Middle Jurassic	<i>Emilia giebelii</i> Ammonite Biozone, early Bajocian, Middle Jurassic	170.3	169.45	[79, 80]
<i>Muiscasaurus catheti</i>	Barremian, Lower Cretaceous	Apitan, Lower Cretaceous	130.77	115.64	[81]
<i>Nanopterygius entekiodon</i>	<i>Aulacostephanus</i> sp. Ammonite Biozone, Kimmeridgean, Upper Jurassic	<i>Aulacostephanus</i> sp. Ammonite Biozone, Kimmeridgean, Upper Jurassic	154.6	149.87	[28]
<i>Ophthalmosaurus icenicus</i>	<i>Kosmoceras jasoni</i> Ammonite Biozone, Callovian, Upper Jurassic	<i>Quenstedtoceras mariae</i> Ammonite Biozone, Oxfordian, Upper Jurassic	165.59	161.39	[82]
<i>Ophthalmosaurus natans</i>	Oxfordian, Late Jurassic	Oxfordian, Late Jurassic	163.5	157.3	[83]
<i>Ophthalmosaurus yasykovi</i>	<i>Epivirgatites nikitini</i> Ammonite Biozone, Tithonian, Upper Jurassic	<i>Craspedites subtilis</i> Ammonite Biozone, Tithonian, Upper Jurassic	147.5	146.4	[84]
<i>Palvennia hoybergeti</i>	<i>Dorsoplantis maximus</i> Ammonite Biozone, Tithonian, Upper Jurassic	<i>Dorsoplantis ilovaiskyi</i> Ammonite Biozone, Tithonian, Upper Jurassic	148.3	147.4	[41]
<i>Paraphthalmosaurus kabanoi</i>	<i>Epivirgatites nikitini</i> Ammonite Biozone, Tithonian, Upper Jurassic	<i>Epivirgatites nikitini</i> Ammonite Biozone, Tithonian, Upper Jurassic	147.5	146.9	[85]
<i>Paravinator wapitiensis</i>	<i>Dorsoplantis panderi</i> Ammonite Biozone, Tithonian, Upper Jurassic	<i>Epivirgatites nikitini</i> Ammonite Biozone, Tithonian, Upper Jurassic	149.6	146.9	[86]
<i>Pervushovisaurus bannovkensis</i>	<i>Subcolumbites</i> Ammonite Biozone, Olenekian, Lower Triassic	<i>Subcolumbites</i> Ammonite Biozone, Olenekian, Lower Triassic	247.7	247.2	[87]
<i>Pervushovisaurus campylodon</i>	Middle Cenomanian, Upper Cretaceous	Middle Cenomanian, Upper Cretaceous	96.24	95.47	[88]
<i>Pessopteryx nisseri</i>	Early Cenomanian, Upper Cretaceous	Early Cenomanian, Upper Cretaceous	100.45	96.5	[89]
	<i>Subcolumbites</i> Ammonite Biozone, Olenekian, Lower Triassic	<i>Subcolumbites</i> Ammonite Biozone, Olenekian, Lower Triassic	247.7	247.2	[65]

Table 3 continued

Taxon	FAD stratigraphy	LAD stratigraphy	FAD (Ma)	LAD (Ma)	References
<i>Phalarodon callawayi</i>	Upper Anisian, Middle Triassic	Upper Anisian, Middle Triassic	243.99	239.1	[90]
<i>Phalarodon fraxsi</i>	Upper Anisian, Middle Triassic	lower Ladinian, Middle Triassic	244.94	237	[91]
<i>Phalarodon major</i>	Upper Anisian, Middle Triassic	Ladinian, Middle Triassic	244.94	237	[92]
<i>Phantonosaurus neubigi</i>	<i>pulcher/robustus</i> Conodont Biozone, Upper Anisian, Middle Triassic	<i>pulcher/robustus</i> Conodont Biozone, Upper Anisian, Middle Triassic	244.94	241.5	[93]
<i>Platypterygius americanus</i>	Upper Albian, Lower Cretaceous	Upper Albian, Lower Cretaceous	107.59	100.5	[94]
<i>Platypterygius australis</i>	Albian, Lower Cretaceous	Albian, Lower Cretaceous	113	100.5	[95, 96]
<i>Platypterygius hauthali</i>	<i>Hatchiceras</i> Ammonite Biozone, Lower Barremian, Lower Cretaceous	<i>Hatchiceras</i> Ammonite Biozone, Lower Barremian, Lower Cretaceous	130.77	129.41	[97]
<i>Platypterygius hiercynicus</i>	Apitan, Lower Cretaceous	Apitan, Lower Cretaceous	125	113	[98]
<i>Platypterygius platypterygius</i>	<i>Hoplites deshayesi</i> Ammonite Biozone, Apitan, Lower Cretaceous	<i>Hoplites deshayesi</i> Ammonite Biozone, Apitan, Lower Cretaceous	125	113	[99]
<i>Platypterygius sachicarum</i>	Barremian, Lower Cretaceous	Apitan, Lower Cretaceous	130.77	113	[100]
<i>Qianichthyosaurus xingyiensis</i>	Ladinian, Middle Triassic	Ladinian, Middle Triassic	241.5	237	[101]
<i>Qianichthyosaurus zhoui</i>	Carnian, Upper Triassic	Carnian, Upper Triassic	237	228.35	[102]
<i>Quasianosteosaurus vikinghoegdai</i>	<i>Subcolumnites</i> Ammonite Biozone, Olenekian, Lower Triassic	<i>Subcolumnites</i> Ammonite Biozone, Olenekian, Lower Triassic	247.7	247.2	[103]
<i>Sclerocormus parviceps</i>	<i>Subcolumnites</i> Ammonite Biozone, Olenekian, Lower Triassic	<i>Subcolumnites</i> Ammonite Biozone, Olenekian, Lower Triassic	247.9	247.7	[104, 105]
<i>Shastasaurus liangae</i>	Upper Carnian, Late Triassic	Upper Carnian, Late Triassic	233.5	228.35	[106]
<i>Shastasaurus pacificus</i>	Upper Carnian, Late Triassic	Upper Carnian, Late Triassic	233.5	228.35	[31]
<i>Shastasaurus sikkaniensis</i>	<i>Epigondolella postera</i> conodont Biozone, <i>Mesohemnavattites columbianus</i> ammonite Biozone, middle Norian, Upper Triassic	<i>Epigondolella postera</i> conodont Biozone, <i>Mesohemnavattites columbianus</i> ammonite Biozone, middle Norian, Upper Triassic	216.9	214.7	[107]
<i>Shonisaurus popularis</i>	Upper Carnian, Late Triassic	Upper Carnian, Late Triassic	233.5	228.35	[108]
<i>Simbirskiaaurus birjukovi</i>	<i>Praeoxyteuthis pugio</i> Belemnite Biozone, lower Barremian, Lower Cretaceous	<i>Praeoxyteuthis pugio</i> Belemnite Biozone, lower Barremian, Lower Cretaceous	130.77	129.41	[88]
<i>Sisteronita seeleyi</i>	Cambridge Greensand Member, early Cenomanian, Upper Cretaceous	Cambridge Greensand Member, early Cenomanian, Upper Cretaceous	100.5	96.24	[109]
<i>Stenopterygius aaleniensis</i>	<i>Leioceras opalinum</i> Ammonite Biozone, Aalenian, Middle Jurassic	<i>Leioceras opalinum</i> Ammonite Biozone, Aalenian, Middle Jurassic	174.1	172.13	[110]
<i>Stenopterygius quadrisissimus</i>	<i>Dactylioceras tenuicostatum</i> Ammonite Biozone, Toarcian, Lower Jurassic	<i>Dactylioceras tenuicostatum</i> Ammonite Biozone, Toarcian, Lower Jurassic	182.7	180.36	[56, 111]
<i>Stenopterygius triscissus</i>	<i>Harpoceras falciferum</i> Ammonite Biozone, Toarcian, Lower Jurassic	<i>Harpoceras falciferum</i> Ammonite Biozone, Toarcian, Lower Jurassic	182.7	180.36	[56, 111]
<i>Stenopterygius uniter</i>	<i>Dactylioceras tenuicostatum</i> Ammonite Biozone, Toarcian, Lower Jurassic	<i>Dactylioceras tenuicostatum</i> Ammonite Biozone, Toarcian, Lower Jurassic	181.7	180.36	[56, 111]
<i>Suevoleviathan disintegr</i>	<i>Dactylioceras tenuicostatum</i> Ammonite Biozone, Toarcian, Lower Jurassic	<i>Dactylioceras tenuicostatum</i> Ammonite Biozone, Toarcian, Lower Jurassic	181.7	180.36	[112, 113]
<i>Suevoleviathan integer</i>	<i>Dactylioceras tenuicostatum</i> Ammonite Biozone, Toarcian, Lower Jurassic	<i>Dactylioceras tenuicostatum</i> Ammonite Biozone, Toarcian, Lower Jurassic	181.7	180.36	[112, 113]
<i>Sveltonectes insolitus</i>	Upper Barremian, Cretaceous	Upper Barremian, Cretaceous	129.6	126.3	[114]

Table 3 continued

Taxon	FAD stratigraphy	LAD stratigraphy	FAD (Ma)	LAD (Ma)	References
<i>Tennodontosaurus azerguensis</i>	<i>Harpoceras bifrons</i> Ammonite Biozone, middle Toarcian, Lower Jurassic	<i>Harpoceras bifrons</i> Ammonite Biozone, middle Toarcian, Lower Jurassic	180.36	178.24	[115]
<i>Tennodontosaurus crassimanus</i>	<i>Dactylioceras tenuicostatum</i> Ammonite Biozone, Toarcian, Lower Jurassic	<i>Harpoceras falciferum</i> Ammonite Biozone, Toarcian, Lower Jurassic	182.7	180.36	[116]
<i>Tennodontosaurus eurycephalus</i>	<i>Arietites bucklandi</i> Ammonite Biozone, lower Sinemurian, Lower Jurassic	<i>Arietites bucklandi</i> Ammonite Biozone, lower Sinemurian, Lower Jurassic	199.3	197.8	[117]
<i>Tennodontosaurus nuertingensis</i>	<i>Uptonia jamesoni</i> Ammonite Biozone, Lower Pliensbachian, Lower Jurassic	<i>Tragophyllocera ibex</i> Ammonite Biozone, Pliensbachian, Lower Jurassic	190.8	187.56	[118, 119]
<i>Tennodontosaurus platyodon</i>	<i>Schlotheimia angulata</i> Ammonite Biozone, Hettangian	<i>Amioceras semicostatum</i> Ammonite Biozone, Sinemurian	200.1	196.31	[117]
<i>Tennodontosaurus trigonodon</i>	<i>Dactylioceras tenuicostatum</i> Ammonite Biozone, Toarcian, Lower Jurassic	<i>Harpoceras falciferum</i> Ammonite Biozone, Toarcian, Lower Jurassic	182.7	180.36	[112, 120]
<i>Thaisaurus chonglakmanii</i>	Lower Triassic	Lower Triassic	250.01	247.2	[121]
<i>Thalattoarchon saurophagus</i>	Nevadiscitulus <i>taylori</i> Ammonite Biozone, Anisian, Middle Triassic	<i>Nevadiscitulus taylori</i> Ammonite Biozone, Anisian, Middle Triassic	247.2	244.6	[122]
<i>Tholodus schmidi</i>	<i>Decurviella decurviella</i> Conodont Biozone, Ladinian (Pelsonian), Middle Triassic	<i>Judicarites/Neoschizodus orbicularis</i> Conodont Biozone, Ladinian (Pelsonian), Middle Triassic	241	237	[123]
<i>Tretocnemus californicus</i>	<i>Metapolygnathus polygnathiformis</i> Conodont Biozone, Carnian, Upper Triassic	<i>Metapolygnathus polygnathiformis</i> Conodont Biozone, Carnian, Upper Triassic	233.5	228.35	[47, 124]
<i>Tretocnemus zittelii</i>	<i>Metapolygnathus polygnathiformis</i> Conodont Biozone, Carnian, Upper Triassic	<i>Metapolygnathus polygnathiformis</i> Conodont Biozone, Carnian, Upper Triassic	233.5	228.35	[47, 124]
<i>Undorosaurus gorodischensis</i>	<i>Epivirgatites nikitinii</i> , Virgatites virgatus Ammonite Biozone, Tithonian, Upper Jurassic	<i>Epivirgatites nikitinii</i> , Virgatites virgatus Ammonite Biozone, Tithonian, Upper Jurassic	147.9	146.9	[85]
<i>Undorosaurus trautscholdi</i>	<i>Epivirgatites nikitinii</i> Ammonite Biozone, Tithonian, Upper Jurassic	<i>Kachipurites fulgens</i> Ammonite Biozone, Tithonian, Upper Jurassic	150	144.1	[125]
<i>Utatsusaurus hataii</i>	<i>Subcolumbites</i> Ammonite Biozone, Olenekian, Lower Triassic	<i>Arnatoceratites</i> Ammonite Biozone, Olenekian, Lower Triassic	247.7	247.2	[126]
<i>Wahlisaurus massarae</i>	Pre-Planorbis beds, Hettangian, Lower Jurassic	Pre-Planorbis beds, Hettangian, Lower Jurassic	201.3	200.85	[127]
<i>Wimanius odontopalatus</i>	Anisian, Middle Triassic	Ladinian, Middle Triassic	241.5	237	[128]
<i>Xiaminiosaurus catactes</i>	<i>Nicarella kockeli</i> Conodont Biozone, Pelsonian, Anisian, Middle Triassic	<i>Nicarella kockeli</i> Conodont Biozone, Pelsonian, Anisian, Middle Triassic	244.94	241.5	[129]

Table 4: **Skull lengths of Ichthyosauriformes included in the analyses.**
Logarithm values are shown to 3 d.p.

Taxon	Skull length (mm)	\log_{10} (Skull length (mm))	References
<i>Acamptonectes densus</i>	1000	3.000	[22]
<i>Aegirosaurus leptospondylus</i>	570	2.756	[23]
<i>Barracudasauroides paxianensis</i>	212	2.326	[26]
<i>Besanosaurus leptorhynchus</i>	510	2.708	[27]
<i>Brachypterygius extremus</i>	1155	3.062	[28]
<i>Callawayia neoscapularis</i>	453	2.656	[93, 130]
<i>Cartorhynchus lenticarpus</i>	55	1.740	[33]
<i>Caypullisaurus bonapartei</i>	1300	3.114	[34]
<i>Chacaicosaurus cayi</i>	980	2.991	[35]
<i>Chaohusaurus geishanensis</i>	117	2.068	[37]
<i>Contectopalatus atavus</i>	130	2.114	[39, 131]
<i>Cryptopterygius kristiansenae</i>	1220	3.086	[33, 37]
<i>Cymbospondylus petrinus</i>	1166	3.067	[31]
<i>Eurhinosaurus longirostris</i>	1250	3.097	[45, 46]
<i>Excalibosaurus costini</i>	1540	3.188	[46, 47]
<i>Guizhouichthysaurus tangae</i>	800	2.903	[52]
<i>Guizhouichthysaurus wolonggangense</i>	645	2.810	[53]
<i>Gulosaurus helmi</i>	87	1.940	[54, 55]
<i>Hauffiopteryx typicus</i>	380	2.580	[56–58]
<i>Hudsonelpidia brevirostris</i>	131	2.117	[60]
<i>Hupehsuchus nanchangensis</i>	126	2.100	[61]
<i>Ichthyosaurus anningae</i>	390	2.591	[62]
<i>Ichthyosaurus breviceps</i>	240	2.380	[63]
<i>Ichthyosaurus communis</i>	256	2.408	[63]
<i>Ichthyosaurus conybeari</i>	216	2.334	[63]
<i>Ichthyosaurus larkini</i>	355	2.550	[64]
<i>Ichthyosaurus somersetensis</i>	438	2.641	[64]
<i>Leptonectes moorei</i>	328	2.516	[66]
<i>Leptonectes solei</i>	1585	3.200	[70]
<i>Leptonectes tenuirostris</i>	523	2.719	[47, 71]
<i>Macgowania janiceps</i>	505	2.703	[60]
<i>Mixosaurus cornalianus</i>	195	2.290	[47, 76]
<i>Mixosaurus kuhnschneyderi</i>	160	2.204	[132]
<i>Mixosaurus xindianensis</i>	223	2.348	[78]
<i>Nannopterygius enthekiodon</i>	600	2.778	[28]
<i>Ophthalmosaurus icenicus</i>	965	2.985	[82]
<i>Ophthalmosaurus natans</i>	1082	3.034	[83]
<i>Palvennia hoybergeti</i>	860	2.934	[41]
<i>Parvinatator wapitiensis</i>	120	2.079	[87]
<i>Phalarodon callawayi</i>	300	2.477	[90]
<i>Phalarodon fraasi</i>	205	2.312	[133, 134]
<i>Phantomosaurus neubigi</i>	550	2.740	[93]
<i>Platypterygius americanus</i>	1250	3.097	[94]
<i>Platypterygius australis</i>	1430	3.155	[95, 96]
<i>Platypterygius hercynicus</i>	1040	3.017	[98]
<i>Platypterygius platydactylus</i>	1170	3.068	[99]
<i>Platypterygius sachicarum</i>	870	2.940	[100]
<i>Qianichthysaurus xingyiensis</i>	270	2.431	[101]

Table 4 continued

Taxon	Skull length (mm)	\log_{10} (Skull length (mm))	References
<i>Qianichthyosaurus zhoui</i>	240	2.380	[102]
<i>Sclerocormus parviceps</i>	100	2.000	[104]
<i>Shastasaurus liangae</i>	750	2.875	[106]
<i>Shastasaurus sikkaniensis</i>	3000	3.477	[107]
<i>Shonisaurus popularis</i>	2750	3.439	[108]
<i>Stenopterygius quadriscissus</i>	625	2.796	[56, 111]
<i>Stenopterygius triscissus</i>	634	2.802	[56, 111]
<i>Stenopterygius uniter</i>	537	2.730	[56, 111]
<i>Suevoleiathan disinteger</i>	860	2.934	[112, 113]
<i>Suevoleiathan integer</i>	690	2.839	[112, 113]
<i>Sveltonectes insolitus</i>	570	2.756	[135]
<i>Temnodontosaurus azerguensis</i>	1700	3.230	[115]
<i>Temnodontosaurus eurycephalus</i>	1020	3.009	[117]
<i>Temnodontosaurus platyodon</i>	1790	3.253	[117]
<i>Temnodontosaurus trigonodon</i>	1090	3.037	[112, 120]
<i>Thalattoarchon saurophagus</i>	1200	3.079	[122]
<i>Utatsusaurus hataii</i>	215	2.332	[126]
<i>Wimanius odontopalatus</i>	250	2.398	[128, 136]
<i>Xinminosaurus catactes</i>	290	2.462	[129]

Supplemental code

Code s1 R code implementing the disparity, principal coordinates, diversity, and discrete character rates analyses. This set of five scripts contains the code used to run the main discrete character analyses in R. Outputs include time-scaled trees, discrete rates of evolution, stratigraphic congruence values; PDF files of all figures produced; CSV files of root ages from the time-scaled trees, stratigraphic congruence tests, and statistical tests (pairwise PERMANOVA between epochs for PCA data and pairwise t -tests of per-bin disparity).

Code s2 Continuous rates analyses in BayesTraits and plotting in R. Rates analyses were run individually on 100 time-scaled trees then combined into consensus trees with branch rates averaged across all runs. Also includes code to create the traitgram of Fig. 4.

References

1. Bapst, D. W. Paleotree: An R Package for Paleontological and Phylogenetic Analysis of Evolution. *Methods in Ecology and Evolution* **3**, 803–807 (2012).
2. Laurin, M. The Evolution of Body Size, Cope’s rule and the origin of amniotes. *Systematic Biology* **53**, 594–622 (2004).
3. Moon, B. C. A New Phylogeny of Ichthyosaurs (Reptilia: Diapsida). *Journal of Systematic Palaeontology* **17**, 129–155 (2017).
4. R Core Team. *R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria* <http://www.R-project.org/> (2014).

5. Lloyd, G. T. Estimating Morphological Diversity and Tempo with Discrete Character-taxon Matrices: Implementation, Challenges, Progress, and Future Directions. *Biological Journal of the Linnean Society* **118**, 131–151 (2016).
6. Close, R. A., Friedman, M., Lloyd, G. T. & Benson, R. B. J. Evidence for a Mid-Jurassic Adaptive Radiation in Mammals. *Current Biology* **25**, 2137–2142 (2015).
7. Wills, M. A., Briggs, D. E. G. & Fortey, R. A. Disparity as an Evolutionary Index: a Comparison of Cambrian and Recent Arthropods. *Paleobiology* **20**, 93–130 (1994).
8. Gower, J. C. A General Coefficient of Similarity and Some of Its Properties. *Biometrics* **24**, 774–786 (1971).
9. Flannery Sutherland, J. T., Moon, B. C., Stubbs, T. L. & Benton, M. J. Does Exception Preservation Distort Our View of Disparity in the Fossil Record? *Proceedings of the Royal Society B: Biological Sciences* **286**, 20190091 (2019).
10. Lehmann, O. E. R., Ezcurra, M. D., Butler, R. J. & Lloyd, G. T. Biases with the Generalized Euclidean Distance Measure in Disparity Analyses with High Levels of Missing Data. *Palaeontology* (2019).
11. Caillez, F. The Analytical Solution of the Additive Constant Problem. *Psychometrika* **48**, 305–308 (1983).
12. Wills, M. A. Crustacean Disparity through the Phanerozoic: Comparing Morphological and Stratigraphic Data. *Biological Journal of the Linnean Society* **65**, 455–500 (1998).
13. Thorne, P. M., Ruta, M. & Benton, M. J. Resetting the Evolution of Marine Reptiles at the Triassic–Jurassic Boundary. *Proceedings of the National Academy of Sciences* **108**, 8339–8344 (2011).
14. Guillerme, T. dispRity: A Modular R Package for Measuring Disparity. *Methods in Ecology and Evolution*, 1755–1763 (2018).
15. Gradstein, F. M., Ogg, J. C., Schmitz, M. D. & Ogg, G. M. *The Geologic Time Scale 2012* 1143 (Elsevier, Oxford, 2012).
16. Reisz, R. R. A Diapsid Reptile from the Pennsylvanian of Kansas. *Special Publication of the Museum of Natural History, University of Kansas* **7**, 74 pp (1981).
17. Falcon-Lang, H. J., Benton, M. J. & Stimson, M. Ecology of the Earliest Reptiles Inferred from Basal Pennsylvanian Trackways. *Journal of the Geological Society, London* **164**, 1113–1118 (2007).
18. Ketchum, H. F. & Barrett, P. M. New Reptile Material from the Lower Triassic of Madagascar: Implications for the Permian–Triassic Extinction Event. *Canadian Journal of Earth Sciences* **41**, 1–8 (2004).
19. Reisz, R. R., Modesto, S. P. & Scott, D. M. A New Early Permian Reptile and Its Significance in Early Diapsid Evolution. *Proceedings of the Royal Society of London B: Biological Sciences* **278**, 3731–3737 (2011).
20. Ruta, M., Cisneros, J. C., Liebrecht, T., Tsuji, L. A. & Müller, J. Amniotes through Major Biological Crises: Faunal Turnover among Parareptiles and the End-Permian Mass Extinction. *Palaeontology* **54**, 1117–1137 (2011).
21. Kissel, R. A., Dilkes, D. W. & Reisz, R. R. *Captorhinus magnus*, a New Captorhinid (Amniota: Eureptilia) from the Lower Permian of Oklahoma, with New Evidence on the Homology of the Astragalus. *Canadian Journal of Earth Sciences* **39**, 1363–1372 (2002).
22. Fischer, V. et al. New Ophthalmosaurid Ichthyosaurs from the European Lower Cretaceous Demonstrate Extensive Ichthyosaur Survival across the Jurassic–Cretaceous Boundary. *PLoS ONE* **7**, e29234 (2012).

23. Bardet, N. & Fernández, M. S. A New Ichthyosaur from the Upper Jurassic Lithographic Limestones of Bavaria. *Journal of Paleontology* **74**, 503–511 (2000).
24. Maxwell, E. E. Generic Reassignment of an Ichthyosaur from the Queen Elizabeth Islands, Northwest Territories, Canada. *Journal of Vertebrate Paleontology* **30**, 403–415 (2010).
25. Druckenmiller, P. S. & Maxwell, E. E. A New Lower Cretaceous (lower Albian) Ichthyosaur Genus from the Clearwater Formation, Alberta, Canada. *Canadian Journal of Earth Sciences* **47**, 1037–1053 (2010).
26. Jiang, D.-Y., Schmitz, L., Hao, W. & Sun, Y.-L. A New Mixosaurid Ichthyosaur from the Middle Triassic of China. *Journal of Vertebrate Paleontology* **26**, 60–69 (2006).
27. Dal Sasso, C. & Pinna, G. *Besanosaurus leptorhynchus* N. Gen. N. Sp., a New Shastasaurid Ichthyosaur from the Middle Triassic of Besano (Lombardy, N. Italy). *Paleontologia Lombarda* **4**, 1–22 (1996).
28. Moon, B. C. & Kirton, A. M. Ichthyosaurs of the British Middle and Upper Jurassic. Part 2, *Brachypterygius*, *Nanopterygius*, *Macropterygius*, and *Taxa Invalida*. *Monograph of the Palaeontographical Society* **172**, 85–176 (2018).
29. Efimov, V. M. The Ichthyosaur *Otschevia pseudoscythica* Gen. Et Sp. Nov. from the Upper Jurassic of Ulyanovsk Volga. *Paleontologicheskii Zhurnal* **1998**, 82–86 (1998).
30. Merriam, J. C. Triassic Ichthyopterygia from California and Nevada. *University of California, Bulletin of the Department of Geology* **3**, 63–108 (1902).
31. Merriam, J. C. Triassic Ichthyosauria, with Special Reference to the American Forms. *Memoirs of the University of California* **1**, 1–252 (1908).
32. McGowan, C. A New Species of *Shastasaurus* (Reptilia: Ichthyosauria) from the Triassic of British Columbia: The Most Complete Exemplar of the Genus. *Journal of Vertebrate Paleontology* **14**, 168–179 (1994).
33. Motani, R. et al. A Basal Ichthyosauriform with a Short Snout from the Lower Triassic of China. *Nature* **517**, 485–488 (2015).
34. Fernández, M. S. A New Ichthyosaur from the Tithonian (Late Jurassic) of the Neuquén Basin, Northwestern Patagonia, Argentina. *Journal of Paleontology* **71**, 479–484 (1997).
35. Fernández, M. S. A New Long-snouted Ichthyosaur from the Early Bajocian of Neuquén Basin (Argentina). *Ameghiniana* **31**, 291–297 (1994).
36. Motani, R., Jiang, D.-Y., Tintori, A., Rieppel, O. C. & Chen, G. Terrestrial Origin of Viviparity in Mesozoic Marine Reptiles Indicated by Early Triassic Embryonic Fossils. *PLoS ONE* **9**, e88640 (2014).
37. Motani, R. et al. Status of *Chaohusaurus chaoxianensis* (chen, 1985). *Journal of Vertebrate Paleontology*, e892011 (2015).
38. Chen, X., Sander, P. M., Cheng, L. & Wang, X. A New Triassic Primitive Ichthyosaur from Yuanan, South China. *Acta Geologica Sinica* **87**, 672–677 (2013).
39. Liu, J. et al. The First Specimen of the Middle Triassic Ichthyosaur *Phalarodon atavus* (Ichthyosauria: Mixosauridae) from South China, Showing Postcranial Anatomy and Peri-tethyan Distribution. *Palaeontology* **56**, 849–866 (2013).
40. Tyborowski, D. A New Ophthalmosaurid Ichthyosaur from the Late Jurassic of Owadów-Brzezinki Quarry, Poland. *Acta Palaeontologica Polonica* **61**, 791–803 (2016).

41. Druckenmiller, P. S., Hurum, J. H., Knutsen, E. M. & Nakrem, H. A. Two New Ophthalmosaurids (Reptilia: Ichthyosauria) from the Agardhfjellet Formation (Upper Jurassic: Volgian/Tithonian), Svalbard, Norway. *Norwegian Journal of Geology* **92**, 311–339 (2012).
42. Sander, P. M. The Large Ichthyosaur *Cymbospondylus buchseri*, Sp. Nov., from the Middle Triassic of Monte San Giorgio (switzerland), with a Survey of the Genus in Europe. *Journal of Vertebrate Paleontology* **9**, 163–173 (1989).
43. Fröbisch, N. B., Sander, P. M. & Rieppel, O. C. A New Species of *Cymbospondylus* (Diapsida, Ichthyosauria) from the Middle Triassic of Nevada and a Re-evaluation of the Skull Osteology of the Genus. *Zoological Journal of the Linnean Society* **147**, 515–538 (2006).
44. Brusatte, S. L. *et al.* Ichthyosaurs from the Jurassic of Skye, Scotland. *Scottish Journal of Geology* **51**, 43–55 (2015).
45. Von Huene, F. F. Ein Neuer Fund Von *Eurhinosaurus longirostris*. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* **93**, 277–284 (1951).
46. McGowan, C. A Putative Ancestor for the Swordfish-like Ichthyosaur *Eurhinosaurus*. *Nature* **322**, 454–456 (1986).
47. McGowan, C. & Motani, R. in *Handbook of Paleoherpetology* (ed Sues, H.-D.) (Verlag Dr. Friedrich Pfeil, Munich, 2003).
48. Paparella, I., Maxwell, E. E., Cipriani, A., Roncaci & Caldwell, M. W. The First Ophthalmosaurid Ichthyosaur from the Upper Jurassic of the Umbrian–Marchean Apennines (Marche, Central Italy). *Geological Magazine* **154**, 837–858 (2016).
49. Arkhangelsky, M. S. On a New Ichthyosaur of the Genus *Otschekia* from the Volgian Stage of the Volga Region near Ulyanovsk. *Paleontological Journal* **35**, 629–635 (2001).
50. Zverkov, N. G., Arkhangelsky, M. S. & Stenshin, I. M. A Review of Russian Upper Jurassic Ichthyosaurs with an Intermedium/humeral Contact. Reassessing *Grendelius* McGowan, 1976. *Proceedings of the Zoological Institute RAS* **319**, 558–588 (2015).
51. Motani, R. Skull of *Grippia longirostris*: No Contradiction with a Diapsid Affinity for the Ichthyopterygia. *Palaeontology* **43**, 1–14 (2000).
52. Pan, X.-R. *et al.* Discussion on *Guizhouichthyosaurus tangae* Cao and Luo in Yin Et Al., 2000 (Reptilia, Ichthyosauria) from the Late Triassic of Guanling County, Guizhou. *Acta Scientiarum Naturalium Universitatis Pekinensis* **42**, 697–703 (2006).
53. Chen, X., Cheng, L. & Sander, P. M. A New Species of *Callawayia* (Reptilia: Ichthyosauria) from the Late Triassic in Guanling, Guizhou. *Geology in China* **34**, 974–982 (2007).
54. Brinkman, D. B., Xijin, Z. & Nicholls, E. L. A Primitive Ichthyosaur from the Lower Triassic of British Columbia, Canada. *Palaeontology* **35**, 465–474 (1992).
55. Cuthbertson, R. S., Russell, A. P. & Anderson, J. S. Cranial Morphology and Relationships of a New Grippidian (Ichthyopterygia) from the Vega-Phroso Siltstone Member (Lower Triassic) of British Columbia, Canada. *Journal of Vertebrate Paleontology* **33**, 831–847 (2013).
56. Maisch, M. W. Revision Der Gattung *Stenopterygius* Jaekel, 1904 Emend. Von Huene, 1922 (Reptilia: Ichthyosauria) Aus Dem Unteren Jura Westeuropas. *Palaeodiversity* **1**, 227–271 (2008).
57. Caine, H. & Benton, M. J. Ichthyosauria from the Upper Lias of Strawberry Bank, England. *Palaeontology* **54**, 1069–1093 (2011).
58. Marek, R. D., Moon, B. C., Williams, M. & Benton, M. J. The Skull and Endocranum of a Lower Jurassic Ichthyosaur Based on Digital Reconstructions. *Palaeontology* **58**, 723–742 (2015).

59. Motani, R., Manabe, M. & Dong, Z. M. The Status of *Himalayasaurus tibetensis* (Ichthyopterygia). *Paludicola* **2**, 174–181 (1999).
60. McGowan, C. A Remarkable Small Ichthyosaur from the Upper Triassic of British Columbia, Representing a New Genus and Species. *Canadian Journal of Earth Sciences* **32**, 292–303 (1995).
61. Carroll, R. L. & Dong, Z. M. *Hupehsuchus*, an Enigmatic Aquatic Reptile from the Triassic of China, and the Problem of Establishing Relationships. *Philosophical Transactions of the Royal Society B: Biological Sciences* **331**, 131–153 (1991).
62. Lomax, D. R. & Massare, J. A. A New Species of *Ichthyosaurus* from the Lower Jurassic of West Dorset, England, U.K. *Journal of Vertebrate Paleontology*, e903260 (2015).
63. McGowan, C. A Revision of the Latipinnate Ichthyosaurs of the Lower Jurassic of England (Reptilia: Ichthyosauria). *Life Sciences Contributions of the Royal Ontario Museum* **100**, 1–30 (1974).
64. Lomax, D. R. & Massare, J. A. Two New Species of *Ichthyosaurus* from the Lowermost Jurassic (Hettangian) of Somerset, England. *Papers in Palaeontology* **3** (ed Benson, R.) 1–20 (2017).
65. Wiman, C. Ichthyosaurier Aus Der Trias Spitzbergens. *Bulletin of the Geological Institute of Upsala* **10**, 124–148 (1910).
66. Roberts, A. J., Druckenmiller, P. S., Saetre, G.-P. & Hurum, J. H. A New Upper Jurassic Ophthalmosaurid Ichthyosaur from the Slottsmøya Member, Agardhfjellet Formation or Central Spitsbergen. *PLoS ONE* **9**, e103152 (2014).
67. Delsett, L. L., Roberts, A. J., Druckenmiller, P. S. & Hurum, J. H. A New Ophthalmosaurid (Ichthyosauria) from Svalbard, Norway, and Evolution of the Ichthyopterygian Pelvic Girdle. *PLoS ONE* **12**, e0169971 (2017).
68. Fischer, V., Arkhangelsky, M. S., Uspensky, G. N., Stenshin, I. M. & Godefroit, P. A New Lower Cretaceous Ichthyosaur from Russia Reveals Skull Shape Conservatism within Ophthalmosaurinae. *Geological Magazine* **151**, 60–70 (2014).
69. McGowan, C. & Milner, A. C. A New Pliensbachian Ichthyosaur from Dorset, England. *Palaeontology* **42**, 761–768 (1999).
70. McGowan, C. A New Species of Large, Long-snouted Ichthyosaur from the English Lower Lias. *Canadian Journal of Earth Sciences* **30**, 1197–1204 (1993).
71. McGowan, C. *Leptopterygius tenuirostris* and Other Long-snouted Ichthyosaurs from the English Lower Lias. *Palaeontology* **32**, 409–427 (1989).
72. McGowan, C. A New and Typically Jurassic Ichthyosaur from the Upper Triassic of British Columbia. *Canadian Journal of Earth Sciences* **33**, 24–32 (1996).
73. Maxwell, E. E. & Caldwell, M. W. A New Genus of Ichthyosaur from the Lower Cretaceous of Western Canada. *Palaeontology* **49**, 1043–1052 (2006).
74. Fischer, V. *et al.* A Basal Thunnosaurian from Iraq Reveals Disparate Phylogenetic Origins for Cretaceous Ichthyosaurs. *Biology Letters* **9**, 20130021 (2013).
75. Maisch, M. W. & Matzke, A. T. *Mikadocephalus gracilirostris* N. Gen., N. Sp., a New Ichthyosaur from the Grenzbitumenzone (Anisian-Ladinian) of Monte San Giorgio (switzerland). *Paläontologische Zeitschrift* **71**, 267–289 (1997).
76. Repossi, E. Il Mixosauro Degli Strati Triasici Di Besano in Lombardia. *Atti della Società Italiana di Scienza Naturali e del Museo Civico di Storia Naturale di Milano* **41**, 361–371 (1902).
77. Brinkmann, W. Die Ichthyosaurier (Reptilia) Aus Der Grenzbitumenzone (mitteltrias) Des Monte San Giorgio (tessin, Schweiz) – Neue Ergebnisse. *Vierteljahrsschrift der Naturforschenden Gesellschaft in Zurich* **143**, 165–177 (1998).

78. Chen, X. & Cheng, L. A New Species of *Mixosaurus* (Reptilia: Ichthyosauria) from the Middle Triassic of Pu'an, Guizhou, China. *Acta Palaeontologica Sinica* **2010**, 251–260 (2010).
79. Fernández, M. S. & Talevi, M. Ophthalmosaurian (Ichthyosauria) Records from the Aalenian–Bajocian of Patagonia (Argentina): An Overview. *Geological Magazine* **151**, 49–59 (2014).
80. Fernández, M. S. A New Ichthyosaur from the Los Molles Formation (Early Bajocian), Neuquén Basin, Argentina. *Journal of Paleontology* **73**, 677–681 (1999).
81. Maxwell, E. E., Dick, D. G., Padilla, S. & Luz Parra, M. A New Ophthalmosaurid Ichthyosaur from the Early Cretaceous of Colombia. *Papers in Palaeontology* **2**, 59–70 (2016).
82. Moon, B. C. & Kirton, A. M. Ichthyosaurs of the British Middle and Upper Jurassic. Part 1, *Ophthalmosaurus*. *Monograph of the Palaeontographical Society* **170**, 1–84 (2016).
83. Gilmore, C. W. Osteology of *Baptanodon* (marsh). *Memoirs of the Carnegie Museum* **2**, 77–129 (1905).
84. Efimov, V. M. Ichthyosaurs of a New Genus *Yasykvia* from the Upper Jurassic Strata of European Russia. *Paleontologicheskii Zhurnal* **1999**, 91–98 (1999).
85. Efimov, V. M. A New Family of Ichthyosaurs, the Undorosauridae Fam. Nov. from the Volgian Stage of the European Part of Russia. *Paleontologicheskii Zhurnal* **1999**, 174–181 (1999).
86. Arkhangelsky, M. S. On a New Genus of Ichthyosaurs from the Lower Volgian Substage of the Saratov, Volga Region. *Paleontologicheskii Zhurnal* **1997**, 87–91 (1997).
87. Nicholls, E. L. & Brinkman, D. B. in *Vertebrate Fossils and the Evolution of Scientific Concepts* Writings in Tribute to Beverly Halstead, by Some of His Many Friends (ed Sarjeant, W. A. S.) 521–535 (Gordon and Breach Publishers, London, 1995). ISBN: 9782881249969.
88. Fischer, V. et al. *Simbirsksiasaurus* and *Pervushovisaurus* Reassessed: Implications for the Taxonomy and Cranial Osteology of Cretaceous Platypterygiine Ichthyosaurs. *Zoological Journal of the Linnean Society* **171**, 822–841 (2014).
89. Fischer, V., Bardet, N., Benson, R. B. J., Arkhangelsky, M. S. & Friedman, M. Extinction of Fish-shaped Marine Reptiles Associated with Reduced Evolutionary Rates and Global Environmental Volatility. *Nature Communications* **7**, 10825 (2016).
90. Schmitz, L., Sander, P. M., Storrs, G. W. & Rieppel, O. C. New Mixosauridae (Ichthyosauria) from the Middle Triassic of the Augusta Mountains (Nevada, USA) and Their Implications for Mixosaur Taxonomy. *Palaeontographica Abteilung A: Paläozoologie—Stratigraphie* **270**, 133–162 (2004).
91. Merriam, J. C. The Skull and Dentition of a Primitive Ichthyosaurian from the Middle Triassic. *University of California, Bulletin of the Department of Geology* **5**, 1–11 (1910).
92. Maisch, M. W. & Matzke, A. T. Observations on Triassic Ichthyosaurs. Part XIV: The Middle Triassic Mixosaurid *Phalarodon major* (v. HUENE, 1916) from Switzerland and a Reconsideration of Mixosaurid Phylogeny. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* **2005**, 597–613 (2005).
93. Sander, P. M. in *Ancient Marine Reptiles* (eds Callaway, J. M. & Nicholls, E. L.) 1st ed., 17–43 (Academic Press, San Diego, California, 1997). ISBN: 9780121552107. <https://www.sciencedirect.com/book/9780121552107/ancient-marine-reptiles> (2019).
94. Romer, A. S. An Ichthyosaur Skull from the Cretaceous of Wyoming. *Contributions to Geology, University of Wyoming* **7**, 27–41 (1968).
95. Wade, M. *Platypterygius australis*, an Australian Cretaceous Ichthyosaur. *Lethaia* **17**, 99–113 (1984).
96. Wade, M. A Review of the Australian Cretaceous Longipinnate Ichthyosaur *Platypterygius*, (Ichthyosauria, Ichthyopterygia). *Memoirs of the Queensland Museum* **28**, 115–137 (1990).

97. Fernández, M. S., Archuby, F., Talevi, M. & Ebner, R. Ichthyosaurian Eyes: Paleobiological Information Content in the Sclerotic Ring of *Caypullisaurus* (Ichthyosauria, Ophthalmosauria). *Journal of Vertebrate Paleontology* **25**, 330–337 (2005).
98. Kolb, C. & Sander, P. M. Redescription of the Ichthyosaur *Platypterygius hercynicus* (Kuhn 1946) from the Lower Cretaceous of Salzgitter (Lower Saxony, Germany). *Palaeontographica Abteilung A: Paläozoologie—Stratigraphie* **288**, 151–192 (2009).
99. Broili, F. Ichthyosaurierreste Aus Der Kreide. *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie. Beilage-Band* **25**, 422–442 (1908).
100. Páramo, M. E. *Platypterygius sachicarum* (Reptilia, Ichthyosauria) Nueva Especie Del Crétacico Do Colombia. *Revista Ingeominas* **6**, 1–12 (1997).
101. Yang, P.-F. et al. A New Species of *Qianichthysaurus* (Reptilia: Ichthyosauria) from Xingyi Fauna (Ladinian, Middle Triassic) of Guizhou. *Acta Scientiarum Naturalium Universitatis Pekinensis* **49**, 1002–1008 (2013).
102. Li, C. Ichthyosaur from Guizhou, China. *Chinese Science Bulletin* **44**, 1329–1333 (1999).
103. Maisch, M. W. & Matzke, A. T. Observations on Triassic Ichthyosaurs. Part XII. a New Early Triassic Ichthyosaur Genus from Spitzbergen. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* **229**, 317–338 (2003).
104. Jiang, D.-Y. et al. A Large Aberrant Stem Ichthyosauriform Indicating Early Rise and Demise of Ichthyosauromorphs in the Wake of the End-Permian Extinction. *Scientific Reports* **6**, 26232 (2016).
105. Motani, R., Jiang, D.-Y., Tintori, A., Ji, C. & Huang, J.-D. Pre- Versus Post-mass Extinction Divergence of Mesozoic Marine Reptiles Dictated by Time-scale Dependence of Evolutionary Rates. *Proceedings of the Royal Society B: Biological Sciences* **284**, 20170241 (2017).
106. Yin, G., Zhou, X., Cao, Z., Yu, Y. & Luo, Y. A Preliminary Study on the Early Late Triassic Marine Reptiles from Guanling, Guizhou, China. *Geology, Geochemistry* **28**, 1–23 (2000).
107. Nicholls, E. L. & Manabe, M. Giant Ichthyosaurs of the Triassic—a New Species of *Shonisaurus* from the Pardonet Formation (Norian: Late Triassic) of British Columbia. *Journal of Vertebrate Paleontology* **24**, 838–849 (2004).
108. Camp, C. L. Large Ichthyosaurs from the Upper Triassic of Nevada. *Palaeontographica Abteilung A: Paläozoologie—Stratigraphie* **170**, 139–200 (1980).
109. Fischer, V., Bardet, N., Guiomar, M. & Godefroit, P. High Diversity in Cretaceous Ichthyosaurs from Europe Prior to Their Extinction. *PLoS ONE* **9**, e84709 (2014).
110. Maxwell, E. E., Fernández, M. S. & Schoch, R. R. First Diagnostic Marine Reptile Remains from the Aalenian (Middle Jurassic): A New Ichthyosaur from Southwestern Germany. *PLoS ONE* **7**, e41692 (2012).
111. Maxwell, E. E. New Metrics to Differentiate Species of *Stenopterygius* (Reptilia: Ichthyosauria) from the Lower Jurassic of Southwestern Germany. *Journal of Paleontology* **86**, 105–115 (2012).
112. Von Huene, F. F. *Die Ichthyosaurier Des Lias Und Ihre Zusammenhänge* 114 pp. (Verlag von Gebrüder Borntraeger, Berlin, 1922).
113. Maisch, M. W. A New Ichthyosaur Genus from the Posidonia Shale (Lower Toarcian, Jurassic) of Holzmaden, SW-Germany with Comments on the Phylogeny of Post-Triassic Ichthyosaurs. *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* **209**, 47–78 (1998).
114. Fischer, V., Masure, E., Arkhangelsky, M. S. & Godefroit, P. A New Barremian (Early Cretaceous) Ichthyosaur from Western Russia. *Journal of Vertebrate Paleontology* **31**, 1010–1025 (2011).

115. Martin, J. E., Fischer, V., Vincent, P. & Suan, G. A Longirostrine *Temnodontosaurus* (Ichthyosauria) with Comments on Early Jurassic Ichthyosaur Niche Partitioning and Disparity. *Palaeontology* **55**, 995–1005 (2012).
116. Melmore, S. A Description of the Type-specimen of Ichthyosaurus Crassimanus, Blake (owen MS.) *Annals and Magazine of Natural History* **6**, 615–619 (1930).
117. McGowan, C. A Revision of the Longipinnate Ichthyosaurs of the Lower Jurassic of England, with Descriptions of Two New Species. *Life Sciences Contributions of the Royal Ontario Museum* **97**, 1–37 (1974).
118. Von Huene, F. F. Neue Ichthyosaurier Aus Württemburg. *Neues Jahrbuch für Mineralogie, Geologie und Paläontologie, Beilage-Band B* **65**, 305–320 (1931).
119. Maisch, M. W. & Hungerbühler, A. Revision of *Temnodontosaurus nuertingensis* (v. Huene, 1931), a Large Ichthyosaur from the Lower Pliensbachian (Lower Jurassic) of Nürtingen, South Western Germany. *Stuttgarter Beiträge zur Naturkunde, Serie B (Geologie und Paläontologie)* **248**, 1–11 (1997).
120. Fraas, E. *Die Ichthyosaurier Der Süddeutschen Trias- Und Jura-ablagerungen* (Verlag der H. Laupp'schenn Buchhandlung, Tübingen, 1891).
121. Mazin, J.-M., Suteethorn, V., Buffetaut, E., Jaeger, J.-J. & Elmcke-Ingavat, R. Preliminary Description of *Thaisaurus chonglakmanii* N. G., N. Sp., a New Ichthyopterygian (Reptilia) from the Early Triassic of Thailand. *Comptes rendus de l'Académie des sciences. Série 2, Mécanique, Physique, Chimie, Sciences de l'univers, Sciences de la Terre* **313**, 1207–1212 (1991).
122. Fröbisch, N. B., Fröbisch, J., Sander, P. M., Schmitz, L. & Rieppel, O. C. Macropredatory Ichthyosaur from the Middle Triassic and the Origin of Modern Trophic Networks. *Proceedings of the National Academy of Sciences* **110**, 1393–1397 (2013).
123. Dalla Vecchia, F. M. First Record of the Rare Marine Reptile *Tholodus schmidi* from the Middle Triassic of the Southern Alps. *Rivista Italiana di Paleontologia e Stratigrafia* **110**, 479–492 (2004).
124. Merriam, J. C. New Ichthyosauria from the Upper Triassic of California. *University of California, Bulletin of the Department of Geology* **3**, 249–263 (1903).
125. Arkhangelsky, M. S. & Zverkov, N. G. On a New Ichthyosaur of the Genus *Undorosaurus*. *Proceedings of the Zoological Institute RAS* **318**, 187–196 (2014).
126. Shikama, T., Kamei, T. & Murata, M. Early Triassic *Ichthyosaurus*, *Utatsusaurus hataii* Gen. Et Sp. Nov., from the Kitakami Massif, Northeast Japan. *Science Reports of Tohoku University (Geology)* **48**, 77–97 (1978).
127. Lomax, D. R. A New Leptonectid Ichthyosaur from the Lower Jurassic (Hettangian) of Nottinghamshire, England, UK, and the Taxonomic Usefulness of the Ichthyosaurian Coracoid. *Journal of Systematic Palaeontology* **15**, 387–401 (2016).
128. Maisch, M. W. & Matzke, A. T. Observations on Triassic Ichthyosaurs. Part II. A New Ichthyosaur with Palatal Teeth from Monte San Giorgio. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* **1998**, 26–41 (1998).
129. Jiang, D.-Y. *et al.* New Primitive Ichthyosaurian (Reptilia, Diapsida) from the Middle Triassic of Panxian, Guizhou, Southwestern China and Its Position in the Triassic Biotic Recovery. *Progress in Natural Science* **18**, 1315–1319 (2008).
130. Nicholls, E. L. & Manabe, M. A New Genus of Ichthyosaur from the Late Triassic Pardonet Formation of British Columbia: Bridging the Triassic–Jurassic Gap. *Canadian Journal of Earth Sciences* **38**, 983–1002 (2001).

131. Maisch, M. W. & Matzke, A. T. The Mixosaurid Ichthyosaur *Contectopalatus* from the Middle Triassic of the German Basin. *Lethaia* **33**, 71–74 (2000).
132. Brinkmann, W. *Sangiorgiosaurus* N. G.—eine Neue Mixosaurier Gattung (Mixosauridae, Ichthyosaura) Mit Quetschzähnen Aus Der Grenzbitumenzone (mitteltrias) Des Monte San Giorgio (schweiz, Kanton Tessin). *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* **207**, 125–144 (1998).
133. Jiang, D.-Y., Schmitz, L., Motani, R., Hao, W. & Sun, Y.-L. The Mixosaurid Ichthyosaur *Phalarodon* Cf. *P. fraasi* from the Middle Triassic of Guizhou Province, China. *Journal of Paleontology* **81**, 602–605 (2007).
134. Økland, I. H., Delsett, L. L., Roberts, A. J. & Hurum, J. H. A *Phalarodon fraasi* (Ichthyosauria: Mixosauridae) from the Middle Triassic of Svalbard. *Norwegian Journal of Geology* **98**, 267–288 (2018).
135. Fischer, V., Guiomar, M. & Godefroit, P. New Data on the Palaeobiogeography of Early Jurassic Marine Reptiles: The Toarcian Ichthyosaur Fauna of the Vocontian Basin (SE France). *Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen* **261**, 111–127 (2011).
136. Maisch, M. W. & Matzke, A. T. Observations on Triassic Ichthyosaurs. Part V. The Skulls of *Mikadocephalus* and *Wimanius* Reconstructed. *Neues Jahrbuch für Geologie und Paläontologie, Monatshefte* **1999**, 345–356 (1999).